On the problem of completeness of QM: von Neumann against Einstein, Podolsky, and Rosen

Andrei Yu. Khrennikov International Center for Mathematical Modelling in Physics and Cognitive Sciences, University of Växjö, S-35195, Sweden Email:Andrei.Khrennikov@msi.vxu.se

November 12, 2018

Abstract

We performed a comparative analysis of the arguments of Einstein, Podolsky and Rosen – EPR, 1935: [1] (against the completeness of QM) and the theoretical formalism of QM (due to von Neumann, 1932: [2]). We found that the EPR considerations do not match at all with the von Neumann's theory. Thus EPR did not criticize the real theoretical model of QM. The root of EPR's paradoxical conclusion on incompleteness of QM is the misuse of von Neumann's projection postulate. EPR applied this postulate to observables with degenerate spectra (which is totally forbidden by the axiomatics of QM).

1 Introduction

Recently I published preprint [3] devoted to misinterpretation of the von Neumann projection postulate in modern QM. I also pointed to the fact that such a misinterpretation played the fundamental role in the EPR arguments against the Copenhagen interpretation. This preprint generated a number of fruitful comments from experts in quantum foundations. In spite of general support of my analysis, it

was frequently pointed out that my considerations are not completely clear. In this paper I would like to make clearer the former presentation (in particular, to escape mathematical calculations [3]).

Moreover, I would like to extend my analysis to the general problem of completeness of QM. The conclusion of such analysis is that this problem was not well possed by Einstein, Podolsky and Rosen (EPR). Therefore it should be reformulated in relation to the real Copenhagen interpretation (and not to its perverse version which was attacked in [1]).

2 The role of the projection postulate in the EPR argument

From comments on my preprint [3] I understood that the role of the projection postulate in the EPR-considerations is practically unknown (even for experts in quantum foundations). The main problem is that not so many people have read the original EPR-paper [1]. Even if one did this, it was not careful reading - since it was easier to understand the EPR-arguments from later books on QM. However the projection postulate is the basis of the EPR-definition of an element of reality. Hence, its use (in fact, misuse) is the main source of dilemma: either incompleteness or nonlocality. We shall see that the right (von Neumann) application of the projection postulate would not generate such a dilemma. In particular, so called "quantum nonlocality" would not at all appear in discussion on completeness of QM (its Copenhagen interpretation).

What was wrong in the EPR-considerations? The crucial point was misuse of reduction of wave function in QM. By speaking about QM one should pay attention both to its mathematical formalism and

¹From the very beginning we emphasize that the EPR-arguments were against QM as a theoretical model (including interpretational part). Thus the EPR story was not about "physical elements of reality", but about their theoretical counterparts in the formalism of QM. We recall that axiomatization of QM was performed by Dirac [4] and von Neumann [2]. Measurement theory was completely formalized in [2]. EPR's arguments are in fact about measurement theory. To be rigorous, they should speak about theoretical counterparts of "elements of reality" in von Neumann's axiomatic model. Unfortunately, EPR did not do this precisely (as we shall see). Instead of speaking about von Neumann's axiomatics, they criticized a QM model which was not rigorously formalized. I think that this absence of rigor was the main root of the "EPR-paradox."

its interpretation. The EPR consideration was not consistent neither with the mathematical formulation (due to von Neumann [2]) nor interpretation (due to Bohr [5]).

We now present the EPR-arguments in detail, since otherwise it would be really impossible to criticize them: details are extremely important. We remind the EPR viewpoint on elements of reality:

"If, without in any way disturbing a system, we can predict with certainty (i.e., with probability equal to unity) the value of a physical quantity then there exists an element of physical reality corresponding to this physical quantity."

We emphasize that the main part of the EPR paper [1] consists of considerations on description of reduction of the wave function in QM. Their aim was to associate elements of reality with elements of the theoretical model of QM. We recall that the EPR critique was against this model (and not at all against some real experimental designs). We shall see that EPR associated their elements of reality with eigenfunctions of corresponding self-adjoint operators. We now present their considerations on reduction.

If ψ is an eigenfunction of the operator \widehat{A} ,

$$\psi' \equiv \widehat{A}\psi = a\psi,\tag{1}$$

where a is a number, and so the physical quantity A has with certainty the value a whenever the particle is in the state ψ . By the criterion of reality, for a particle in the state given by ψ for which (1) holds there is an element of physical reality corresponding to the physical quantity A. For example,

$$\psi = e^{(i/\hbar)p_0x},\tag{2}$$

where p_0 is some constant number, and x the independent variable. Since the operator corresponding to the momentum of the particle is

$$\widehat{p} = \frac{\hbar}{i} \frac{\partial}{\partial x},\tag{3}$$

we obtain

$$\psi' = \widehat{p}\psi = \frac{\hbar}{i}\frac{\partial}{\partial x}\psi = p_0\psi. \tag{4}$$

Thus in the state given by (2) the momentum has certainly the value p_0 . It thus has meaning to say that the momentum of the particle in the state given by (2) is real.

On the other hand, if (1) does not hold we can no longer speak of the physical quantity A having a particular value. This is the case, for example, with the coordinate of the particle. The operator corresponding to it, say \widehat{q} , is the operator of multiplication by the independent variable. Thus

$$\widehat{q}\psi = x\psi \neq a\psi. \tag{5}$$

In accordance with quantum mechanics we can only say that the relative probability that a measurement of the coordinate will give a result lying between a and b is

$$\mathbf{P}_{\psi}([a,b]) = \int_{a}^{b} \psi \bar{\psi} dx = \int_{a}^{b} dx = b - a. \tag{6}$$

Since this probability depends upon the difference b-a, we see that all values of the coordinate are equally probable.

More generally, if the operators corresponding to two physical quantities, say A and B, do not commute, that is, if $[\widehat{A}, \widehat{B}] = \widehat{A}\widehat{B} - \widehat{B}\widehat{A} \neq 0$, then the precise knowledge of one of them precludes such a knowledge of the other (since there are no common eigenvectors for A and B). Furthermore, any attempt to determine the latter experimentally will alter the state of the system in such a way as to destroy the knowledge of the first.

From this it follows that: either

a) the quantum mechanical description of reality given by the wave function is not complete;

or

b) when the operators corresponding to two physical quantities do not commute the two quantities cannot have simultaneous reality.

For if both of them had simultaneous reality—and thus definite values—these values would enter into the complete description, according to the condition of completeness. If then the wave function provided such a complete description of reality, it would contain these values; these would be predictable.

By the Copenhagen interpretation of quantum mechanics it is assumed that the wave function does contain a complete description of the physical reality of the system in the state to which it corresponds.

Let us suppose that we have two systems S_1 and S_2 which we permit to interact from the time t = 0 to t = T, after which time we suppose that there is no longer any interaction between the two parts.

We further suppose that the states of the two systems before t = 0 were known. We can then calculate, with the help of the Schrödinger equation, the state of the combined system $S_1 + S_2$ at any subsequent time; in particular, for any t > T.

Let us designate the corresponding wave function (calculated with the aid of the Schrödinger equation) by Ψ . This is the function of the two variables x_1 and x_2 corresponding to the systems S_1 and S_2 respectively, $\Psi = \Psi(x_1, x_2)$. We cannot, however, calculate the state in which either one of the two systems is left after the interaction. This, according to quantum mechanics, can be done with the help of the further measurements by a process known as the reduction of the wave function. Let us consider the essentials of this process.

Let $a_1, a_2, a_3, ...$ be the eigenvalues of an operator A corresponding to some physical quantity A pertaining to the system S_1 and $u_1(x_1), u_2(x_1), u_3(x_1), ...$ the corresponding eigenfunctions. Then Ψ , considered as a function of x_1 , can be expressed as

$$\Psi(x_1, x_2) = \sum_{n=1}^{\infty} u_n(x_1)\psi_n(x_2)$$
 (7)

Here the $\psi_n(x_2)$ are to be regarded merely as the coefficients of the expansion of $\Psi(x_1, x_2)$ into a series of orthogonal functions $u_n(x_1)$. Suppose now that the quantity A is measured and is found to have the value a_k . It is then concluded that after the measurement the first system is left in the state given by the wave function $u_k(x_1)$, and the second system is left in the state given by the wave function $\psi_k(x_2)$. This is the process of reduction of the wave function; the wave function given by the infinite series (7) is reduced to a single term $u_k(x_1)\psi_k(x_2)$.

The set of functions $u_n(x_1)$ is determined by the choice of the physical quantity A. If, instead of this, we had chosen another quantity, say B, with the operator \widehat{B} having the eigenvalues $b_1, b_2, b_3, ...$ and eigenfunctions $v_1(x_1), v_2(x_1), v_3(x_1), ...$ we should have obtained, instead of (7), the expansion

$$\Psi(x_1, x_2) = \sum_{s=1}^{\infty} v_s(x_1)\phi_s(x_2), \tag{8}$$

where ϕ_s are the new coefficients. If the quantity B is now measured and is found to have the value b_r , we conclude that after the measurement the system S_2 is left in the state given by $\phi_r(x_2)$.

Let us now go back to the consideration of the quantum state Ψ . As we have seen, as a consequence of two different measurements performed upon the first system S_1 (for the quantities A and B) the second system may be left in states with two different wave functions $-\psi_k(x_2)$ and $\phi_r(x_2)$. On the other hand, since at the time of measurement the two systems no longer interact, no real change can take place in the second system as a consequence of anything that may be done to the first system. This is, of course, merely a statement of what is meant by the absence of an interaction between the two systems. Thus it is possible to assign two different wave functions (in our example ψ_k and ϕ_r) to the same reality (the second system after the interaction with the first).

Now, it may happen that the two wave functions ψ_k and ϕ_r are eigenfunctions of two non-commuting operators corresponding to some physical quantities P and Q, respectively. That this may actually be the case can best be shown by an example, see [1].

3 The von Neumann projection postulate

In von Neumann's book [2] the cases of observables with nondegenerate and degenerate spectra were sharply distinguished. The postmeasurement state is well defined (and given by the corresponding eigenvector) only for observables with nondegenerate spectra. Only in this case EPR might say that one could assign the wave function with the physical system (after the measurement). However, if spectrum is degenerate, then by the von Neumann axiomatics of QM the post-measurement state is not determined.

Thus one could not assign the definite wave function with the physical system (after measurement).

It is amazing that EPR did not pay attention to this crucial point. I could not exclude that they even did not read von Neumann's book. In their paper the projection postulate is applied for observables with degenerate spectra, but in such a way as if they were observables with nondegenerate spectra.

By considering partial measurements on subsystems of composite systems one immediately moves to the domain of degenerate measurements. Those operators A and B considered by EPR have degenerate spectra. Therefore by measuring e.g. A one would not determine the

state of a composite system $S_1 + S_2$. Hence, the state of S_2 is not determined by A-measurement on S_1 . The wave function $\psi_k(x_2)$ could not be assigned with S_2 . It is impossible to proceed as EPR did at the very end of their general considerations on measurements on composite systems. Since even one wave function, $\psi_k(x_2)$, could not be assigned with S_2 , it is totally meaningless to write about assigning of two different wave functions to the same reality.

Conclusion. EPR did not prove that QM is incomplete. They did mistake by assuming that by measurement of observable A (respectively, B) on S_1 the linear combination (7) (respectively, (8)) is reduced to a single summand.

4 EPR is about precise correlations

My correspondence with readers of preprint [3] demonstrated that considerations of EPR on reduction of the wave function (which were presented in section 2) were never discussed seriously. This part of EPR's paper (two of totally four pages) is practically ignored. Instead of this,l people have always been concentrated on the last page of the paper containing the discussion on precise correlations for the position and momentum. As e.g. Elena Loubentz and Joachim Kupsch pointed out in E-mails to me, the EPR paper is not about the projection postulate, but about measurements for states with precise correlations. The essence of the EPR conclusions is presented in short on page 780:

"Returning now to the general case contemplated in Eqs. (7) and (8), we assume that ψ_k and ϕ_r are indeed eigenfunctions of some non-commuting operators P and Q, corresponding to the eigenvalues p_k and q_r , respectively. Thus by measuring either A or B we are in a position to predict with certainty, and without in any way disturbing the second system, whether the value of the quantity P (that is p_k) or the value of the quantity Q (that is q_r). In accordance with our criterion of reality, in the first case we must consider the quantity P as being an element of reality, in the second case the quantity Q is an element of reality."

As I understood, the last sentence has always been considered as the very end of the story. However, (by some reason) EPR continued:

"But, as we have seen, both wave functions ψ_k and ϕ_r , belong to the same reality."

Opposite to the majority of readers of their paper or (and it was more common) some texts about their paper, EPR were not able to get the complete satisfaction via producing of elements of reality for the second particle via A and B measurements on the first one. They had to come back to their rather long story (pages 788-789) on reduction of the wave function. I think that this EPR's comeback to reduction is the crucial point of their argument. Why did they need do this? I think that by the following reason. It is impossible to associate simultaneously two "experimental elements of reality" with S_2 on the basis of measurement on S_1 , since (as everybody understood well) either A or B measurement could be performed on S_1 (but not both A and B). Therefore EPR were able to associate with S_2 only "theoretical elements of reality" represented by the wave functions $\psi_k(x_2)$ and $\phi_r(x_2)$ - eigenfunctions of the two non-commuting operators P and Q (for the second particle).

And it was enough for their purpose, since they wanted to prove incompleteness of QM as a *theoretical* model.

Conclusion. EPR were not able to proceed without appealing to the projection postulate (with all consequences of its misuse).

5 Refinement measurements

However, according to von Neumann by obtaining a fixed value, say $A = \alpha$, for measurement on S_1 , one does not determine the state of $S_1 + S_2$ (and, hence, neither the state of S_2).

To determine the state of $S_1 + S_2$, one should perform some refinement measurement. In QM it is represented by an operator commuting with $A \otimes I$ and eliminating degeneration². Since any operator of the form $I \otimes C$ commutes with $A \otimes I$, it is natural to consider refinement observable corresponding to measurement on S_2 . The position Q and momentum P operators considered by EPR give examples of von Neumann's refinement measurements. Each of them determine the state of $S_1 + S_2$ (and hence S_2) uniquely.

Moreover, for any operator with degenerate spectrum its measurement is ambiguous [2]. Thus in the EPR case measurement of A could not at all be considered as measurement on $S_1 + S_2$. It is just measurement on S_1 .

²Here $A: L_2(\mathbf{R}^3) \to L_2(\mathbf{R}^3), A \otimes I: L_2(\mathbf{R}^3) \otimes L_2(\mathbf{R}^3) \to L_2(\mathbf{R}^3) \otimes L_2(\mathbf{R}^3).$

However, for EPR the story about so called EPR-states was not simply the standard story about von Neumann's refinement measurements.

6 The EPR paper as the source of the idea about quantum nonlocality

At the very end of their paper EPR discussed a problem which later became known as the problem of *quantum nonlocality*:

"One could object to this conclusion on the grounds that our criterion of reality is not sufficiently restrictive. Indeed, one would not arrive at our conclusion if one insisted that two or more physical quantities can be regarded as simultaneous elements of reality only when they can be simultaneously measured or predicted. On this point of view, since either one or the other, but not both simultaneously, of the quantities P and Q can be predicted, they are not simultaneously real. This makes the reality of P and Q depend upon the process of measurement carried out on the first system, which does not disturb the second system in any way. No reasonable definition of reality could be expected to permit this."

Later nonlocality was coupled to the von Neumann projection postulate in the following way. To escape incompleteness of QM, one should not assign the wave function $\psi_k(x_2)$ with S_2 before the Ameasurement on S_1 . One might say that the A-measurement on S_1 produces instantaneous action on S_2 and its state is collapsed into $\psi_k(x_2)$. For example, one can find an example of such a reasoning in the paper of Alain Aspect [6].

This form of reasoning has nothing to do with QM. By the same von Neumann's projection postulate the state of S_2 is NOT determined by measurement on S_1 . There is no even trace of action at the distance!

Conclusion. "Quantum nonlocality" appeared as a consequence of misuse of the projection postulate. We also emphasize that EPR considered quantum nonlocality as a totally absurd alternative to their arguments in favor of incompleteness of QM.

7 Nonlocality of the experiment design as opposed to EPR state nonlocality

7.1 Quantum theory and joint measurements of compatible observables

We have already discussed that from the QM-viewpoint (based on von Neumann's axiomatics) the whole EPR story is about refinement measurements for operators with degenerate spectra. It would be useful to analyse (by using the conventional QM-framework) the procedure of joint measurement of two compatible observables, say A and Q: [A, Q] = 0.

The crucial point is that by von Neumann, to design joint measurement of A and Q, one should design measurement of third observable, say C, such that A = f(C) and Q = g(C), where $f, g : \mathbf{R} \to \mathbf{R}$ are some functions. In the EPR case we want to have C with nondegenerate spectrum and A is observable on S_1 and Q on S_2 .

Since A and Q are measured in different domains of spacetime, the design of measurement of C should be nonlocal. It is an extremely important point.

What does it mean "nonlocal design"?

In particular, it means that one should perform the *time synchro-nization* between results of measurement of A and Q. It is important to be totally sure that clicks of the A-detector (giving the result of measurement on S_1) and the Q-detector (giving the result of measurement on S_2) match each other. We emphasize that in the real experimental setup for the EPR-Bohm experiment for photon polarization, see e.g., [7], [8], such a time synchronization is really realized via the nonlocal experimental design - via using the *time window*. The time window constraint

$$|t_i^A - t_i^Q| < \Delta$$

is evidently nonlocal. We also point out to the *synchronization of space* frames. Orientations of polarization beam splitters are chosen in one fixed space frame (in the complete accordance with Bohr's ideology [5]).

7.2 The EPR state nonlocality

If one proceeds with so called quantum nonlocality induced by the misuse of the projection postulate, then he should take such a nonlocality very seriously. It would be real physical nonlocality of states. We again recall that EPR considered such a nonlocality as totally absurd.

Conclusion. The correct application of the projection postulate implies the nonlocal experimental design of the EPR-type experiments; in particular, the time synchronization (e.g., via the time window) as well as the choice of the fixed space frame. This experimental design nonlocality has nothing to do with so called "quantum nonlocality".

8 Bohr's reply to Einstein

It is typically emphasized that Bohr's reply [5] is very difficult for understanding. I totally agree with such a common viewpoint. I was able to understand Bohr only on the basis of previous considerations on the role of the projection postulate in the EPR considerations. Unfortunately, in Bohr's reply there was no even trace of von Neumann's axiomatization of QM³. Consequently Bohr did not pay any attention to the role of the projection postulate in the EPR considerations. He missed the EPR-trick with assigning to S_2 two wave functions, $\psi_k(x_2)$ and $\phi_r(x_2)$, which are eigenfunctions of two noncommutative observables, say P and Q. It is very important in the EPR considerations that these wave functions and not measurements by themselves represent "elements of reality" in QM (as a theoretical model). Thus, instead of analyzing this tricky point in the EPR paper, Bohr proceeded in the purely experimental framework. He simply recalled his ideas on complementarity of various measurement setups in relation to the EPR-considerations. In short his message was that since one could not combine two measurement setups for S_1 related to incompatible quantities, it is impossible to assign two corresponding elements of reality to S_2 . Bohr concluded that the EPR notion of an element of reality was ambiguous.

The problem was that EPR "proved" that QM is incomplete as a theoretical model, but Bohr replied by supporting his old thesis that

³I strongly suspect that neither Einstein nor Bohr had read von Neumann's book at that time.

QM is complete as an experimental methodology. It seems that the resulting common opinion was not in favor of Bohr's reply. And it is clear why. If EPR really were able to prove that the formalism of QM implies assigning to S_2 of two wave functions, $\psi_k(x_2)$ and $\phi_r(x_2)$, corresponding to two noncommuting operators Q and P, I would (and I was!) on their side. The point (presented in this paper) is that they were not able to do this by using the QM formalism in the proper way.

Conclusion. Bohr's reply in spite correctness of his arguments, did not contain the analysis of the real roots of the "EPR paradox". It induced a rather common impression that EPR's argument is not trivially reduced to the old problem of complementarity. It was commonly accepted that the only possibility to escape assigning "elements of reality" corresponding to incompatible observables to the same particle is to accept quantum nonlocality.

9 Concluding remarks

It seems that the "EPR-paradox" was finally resolved in this paper. I hope that it would stimulate people to look for various ways beyond QM. By von Neumann's axiomatics of QM [2] the notion of measurement of observable A with degenerate spectrum is ambiguous. It is well defined only via refinement measurement given by observable Cwith nongenerate spectrum such that A = f(C). Since any observable A on the subsystem S_1 of a composite system $S = S_1 + S_2$ has degenerate spectrum in the tensor Hilbert space of S-states, it is totally meaningless to discuss (as EPR did) its measurement without fixing a refinement measurement on S_2 . If such a refinement is not fixed from the very beginning, then A-measurement has nothing to do with measurements on the composite systems S. It could not change the S-state and, hence, the S_2 -state. Bohr's reply [5] to Einstein could be interpreted in the same way. Thus the EPR-attack against QM was not justified. Unfortunately, this attack was the source of naive Einsteinian realism (assigning to the same system S_2 of two wave functions $\psi_k(x_2)$ and $\phi_r(x_2)$ corresponding to noncommutative operators) and quantum nonlocality. We also point out to practically unknown fact that so called EPR states were studied in detail by von Neumann [2], pp. 434-435. But he was able to proceed without assigning two wave functions (corresponding to noncommuting operators) to the same system. Consequently, no traces of incompleteness of QM or its

nonlocality could be found in [2].

This paper was written during my short visit to University of Oxford, 23-27 March. 2008. Unexpected shift of period of the Conference "Quantum Interactions" (due to Ouster) gives me a possibility to formalize my views elaborated during last ten years. I would like to thank Bob Coecke for hospitality and discussions. I also would like to thank Sven Aerts for an interesting conversation on probabilistic incompatibility. The talk of Basil Hiley (in which he claimed that Bohr's reply is totally un-understandable and nonlocality is the only reasonable choice) was extremely stimulating for me: I read Bohr's reply once again. I would like to thank K. Hess, A. Plotnitsky, E. Loubentz, J. Kupsch, A. Majewski for critical comments on my preprint [3].

References

- [1] A. Einstein, B. Podolsky, N. Rosen, Phys. Rev. 47, 777–780 (1935).
- [2] J. von Neumann, Matematische Grundlagen der Quantenmechanik, Springer, Berlin, 1932.
- [3] A. Khrennikov, The role of von Neumann and Lders postulates in the EPR-Bohm-Bell considerations: Did EPR make a mistake? http://arxiv.org/abs/0801.0419
- [4] P. A. M. Dirac, The Principles of Quantum Mechanics1930.
- [5] N. Bohr, Phys. Rev. 48, 696 (1935).
- [6] A. Aspect, Bell's Theorem: The Naive View of an Experimentalist. http://arxiv.org/abs/quant-ph/0402001.
- [7] G. Weihs, T. Jennewein, C. Simon, H. Weinfurter, and A. Zeilinger, *Phys. Rev. Lett.*, vol. 81, pp. 5039-5042, 1998.
- [8] G. Weihs, "A test of Bell's inequality with spacelike separation," in *Proc. Conf. Foundations of Probability and Physics-4*, Melville, NY: American Institute of Physics, Ser. Conference Proceedings, vol. 889, pp. 250-262, 2007.