

Microphysical analogues of flyby anomalies

Karl Svozil*

Institute for Theoretical Physics, Vienna University of Technology,

Wiedner Hauptstraße 8-10/136, A-1040 Vienna, Austria

Abstract

We discuss Doppler shift and interferometric measurements in analogy to the recently reported macroscopic flyby anomalies.

PACS numbers: 04.80.Cc, 03.65.Nk

Keywords: flyby anomaly, dynamical anomaly, interference




FIG. 1: A particle beam flies by a heavy rotating (magnetic) object. The flyby should be along a path parallel to the rotation axis.

The recently reported anomalous orbital-energy changes observed during six Earth flybys by the Galileo, NEAR, Cassini, Rosetta, and MESSENGER spacecraft documented an unexpected frequency increase in the post encounter radio Doppler data generated by stations of the NASA Deep Space Network [1, 2]. These observations deserve great attention, mainly because of the possible consequences for theories of gravitation and electromagnetism [3].

In what follows we shall investigate microscopic analogues of macroscopic flyby configurations for which anomalies have been noted. We first start with a straightforward discussion of flyby configurations and later concentrate on interferometric setups.

In a first approach consider a beam of neutral particles, such as neutrons or photons, being deflected by a heavy rotating (magnetic) object. The flyby should be along a path parallel to the rotation axis. This configuration is depicted in Fig. 1.

For the prediction of the frequency shift, we take formula (2) of Ref. [1]

$$\Delta\omega = 2K\omega(\cos\delta_i - \cos\delta_o),$$

with δ_i and δ_o as the declinations of the incoming and outgoing osculating asymptotic velocity vectors, and $K = 2\Omega R/c$, where Ω is the angular rotational velocity and R the mean radius of the heavy object. c stands for the speed of light in vacuum. In our case, as $\delta_i = 0$ and $\delta_o = \pi$, we




FIG. 2: Mach-Zehnder interferometer, in which one beam path is parallel to the rotation axis of a heavy rotating (magnetic) object O .

obtain as an estimate for the frequency shift

$$\Delta\omega = \frac{8\Omega R}{c} \omega. \quad (1)$$

One straightforward way to realize heavy (magnetic) rotating objects are stacks of data disks which operate at about 7200 revolutions per minute, corresponding to 120 Hz or 754 rad/s. This is about seven orders of magnitude greater than Earth's angular rotation velocity. Typical disks have a radius of about 0.05 m, which is about eight orders of magnitude smaller than Earth's mean radius. Thus, in this case,

$$K = 2.513 \times 10^{-7} \quad (2)$$

is only about a factor of ten smaller than for Earth.

Another possibility to measure possible frequency shifts of microscopic particles and light is by comparing the phases acquired along two different paths. Consider, for the sake of demonstration, a Mach-Zehnder device depicted in Fig. 2, in which heavy rotating (magnetic) mass is placed along one beam path, whereas the other beam path is left unperturbed. As usual [4, 5], the computation

proceeds by successive substitution (transition) of states; i.e., if all components are lossless, then

$$\begin{aligned}
S_1 : a &\rightarrow (b + ic)/\sqrt{2} \quad , \\
O : b &\rightarrow be^{i\Delta\omega} \quad , \\
S_2 : b &\rightarrow (e + id)/\sqrt{2} \quad , \\
S_2 : c &\rightarrow (d + ie)/\sqrt{2} \quad .
\end{aligned}
\tag{3}$$

The resulting transition is

$$a \rightarrow \psi = i \left(\frac{e^{i\Delta\omega} + 1}{2} \right) d + \left(\frac{e^{i\Delta\omega} - 1}{2} \right) e \quad .
\tag{4}$$

As can be expected, for $\Delta\omega = 0$ Eq. (4) reduces to $a \rightarrow id$, and an emitted quant is detected only by D_1 . If $\Delta\omega = \pi$, Eq. (4) reduces to $a \rightarrow -e$, and an emitted quant is detected only by D_2 . If one varies the phase shift $\Delta\omega$, one obtains the following detection probabilities:

$$\begin{aligned}
P_{D_1}(\Delta\omega) &= |(d, \psi)|^2 = \cos^2\left(\frac{\Delta\omega}{2}\right) = \cos^2\left(\frac{4\Omega R}{c} \omega\right), \\
P_{D_2}(\Delta\omega) &= |(e, \psi)|^2 = \sin^2\left(\frac{\Delta\omega}{2}\right) = \sin^2\left(\frac{4\Omega R}{c} \omega\right).
\end{aligned}
\tag{5}$$

Taking the estimate for the hard disk configuration as before, the frequency shifts are of the order of 5×10^{-7} times the frequency of light or of the matter waves.

In summary, we have suggested to test the flyby of microscopic particles along heavy (magnetic) rotating objects whose rotation axis are parallel to the beam path. Because of their sensitivity, interferometric setups appear to be particularly promising.

* Electronic address: svozil@tuwien.ac.at; URL: <http://tph.tuwien.ac.at/~svozil>

- [1] J. D. Anderson, J. K. Campbell, J. E. Ekelund, J. Ellis, and J. F. Jordan, "Anomalous Orbital-Energy Changes Observed during Spacecraft Flybys of Earth," *Physical Review Letters* **100**, 091 102 (2008).
<http://dx.doi.org/10.1103/PhysRevLett.100.091102>
- [2] J. D. Anderson, J. K. Campbell, and M. M. Nieto, "The energy transfer process in planetary flybys," *New Astronomy* **12**, 383–397 (2007).
<http://dx.doi.org/10.1016/j.newast.2006.11.004>
- [3] H. Dittus, C. Lmmerzahl, and S. G. Turyshev, *Lasers, Clocks and Drag-Free Control Exploration of Relativistic Gravity in Space*, Astrophysics and Space Science Library, Vol. 349 (Springer, Heidelberg, 2006).

[4] D. M. Greenberger, M. A. Horne, and A. Zeilinger, "Multiparticle interferometry and the superposition principle," *Physics Today* **46**, 22–29 (1993).

[5] K. Svozil, "Noncontextuality in multipartite entanglement," *J. Phys. A: Math. Gen.* **38**, 5781–5798 (2005).

<http://dx.doi.org/10.1088/0305-4470/38/25/013>