

**INVERSE INDEFINITE SPECTRAL PROBLEM FOR SECOND
ORDER DIFFERENTIAL OPERATOR PENCIL WITH
COMPLEX PERIODIC COEFFICIENTS**

R.F. Efendiev

**Institute Applied Mathematics, Baku State University,
Z.Khalilov, 23, AZ1148, Baku, Azerbaijan, rakibaz@yahoo.com**

ABSTRACT.

The inverse problem for the differential operator pencil with complex periodic potential and discontinuous coefficients on the axis is studied. Main characteristics of the fundamental solutions are investigated, the spectrum of the operator is studied. We give formulation of the inverse problem, prove a uniqueness theorem and provide a constructive procedure for the solution of the inverse problem.

Key words:

Discontinuous equations; Tuning points; Spectral singularities; Inverse spectral problems; Continuous spectrum;

MSC:

34A36; 34M60; 34L05; 47A10; 47A70:

INTRODUCTION.

Let L be an operator generated by the operation $\frac{1}{\rho(x)} \left\{ -\frac{d^2}{dx^2} + 2\lambda p(x) + q(x) \right\}$ in the space $L_2(-\infty, +\infty, \rho(x))$.

Here λ is a complex number, coefficients $p(x)$ and $q(x)$ are

$$p(x) = \sum_{n=1}^{\infty} p_n e^{inx}; \sum_{n=1}^{\infty} n \cdot |p_n| < \infty \quad ; \quad q(x) = \sum_{n=1}^{\infty} q_n e^{inx}; \sum_{n=1}^{\infty} |q_n| < \infty \quad (1)$$

and

$$\rho(x) = \begin{cases} 1 & \text{for } x \geq 0, \\ -\beta^2 & \text{for } x < 0. \end{cases} \quad (2)$$

If $\rho(x) = 1$, then equation $Ly = \lambda^2 y$ described wave propagation in a one dimensional nonconservative medium in a frequency domain, where λ is the wave number (momentum), λ^2 is energy, $p(x)$ describes the joint effect of absorption and generation of energy, $q(x)$ describes the regeneration of the force density. Because of important applications in quantum mechanics, it is interesting to investigate the spectral characteristics of the operator pencil L , and at least one of the colliding particles is a fermion, it is relevant to solve the inverse problem in the presence of central and spin orbital potentials.

Our investigation was stimulated by M.G.Casymov's paper [1] where he first considered this potential. Especially, we would like to indicate the paper V. Guillemin, A. Uribe [2] where the potential $q(x) = \sum_{n=1}^{\infty} q_n e^{inx}$ plays a vital part for solving the KdV equation. The inverse problems connected with potentials of the form (1) were considered in [3-4].

REPRESENTATION OF FUNDAMENTAL SOLUTIONS.

Here we study the solutions of the main equation

$$-y''(x) + 2\lambda p(x)y(x) + q(x)y(x) = \lambda^2 \rho(x) y(x)$$

that will be convenient in future.

We first consider the solutions $f_1^+(x, \lambda)$ and $f_2^+(x, \lambda)$, determined by the conditions at infinity

$$\lim_{Imx \rightarrow \infty} f_1^+(x, \lambda) e^{-i\lambda x} = 1, \quad \lim_{Imx \rightarrow \infty} f_2^+(x, \lambda) e^{-\beta \lambda x} = 1.$$

We can prove the existence of these solutions if the condition (1) is fulfilled for the potential. This will be unique restriction on the potential and later on we'll consider it to be fulfilled.

Theorem 1. Let $q(x)$ and $p(x)$ be of the form (1) and $\rho(x)$ satisfy condition (2). Then equation $Ly = \lambda^2 y$ has special solutions of the form

$$f_1^+(x, \lambda) = e^{i\lambda x} \left(1 + \sum_{n=1}^{\infty} v_n^+ e^{inx} + \sum_{n=1}^{\infty} \sum_{\alpha=n}^{\infty} \frac{v_{n\alpha}^+}{n+2\lambda} e^{i\alpha x} \right) \quad \text{for } x \geq 0, \quad (3)$$

$$f_2^+(x, \lambda) = e^{\lambda \beta x} \left(1 + \sum_{n=1}^{\infty} v_n^+ e^{inx} + \sum_{n=1}^{\infty} \sum_{\alpha=n}^{\infty} \frac{v_{n\alpha}^+}{n-2i\lambda\beta} e^{i\alpha x} \right) \quad \text{for } x < 0 \quad (4)$$

where the numbers v_n^+ and $v_{n\alpha}^+$ are determined from the following recurrent relations

$$\alpha^2 \nu_{\alpha}^+ + \alpha \sum_{n=1}^{\alpha} \nu_{n\alpha}^+ + \sum_{s=1}^{\alpha-1} \left(q_{\alpha-s} \nu_s^+ - p_{\alpha-s} \sum_{n=1}^s \nu_{ns}^+ \right) + q_{\alpha} = 0, \quad (5)$$

$$\alpha(\alpha-n) \nu_{n\alpha}^+ + \sum_{s=n}^{\alpha-1} (q_{\alpha-s} - n \cdot p_{\alpha-s}) \nu_{ns}^+ = 0, \quad (6)$$

$$\alpha \nu_{\alpha}^+ + \sum_{s=1}^{\alpha-1} \nu_s^+ p_{\alpha-s} + p_{\alpha} = 0. \quad (7)$$

and the series (3-4) admits double termwise differentiation.

Remark1: If $\lambda \neq -\frac{\alpha}{2}$ and $Im\lambda \geq 0$, then $f_1^+(x, \lambda) \in L_2(0, +\infty)$.

Remark2: If $\lambda \neq -\frac{i\alpha}{2\beta}$ and $Re\lambda \geq 0$, then $f_2^+(x, \lambda) \in L_2(-\infty, 0)$.

Extending $f_1^+(x, \lambda)$ and $f_2^+(x, \lambda)$ as solutions of equation $Ly = \lambda^2 y$ on $x < 0$ and $x \geq 0$ respectively and using the conjunction conditions

$$\begin{aligned} y(0+) &= y(0-), \\ y'(0+) &= y'(0-), \end{aligned}$$

we can prove the following lemma.

Lemma 1: $f_1^+(x, \lambda)$ and $f_2^+(x, \lambda)$ may be extended as solutions of equation $Ly = \lambda^2 y$ on $x < 0$ and $x \geq 0$, respectively. Then we get

$$f_2^+(x, \lambda) = C_{11}(\lambda) f_1^+(x, \lambda) + C_{12}(\lambda) f_1^-(x, \lambda) \quad \text{for } x \geq 0,$$

$$f_1^+(x, \lambda) = C_{22}(\lambda) f_2^+(x, \lambda) + C_{21}(\lambda) f_2^-(x, \lambda), \quad \text{for } x < 0,$$

where

$$f_1^-(x, \lambda) = e^{-i\lambda x} \left(1 + \sum_{n=1}^{\infty} v_n^- e^{inx} + \sum_{n=1}^{\infty} \sum_{\alpha=n}^{\infty} \frac{v_{n\alpha}^-}{n-2\lambda} e^{i\alpha x} \right) \quad \text{for } x \geq 0,$$

$$f_2^-(x, \lambda) = e^{-\lambda\beta x} \left(1 + \sum_{n=1}^{\infty} v_n^- e^{inx} + \sum_{n=1}^{\infty} \sum_{\alpha=n}^{\infty} \frac{v_{n\alpha}^-}{n+2i\lambda\beta} e^{i\alpha x} \right) \quad \text{for } x < 0.$$

The numbers v_n^- , $v_{n\alpha}^-$ are determined from the following recurrent relations

$$\alpha^2 v_\alpha^- + \alpha \sum_{n=1}^{\alpha} v_{n\alpha}^- + \sum_{s=1}^{\alpha-1} \left(q_{\alpha-s} v_s^- - p_{\alpha-s} \sum_{n=1}^s v_{ns}^- \right) + q_\alpha = 0,$$

$$\alpha(\alpha-n) v_{n\alpha}^- + \sum_{s=n}^{\alpha-1} (q_{\alpha-s} + n \cdot p_{\alpha-s}) v_{ns}^- = 0,$$

$$\alpha v_\alpha^- - \sum_{s=1}^{\alpha-1} v_s^- p_{\alpha-s} - p_\alpha = 0,$$

and

$$C_{11}(\lambda) = \frac{W[f_2^+(0, \lambda), f_1^-(0, \lambda)]}{2i\lambda}, \quad C_{12}(\lambda) = \frac{W[f_1^+(0, \lambda), f_2^+(0, \lambda)]}{2i\lambda},$$

$$C_{22}(\lambda) = \frac{i}{\beta} C_{11}(-\lambda), \quad C_{21}(\lambda) = -\frac{i}{\beta} C_{12}(\lambda).$$

Lemma 2. L has no eigenvalues for real and pure imaginary λ . It's continuous

spectra consist of axes $Re\lambda = 0$ and $Im\lambda = 0$ on which there may exist spectral singularities coinciding with the numbers $\frac{in}{2\beta}$, $\frac{n}{2}$, $n = \pm 1, \pm 2, \pm 3, \dots$

Lemma 3. The eigenvalues of operator L are finite and coincide with zeros of the functions $C_{12}(\lambda)$, $C_{11}(-\lambda)$, $C_{12}(-\lambda)$, $C_{11}(\lambda)$ from sectors $S_k = \{k\pi/2 < \arg \lambda < (k+1)\pi/2\}$, $k = \overline{0, 3}$ respectively.

Definition 1. The data $\{\lambda_n, C_{11}(\lambda), C_{12}(\lambda)\}$ are called the spectral data of L .

INVERSE PROBLEM. Given the spectral data $\{\lambda_n, C_{11}(\lambda), C_{12}(\lambda)\}$ construct the potential $q(x), p(x)$ and β .

Let

$$f_n^\pm(x) = \lim_{\lambda \rightarrow \mp \frac{n}{2}} (n \pm 2\lambda) f_1^\pm(x, \lambda) = \sum_{\alpha=n}^{\infty} v_{n\alpha}^\pm e^{i\alpha x} e^{-i\frac{n}{2}x} \quad (8)$$

Then holds the relation

$$f_n^\pm(x) = v_{nn}^\pm f_1^\mp \left(x, \mp \frac{n}{2} \right) \quad (9)$$

Using the results obtained above we arrive at the following procedure for solution of the inverse problem.

1. Taking into account (8-9) it is easy to check that

$$\lim_{\lambda \rightarrow n/2} (n - 2\lambda) \frac{C_{11}(\lambda)}{C_{12}(\lambda)} = v_{nn}^-, \quad \lim_{\lambda \rightarrow -n/2} (n + 2\lambda) \frac{C_{12}(\lambda)}{C_{11}(\lambda)} = v_{nn}^+$$

consequently we find all numbers v_{nn}^\pm , $n = 1, 2, \dots$

2. Taking into account (9) we get

$$\sum_{\alpha=B}^{\infty} v_{n\alpha}^\pm e^{i\alpha x} e^{-i\frac{n}{2}x} = v_{nn}^\pm \cdot e^{i\frac{n}{2}x} \left(1 + \sum_{n=1}^{\infty} v_n^\mp e^{inx} + \sum_{n=1}^{\infty} \sum_{\alpha=n}^{\infty} \frac{v_{n\alpha}^\mp}{n+n} e^{i\alpha x} \right)$$

or

$$v_{m, \alpha+m}^\pm = v_{mm}^\pm \left(v_\alpha^\mp + \sum_{n=1}^{\alpha} \frac{v_{n\alpha}^\mp}{n+m} \right)$$

from which all numbers $v_{n\alpha}^\pm$, $\alpha = 1, 2, \dots$, $n = 1, 2, \dots, n < \alpha$ are defined.

3. Then from recurrent formula (5-7), find all numbers p_n and q_n .

4. The number β is defined from equality

$$\beta = iC_{11}(\lambda_n) C_{11}(-\lambda_n).$$

Really using Lemma 1 we derive for $\lambda_n \in S_0$ the true relation

$$C_{11}(\lambda_n) = \frac{f_2^+(x, \lambda_n)}{f_1^+(x, \lambda_n)}, \quad C_{22}(\lambda_n) = \frac{f_1^+(x, \lambda_n)}{f_2^+(x, \lambda_n)} \text{ i.e. } C_{11}(\lambda_n) C_{22}(\lambda_n) = 1.$$

Then from Lemma 1 we get $\beta = iC_{11}(\lambda_n) C_{11}(-\lambda_n)$.

So inverse problem has a unique solution and the numbers q_n, p_n and β are defined constructively by the spectral data.

Theorem 2. The specification of the spectral data uniquely determines potentials $p(x)$, $q(x)$ and β .

REFERENCES.

1. Gasymov M.G. Spectral analysis of a class non-self-adjoint operator of the second order. Functional analysis and its appendix. (In Russian) 1980, V34.1.pp.14-19
2. V.Guillemin, A. Uribe Hardy functions and the inverse spectral method. Comm.In Partial Differential equations,8(13),1455-1474(1983)
3. Efendiev, R.F. The Characterization Problem for One Class of Second Order Operator Pencil with Complex Periodic Coefficients. Moscow Mathematical Journal, 2007, Volume 7, Number 1, pp.55-65.
4. Efendiev, R. F. Complete solution of an inverse problem for one class of the high order ordinary differential operators with periodic coefficients. Zh. Mat. Fiz. Anal. Geom. 2006, V. 2, no. 1, 73–86,

**SPECTRAL ANALYSIS FOR ONE CLASS OF SECOND-ORDER
INDEFINITE NON-SELF-ADJOINT DIFFERENTIAL
OPERATOR PENCIL**

R.F. Efendiev

Institute of Applied Mathematics, Baku State University,
Z.Khalilov, 23, AZ1148, Baku, Azerbaijan,

efendievrakib@bsu.az

ABSTRACT

The inverse problem for the Schrodinger operator with complex periodic potentials and discontinuous coefficient on the axis is studied. Main characteristics of the fundamental solutions are investigated, the spectrum of the operator is studied. We give the formulation of the inverse problem, prove a uniqueness theorem and provide a constructive procedure for the solution of the inverse problem.

Key words: Discontinuous equation, Spectral singularities, Inverse spectral problem, Continuous spectrum.

MSC: 34A36, 34L05, 47A10, 47A70

INTRODUCTION

In this paper we propose a method for solving the inverse spectral problem of wave propagation with discontinuous wave speed in one-dimensional layered-inhomogeneous medium in the frequency domain which is described by Schrodinger equation

$$-y''(x) + 2\lambda p(x)y(x) + q(x)y(x) = \lambda^2 \rho(x)y(x) \quad (1)$$

where R - real axis, λ - wave number (momentum), λ^2 - energy, $p(x)$ - describes the joint effect of absorption and generation of energy, $q(x)$ - describes the regeneration of the force density, $\rho^{-1}(x)$ - wave speed.

An inverse problem consists in inferring some properties of given problem from a complete or partial knowledge of its solution. The definition is rather vague: nevertheless it points out that the difference between "direct problem" and an "inverse problem" is, to some extent, arbitrary and its rational is to be found in the historical development of that particular problem. At this time, there is no general inverse spectral theory-even the simplest cases require considerable ingenuity for their resolution and none of inverse problems can be really posed unless a class of coefficients is specified in advance.

The direct and inverse scattering problems for the classical Schrodinger operators have been studied extensively (see [1-7] and references therein).

For the Sturm-Liouville operator with different singularities (i.e. on the half-line having an arbitrary number of turning points, having singularities and turning points at the end-points of the interval) the determination of the spectral

function or normalizing constants has been studied by G. Freiling and V.Yurko [8-11] These results are mainly based on Volterra operator transformation and contour integrations.

The string equation or wave equation in layered medium was investigated by M.G. Krein [12], A.S. Blagoveshchensky [13], N.I.Grinberg [14], and others.

M.I.Belishev [15] first considered (1) with $p(x) = 0, q(x) = 0$ and solved the inverse problem of reconstructing the $\rho(x)$ of a finite inhomogeneous string from the frequencies and energies of its normalized characteristic vibrations in the case where $\rho(x)$ can change sign (is indefinite).

The inverse spectral problem of recovering pencils of second-order differential operators on the half-line with turning points was studied by V.Yurko in [12], where it was established properties of spectral characteristics, gave formulation of the inverse problem and prove a uniqueness theorem for solution of the inverse problem.

The considered problem is of interest not only for its own sake, but also because there are other inverse problems in physics which can be reduced to it. For example, kinetic equation after separation of variables reduce Sturm-Liouville eigenvalue problems with an indefinite function.

In neutron transport one may deal with groups of neutrons of different speeds, anisotropic interactions and interactions between different groups of neutrons, which in general leads to nonselfadjoint operator.

In present work we suggested that layered-inhomogeneous medium can also absorption and emit an energy. Therefore, the potentials $p(x), q(x)$ are complex valued.

Complex potentials have been used in the past to model neutron absorption [16,17] and thus the imaginary potential represents emission or absorption. For example, in the crystal model the complex potential is viewed as representing a "prepared" material. The material is assumed to be capable of absorption or emission.

Now we begin to study the inverse problem for a pencil L of non-self-adjoint differential operators pencil generated by formal differential expression

$$l\left(\frac{d}{dx}, \lambda\right) \equiv \frac{1}{\rho(x)} \left\{ -\frac{d^2}{dx^2} + 2\lambda p(x) + q(x) \right\} \quad (2)$$

in the space $L_2(-\infty, \infty, \rho(x))$, λ is complex, and the coefficients $p(x), q(x)$ and $\rho(x)$ are

$$p(x) = \sum_{n=1}^{\infty} p_n e^{inx}; \sum_{n=1}^{\infty} n \cdot |p_n| < \infty \quad (3)$$

$$q(x) = \sum_{n=1}^{\infty} q_n e^{inx}; \sum_{n=1}^{\infty} |q_n| < \infty \quad (4)$$

$$\rho(x) = \begin{cases} 1 & \text{for } x \geq 0, \\ -\beta^2 & \text{for } x < 0. \end{cases} \quad (5)$$

Indefinite differential pencils (1) with coefficients (3-5), produce qualitative modifications in the investigations of the inverse problem.

The feature of our case is the fact that an operator of transformation is absent and because the function $\rho(x)$ is nonsmooth, the wave equation is cannot be reduced to the Schrodinger equation by change of variables.

Firstly potential (4) was considered by M.G.Gasymov[18]. As a final remark, we mention some other works related to the potentials (3)-(4)[20-23].

Note that at $p(x) = 0$ given problem was investigated in the work [23].

By taking into account these point our aim in this paper is study of the spectrum and also to solve the inverse problem for singular non-self-adjoint operators by transmission coefficient and normalizing numbers corresponding to quasi-eigenfunctions of the Sturm- Liouville operator with complex periodic potential and discontinuous coefficients on the axis.

1 We will investigate the representation of fundamental solutions and some relations between these solutions which will be very useful for solving of an inverse problem. In our investigations the basic role will be played a good representation of the solutions of the basic equation (1). As the potential allows bounded analytic continuation to the upper half of the complex plane , we can conduct detailed analysis of problem (2)-(5).

2 We will study the spectrum of the problem (2)-(5). As our solutions have an obvious form that kernel of the resolvent will have a simple representation and consequently it will be possible to study its analytical properties in detail.

3 We will put and solve the inverse problem. As it was pointed out in the introduction some of the authors for solving of the inverse problems employed operator transformations and studied solvability of the obtained Gelfand-Levitan type integral equations. In our case, to solve the inverse problem we have to solve a system of algebraic equations. So inverse problem has a unique solution and the numbers q_n, p_n and β are defined constructively by the spectral data.

REPRESENTATION OF FUNDAMENTAL SOLUTIONS.

Here we study the solutions of the main equation

$$-y''(x) + 2\lambda p(x)y(x) + q(x)y(x) = \lambda^2 \rho(x) y(x)$$

that will be convenient in future. We can prove the existence of these solutions if the condition (3)-(5) is fulfilled for the potential. This will be unique restriction on the potential and later on we'll consider it to be fulfilled.

Theorem 1. Let $q(x)$ and $p(x)$ be of the form (3), (4) and $\rho(x)$ satisfy condition (5). Then equation $Ly = \lambda^2 y$ has special solutions of the form

$$f_1^\pm(x, \lambda) = e^{\pm i\lambda x} \left(1 + \sum_{n=1}^{\infty} v_n^\pm e^{inx} + \sum_{n=1}^{\infty} \sum_{\alpha=n}^{\infty} \frac{v_{n\alpha}^\pm}{n \pm 2\lambda} e^{i\alpha x} \right) \text{ for } x \geq 0 \quad (6)$$

and

$$f_2^\pm(x, \lambda) = e^{\pm\lambda\beta x} \left(1 + \sum_{n=1}^{\infty} v_n^\pm e^{inx} + \sum_{n=1}^{\infty} \sum_{\alpha=n}^{\infty} \frac{v_{n\alpha}^\pm}{n \mp 2i\lambda\beta} e^{i\alpha x} \right) \text{ for } x < 0 \quad (7)$$

where the numbers v_n^\pm and $v_{n\alpha}^\pm$ are determined from the following recurrent relations

$$\alpha^2 v_\alpha^\pm + \alpha \sum_{n=1}^{\alpha} v_{n\alpha}^\pm + \sum_{s=1}^{\alpha-1} \left(q_{\alpha-s} v_s^\pm - p_{\alpha-s} \sum_{n=1}^s v_{ns}^\pm \right) + q_\alpha = 0, \quad (8)$$

$$\alpha(\alpha - n) v_{n\alpha}^\pm + \sum_{s=n}^{\alpha-1} (q_{\alpha-s} \mp n \cdot p_{\alpha-s}) v_{ns}^\pm = 0, \quad (9)$$

$$\alpha v_\alpha^\pm \pm \sum_{s=1}^{\alpha-1} v_s^\pm p_{\alpha-s} \pm p_\alpha = 0. \quad (10)$$

and the series (6,7) admits double termwise differentiation.

Remark1: If $\lambda \neq -\frac{n}{2}$ and $Im\lambda > 0$, then $f_1^+(x, \lambda) \in L_2(0, +\infty)$.

Remark2: If $\lambda \neq -\frac{in}{2\beta}$ and $Re\lambda > 0$, then $f_2^+(x, \lambda) \in L_2(-\infty, 0)$.

By analogy [22], it is easy to see that equation (1) for $\lambda \neq 0, \lambda \neq \pm\frac{n}{2}, \lambda \neq \pm\frac{in}{2\beta}$ has fundamental solutions $f_1^+(x, \lambda), f_1^-(x, \lambda)$ ($f_2^+(x, \lambda), f_2^-(x, \lambda)$) for which

$$W[f_1^+(x, \lambda), f_1^-(x, \lambda)] = 2i\lambda,$$

$$W[f_2^+(x, \lambda), f_2^-(x, \lambda)] = -2i\lambda\beta$$

(where $W[f, g] = f'g - fg'$) is satisfied

Then use the conjunction conditions

$$y(0+) = y(0-),$$

$$y'(0+) = y'(0-)$$

we get that each solution of equation (1) may be represented as linear combinations of these solutions

$$f_2^+(x, \lambda) = C_{11}(\lambda) f_1^+(x, \lambda) + C_{12}(\lambda) f_1^-(x, \lambda), \text{ for } x \geq 0, \quad (11)$$

$$f_1^+(x, \lambda) = C_{22}(\lambda) f_2^+(x, \lambda) + C_{21}(\lambda) f_2^-(x, \lambda), \text{ for } x < 0, \quad (12)$$

were

$$C_{11}(\lambda) = \frac{W[f_2^+(0, \lambda), f_1^-(0, \lambda)]}{2i\lambda},$$

$$C_{12}(\lambda) = \frac{W[f_1^+(0, \lambda), f_2^+(0, \lambda)]}{2i\lambda},$$

$$C_{22}(\lambda) = \frac{W[f_1^+(0, \lambda), f_2^-(0, \lambda)]}{2\beta\lambda}, \quad (13)$$

$$C_{21}(\lambda) = \frac{W[f_2^+(0, \lambda), f_1^+(0, \lambda)]}{2\beta\lambda}. \quad (14)$$

By (13) (14) the coefficients $C_{22}(\lambda)$, $C_{21}(\lambda)$ can be predetermined and at $x \geq 0$ we get

$$C_{22}(\lambda) = \frac{i}{\beta}C_{11}(-\lambda), C_{21}(\lambda) = -\frac{i}{\beta}C_{12}(\lambda). \quad (15)$$

Let

$$f_n^\pm(x) = \lim_{\lambda \rightarrow \mp \frac{n}{2}} (n \pm 2\lambda) f_1^\pm(x, \lambda) = \sum_{\alpha=n}^{\infty} v_{n\alpha}^\pm e^{i\alpha x} e^{-i\frac{n}{2}x}, \quad (16)$$

It follows from relation (6) that $f_n^\pm(x) \neq 0$ is valid for $v_{n\alpha}^\pm \neq 0$.

From this we obtained that $W[f_n^\pm(x), f_1^\mp(x, \mp \frac{n}{2})] = 0$ and consequently the functions $f_n^\pm(x), f_1^\mp(x, \mp \frac{n}{2})$ are linear dependent.

Therefore

$$f_n^\pm(x) = S_n^\pm f_1^\mp\left(x, \frac{n}{2}\right). \quad (17)$$

Comparing these formulas we get that $S_n^\pm = v_{nn}^\pm$.

SPECTRUM OF THE OPERATOR L

To study the spectrums of the operator L at first we calculate the kernel of the resolvent of the operator $(L - \lambda^2 I)$. Divided the plane λ into sectors

$$S_k = \{k\pi/2 < \arg \lambda < (k+1)\pi/2\}, k = \overline{0, 3}.$$

By means of general methods for the kernel of the resolvent, we get

$$R_{11}(x, t, \lambda) = \frac{1}{W[f_1^+, f_2^+]} \begin{cases} f_1^+(x, \lambda) f_2^+(t, \lambda) & \text{for } t < x \\ f_1^+(t, \lambda) f_2^+(x, \lambda) & \text{for } t > x \end{cases} \quad \lambda \in S_0, \quad (18)$$

$$R_{12}(x, t, \lambda) = \frac{1}{W[f_1^+, f_2^-]} \begin{cases} f_1^+(x, \lambda) f_2^-(t, \lambda) & \text{for } t < x \\ f_1^+(t, \lambda) f_2^-(x, \lambda) & \text{for } t > x \end{cases} \quad \lambda \in S_1, \quad (19)$$

$$R_{21}(x, t, \lambda) = \frac{1}{W[f_1^-, f_2^-]} \begin{cases} f_1^-(x, \lambda) f_2^-(t, \lambda) & \text{for } t < x \\ f_1^-(t, \lambda) f_2^-(x, \lambda) & \text{for } t > x \end{cases} \quad \lambda \in S_2, \quad (20)$$

$$R_{22}(x, t, \lambda) = \frac{1}{W[f_1^-, f_2^+]} \begin{cases} f_1^-(x, \lambda) f_2^+(t, \lambda) & \text{for } t < x \\ f_1^-(t, \lambda) f_2^+(x, \lambda) & \text{for } t > x \end{cases} \quad \lambda \in S_3. \quad (21)$$

Remark 3 : $R_\lambda = (L - \lambda^2 I)^{-1}$ exists and bounded for all λ out of $\{Re\lambda = 0\} \cup \{Im\lambda = 0\}$ and $C_{11}(\pm\lambda) \neq 0$, $C_{12}(\pm\lambda) \neq 0$.

Theorem 2. L has no eigenvalues for real and pure imaginary λ . It's continuous spectra consist of axes $Re\lambda = 0$ and $Im\lambda = 0$ on which there may exist spectral singularities coinciding with the numbers $\frac{in}{2\beta}$, $\frac{n}{2}$, $n = \pm 1, \pm 2, \pm 3, \dots$

Proof:

We note that equation (1) has fundamental solutions $f_1^+(x, \lambda), f_1^-(x, \lambda), f_2^+(x, \lambda), f_2^-(x, \lambda)$ on $|Im\lambda| < \frac{\varepsilon}{2}$ ($|Re\lambda| < \frac{\varepsilon}{2}$).

Then for $Im\lambda = 0$ solution of equation (1) can be written in the form

$$y(x, \lambda) = C_1 f_1^+(x, \lambda) + C_2 f_1^-(x, \lambda)$$

In case $Im\lambda = 0$ the solution $f_1^\pm(x, \lambda)$ has the form

$$f_1^\pm(x, \lambda) = e^{\pm i Re\lambda x} \left(1 + \sum_{n=1}^{\infty} v_n^\pm e^{inx} + \sum_{n=1}^{\infty} \frac{1}{n \pm 2\lambda} \sum_{\alpha=n}^{\infty} V_{n\alpha} e^{i\alpha x} \right)$$

then $y(x, \lambda) \in L_2(-\infty, +\infty)$ except when $C_1 = C_2 = 0$.

Analogously we can prove case $Re\lambda = 0$.

Since in $|Re\lambda| < \frac{\varepsilon}{2}$ the functions $f_2^+(x, \lambda), f_2^-(x, \lambda)$ form fundamentals solutions, then

$$y(x, \lambda) = C_3 f_2^+(x, \lambda) + C_4 f_2^-(x, \lambda) .$$

If $Re\lambda = 0$ then the solution $f_2^\pm(x, \lambda)$ has the form

$$f_2^\pm(x, \lambda) = e^{\pm i Im\lambda \beta x} \left(1 + \sum_{n=1}^{\infty} v_n^\pm e^{inx} + \sum_{n=1}^{\infty} \frac{1}{n \pm 2Im\lambda\beta} \sum_{\alpha=n}^{\infty} V_{n\alpha} e^{i\alpha x} \right)$$

then $y(x, \lambda) \in L_2(-\infty, +\infty)$ except when $C_3 = C_4 = 0$.

Taking into account Remark 2, we'll investigate the function

$$R(x, t, \lambda) = \begin{cases} R_{11}(x, t, \lambda) & , \quad \lambda \in S_0 \\ R_{12}(x, t, \lambda) & , \quad \lambda \in S_1 \\ R_{21}(x, t, \lambda) & , \quad \lambda \in S_2 \\ R_{22}(x, t, \lambda) & , \quad \lambda \in S_3 \end{cases}$$

in the neighborhood of poles λ^* from $\{Re\lambda = 0\} \cup \{Im\lambda = 0\}$. Without loss of generality we can investigate $R(x, t, \lambda)$ in the neighborhood of poles λ_0^* from $[0, \infty)$. Then the number λ_0^* coincides with one of the numbers $(\frac{n}{2})$, $n \in N$. From (18)-(21) it follows that the limit $\lim_{\lambda \rightarrow \lambda_0^*} (\lambda - \lambda_0^*) R(x, t, \lambda) = R_0(x, t)$

exists and $R_0(x, t)$ is a bounded function with respect to all the variables.

Let $\theta(x)$ be an arbitrary finite function. Then $\varphi(x) = \int_{-\infty}^{+\infty} R_0(x, t)\theta(t) dt$ is a

bounded solution of equation (1) for $\lambda = \lambda_0^*$. Therefore $\varphi(x) = C_0 f_1^+(x, \lambda_0^*)$. Comparison of the last relation with formulae (18)-(21) shows that if $\lambda_0^* \neq \frac{n}{2}, n \in N$ then $C_0 = 0$ and so the kernel of the resolvent has removable singularity at the point λ_0^* . So there is a case $\lambda_0^* = \frac{n}{2}, n \in N$ where the kernel of resolvent has poles of the first order. Since $f_1^+(x, \lambda_0^*) \notin L_2(-\infty, +\infty)$ then $\lambda_0^* = (\frac{n}{2})$ is a spectral singularity of the operator L in the sense of M.N.Naimark [24, p.450].(Analogously we can show that the resolvent has poles of the first order at the points $\frac{in}{2\beta}, \frac{n}{2}, n = \pm 1, \pm 2, \pm 3, \dots$ which are spectral singularities of the operator L).

In order to show that all numbers from $\{Re\lambda = 0\} \cup \{Im\lambda = 0\}$ belong to the continuous spectra we introduce the Dirichlet condition at $x = 0$, the resulting operator L^0 splits into the direct sum of corresponding Dirichlet operators L^-, L^+ on the negative and positive semi-axes. Therefore the essential spectrum of L^0 is $\{Re\lambda = 0\} \cup \{Im\lambda = 0\}$, and the initial operator L has the same essential spectrum. Since eigenvalues of infinite multiplicities are not possible, the essential spectrum of L coincide with the continuous one. The Theorem is proved.

Lemma 1. The eigenvalues of operator L are finite and coincide with zeros of the functions $C_{12}(\lambda), C_{11}(-\lambda), C_{12}(-\lambda), C_{11}(\lambda)$ from sectors $S_k = \{k\pi/2 < \arg \lambda < (k+1)\pi/2\}, k = \overline{0, 3}$ respectively.

Definition 1. The data $\{C_{11}(\lambda), C_{12}(\lambda)\}$ are called the spectral data of L .

INVERSE PROBLEM

From the representation (18)-(21) it also follows that for each x and t from $(-\infty, +\infty)$ the kernel $R(x, t, \lambda)$ admits a meromorphic continuation from the sectors $S_k, k = 0, 1, 2, 3$ and may have poles at the points $(\pm \frac{n}{2})$ and $(\pm \frac{in}{2\beta})$, $n \in N$, outside of $S_k, k = 0, 1, 2, 3$. Such poles of the resolvent are called quasi-stationary states of the operator L . Thus the quasi-stationary states of the operator L are the numbers $(\pm \frac{n}{2})$ and $(\pm \frac{in}{2\beta})$, $n \in N$. The numbers $V_{nn}, n \in N$ play a part of normalizing numbers corresponding to quasi-eigenfunction of the operator L . So, it is make natural the formulation of the inverse problem about reconstruction of the potential of the equation (1) and the number β .

Let's study the inverse problem for the problem (1-3). The inverse problem is formulated as follows.

Given the spectral data $\{C_{11}(\lambda), C_{12}(\lambda)\}$ construct the potential $q(x), p(x)$ and β .

Using the results obtained above we arrive at the following procedure for solution of the inverse problem.

1. Taking into account (16) it is easy to check that

$$\lim_{\lambda \rightarrow n/2} (n-2\lambda) \frac{C_{11}(\lambda)}{C_{12}(\lambda)} = v_{nn}^-, \quad \lim_{\lambda \rightarrow -n/2} (n+2\lambda) \frac{C_{12}(\lambda)}{C_{11}(\lambda)} = v_{nn}^+$$

consequently we find all numbers v_{nn}^\pm , $n = 1, 2, \dots$

2. Taking into account (16) we get

$$\sum_{\alpha=B}^{\infty} v_{n\alpha}^\pm e^{i\alpha x} e^{-i\frac{n}{2}x} = v_{nn}^\pm \cdot e^{i\frac{n}{2}x} \left(1 + \sum_{n=1}^{\infty} v_n^\mp e^{in x} + \sum_{n=1}^{\infty} \sum_{\alpha=n}^{\infty} \frac{v_{n\alpha}^\mp}{n+n} e^{i\alpha x} \right)$$

or

$$v_{m,\alpha+m}^\pm = v_{mm}^\pm \left(v_\alpha^\mp + \sum_{n=1}^{\alpha} \frac{v_{n\alpha}^\mp}{n+m} \right)$$

from which all numbers $v_{n\alpha}^\pm$, $\alpha = 1, 2, \dots$, $n = 1, 2, \dots, n < \alpha$ are defined.

3. Then from recurrent formula (8-10), find all numbers p_n and q_n .

4. The number β is defined from equality

$$\beta = iC_{11}(\lambda_n) C_{11}(-\lambda_n).$$

Really from (11-12) we derive for $\lambda_n \in S_0$ which is a root of the equation $C_{12}(\lambda) = 0$ the true relation

$$C_{11}(\lambda_n) = \frac{f_2^+(x, \lambda_n)}{f_1^+(x, \lambda_n)}, \quad C_{22}(\lambda_n) = \frac{f_1^+(x, \lambda_n)}{f_2^+(x, \lambda_n)} \text{ i.e. } C_{11}(\lambda_n) C_{22}(\lambda_n) = 1.$$

Then we get $\beta = iC_{11}(\lambda_n) C_{11}(-\lambda_n)$.

So inverse problem has a unique solution and the numbers q_n, p_n and β are defined constructively by the spectral data.

Theorem 3. The specification of the spectral data uniquely determines potentials $p(x)$, $q(x)$ and β .

REFERENCES.

- 1 T. Aktosun, M. Klaus, and C. van der Mee, Wave scattering in one dimension with absorption, J. Math. Phys. 39, 1957-1992, 1998.
- 2 T. Aktosun, M. Klaus, and C. van der Mee, Inverse scattering in one-dimensional nonconservative media, Integral Equations and Operator Theory 30, 297-316, 1998
3. Jaulent M. and C. Jean, The inverse s-wave scattering problem for a class of potentials depending on energy, Commun. Math. Phys. 28, 177-220, 1972.
4. Jaulent M. and C. Jean, The inverse problem for the one-dimensional Schrodinger equation with an energy-dependent potential. I, Ann. Inst. Henri Poincare, Sect. A 25, 105-118, 1976.
5. Jaulent M and C. Jean, The inverse problem for the one-dimensional Schrodinger equation with an energy-dependent potential. II, Ann. Inst. Henri Poincare, Sect. A 25, 119-137, 1976.
6. Jaulent M., Inverse scattering problems in absorbing media, J. Math. Phys. 17, 1351-1360, 1976.

7. Pivovarchik V.N. Reconstruction of the potential of the Sturm-Liouville equation from three spectra of - Functional Analysis and Its Applications, 1999
8. Freiling G, Yurko V Inverse problems for differential equations with turning points - Inverse problems, 1997
9. Eberhard W. and Freiling G., The distribution of the eigenvalues for second order eigenvalue problems in the presence of an arbitrary number of turning points, Results in Mathematics, 21, 24-41, 1992
10. Freiling G. and Yurko V.A. , Inverse Sturm-Liouville Problems and their Applications, NOVA Science Publishers, New York, 2001.
11. Yurko V Inverse spectral problems for differential pencils on the half-line with turning points - Journal of Mathematical Analysis and Applications, 2006
12. Krein MG On determination of the potential of a particle from its S-function - Dokl. Akad. Nauk SSSR, 1955
13. Blagoveshchensky A.S., A one-dimensional inverse boundary value problem for a second order hyperbolic equation. (Russian) Zap. Nauch. Sem. Leningrad. Otdel. Mat. Inst. Steklov. (LOMI) 15 1969 85–90.
14. Grinberg N. I. 1991 The one-dimensional inverse scattering problem for the wave equation, Math. USSR Sb. 70, 557-572, [Mat. Sb. 181 (8), 1114-1129 (1990) (Russian)].
15. Belishev MI Inverse spectral indefinite problem for the equation $y'' + \lambda p(x)y = 0$ on an interval - Functional Analysis and Its Applications, 1987
16. Bethe H.A A continuum theory of the compound nucleus - Physical Review, 57, 1125, 1940
17. Feshbach H , Porter CE, Weisskopf VF Model for nuclear reactions with neutrons - Physical Review, 1954
18. Gasymov M.G. Spectral analysis of a class non-self-adjoint operator of the second order. Functional analysis and its applications. (In Russian), V34.1. pp.14-19, 1980
19. Gasymov MG., Gusejnov GS Determination of diffusion operators according to spectral data- Dokl., Akad. Nauk Az. SSR, (In Russian) 1981
20. Efendiev, R.F. The Characterization Problem for One Class of Second Order Operator Pencil with Complex Periodic Coefficients. Moscow Mathematical Journal, 2007, Volume 7, Number 1, pp.55-65.
21. Efendiev, R. F. Complete solution of an inverse problem for one class of the high order ordinary differential operators with periodic coefficients. Zh. Mat. Fiz. Anal. Geom. 2006, V.2, no. 1, 73–86,
22. Efendiev, R.F. Spectral analysis of a class of non-self-adjoint differential operator pencils with a generalized function. Teoreticheskaya i Matematicheskaya Fizika, 2005, Vol.145, 1. pp.102-107, October (in Russian). Theoretical and Mathematical Physics, 145(1):1457-461, (English).
23. Efendiev, R. F Inverse indefinite spectral problem for second order differential operator with complex periodic coefficients- Proceeding of Baku State University, (In Russian), 2008, 4, 17-22
24. Naimark, M.A. Linear differential operators. Moscow, Nauka, 1969 (Russian).