

Consistent classical and quantum mixed dynamics

Michael J. W. Hall
Theoretical Physics, IAS,
Australian National University,
Canberra ACT 0200, Australia

A recent proposal for mixed dynamics of classical and quantum ensembles is shown, in contrast to other proposals, to satisfy the minimal algebraic requirements proposed by Salcedo for any consistent formulation of such dynamics. It is further shown that additional desirable requirements, related to separability, may be satisfied by imposing a weak ‘independence’ constraint on the class of physical ensembles. Although the mixed dynamics is formulated in terms of ensembles on configuration space, thermodynamic mixtures of such ensembles may be defined which are equivalent to canonical phase space ensembles on the classical sector. Hence, the formulation appears to be both consistent and physically complete.

PACS numbers: 03.65.Ta

I. INTRODUCTION

Many proposals have been made for mixing classical and quantum dynamics [1, 2]. These proposals may be broadly classified into mean-field, phase-space and trajectory categories, and all fail some important criterion - such as conservation of energy and probability, back-reaction by the quantum component on the classical component, positivity of probabilities, correct equations of motion in the limit of no interaction, and describing all interactions of physical interest [2, 3, 4, 5, 6, 7, 8, 9, 10, 11]). Hence, while often of practical interest for calculations in molecular dynamics [10, 11, 12], the above proposals have not led to a formulation of mixed classical and quantum dynamics that may be regarded as physically fundamental. Such a formulation would not only be of interest for making physically-consistent numerical calculations, but also for modelling ‘classical’ measurement apparatuses and coupling quantum systems to classical spacetime metrics [4].

Recently, a new proposal has been made that falls outside the above categories [13]. It is based on the description of physical systems by ensembles on configuration space, and satisfies all of the abovementioned criteria. Moreover, it has been successfully applied to discuss position and spin measurements, interacting classical and quantum oscillators, and the coupling of quantum fields to classical spacetime [13, 14]. However, while this ‘configuration-ensemble’ approach is therefore a promising formulation of mixed dynamics, only particular observable properties, such as position, momentum, energy and angular momentum, have been discussed in any detail. Hence, further investigation is necessary to determine whether it is fully self-consistent.

In this regard, Salcedo has recently specified two “*minimal requirements for a consistent classical-quantum formulation*” [10]:

- (i) a Lie bracket may be defined on the set of observables, and
- (ii) the Lie bracket is equivalent to the classical Poisson

bracket for any two classical observables, and to $(i\hbar)^{-1}$ times the quantum commutator for any two quantum observables.

These have been previously justified on physical grounds by Caro and Salcedo [6], and are highly nontrivial: none of the abovementioned proposals, in those cases where they have been sufficiently developed to identify the general ‘classical’ and ‘quantum’ observables of the theory, satisfy both requirements [6, 10, 11]. These minimal requirements therefore provide a critical test for the configuration-ensemble approach to mixed dynamics.

In Sec. II it is shown that the configuration-ensemble formulation, unlike other proposals, *does* pass the above test. In particular, a Lie algebra of observables may be defined that satisfies requirements (i) and (ii) above.

Moreover, it is shown in Sec. III that the additional reasonable requirement

- (iii) The Lie bracket of any quantum observable with any classical observable vanishes,

may also be satisfied, by imposing a weak ‘independence’ constraint on the set of physical ensembles. This requirement has been suggested previously by Caro and Salcedo [6] (see equation (18) thereof), and implies, for example, that the classical and quantum components of a mixed ensemble will evolve independently under a Hamiltonian that is the sum of a classical and a quantum Hamiltonian. The consistent description of measurement interactions is also discussed in Sec. III.

Finally, in Sec. IV it is shown that the configuration-ensemble approach is consistent with thermodynamics. In particular, a generalised ‘canonical ensemble’ may be defined as a suitable mixture of distinguishable stationary ensembles on configuration space, which is equivalent to the usual canonical ensemble on phase space for an ergodic classical system, and to the usual canonical ensemble on Hilbert space for a quantum system. This demonstrates that the formulation of the approach on configuration space is not a barrier to describing all physically relevant systems (and also implies classical statistical mechanics may be given a Hamilton-Jacobi formulation).

II. OBSERVABLES IN THE CONFIGURATION-ENSEMBLE APPROACH

A. General observables

The description of physical systems by ensembles on configuration space may be introduced at quite a fundamental and generic level [13, 15]. The starting point is simply a probability density P on the configuration space of the system, the dynamics of which is assumed to satisfy an action principle. Thus, there is a canonically conjugate quantity, S , on the configuration space, and an ‘ensemble’ Hamiltonian, $\tilde{H}[P, S]$, such that

$$\frac{\partial P}{\partial t} = \frac{\delta \tilde{H}}{\delta S}, \quad \frac{\partial S}{\partial t} = -\frac{\delta \tilde{H}}{\delta P}. \quad (1)$$

Here $\delta/\delta f$ denotes the appropriate variational derivative on the configuration space. For the particular case of a continuous configuration space, indexed by ‘position coordinate’ ξ , and a functional $L[f]$ of the form $L[f] = \int d\xi F(f, \nabla f, \xi)$, one has the useful formula [16]

$$\delta L/\delta f = \partial F/\partial f - \nabla \cdot [\partial F/\partial(\nabla f)].$$

As a simple example, the ensemble Hamiltonian

$$\tilde{H}_Q = \int dq P \left[\frac{|\nabla S|^2}{2m} + \frac{\hbar^2}{8m} \frac{|\nabla P|^2}{P^2} + V(q) \right]$$

describes a quantum spin-zero particle of mass m , moving under a potential $V(q)$. In particular, the equations of motion (1) reduce in this case to the real and imaginary parts of the Schrödinger equation

$$i\hbar(\partial\psi/\partial t) = [-\hbar^2/2m)\nabla^2 + V] \psi,$$

where $\psi(q) := P^{1/2}e^{iS/\hbar}$. Moreover, if the limit $\hbar \rightarrow 0$ is taken in the ensemble Hamiltonian \tilde{H}_Q , the equations of motion for P and S reduce to the Hamilton-Jacobi and continuity equations for an ensemble of classical particles [13]. Thus, in the configuration-ensemble approach, the primary difference between quantum and classical mechanics lies in the choice of the ensemble Hamiltonian.

While a number of properties and applications of the configuration ensemble formalism have been previously considered, the description of ‘observables’ has only been briefly alluded to, with emphasis on particular quantities such as position, momentum, spin and energy [13]. The general description is therefore addressed here, to enable the properties of observables for mixed configuration ensembles to be compared against the two minimal requirements discussed in the Introduction.

Note first that the conjugate pair (P, S) allows a Poisson bracket to be defined for any two functionals $A[P, S]$ and $B[P, S]$, via [16]

$$\{A, B\} := \int d\xi \left(\frac{\delta A}{\delta P} \frac{\delta B}{\delta S} - \frac{\delta B}{\delta P} \frac{\delta A}{\delta S} \right) \quad (2)$$

(where integration is replaced by summation over any discrete parts of the configuration space). Thus, the equations of motion (1) may be written as $\partial P/\partial t = \{P, \tilde{H}\}$ and $\partial S/\partial t = \{S, \tilde{H}\}$, and more generally it follows that

$$\partial A/\partial t = \{A, \tilde{H}\}$$

for any functional $A[P, S]$. The Poisson bracket is well known to be a Lie bracket [16], and in particular is linear, antisymmetric, and satisfies the Jacobi identity. Hence, the first minimal consistency requirement, given in Sec. I above, is automatically satisfied by choosing the observables to be any set of functionals of P and S that is closed with respect to the Poisson bracket.

It is important to note that an *arbitrary* functional $A[P, S]$ will not be allowable as an observable in general. This is analogous to the restriction of expectation values to bilinear forms of linear Hermitian operators in standard quantum mechanics, even though a commutator bracket can be more generally defined [17]. Thus, for example, the infinitesimal canonical transformation

$$P \rightarrow P + \epsilon \delta A/\delta S, \quad S \rightarrow S - \epsilon \delta A/\delta P$$

generated by any observable A must preserve the interpretation of P as a probability density, i.e., the normalisation and positivity of P must be preserved. This imposes the respective fundamental conditions [13]

$$A[P, S + c] = A[P, S], \quad \delta A/\delta S = 0 \text{ if } P(\xi) = 0, \quad (3)$$

where c is an arbitrary constant.

Note that each of the above conditions is consistent with the Poisson bracket. First, defining $I[P, S] := \int d\xi P$, the normalisation condition is simply the requirement that I is invariant under allowed canonical transformations, i.e., that $\delta I = \epsilon\{I, A\} = 0$. Hence, if it holds for two observables A and B , then it automatically holds for $\{A, B\}$ via the Jacobi identity, since

$$\{I, \{A, B\}\} = -\{A, \{B, I\}\} - \{B, \{I, A\}\} = 0.$$

Similarly, the positivity condition may be rewritten as $\delta P = \epsilon\{P, A\} = 0$ whenever $P(\xi) = 0$ (otherwise $P(\xi)$ can be decreased below 0 by choosing the sign of ϵ appropriately), which again holds for $\{A, B\}$, if it holds for A and B , as a consequence of the Jacobi identity.

It follows that the observables corresponding to a configuration-ensemble description should be chosen as some set of functionals satisfying the normalisation and positivity constraints (3), that is closed with respect to the Poisson bracket in Eq. (2). Under any such choice, the first minimal requirement (i) in Sec. I is automatically satisfied, where the Lie bracket is identified with the Poisson bracket.

B. Classical and quantum observables

To determine whether the second minimal requirement (ii) in Sec. I is also satisfied, it is necessary to

first define the ‘classical’ and ‘quantum’ observables of a mixed quantum-classical ensemble.

Consider, therefore, a mixed quantum-classical ensemble, indexed by the joint configuration $\xi = (q, x)$, where q labels the quantum configuration and x labels the classical configuration. For example, q may refer to the position of a quantum system, or, more generally, label some complete set of kets $\{|q\rangle\}$ [15]. In contrast, x will always be taken here to refer to some continuous set of coordinates on a classical configuration space (to enable the discussion of classical phase space relations). Hence, the mixed ensemble is described by two conjugate quantities $P(q, x, t)$ and $S(q, x, t)$.

First, for any real classical phase space function $f(x, k)$, define the corresponding classical observable C_f by

$$C_f := \int dq dx P f(x, \nabla_x S). \quad (4)$$

(where integration with respect to q is replaced by summation over any discrete portions of the quantum configuration space). This is similar in form to a classical average, and hence the numerical value of C_f will be identified with the predicted expectation value of the corresponding function $f(x, k)$. It is easily checked that C_f satisfies the required normalisation and positivity conditions (3). Note that, for classical observables, $\nabla_x S$ plays the role of a momentum associated with the configuration (q, x) .

The Poisson bracket of any two classical observables C_f and C_g follows, using Eq. (2) and integration by parts with respect to x , as

$$\begin{aligned} \{C_f, C_g\} &= \int dq dx [-f \nabla_x \cdot (P \nabla_k g) + g \nabla_x \cdot (P \nabla_k f)] \\ &= \int dq dx P (\nabla_x f \cdot \nabla_k g - \nabla_x g \cdot \nabla_k f) \\ &= C_{\{f, g\}}, \end{aligned} \quad (5)$$

where all quantities in the integrands are evaluated at $k = \nabla_x S$, and $\{f, g\}$ denotes the usual Poisson bracket for phase space functions. Hence, the Lie bracket for classical observables is equivalent to the usual phase space Poisson bracket, as required. Given that the observables are evaluated on a configuration space, rather than on a phase space, this is a somewhat remarkable result.

Second, for any Hermitian operator M acting on the Hilbert space spanned by the kets $\{|q\rangle\}$, define the corresponding quantum observable Q_M by

$$\begin{aligned} Q_M &:= \int dq dx \psi^*(q, x) M \psi(q, x) \\ &= \int dq dq' dx (PP')^{1/2} e^{i(S-S')/\hbar} \langle q' | M | q \rangle, \end{aligned} \quad (6)$$

where $\psi(q, x) := P(q, x)^{1/2} e^{iS(q, x)/\hbar}$, $P = P(x, q)$, $P' = P(x, q')$, etc (and where integration with respect to q and q' is replaced by summation over any discrete portions of the quantum configuration space). This is similar in form

to a quantum average, with respect to the ‘hybrid wavefunction’ $\psi(q, x)$, and hence the numerical value of Q_M will be identified with the predicted expectation value of the corresponding operator M . It follows immediately from the second equality in Eq. (6) that Q_M satisfies the normalisation and positivity conditions (3).

To evaluate the Poisson bracket of any two quantum observables Q_M and Q_N , it is convenient to first express the Poisson bracket in terms of the hybrid wavefunction $\psi(q, x)$ and its complex conjugate $\psi^*(q, x)$. One has in particular for any real functional $A[P, S]$ that

$$\frac{\delta A}{\delta P} = \frac{\partial \psi}{\partial P} \frac{\delta A}{\delta \psi} + \frac{\partial \psi^*}{\partial P} \frac{\delta A}{\delta \psi^*} = \frac{1}{\psi^* \psi} \operatorname{Re} \left\{ \psi \frac{\delta A}{\delta \psi} \right\},$$

$$\frac{\delta A}{\delta S} = \frac{\partial \psi}{\partial S} \frac{\delta A}{\delta \psi} + \frac{\partial \psi^*}{\partial S} \frac{\delta A}{\delta \psi^*} = -\frac{2}{\hbar} \operatorname{Im} \left\{ \psi \frac{\delta A}{\delta \psi} \right\},$$

and hence, noting $-ad + bc = \operatorname{Im}\{(a + ib)(c - id)\}$, that

$$\{A, B\} = \frac{2}{\hbar} \operatorname{Im} \left\{ \int dq dx \frac{\delta A}{\delta \psi} \frac{\delta B}{\delta \psi^*} \right\}. \quad (7)$$

Recalling that M and N are Hermitian, so that $\psi^* M \psi$ may be replaced by $(M\psi)^* \psi$ in equation (6), it immediately follows that

$$\{Q_M, Q_N\} = \frac{2}{\hbar} \operatorname{Im} \left\{ \int dq dx (M\psi)^* N \psi \right\} = Q_{[M, N]/(i\hbar)}, \quad (8)$$

where $[M, N]$ denotes the usual quantum commutator $MN - NM$. Hence, the Lie bracket for quantum observables is equivalent to the usual quantum commutator, as required.

Eqs. (2), (5) and (8) are the main results of this section. They imply that any set of observables containing the quantum observables C_f and the quantum observables Q_M , that is closed under the Poisson bracket, will satisfy both of the minimal requirements (i) and (ii) in Sec. I for a consistent mixed quantum-classical formulation. In the following sections, it will be shown that further desirable properties can also be accommodated within the configuration-ensemble formulation.

III. INTERACTION AND MEASUREMENT

In contrast to other proposals for mixed dynamics, the configuration-ensemble approach has been shown to pass the critical test of meeting the two minimal consistency requirements (i) and (ii) in Sec. I. However, while the approach thereby gains a special status, there are a number of other requirements that may reasonably be expected of a physical theory. Some of these, related to local aspects of interaction and measurement, are discussed below.

A. An independence constraint

One natural requirement for a mixed quantum-classical formulation is that it is able to consistently describe non-interacting classical and quantum systems. For example, in the case of interacting classical and quantum spin-zero particles having masses m and m' respectively, consider the ensemble Hamiltonian [13]

$$\tilde{H} = \tilde{H}_C + \tilde{H}_Q + \tilde{H}_I, \quad (9)$$

comprising classical and quantum kinetic terms

$$\tilde{H}_C = \frac{1}{2m} \int dq dx P |\nabla_x S|^2,$$

$$\tilde{H}_Q = \frac{1}{2m'} \int dq dx P \left[|\nabla_q S|^2 + \frac{\hbar^2}{4} \frac{|\nabla_q P|^2}{P^2} \right],$$

and an interaction term

$$\tilde{H}_I = \int dq dx P V(q, x, t)$$

for some potential function $V(q, x, t)$. Note that \tilde{H}_C and \tilde{H}_Q correspond to $f(x, k) = |k|^2/(2m)$ and $M = -\hbar^2 |\nabla_q|^2/(2m')$ in Eqs. (4) and (6), respectively, and that the normalisation and positivity conditions (3) are trivially satisfied. One expects that the particular choice $V \equiv 0$ should lead to independent evolution of the classical and quantum components, corresponding to non-interaction between the components. However, a straightforward calculation yields the Poisson bracket

$$\{\tilde{H}_C, \tilde{H}_Q\} = 2\hbar^2 \int dq dx P (\nabla_x S) \cdot \nabla_x (P^{-1/2} \nabla_q^2 P^{1/2}),$$

which does not vanish identically for arbitrary P and S .

There are two possible ways to resolve the above issue. The first is to permit a description in which the coupling between quantum and classical systems can never be turned off. This might be used to advance, for example, a new mechanism for physical decoherence, that relies purely on the existence of classical as well as quantum ensembles. However, this ‘interaction without interaction’ option will not be considered further here. The second possibility is to impose the dynamical constraint that the above Poisson bracket must indeed vanish for all *physical* ensembles. This is closely analogous to the imposition of ‘weak constraints’ in quantum field theory, which similarly determine a class of physical states [18]. It is this second option that will be investigated here.

In particular, consider the assumption that the canonical transformation generated by any classical observable, C_f , when applied to any physical configuration ensemble, (P, S) , cannot alter any quantum observable, Q_M , and vice versa. This implies the weak constraint

$$\{C_f, Q_M\} \approx 0, \quad (10)$$

where \approx is used to denote equality only for the class of *physical* ensembles, similarly to the notation used in quantum constraint theory [18]. Note that this constraint corresponds to requirement (iii) in Sec. I being satisfied for all physical ensembles.

The set of ensembles compatible with the independence constraint (10) is in fact very large, and includes all ensembles likely to appear in any calculation involving the interaction of classical and quantum ensembles. For example, as shown below, it includes any configuration ensemble (P, S) for which the classical and quantum components are *independent* at some time, i.e., for which there is some time t_0 such that [13]

$$P(q, x, t_0) = P_Q(q) P_C(x), \quad S(q, x, t_0) = S_Q(q) + S_C(x). \quad (11)$$

Thus, at time t_0 the ensemble is fully described by two conjugate pairs (P_Q, S_Q) and (P_C, S_C) , corresponding to the quantum and classical components respectively. Note that such independent initial conditions are generally assumed in *any* model of classical-classical and quantum-quantum interactions, including measurement interactions, and thermal interactions between a given system and a heat reservoir.

To show that all ensembles satisfying Eq. (11) automatically satisfy constraint (10), it is convenient to use the representation of the Poisson bracket in Eq. (7). First, from Eq. (4) one has

$$C_f = \int dq dx \psi^* \psi f(x, k), \quad k = \frac{\hbar}{2i} \left(\frac{\nabla_x \psi}{\psi} - \frac{\nabla_x \psi^*}{\psi^*} \right),$$

and hence that

$$\begin{aligned} \frac{\delta C_f}{\delta \psi} &= \psi^* f + \psi^* \psi (\nabla_k f) \cdot \frac{\partial k}{\partial \psi} \\ &\quad - \nabla_x \cdot \left(\psi^* \psi (\nabla_k f) \cdot \frac{\partial k}{\partial (\nabla_x \psi)} \right) \\ &= \psi^* f - \frac{\hbar}{2i} \frac{\psi^*}{\psi} (\nabla_k f \cdot \nabla_x \psi) - \frac{\hbar}{2i} \nabla_x \cdot (\psi^* \nabla_k f). \end{aligned}$$

Second, from Eq. (6) one has $\delta Q_M / \delta \psi^* = M \psi$. Now, Eq. (11) is equivalent to a factorisation $\psi(q, x, t_0) = \psi_Q(q) \psi_C(x)$ of the hybrid wavefunction, implying that $f(x, k)$ is independent of q . Hence, at time t_0 , one obtains

$$\begin{aligned} \int dq dx \frac{\delta C_f}{\delta \psi} \frac{\delta Q_M}{\delta \psi^*} &= \int dq \psi_Q^* M \psi_Q \left\{ \int dx \psi_C^* \psi_C f \right. \\ &\quad \left. - \frac{\hbar}{2i} \int dx [\psi_C^* (\nabla_k f \cdot \nabla_x \psi_C) - \psi_C \nabla_x \cdot (\psi_C^* \nabla_k f)] \right\} \\ &= \int dq \psi_Q^* M \psi_Q \int dx \psi_C^* \psi_C f, \end{aligned}$$

where integration by parts has been used to obtain the final result. This expression is clearly real, implying immediately from Eq. (7) that

$$\{C_f, Q_M\} = 0$$

at time t_0 . Hence, since Poisson brackets are preserved under Hamiltonian evolution, this holds for all times, and the independence constraint (10) is satisfied for $P(q, x, t)$ and $S(q, x, t)$ as claimed.

More generally, it also follows that the constraint is satisfied by any ensemble related to an independent ensemble via some canonical transformation.

It is concluded that requirement (iii) in Sec. I can consistently be accommodated within the configuration-ensemble approach, as a weak constraint on the set of physical ensembles as per Eq. (10).

B. Measurement aspects

Examples of measurements of position and spin on quantum ensembles, via interaction with an ensemble of classical pointers, and the corresponding decoherence of the quantum component relative to the classical component, have been described previously within the configuration-ensemble approach [13]. Here it is noted that a simple model exists for describing the measurement of any quantum observable via a classical measuring apparatus, and that this model is consistent with the weak independence constraint (10).

In particular, the measurement of an arbitrary quantum observable, Q_M , may be modelled by the ensemble Hamiltonian

$$\tilde{H} := \tilde{H}_0 + \kappa(t) \int dq dx \psi^*(q, x) \left(\frac{\hbar}{i} \frac{\partial}{\partial x} \right) M \psi(q, x),$$

where $\kappa(t)$ vanishes outside the measurement period, and x denotes the position of a one-dimensional classical pointer (with integration over q replaced by summation over any discrete values). Note that the interaction term satisfies the normalisation and positivity constraints (3). As will be shown, this term correlates the eigenvalues of M with the position of the pointer, in a manner rather similar to purely quantum models of measurement.

It is convenient to assume that the measurement takes place over a sufficiently short time period, $[0, T]$, such that \tilde{H}_0 can be ignored during the measurement. The equations of motion during the interaction then follow via Eqs. (1) and (7) as being equivalent to the hybrid Schrödinger equation

$$i\hbar \frac{\partial \psi}{\partial t} = \kappa(t) \left(\frac{\hbar}{i} \frac{\partial}{\partial x} \right) M \psi.$$

For an initially independent ensemble at time $t = t_0 = 0$, as per Eq. (11), this equation may be trivially integrated to give

$$\psi(q, x, T) = \sum_n c_n \psi_C(x - K\lambda_n) \langle q|n \rangle \quad (12)$$

at the end of the measurement interaction, where $K = \int_0^T dt \kappa(t)$, $|n\rangle$ denotes the eigenstate corresponding to

eigenvalue λ_n of M , and $c_n = \langle n|\psi_Q\rangle$ (with $|\psi_Q\rangle$ defined via $\langle q|\psi_Q\rangle := \psi_Q(q)$). It has been assumed for simplicity here that M is nondegenerate (the degenerate case is considered further below).

The pointer probability distribution after measurement follows immediately from Eq. (12) as

$$P(x, T) = \int dq \psi^* \psi = \sum_n |c_n|^2 P_C(x - K\lambda_n). \quad (13)$$

Hence, the initial pointer distribution is displaced by an amount $K\lambda_n$ with probability $|c_n|^2$, thus correlating the position of the pointer with the eigenvalues of M . In particular, choosing a sufficiently narrow initial distribution $P_C(x)$ (eg, a delta-function), the displaced distributions will be nonoverlapping (corresponding to a ‘good’ measurement), and eigenvalue λ_n will be perfectly correlated with the measured pointer position.

The above shows that the measurement of any quantum observable may be modelled via interaction with a strictly classical measuring apparatus. Note that ‘collapse’ of the quantum component of the ensemble can also be modelled, *if* desired. Suppose in particular that the ‘real’ position of the pointer is determined to be $x = a$. This must correspond to just one of the nonoverlapping distributions $P_C(x - K\lambda_n)$, and updating $P(q, x, T)$ via Bayes theorem implies, via Eqs. (12) and (13), that

$$P_a(q, x, T) = \delta(x-a) P(q, a, T) / P(a, T) = \delta(x-a) |\langle q|n\rangle|^2.$$

Moreover, it is natural to update the conjugate quantity $S(q, x, t)$ via the minimal substitution

$$S_a(q, x, t) = S(q, a, T).$$

Note that the ‘collapsed’ ensemble after measurement thus has the form of Eq. (11) (with quantum component described by $\psi_a(q) = \langle q|n\rangle$ up to a phase factor), implying that collapse is compatible with the independence constraint (10).

Finally, for a degenerate operator M with eigenvalue decomposition $\sum_n \lambda_n E_n$, similar results are obtained, but with $c_n \langle q|n\rangle$ in Eq. (12) replaced by $\langle q|E_n|\psi_Q\rangle$, $|c_n|^2$ by $p_n = \langle \psi_Q|E_n|\psi_Q\rangle$, and the ‘collapsed’ quantum component by $\psi_a(q) = (p_n)^{-1/2} \langle q|E_n|\psi_Q\rangle$.

IV. MIXTURES AND THERMODYNAMICS

It has been demonstrated above that the configuration-ensemble approach provides a consistent formulation of mixed quantum and classical dynamics. However, given that this approach describes classical ensembles via a configuration space, rather than a phase space, this raises a potential completeness issue: do all classical phase space ensembles have a counterpart in this approach? Further, given the lack of a natural phase space entropy, can the configuration-ensemble approach deal with thermal ensembles in a manner that is compatible with both

classical and quantum thermodynamics? It is shown briefly below that both these questions have positive answers.

The central concept required is that of a *mixture* of configuration ensembles. In particular, if a physical system is described by the configuration ensemble (P_j, S_j) with prior probability p_j , then it may be said to correspond to the mixture $\{(P_j, S_j); p_j\}$. For quantum ensembles, such mixtures are conveniently represented by density operators. More generally, however, there is no similarly convenient representation. The average of any observable $A[P, S]$ over a mixture is given by

$$\langle A \rangle = \sum_j p_j A[P_j, S_j]. \quad (14)$$

The first question posed above can now easily be answered, using the fact that any classical phase space point, $\gamma = (x', k')$, may be described by a classical configuration ensemble (P_γ, S_γ) , defined by

$$P_\gamma(x) := \delta(x - x'), \quad S_\gamma(x) := k' \cdot x.$$

In particular, the value of any classical observable C_f , corresponding to the average value of f , follows via Eq. (4) as

$$C_f[P_\gamma, S_\gamma] = \langle f \rangle_\gamma = f(x', k'). \quad (15)$$

It follows immediately that *any* classical phase space ensemble, represented by some phase space density $p(x', k')$, may equivalently be described by the mixture $\{(P_\gamma, S_\gamma); p\}$ of classical configuration ensembles.

To address the second question above, one further requires the notions of ‘stationary’ and ‘distinguishable’ configuration ensembles. First, stationary ensembles are those for which all observable quantities are time-independent, and are characterised by the property [13]

$$\partial P / \partial t = 0, \quad \partial S / \partial t = -E, \quad (16)$$

for some constant E . Second, two configuration ensembles are defined to be distinguishable if there is some observable which can distinguish unambiguously between them, i.e., the ranges of the observable for each ensemble do not overlap. Thus, for example, two quantum ensembles are distinguishable if the corresponding wavefunctions are orthogonal, while two classical ensembles are distinguishable if the ranges of (x, k) over the supports of the ensembles are nonoverlapping (with $k = \nabla_x S$).

A *thermal mixture* may now be defined as a mixture of distinguishable stationary ensembles $\{(P, S); p(P, S|\tilde{H})\}$ such that

$$p(P, S|\tilde{H}) \sim e^{-\beta \tilde{H}[P, S]}, \quad \beta > 0. \quad (17)$$

This definition is, for present purposes, justified by its consequences, but it may also be motivated by appealing to properties of two distinct noninteracting systems in

thermal equilibrium, described by joint ensemble Hamiltonian \tilde{H}_T , for which one expects

$$p(P, S|\tilde{H}_T) = p(P, S|\tilde{H}) p(P', S'|\tilde{H}'),$$

for pairs (P, S) , (P', S') of stationary ensembles.

For a quantum system with Hamiltonian operator H , the above definition immediately leads to the usual quantum canonical ensemble represented by the density operator proportional to $e^{-\beta H}$. It will now be shown that for an ergodic classical system with phase space Hamiltonian $H(x, k)$, the corresponding thermal mixture of configuration ensembles is equivalent to the classical canonical ensemble with phase space density proportional to $e^{-\beta H}$.

In particular, the ensemble Hamiltonian is given by

$$\tilde{H}[P, S] = C_H = \int dx P H[x, \nabla S],$$

and it follows from Eqs. (1) and (16) that the stationary ensembles are then given by the solutions of the continuity and Hamilton-Jacobi equations

$$\nabla_x \cdot [P \nabla_k H(x, \nabla_x S)] = 0, \quad H(x, \nabla_x S) = E.$$

Now, since H is time-independent, the general solution to the Hamilton-Jacobi equation for S is of the form $W(x) - Et$, and generates a canonical transformation on phase space from (x, k) to a set of constants of the motion, which may be chosen as the initial values (x_0, k_0) at some fixed time [16]. One then has $E = H(x_t, k_t) = H(x_0, k_0)$, and a corresponding set of solutions $S_{x_0, k_0}(x, t) := W_{k_0}(x) - H(x_0, k_0)t$ of the Hamilton-Jacobi equation (where $x_0 = \nabla_{k_0} W_{k_0}$ [16]). Further, recalling that the classical velocity is $\dot{x}_t = \nabla_k H$, the above continuity equation simply requires that $P(x)|\dot{x}_t|$ is constant along any given trajectory (x_t, k_t) . One may therefore define a corresponding set of solutions by $P_{x_0, k_0}(x) \sim |\dot{x}_t|^{-1}$ when x lies on the particular trajectory having initial values (x_0, k_0) , and vanishing elsewhere.

The stationary ensembles $(P_{x_0, k_0}, S_{x_0, k_0})$ are all distinguishable, since they correspond to a set of nonoverlapping trajectories defined by their initial values (x_0, k_0) . Hence, they are suitable for defining a thermal mixture. Further, by construction, one has via definition (4) that

$$C_f[P_{x_0, k_0}, S_{x_0, k_0}] = \int dx_t |\dot{x}_t|^{-1} f(x_t, k_t) / \int dx_t |\dot{x}_t|^{-1},$$

where integration is along the trajectory defined by initial point (x_0, k_0) . Changing the variable of integration to t then gives

$$C_f[P_{x_0, k_0}, S_{x_0, k_0}] = \lim_{T \rightarrow \infty} \frac{1}{T} \int_0^T dt f(x_t, p_t), \quad (18)$$

which is always well defined for ergodic systems. Further, for such systems the righthand side is simply the microcanonical ensemble average of $f(x, k)$, corresponding

to constant energy $E = H(x_0, k_0)$ [19]. Finally, noting $\tilde{H}[P_{x_0, k_0}, S_{x_0, k_0}] = H[x_0, k_0]$ by construction, the associated thermal mixture in Eq. (17) is characterised by $p(x_0, k_0) \sim e^{-\beta H[x_0, k_0]}$, and it immediately follows via Eqs. (14) and (18) that the average of f over the mixture is the canonical ensemble average, as desired.

V. DISCUSSION

The configuration-ensemble approach satisfies the two minimal requirements for a consistent formulation of mixed dynamics, as shown in Sec. II. No other formulation appears to be known which passes this critical test.

Further, an additional ‘independence’ requirement may also be satisfied by imposing a weak constraint on the class of physical ensembles, analogous to the use of gauge constraints in quantum field theory, as shown in Sec. III. This constraint is satisfied, for example, by any mixed ensemble that is related to some pair of independent quantum and classical ensembles by a canonical transformation.

More generally, it should be noted that the configuration-ensemble approach has a number of interesting applications outside the domain of mixed dynamics, providing a basis for, eg, the derivation of classical and quantum equations of motion [20], a generalisation of quantum superselection rules [15], and, as seen in Sec. IV above, a ‘Hamilton-Jacobi’ approach to classical statistical mechanics. Hence, overall, the approach appears to be valuable in providing a fundamental tool for describing physical systems, and merits further investigation.

It is of interest to remark on how the configuration-ensemble approach is able to avoid various ‘no-go’ theorems on mixed dynamics in the literature [5, 6, 7, 9]. This is essentially due to such theorems requiring a formal assumption that the set of observables can be extended to form a product algebra, where the product $A * B$ is assumed to satisfy $C_f * C_g = C_{fg}$, $Q_M * Q_N = Q_{MN}$, and some further property such as the Leibniz rule $\{A, B * C\} = \{A, B\} * C + B * \{A, C\}$. However, the assumption of such a product algebra clearly goes beyond the domain of observable quantities (eg, the product of two Hermitian operators is not a Hermitian operator), and hence cannot be justified on physical grounds. Thus, any import of such ‘no-go’ theorems, for the configuration-ensemble approach, where no such product is defined or required, is purely formal in nature.

Note that for application to mixed dynamics, the set of observables must be chosen such that it contains the classical and quantum observables defined in Eqs. (4) and (6), with all members satisfying the normalisation and positivity conditions in Eq. (3). It must also, of course, contain the Poisson bracket of any two of its members - however, this can always be assured by replacing a given set by its closure under the Poisson bracket operation. It is of interest to consider what further physical conditions might be imposed on the set of observables. For

example, for ensembles of interacting classical and quantum nonrelativistic particles, it is reasonable to require this set contains ensemble Hamiltonians of the form defined in Eq. (9), with the potential function V restricted to be of the form $V(q - x, t)$. The equations of motion will then be invariant under Galilean transformations, although with the interesting property that the centre of mass and relative motions do not decouple [13].

A more general condition that might be imposed on observables is that they are homogenous of degree unity with respect to the probability density P , i.e.,

$$A[\lambda P, S] = \lambda A[P, S] \quad (19)$$

for all $\lambda \geq 0$. Note that if this condition holds for two observables A and B , then it holds for the Poisson bracket $\{A, B\}$, as may be checked by direct substitution into Eq. (2). It is also easily verified to hold for the classical and quantum observables defined in Eqs. (4) and (6). Differentiating Eq. (19) on both sides with respect to λ and choosing $\lambda = 1$ yields the numerical identity

$$A[P, S] = \int d\xi P (\delta A / \delta P) =: \langle \delta A / \delta P \rangle, \quad (20)$$

i.e., A can be calculated by integrating over a local density on the configuration space. Thus, the homogeneity condition consistently allows observables to be interpreted both as generators of canonical transformations and as expectation values.

Finally, while the question of decoherence has not been addressed in any detail here, it is worth noting that the configuration-ensemble approach provides several possibilities in this regard. First, for any mixed quantum-classical ensemble, one may define a conditional quantum wavefunction $\psi_x(q)$ which describes the conditional decoherence of the quantum component relative to the classical component [13]. Second, as noted briefly in Sec. III A, if the weak independence constraint (10) is not imposed, then the very existence of a configuration-ensemble not satisfying this constraint may lead to ‘interaction without interaction’ and thus to intrinsic decoherence of the quantum component. Third, while at any time the hybrid wavefunction $\psi(q, x, t)$ describing a mixed quantum-classical system always has a decomposition of the form

$$\psi(q, x, t) = \sum_n \sqrt{p_n(t)} \psi_{C,n}(x, t) \psi_{Q,n}(q, t)$$

(eg, a Schmidt decomposition), only at *particular* times (if at all, depending on the ensemble Hamiltonian), can it have such a decomposition for which (i) the classical ensembles corresponding to $\psi_{C,n}(x, t)$ are *classically* distinguishable (see Sec. IV), and (ii) the quantum ensembles corresponding to $\psi_{Q,n}(q, t)$ are mutually orthogonal. Moreover, unlike a Schmidt decomposition, such a decomposition would be unique even for equal $p_n(t)$. Hence, decoherence could be modelled by imposing a spontaneous ‘collapse’ of the ensemble at such well-defined times, similarly to the collapse model in Sec. III B.

Acknowledgments

I am grateful to Marcel Reginatto for helpful discussions, and for pointing out an error in an early draft.

-
- [1] For example, E.C.G. Sudarshan, *Pramana* **6**, 117 (1976); I.V. Aleksandrov, *Z. Naturforsch. Teil A* **36** 902 (1981); A. Anderson, *Phys. Rev. Lett.* **74** 621 (1995); L. Diosi and J.J. Halliwell, *Phys. Rev. Lett.* **81**, 2846 (1998); L. Diosi, N. Gisin, and W.T. Strunz, *Phys. Rev. A* **61**, 022108 (2000); E. Gindensperger, C. Meier, and J.A. Beswick, *J. Chem. Phys.* **113** 9369 (2000); O.V. Prezdho and C. Brooksby, *Phys. Rev. Lett.* **86**, 3215 (2001); I. Burghardt and G. Parlant, *J. Chem. Phys.* **120**, 3055 (2004); V.V. Kisil, *Europhys. Lett.* **72**, 873 (2005); O.V. Prezdho, *J. Chem. Phys.* **124** 201104 (2006).
- [2] L. Diosi, N. Gisin, and W.T. Strunz, *Phys. Rev. A* **61**, 022108 (2000).
- [3] T.N. Sherry and E.C.G. Sudarshan, *Phys. Rev. D* **20**, 857 (1979).
- [4] W. Boucher and J. Traschen, *Phys. Rev. D* **37**, 3522 (1988);
- [5] L.L. Salcedo, *Phys. Rev. A* **54**, 3657 (1996).
- [6] J. Caro and L.L. Salcedo, *Phys. Rev. A* **60** 842 (1999).
- [7] A. Peres and D.R. Terno, *Phys. Rev. A* **63**, 022101 (2001).
- [8] L.L. Salcedo, *Phys. Rev. Lett.* **90**, 118901 (2003).
- [9] D. Sahoo, *J. Phys. A* **37**, 997 (2004).
- [10] L.L. Salcedo, *J. Chem. Phys.* **126** 057101 (2007).
- [11] F. Agostini, S. Caprara and G. Ciccotti, *Europhys. Lett.* **78**, 30001 (2007).
- [12] See for example, N. Makri, *Annu. Rev. Phys. Chem.* **50**, 167 (1999); O. Prezdho and C. Brooksby, *Phys. Rev. Lett.* **90**, 118902 (2003); O.V. Prezdho, *J. Chem. Phys.* **126** 057102 (2007); and references therein.
- [13] M.J.W. Hall and M. Reginatto, *Phys. Rev. A* **72** 062109 (2005).
- [14] M. Albers, C. Kiefer and M. Reginatto, eprint arXiv:0802.1978v1 [gr-qc].
- [15] M.J.W. Hall, *J. Phys. A* **37** 7799 (2004).
- [16] H. Goldstein, *Classical Mechanics* (Addison-Wesley, New York, 1950), Chaps. 8, 9 and 11.
- [17] S. Weinberg, *Ann. Phys. N.Y.* **194** 336 (1989).
- [18] M. Henneaux and C. Teitelbohm, *Quantization of Gauge Systems* (Princeton University Press, New Jersey, 1992), Chaps. 1 and 2.
- [19] G. Morandi, F. Napoli and E. Ercolessi, *Statistical Mechanics*, 2nd edition (World Scientific, Singapore, 2001), Chap. 2.
- [20] M.J.W. Hall and M. Reginatto, *J. Phys. A* **35**, 3289 (2002); M.J.W. Hall, K. Kumar and M. Reginatto, *J. Phys. A* **36**, 9779 (2003); M.J.W. Hall, *Gen. Rel. Gravitat.* **37** 1505 (2005).