Partial teleportation of entangled atomic states

W. B. Cardoso^{1, *} and N. G. de Almeida²

¹Instituto de Física, Universidade Federal de Goiás, 74.001-970, Goiânia (GO), Brazil. ²Núcleo de Pesquisas em Física, Universidade Católica de Goiás, 74.605-220, Goiânia (GO), Brazil.

In this paper we propose a scheme for partially teleporting entangled atomic states. By partial teleportation we mean that teleportation will occur by changing one of the partner of the entangled state to be teleported. Our scheme can be implemented using only four two-level atoms interacting either resonantly or off-resonantly with a single cavity-QED. The estimative of losses occurring during this partial teleportation process is accomplished through the phenomenological operator approach technique.

PACS numbers: 03.67.Hk; 42.50.Dv

I. INTRODUCTION

Quantum teleportation, first suggested by Bennett et al. [1], is one of the cornerstones of quantum information and computation [2, 3, 4]. The crucial ingredient characterizing this phenomenon is the transfer of information between noninteracting systems at the expense of a quantum channel. This issue has received great attention since its pioneer proposal, mainly after its experimental realizations from 1997 onwards [5, 6, 7]. In the meantime, various proposals have been suggested for implementing teleportation, for instance, teleportation of trapped wave fields inside high-Q microwave cavities [8], teleportation of running wave fields [9, 10, 11], teleportation by particle-hole annihilation in the Fermi sea [14], teleportation in noninertial frames [15], teleportation of electronic many-qubit states encoded in the electron spin of quantum dots [16], teleportation of single-mode thermal state of light field [17], teleportation of a macroscopic atomic ensemble [18], teleportation between nanomechanical modes in an all solid-state circuit [19], teleportation of trapped field states inside a single bimodal cavity [20], nonprobabilistic teleportation of a field state via cavity QED [21], and teleportation of the angular spectrum of a single-photon field [22], among others.

Since the pioneering work by Bennet et al.[1], several schemes for teleportation differing from this original protocol (OP) have appeared in the literature. Actually, in the experiment of Ref.[6], Boschi et al. explored both the polarization and the state of the photon through two distinct paths to demonstrate teleportation. Braunstein in Ref. [23] and Nielsen and Caves in Ref. [24] showed that teleportation is possible in a reversible manner, and a protocol for its implementation in the optical lattice domain was presented in Ref. [25]. In Ref. [26], de Almeida et al. used Greenberger-Horne-Zeilinger (GHZ) states as the nonlocal channel, instead of the standard Einstein-Podolsky-Rosen (EPR) states. In Ref.[27], Karlsson and Bourennane, also using a GHZ channel, showed how to accomplish teleportation controlled by a third party. Controlled teleportation involving many agents were also considered in Ref. [28]. In Ref. [29], Popescu substituted the nonlocal channel by the mixed Werner states. Entangled mixed states as nonlocal channel was further considered in Refs.[30], and in Ref. [31], Christian Trump et al. substituted the Bell base in the teleportation process by a partially entangled base. In Ref. [32], Vaidman considered both a spin state and a system with continuous variable, and presented a "cross measurement" method to achieve a two-way teleportation. Later on, Moussa [33] showed how to implement this "cross measurement" method in the context of QED cavity, named by him as teleportation with *identity interchange* of quantum states. In Ref.[34], Zheng refers to the approximated teleportation without Bell states measurements of the superposition of zero- and one-photon states from one high -Q cavity to another, with fidelity near 99% [35]. In Ref. [11], H. W. Lee and J. Kim showed how to partially teleport an entanglement of zero and one photon state in the running wave domain. Also, the term partial teleportation is used in literature to deal with teleportation of an unknown state with the generation of its clone [12, 13]. Here we use the term "partial Teleportation" in the same sense as that in [11].

In this paper we present a scheme for partial teleportation in the cavity QED domain. Our scheme uses four two-level atoms interacting either on or off resonantly with a single mode of a high-Q cavity, Ramsey zones, and selective atomic state detectors. To simplify the estimative of losses occurring during the partial teleportation process, we used the phenomenological operator approach technique (POA) [36].

II. IDEAL TELEPORTATION PROCESS

We assume atom 1 previously entangled with atom 2 in the following state, which is the state we want to teleport

$$|\phi\rangle_{12} = C_0 |g\rangle_1 |e\rangle_2 + C_1 |e\rangle_1 |g\rangle_2,$$
 (1)

where C_0 and C_1 are unknown coefficients obeying $|C_0|^2 + |C_1|^2 = 1$, and $|g\rangle$ ($|e\rangle$) is the atomic ground (excited) state. This state can be prepared, for instance, by the method presented in Ref. [37], where two two-level atoms interacts simultaneously with a single mode of the cavity-field.

The Hamiltonian describing the atom-field interaction, in the interaction picture, is

$$H_I = \hbar \lambda \left(a^{\dagger} \sigma^- + a \sigma^+ \right), \tag{2}$$

^{*}Electronic address: wesleybcardoso@gmail.com



FIG. 1: Scheme for accomplishment of partial teleportation in cavity QED.

when the atom-field interaction is resonant, and

$$H_I = \frac{\hbar \lambda^2}{\delta} a^{\dagger} a \sigma^+ \sigma^-, \qquad (3)$$

when the atom-field interaction is off-resonant. This condition is valid provided that $\overline{n}\lambda^2 \ll \delta^2 + \gamma^2$, where \overline{n} is the mean photon number and γ is the damping rate for the cavity-field. Here a^{\dagger} and a are the creation and annihilation operators for the cavity field mode, σ^+ and σ^- are the raising and lowering operators for the atom, λ is the atom-field coupling constant, and $\delta = \omega - \omega_0$ is the detuning between the cavity field frequency ω and the atomic frequency ω_0 .

To compose the nonlocal channel, a third atom, initially prepared in the excited state $|e\rangle_3$, interacts resonantly with the cavity field mode A. Adjusting the atom-field interaction time to $t = \pi/4\lambda$, the nonlocal channel will be given by

$$|\psi\rangle = \frac{1}{\sqrt{2}} \left(|e\rangle_3 \left|0\rangle_A - i\left|g\rangle_3 \left|1\rangle_A\right)\right.$$
(4)

The state of the whole system composed by the two-level atoms and the cavity mode field is

$$\begin{split} |\psi\rangle_{total} &= \frac{1}{2} \left[\left| \Psi^{+} \right\rangle_{1A} \left(C_{0} \left| g \right\rangle_{3} \left| e \right\rangle_{2} + C_{1} \left| e \right\rangle_{3} \left| g \right\rangle_{2} \right) \\ &+ \left| \Psi^{-} \right\rangle_{1A} \left(C_{0} \left| g \right\rangle_{3} \left| e \right\rangle_{2} - C_{1} \left| e \right\rangle_{3} \left| g \right\rangle_{2} \right) \\ &+ \left| \Phi^{+} \right\rangle_{1A} \left(C_{0} \left| e \right\rangle_{3} \left| e \right\rangle_{2} - C_{1} \left| g \right\rangle_{3} \left| g \right\rangle_{2} \right) \\ &+ \left| \Phi^{-} \right\rangle_{1A} \left(C_{0} \left| e \right\rangle_{3} \left| e \right\rangle_{2} + C_{1} \left| g \right\rangle_{3} \left| g \right\rangle_{2} \right) \right], \quad (5) \end{split}$$

where, for convenience, we have defined the Bell states $|\Psi^{\pm}\rangle_{1A}$ and $|\Phi^{\pm}\rangle_{1A}$ as

$$\left|\Psi^{\pm}\right\rangle_{1A} = \frac{1}{\sqrt{2}} \left(-i\left|g\right\rangle_{1}\left|1\right\rangle_{A} \pm \left|e\right\rangle_{1}\left|0\right\rangle_{A}\right),\tag{6}$$

$$\left|\Phi^{\pm}\right\rangle_{1A} = \frac{1}{\sqrt{2}} \left(\left|g\right\rangle_{1}\left|0\right\rangle_{A} \pm i\left|e\right\rangle_{1}\left|1\right\rangle_{A}\right). \tag{7}$$

As in the OP, the teleportation is completed after Alice measuring on its particles (cavity mode field A and atom 1) and sending her result to Bob, whom will know which unitary operation to accomplish on its particles to recover the entangled state that Alice wanted to teleport. Note, however, that different from the OP, when comparing the teleported state, resulting from Eq.(5), with the state to be teleported, Eq.(1), we see that the partner 1 was substituted by the particle 3. This explain the "partial teleportation" term used. Next, we show how Alice must proceed in order to perform the Bell state measurements.

A. Bell State Measurements

First, the atom 1 crosses a Ramsey zone R, adjusted to produce the following evolutions $|e\rangle_1 \rightarrow \frac{1}{\sqrt{2}} (|g\rangle_1 + |e\rangle_1)$ and $|g\rangle_1 \rightarrow \frac{1}{\sqrt{2}} (|g\rangle_1 - |e\rangle_1)$. Then, atom 1 crosses the cavity interacting off-resonantly with mode A, with the interaction time adjusted to $\chi t = \pi$, resulting in the following evolutions $|g\rangle_1 |0\rangle_A \rightarrow |g\rangle_1 |0\rangle_A$, $|g\rangle_1 |1\rangle_A \rightarrow |g\rangle_1 |1\rangle_A$, $|e\rangle_1 |0\rangle_A \rightarrow |e\rangle_1 |0\rangle_A$, $|e\rangle_1 |1\rangle_A \rightarrow -|e\rangle_1 |1\rangle_A$. Then, the atom 1 crosses another Ramsey zone R' adjusted like the first R. As a consequence, the states of the Bell bases evolve as

$$\left|\Psi^{\pm}\right\rangle_{1A} \to \frac{1}{\sqrt{2}} \left|g\right\rangle_{1} \left(-i\left|1\right\rangle_{A} \pm \left|0\right\rangle_{A}\right),\tag{8}$$

$$\left|\Phi^{\pm}\right\rangle_{1A} \to \frac{1}{\sqrt{2}} \left|e\right\rangle_1 \left(\left|0\right\rangle_A \pm i \left|1\right\rangle_A\right). \tag{9}$$

Thus, by selectively detecting the state of the atom 1 it is possible to decide if the joint state was $|\Psi^{\pm}\rangle_{1A}$ or $|\Phi^{\pm}\rangle_{1A}$. Next, we have to discern the phase (\pm) . This is done by sending another two-level atom 4 in the ground state $|g\rangle_4$ resonantly with mode A. The Atom $|g\rangle_4$, after interacting resonantly with mode A as indicated by Eq.(2), crosses R resulting

$$\left|\Psi^{\pm}\right\rangle_{1A}\left|g\right\rangle_{4} \rightarrow \begin{cases} \left|g\right\rangle_{1}\left|e\right\rangle_{4}\left|0\right\rangle_{A} & \text{if } (+) \\ \left|g\right\rangle_{1}\left|g\right\rangle_{4}\left|0\right\rangle_{A} & \text{if } (-) \end{cases}, \quad (10)$$

$$\left|\Phi^{\pm}\right\rangle_{1A}\left|g\right\rangle_{4} \to \begin{cases} \left|e\right\rangle_{1}\left|g\right\rangle_{4}\left|0\right\rangle_{A} & \text{if (+)} \\ \left|e\right\rangle_{1}\left|e\right\rangle_{4}\left|0\right\rangle_{A} & \text{if (-)} \end{cases}$$
(11)

Thus, whenever the atoms 1 and 4 are measured both in its ground or excited states the phase is of (-), otherwise, the phase is (+). The perfect discrimination between the four states composing the Bell base can be accomplished by Alice through the detection of the atoms 1 and 4, separately. Therefore, Alice sends a sign to Bob, whom accomplishes an appropriate rotation in the states of the atoms 2 and 3 to complete the partial teleportation with 100% of fidelity and success probability, in the ideal case. The unitary operations required by Bob are summarized in Table 1.

| BSM | $ \psi angle_{32}$ | Unitary operation |
|------------------------------------|---|------------------------------------|
| $\left \Psi^{+}\right\rangle_{1A}$ | $C_0 g\rangle_3 e\rangle_2+C_1 e\rangle_3 g\rangle_2$ | $\mathbb{I}_3\otimes\mathbb{I}_2$ |
| $\left \Psi^{-}\right\rangle_{1A}$ | $C_0 g\rangle_3 e\rangle_2-C_1 e\rangle_3 g\rangle_2$ | $\sigma_{3z}\otimes\mathbb{I}_2$ |
| $\left \Phi^{+}\right\rangle_{1A}$ | $C_0 e\rangle_3 e\rangle_2 - C_1 g\rangle_3 g\rangle_2$ | $\sigma_{3y}\otimes\mathbb{I}_2$ |
| $\left \Phi^{-}\right\rangle_{1A}$ | $C_0 e angle_3 e angle_2+C_1 g angle_3 g angle_2$ | $\sigma_{3x} \otimes \mathbb{I}_2$ |

TABLE I: The results of the teleportation scheme. BSM denotes the resulting measurement on atom 1 and field mode A. Unitary operation denotes the required operation by Bob after Alice communicating her results. The σ_{3j} is the Pauli operator σ_j acting on atom 3.

III. DECAY OF THE FREE ATOMIC EXCITED STATE

A. Phenomenological operator approach (POA)

Here we observe that the coupling of the atomic states to a surrounding environment \mathcal{E} can be described by the relations [36]

$$g\rangle|\mathcal{E}\rangle \xrightarrow{U_t} |g\rangle\hat{\mathcal{T}}_0|\mathcal{E}\rangle,$$
 (12)

$$|e\rangle|\mathcal{E}\rangle \xrightarrow{U_t} |e\rangle\hat{\mathcal{T}}_e^{\dagger}|\mathcal{E}\rangle + |g\rangle\hat{\mathcal{T}}_g^{\dagger}|\mathcal{E}\rangle,$$
 (13)

where $|\mathcal{E}\rangle$ denotes the initial state of the environment, the operators $\hat{\mathcal{T}}$, acting on this state, account for the atomenvironment coupling, and U_t denotes an unitary operation mixing the atom to its environment. We will assume the environment $|\mathcal{E}\rangle$ in the vacuum state, which is an excellent approximation for high-Q cavities in the microwave domain [38]. Accordingly, we assume that $\hat{\mathcal{T}}_0^{\dagger} = \mathbf{1}$, $\hat{\mathcal{T}}_e^{\dagger} = \mathbf{f}(t) = e^{-\kappa t}\mathbf{1}$, $\hat{\mathcal{T}}_g^{\dagger} = \sum_j \mathbf{g}_j(t)\hat{b}_j^{\dagger}$, with $\sum_j |\mathbf{g}_j(t)|^2 = 1 - e^{-2\kappa t}$, κ denoting the spontaneous decay rate of the atomic excited state, $\mathbf{1}$ is the identity operator, b_j^{\dagger} is the creation operator, having a corresponding annihilation operator b_j , of the *j*th oscillator mode of the environment, and *t* is the time elapsing after the atom suffering a given excitation. With these assumptions, it is straightforward to verify that the superposition $(|g\rangle + |e\rangle)/\sqrt{2}$ leads to the reduced density operator

$$\rho = \frac{1}{2} \left\{ \exp(-2\kappa t) |e\rangle \langle e| + [2 - \exp(-2\kappa t)] |g\rangle \langle g| \right. \\ \left. + \left. \exp(-\kappa t) \left(|e\rangle \langle g| + |g\rangle \langle e| \right) \right\}.$$
(14)

Note that the evolution (12) and (13) are consistent with the well-known result, following from the Weisskopf-Wigner approximation, that an unstable atomic state decays exponentially. In this case, the phenomenological-operator evolution leads to the same atomic density operator as the one we obtain using an *ab-initio* master equation approach. Moreover, due to recent advances in high-Q cavities [40], we will neglect the damping time of the mode A.

B. Decay of the teleported state

To simplify our estimative of the losses, we assume the whole state starting to decay after the preparation of the quantum channel. In the first step, the phenomenological operators used to introduce damping effects, Eqs.(12) and (13), are applied to the whole system until the time t. Then, for each excitement suffered by the atoms during the teleportation process, a new phenomenological operator is included, which modify the decay probability of the atomic states, and as a consequence, the fidelity of the whole teleportation process. Summarizing the applications of the phenomenological operators from the beginning, *i.e.*, since the preparation of the quantum channel, until the end of our teleportation protocol, which occurs at the instant that the fourth atom is detected, we

have to apply them soon after i) the atom 1 crossing the first Ramsey zone (t_1) ; *ii*) the atom 1 crossing the second Ramsey zone (t_2) ; *iii*) the atom 4 interacting resonantly with the mode field cavity (t_3) ; iv) the atom 4 crossing the Ramsey zone (t_4) . After the inclusion of the decay via POA, the state of the whole system by the time the teleportation is concluded becomes a mixture, being represented by a reduced operator density when the reservoir is traced out. In our estimative, we take the case of the teleported state in Bob hands when Alice measures the Bell state $|\Psi^+\rangle_{1A}$ $(|g\rangle_1 |e\rangle_4)$. The corresponding fidelity is shown in Fig. 1. Note that at the time the teleportation is completed (t_4) the fidelity rounds 1, indicating that we can safely neglect losses occurring during the teleportation process. In fact, taking $t_1 = 2\mu s$, as reported in [41], we will have $t_2 \simeq 5 \times 10^{-4} s + 2\mu s + t_1$, which is the necessary time for the atom 1 to interact dispersively with the cavity field and to cross the Ramsey zone, $t_3 = 10^{-4}s + t_2$, which is the necessary time for the atom-field resonant interaction, and $t_4 = 2\mu s + t_3 \simeq 6,06 \times 10^{-4} s$, which is much shorter than the atomic decay $\kappa^{-1} \cong 10^{-2}s$, being the fidelity at this time 0.99 as can be seen from Fig.1. However, as the time goes on, the decay becomes faster and the fidelity is reduced to 2/3, corresponding to the no-cloning limit [42], at the instant $t_f = 5,78 \times 10^{-3}s$. Therefore, the effective time during which the teleported state is at our disposal for further operations is $t_f - t_4 = 5, 17 \times 10^{-3} s$.



FIG. 2: Decay effects of the teleported state. In (a) we see the behavior of the fidelity and its dependence with both the life-time of the atomic state and the value of the coefficients of the state to be teleported. In (b) we see the behavior of the fidelity for the fixed values of the coefficients $C_0 = C_1 = 1/\sqrt{2}$.

IV. COMMENTS AND CONCLUSION

Since the teleportation protocol by Bennett *et al.*[1], several other proposals have appeared, modifying slightly or substantially the original protocol. In this paper we have explored a kind of teleportation named by *partial teleportation* [11]. In our scheme, Alice has an atomic state to be teleported, given by an entanglement of the particles 1 and 2, and shares with Bob a nonlocal channel, composed by the joint state of the particles A (a single mode of a high Q cavity) and 3 (another atom). Performing a Bell measurement on the states of particle A and 1, and informing Bob her result, the following interesting result emerges, after the usual rotation performed

- C. H. Bennett, G. Brassard, C. Crepeau, R. Jozsa, A. Peres, W. K. Wootters, Phys. Rev. Lett. 70 (1993) 1895.
- [2] G. Brassard, S. L. Braunstein, R. Cleve, Physica D 120 (1998) 43.
- [3] D. Gottesman, I. L. Chuang, Nature 402 (1999) 390; Knill E, R. Laamme, G. J. Milburn, Nature 409 (2001) 46.
- [4] M. A. Nielsen, I. L. Chuang, *Quantum Computation and Quantum Information* (Cambridge University Press, Cambridge, England, 2000).
- [5] D. Bouwmeester et al., Nature 390 (1997) 575.
- [6] D. Boschi et al., Phys. Rev. Lett. 80 (1998) 1121.
- [7] A. Furusawa et al., Science 282 (1998) 706; E. Lombardi et al., Phys. Rev. Lett. 88 (2002) 070402; H. de Riedmatten et al., Phys. Rev. Lett. 92 (2004) 047904; Y. F. Huang et al., Phys. Rev. Lett. 93 (2004) 240501; M. D. Barrett et al., Nature 429 (2004) 737; M. Riebe et al., Nature 429 (2004) 734; D. Fattal et al., Phys. Rev. Lett. 92 (2004) 037904; N. Takei et al., Phys. Rev. Lett. 94 (2005) 220502; G. Y. Xiang, J. Li, G. C. Guo, Phys. Rev. A 71 (2005) 044304; Z. Zao et al., Phys. Rev. Lett. 95 (2006) 030502; J. F. Sherson et al., Nature 443 (2006) 557; J. A. W. van Houwellingen et al., Phys. Rev. A 74 (2006) 022303; H. Yonezawa, S. L. Braunstein, A. Furusawa, Phys. Rev. Lett. 99 (2007) 110503.
- [8] L. Davidovich et al., Phys. Rev. A 50 (1994) R895; J. I. Cirac, A. S. Parkins, Phys. Rev. A 50 (1994) R4441; T. Sleator, H. Weinfurter, Ann. N. Y. Acad. Sci. 755 (1995) 715; M. H. Y. Moussa, Phys. Rev. A 54 (1996) 4661; M. Ikram, S. Y. Zhu, M. S. Zubairy, Phys. Rev. A 62 (2000) 022307; N.G. de Almeida, R. Napolitano, M.H.Y. Moussa, Phys. Rev A. 62 (2000) 010101(R); G. Pires et al., Phys. Rev. A 70 (2004) 025803; P. P. Munhoz et al., J. Phys. B: At. Mol. Opt. Phys. 38 (2005) 3875; Z. B. Yang, J. Phys. B: At. Mol. Opt. Phys. 39 (2006) 603; W. B. Cardoso et al., J. Phys. B: At. Mol. Opt. Phys. 40 (2007) 1089; Z. B. Yang, J. Phys. B: At. Mol. Opt. Phys. 40 (2007) 1093; Z. B. Yang, Chinese Phys. 16 (2007) 0329; W. B. Cardoso, Int. J. Theor. Phys. DOI 10.1007/s10773-007-9524-x; C. J. Shan et al., Int. J. Quant. Inf. 5 (2007) 359; C. R. Carvalho et al., J. Phys. B: At. Mol. Opt. Phys. 40 (2007) 1271; W. B. Cardoso et al., Chinese Physics B 17 (2008) 60.
- [9] S. L. Braunstein, H. J. Kimble, Phys. Rev. Lett. 80 (1998) 869.
- [10] C. J. Villas-Bôas, N. G. de Almeida, M. H. Y. Moussa, Phys. Rev. A 60 (1999) 2759.
- [11] H. W. Lee, J. Kim, Phys. Rev. A 63 (2000) 012305.
- [12] R. Filip, Phys. Rev. A 69 (2004) 052301.
- [13] Z. Zhao et al., Phys. Rev. Lett. 95 (2005) 030502.

by Bob: particle 3 takes exactly the role of particle 1 in the entanglement addressed to Alice, but in Bob location. As the entanglement between the particles 1 and 2 is broken and a new entanglement between the particles 3 and 2 is created in a different place, this characterizes a partial teleportation. To estimate the losses occurring during and after the teleportation process, we have used the method of the phenomenological operator approach (POA) introduced in Ref.[36]. Through the POA, we showed an estimative indicating a high fidelity by the time the teleportation is completed and the precise time that the fidelity achieve the no-cloning limit of 2/3.

We thanks CNPq and CAPES, Brazilian Agency, and PROPE/UCG, for support this work.

- [14] C. W. J. Beenaker, M. Kindermann, Phys. Rev. Lett. 92 (2004) 056801.
- [15] P. M. Alsing, D. McMahon, G. J. Milburn, J. Opt. B: Quantum Semiclass. Opt. 6 (2005) S834.
- [16] M. N. Leuemberger, M. E. Flatté, Phys. Rev. Lett. 94 (2005) 107401.
- [17] J. Zhang, C. Xie, K. Peng, J. Opt. B: Quantum Semiclass. Opt. 7 (2005) 313.
- [18] A. Dantan et al., Phys. Rev. Lett. 94 (2005) 050502.
- [19] L. Tian, S. M. Carr, Phys. Rev. B 74 (2006) 125314.
- [20] G. Pires *et al.*, Phys. Rev. A 71 (2005) 060301; Iara P. de Queirós *et. al.*, Phys. Rev. A 76 (2007) 034101.
- [21] C. R. Carvalho *et al.*, J. Phys. B: At. Mol. Opt. Phys. 40 (2007) 1271.
- [22] S. P. Walborn et al., Phys. Rev. A 76 (2007) 033801.
- [23] S. L. Braunstein, Phys. Rev. A 53 (1996) 1900.
- [24] M. A. Nielsen, C. M. Caves, Phys. Rev. A 55 (1997) 2547.
- [25] L. Santos, D. Bru β , J. Phys. A: Math. Gen. 34 (2001) 7003.
- [26] N. G. de Almeida et al., Phys. Lett A. 241 (1998) 213.
- [27] A. Karlsson, M. Bourennane, Phys. Rev. A 58 (1998) 4394.
- [28] C. P. Yang, S. I. Chu, S. Han, Phys. Rev. A 70 (2004) 022329;
 Z. J. Zhang, Z. X. Man, Phys. Lett. A 341 (2005) 55; Z. X. Man *et al.*, J. Phys. B: At. Mol. Opt. Phys. 40 (2007) 1767.
- [29] S. Popescu, Phys. Rev. Lett. 72 (1994) 797.
- [30] S. B. Zheng, Chin. Phys. Lett. 23 (2006) 2356; S. Bandyopadhyay, B. Sanders, Phys. Rev. A 74 (2006) 32310.
- [31] C. Trump, D. Bru β , M. Lewenstein, Phys. Lett A. 279 (2001) 7.
- [32] L. Vaidman, Phys. Rev. A 49 (1994) 1473.
- [33] M. H. Y. Moussa, Phys. Rev A. 55 (1997) R3287.
- [34] S. B. Zheng, Phys. Rev. A 69 (2001) 064302.
- [35] This nonunity fidelity is a consequence of the imperfect discrimination of the Bell states, as required by the OP. See R. W. Chhajlany, A. Wójcik, Phys. Rev. A 73 (2006) 016302.
- [36] N. G. de Almeida, R. Napolitano, M. H. Y. Moussa, Phys. Rev. A 62 (2000) 033815.
- [37] S. B. Zheng and G. C. Guo, Phys. Rev. Lett. 85 (2000) 2392.
- [38] M. Brune et al., Phys. Rev. Lett. 77 (1996) 4887.
- [39] G. Rempe, F. Schmidt-Kaler, H. Walther, Phys. Rev. Lett. 46 (1990) 2783.
- [40] S. Gleyzes et. al, Nature 446 (2007) 297.
- [41] J. M. Raimond, M. Brune, S. Haroche, Rev. Mod. Phys. 73 (2001) 565.
- [42] F. Grosshans, P. Grangier, Phys. Rev. A 64 (2001) R010301.