Spin-Selective Localization Due to Intrinsic Spin-Orbit Coupling

Yaroslav Tserkovnyak and Shimul Akhanjee

Department of Physics and Astronomy, University of California, Los Angeles, California 90095, USA

(Dated: April 23, 2019)

We study spin-dependent diffusive transport in the presence of a tunable spin-orbit (SO) interaction in a two-dimensional (2D) electron system. The spin precession in the SO coupling field affects the interference between different electronic trajectories that is governed by a covariant curvature associated with the SO interaction. Controlling this curvature field by modulating the SO coupling strength and its gradients by, e.g., electric or elastic means, opens intriguing possibilities for exploring spin-selective localization physics. In particular, applying a weak magnetic field allows the control of the electron localization independently for two spin directions, with the spin-quantization axis "engineered" by appropriate SO interaction gradients.

PACS numbers: 72.25.Dc,73.20.Fz,71.70.Ej,85.75.-d

Recent years have seen a growing interest in spin-orbit (SO) interactions in metals, semiconductors, and topological insulators, with significant advances in the physics of spin Hall effects and other phenomena, where SO coupling profoundly modifies the topology of electronic transport [1, 2, 3, 4]. The interest in these problems is also fueled by the desire to develop spintronic functionality in alternative to magnetic systems.

Spin-transport phenomena are in general associated with a gauge freedom of position-dependent spin rotation, with a natural choice of spin gauge often dictated by the geometry of a measuring setup. For a basic theoretical understanding, however, the primary focus is generally devoted to physical quantities that can be expressed in terms of a gauge-covariant curvature field. In particular, a vanishing curvature would imply that the SO coupling can be formally eliminated, while it could appear experimentally as a "persistent spin structure" [5].

The SU(2) gauge symmetry and the associated Yang-Mills curvature field were recently discussed in the contexts of the spin Hall physics, the Aharonov-Casher effect, and other spin-transport phenomena [6]. Additionally, there have been parallel studies on laser-induced non-Abelian gauge fields in cold-atom optical lattices [7], that discussed the Hofstadter "moth" spectrum, the Anderson localization transition, and the possibility of realizing non-Abelian interferometry. In this paper, we study this SU(2) gauge field in the context of disordered two-dimensional (2D) electron systems, utilizing special "nearly Abelian" limit of the non-Abelian Stoke's theorem. To illustrate our general ideas by a practical example, we consider quantum transport corrections in a disordered 2D conductor with intrinsic SO coupling. In particular, we envision a tunable spin-selective localization, controlled by electric gates or by elastic strain, which can spatially modulate the strength of the SO interaction.

Our formalism helps to unravel the precise geometric nature of the precession of electrons in SO fields, which is governed by the SU(2) curvature field, according to the non-Abelian Stoke's theorem. In the context of the weak-localization physics, we are interested in the interference of the closed time-reversed paths, which we show to be described by a simplified version of the Stoke's theorem. The latter applies to smaller loops with a moderate net spin precession, while a large spin precession is responsible for the destruction of the phase coherence of longer time-reversed loops. Our discussion should be relevant to recent experimental studies of weak (anti)localization in 2D heterostructures with strong intrinsic SO coupling [8], and is complementary to the established theoretical framework for disordered systems [9], by explicitly revealing the geometric structure and exploring the role of SO gradients. As a side comment, we note that weak-localization transport corrections (corresponding to maximally crossed diagrams in the Kubo formalism) have subtle but important consequences for the spin Hall physics in disordered 2D Rashba systems [10]. New perspective in the problem of spin-dependent interference and localization effects may thus benefit other areas of spintronics.

In order to illustrate the essential physics, let us consider the following time-independent Hamiltonian:

$$\hat{H} = \frac{1}{2m} \left[-i\hbar \nabla + \frac{e}{c} \hat{\mathbf{A}}(\mathbf{r}) \right]^2 - eV(\mathbf{r}), \qquad (1)$$

where *m* is the effective mass and -e is the electron charge. The scalar potential $V(\mathbf{r})$ describes a quenched disorder and the 2×2 "vector potential" $\hat{\mathbf{A}}$ is responsible for the SO coupling. The 2×2 electron propagator from \mathbf{r}_i to \mathbf{r}_f along a certain contour *c* in time *t* is given by

$$\hat{K}(\mathbf{r}_f, \mathbf{r}_i; t) = T_c e^{-(2\pi i/\phi_0) \int_c d\mathbf{r} \cdot \mathbf{A}(\mathbf{r})} K(\mathbf{r}_f, \mathbf{r}_i; t) \,. \tag{2}$$

Here, $\phi_0 = hc/e$ is the magnetic flux quantum, $K(\mathbf{r}_f, \mathbf{r}_i; t)$ is the spinless propagator corresponding to Eq. (1) with $\hat{\mathbf{A}} \equiv 0$, and T_c is the path-ordering operator. If we are interested in the interference along different trajectories, the basic building block for describing SO coupling effects is provided by the Wilson-loop integral

$$\hat{W}_c(\mathbf{r}) = T_c e^{-(2\pi i/\phi_0) \oint_c d\mathbf{r}' \cdot \hat{\mathbf{A}}(\mathbf{r}')}, \qquad (3)$$

$$\hat{W}_c(\mathbf{r}) = e^{-2\pi i \phi/\phi_0} \quad \text{(Abelian)}, \qquad (4)$$

where $\phi = \int_{s} d\mathbf{S} \cdot (\nabla \times \mathbf{A})$ is the magnetic flux integrated over the surface area $d\mathbf{S}$. Note that Eq. (4) is gauge invariant, since \hat{W}_{c} is now expressed in terms of the magnetic field $\mathbf{B} = \nabla \times \mathbf{A}$.

For the general case, when the connection $\hat{\mathbf{A}}$ has noncommuting matrix components, the non-Abelian Stoke's theorem relates the exponentiated loop integral (3) over $\hat{\mathbf{A}}$ to the curvature described by the covariant tensor field

$$\hat{\mathcal{F}}_{\mu\nu} = \partial_{\mu}\hat{A}_{\nu} - \partial_{\nu}\hat{A}_{\mu} + \frac{2\pi i}{\phi_0}[\hat{A}_{\mu}, \hat{A}_{\nu}], \qquad (5)$$

where μ and ν label the spatial components [11]:

$$\hat{W}_c(\mathbf{r}) = T_s e^{-(\pi i/\phi_0) \int_s dr'^{\mu} \wedge dr'^{\nu} \tilde{\mathcal{F}}_{\mu\nu}(\mathbf{r}')} \,. \tag{6}$$

Here, T_s is the surface-ordering operator [11] and $\tilde{\mathcal{F}}_{\mu\nu}$ is the path-dependent curvature defined by

$$\tilde{\mathcal{F}}_{\mu\nu}(\mathbf{r}') = \hat{K}(\mathbf{r}, \mathbf{r}') \hat{\mathcal{F}}_{\mu\nu}(\mathbf{r}') \hat{K}(\mathbf{r}', \mathbf{r}) .$$
(7)

Clearly the mathematical structure for non-Abelian fields requires a more elaborate approach than the more familiar Abelian version. Let us, therefore, start by including only the leading noncommutative corrections to the Stoke's theorem. Namely, if the loop c encloses a sufficiently small area $S \sim r^2$, we can approximate

$$\hat{W}_c(\mathbf{r}) \approx e^{-2\pi i \phi/\phi_0 + \mathcal{O}(r^3)}, \qquad (8)$$

where

$$\hat{\phi} = \frac{1}{2} \int_{s} dr'^{\mu} \wedge dr'^{\nu} \hat{\mathcal{F}}_{\mu\nu}(\mathbf{r}') = \int_{s} dS \, \hat{\mathcal{F}}_{12}(\mathbf{r}') \qquad (9)$$

is the "plain" flux of the field $\hat{\mathcal{F}}$ through the loop in 2D, without any surface ordering or path dependence. The relative corrections to the approximation (8) scale as $r/l_{\rm so}$, where $l_{\rm so} \sim \phi_0/|\mathbf{A}|$ is the spin-precession length.

As a specific example, let us consider the Rashba Hamiltonian, corresponding to

$$\hat{\mathbf{A}}(\mathbf{r}) = \lambda(\mathbf{r}) \, \mathbf{z} \times \hat{\boldsymbol{\sigma}} \,, \tag{10}$$

where \mathbf{r} lies in the xy plane and \mathbf{z} is a normal unit vector. $\hat{\boldsymbol{\sigma}}$ is a vector of the Pauli matrices, which are generators of the SU(2) group. In Ref. [3], the problem of the "boundary spin Hall effect" was discussed where $\lambda(\mathbf{r})$ changes abruptly from some constant value λ_0 to zero.

This could model, for instance, a lateral Hall contact for spin injection by a Rashba system [12]. Such a variation of $\lambda(\mathbf{r})$ results in the fictitious magnetic field

$$\hat{\mathbf{B}} = \boldsymbol{\nabla} \times \hat{\mathbf{A}} = (\hat{\boldsymbol{\sigma}} \cdot \boldsymbol{\nabla} \lambda) \mathbf{z}, \qquad (11)$$

directed along the z axis, which has opposite signs for spins up and down along the gradient $\nabla \lambda$. If the SO strength λ variation is abrupt on the scale of $l_{\rm so}$, the fictitious field (11) dominates the boundary physics, resulting in a spin-dependent (ordinary) Hall effect. The noncommutative contribution to the gauge field (5) in this case is small. However, in the opposite limit of a homogeneous SO coupling, the curvature field is determined entirely by the commutator: $\hat{\mathcal{F}}_{12} \propto i[\hat{A}_1, \hat{A}_2] \propto \hat{\sigma}_z$. Let us notice one qualitative difference between the limits of the homogeneous and strongly inhomogeneous SO interactions: In the former case, the spin quantization set by the curvature field is along the normal z axis, while in the latter it is along the in-plane direction determined by the gradient $\nabla \lambda$ [4]. In the very special case of a combination of the linear Dresselhaus and Rashba SO interactions of equal strength, we have a trivial (Abelian) SU(2) field, whose curvature vanishes in homogeneous systems [5].

The Wilson-loop integral (3), in principle contains all of the information about the spin precession and interference in the presence of the SO coupling. The problem simplifies either in the presence of strong inhomogeneities, making the theory effectively Abelian to the leading order in the SO strength [3], or in the case of a simple geometry, such as a single-loop Aharonov-Casher effect [4]. A more general 2D theory would require evaluating Eq. (6) over many possible closed trajectories, which is a formidable task without a simple representation for the surface integral on the right-hand side of the equation. Furthermore, it is not a priori clear how to express various physical quantities of interest in terms of W_c . We will argue that the physics of weak localization steers clear of these issues. First of all, the integral (3) enters naturally in the semiclassical path-integral construction of the quantum transport corrections [13]. Furthermore, we will see that Eq. (8) gives an adequate approximation for many cases of interest.

Let us recall the quantum correction to the classical conductivity in disordered systems [13]:

$$\sigma_Q = -\frac{4e^2}{h} D \int_{\tau} dt \tilde{R}(t) e^{-t/\tau_{\varphi}} , \qquad (12)$$

where $D = v_F^2 \tau/2$ is the semiclassical diffusion coefficient and $dt\tilde{R}(t)$ is the return probability associated with the interference of all classical paths with their time-reversed counterparts within the time interval (t, t+dt). The elastic scattering time τ serves as the lower limit of integration, while the upper limit is effectively set by the phasecoherence time $\tau_{\varphi} \gg \tau$, according to the factor $e^{-t/\tau_{\varphi}}$.

The classical probability of return can be formally written as $R(t) = \int d[\mathbf{r}(t')] P_t[\mathbf{r}(t')]$, for closed paths in a neighborhood of the trajectory $\mathbf{r}(t')$. The probability for the realization of a (coarse-grained) Boltzmannian path $\mathbf{r}(t')$ is given by the Wiener measure:

$$P_t[\mathbf{r}(t')] \propto e^{-\int_0^t dt' \dot{\mathbf{r}}(t')^2/4D} \,. \tag{13}$$

In the presence of a SO interaction (1), the interference between counter-propagating closed trajectories acquires an SU(2) phase correction (3). For a sufficiently weak SO interaction, the trajectories are assumed to be otherwise unaffected. (We will specify the exact condition below.) The interference between the counter-propagating closed trajectories is affected by the spin precession as follows [13]:

$$\hat{R}(t) = \int d[\mathbf{r}(t')] P_t[\mathbf{r}(t')] \hat{W}_c[\mathbf{r}(t')] \hat{W}_c^{\dagger}[\bar{\mathbf{r}}(t')], \qquad (14)$$

where $\bar{\mathbf{r}}(t')$ is the time-reversed trajectory. The return probability \tilde{R} entering Eq. (12) has to be appropriately spin-averaged: $\tilde{R} = \text{Tr}[\hat{R}]/2$. Let us notice that $\hat{W}_c^{\dagger}[\bar{\mathbf{r}}(t')] = \hat{W}_c[\mathbf{r}(t')]$ and use the approximation (8) to find for the spin return interference:

$$\hat{W}_c[\mathbf{r}(t')]\hat{W}_c^{\dagger}[\bar{\mathbf{r}}(t')] \approx e^{-4\pi i\hat{\phi}/\phi_0} \,. \tag{15}$$

We will return later to discussing the range of validity of this approximation in the context of weak localization. Tracing this over spin to get the quantum conductivity correction (12), we see that the calculation now reduces to finding the eigenstates $\pm \phi$ of the traceless Hermitian matrix $\hat{\phi}$ [in the case of an SU(2) field]. For the more general SU(2)×U(1) field (e.g., Rashba SO plus the electromagnetic vector potential), $\hat{\phi}$ may acquire a finite trace, with the two eigenstates ϕ_{\pm} becoming

$$\phi_{\pm} = \phi_m \pm \phi, \quad \hat{\phi} = \phi_m + \phi \,\hat{\boldsymbol{\sigma}} \cdot \mathbf{n} \,. \tag{16}$$

Here, ϕ_m is the ordinary magnetic flux through the loop, ϕ is the SO contribution, and **n** is a unit vector defining spin-quantization axis. In general, by not relying upon the approximation (8), the same composition is upheld, although the Hermitian matrix $\hat{\phi}$ that determines spin precession is not given simply by the flux (9) [4]. Note that we have neglected Zeeman splitting, focusing on the case when the magnetic field is oriented perpendicular to the plane of the electron motion [14].

One intriguing consequence of Eq. (16) is the possibility to delocalize electrons spin-selectively by an appropriate combination of an applied magnetic field and an adjustable SO interaction. The relevant spin-quantization axis **n** can be chosen to be along the in-plane gradient of a strongly inhomogeneous SO strength, Eq. (11). This would provide a pragmatic scenario, since we could have large fictitious fields with moderate magnitudes of the SO interaction, while allowing for some freedom in choosing the spin-quantization axis. In practice, however, this hinges on the ability to control large gradients of the SO strength with elastic strain or electric gates, for instance.

The quantum conductivity correction in the presence of an effective flux (16) is given in 2D for each spin by [13]

$$\sigma_Q^{\pm} = -\psi \left(\frac{1}{2} + \frac{\phi_0}{8\pi |\phi_{\pm}|} \frac{\tau_{\varphi}}{\tau}\right) + \psi \left(\frac{1}{2} + \frac{\phi_0}{8\pi |\phi_{\pm}|}\right), \quad (17)$$

in terms of the digamma function ψ , in units of $e^2/2\pi h$. When $|\phi_{\pm}| \ll \phi_0$, this approximates to

$$\sigma_Q^{\pm} \approx -\ln\frac{\tau_{\varphi}}{\tau} + \frac{2}{3} \left(\frac{2\pi\phi_{\pm}}{\phi_0}\right)^2 \quad (|\phi_{\pm}| \ll \phi_0) \,. \tag{18}$$

 ϕ_{\pm} entering these equations is given by the flux of the effective field $B_{\pm} = B_m \pm B$ through the area $l_{\varphi}^2 = D\tau_{\varphi}$. B_m is the out-of-plane magnetic field, and the fictitious field $\pm B$ is given by the eigenvalues of the curvature

$$\hat{\mathcal{F}}_{12} = \hat{\boldsymbol{\sigma}} \cdot \boldsymbol{\nabla} \lambda - (4\pi i \lambda^2 / \phi_0) \hat{\sigma}_z , \qquad (19)$$

assuming for simplicity a constant gradient $\nabla \lambda$, on the scale of l_{φ} . Notice that because of the covariance of the Yang-Mills curvature $\hat{\mathcal{F}}$, B must be gauge invariant. In the opposite limit of $|\phi_{\pm}| = |B_{\pm}|l_{\varphi}^2 \gg \phi_0$ (when, e.g., $\tau_{\varphi} \to \infty$), Eq. (17) approximates to

$$\sigma_Q^{\pm} \approx -\ln \frac{\phi_0}{8\pi |B_{\pm}| D\tau} \quad (|\phi_{\pm}| \gg \phi_0) \,. \tag{20}$$

The above limits are obtained easily by using the asymptotic behavior of the digamma function: $\psi(1/2 + z) = \ln z + 1/24z^2 + \mathcal{O}(1/z^4)$ at $z \to \infty$ and $\psi(1/2 + z) = \mathcal{O}(z)$ at $z \to 0$. We only need to note that our semiclassical treatment based on Eqs. (13)-(15) requires that $|B_{\pm}| \ll \phi_0/l^2$ (where $l = v_F \tau$ is the mean free path), so that we always have $(\phi_0/|\phi_{\pm}|)(\tau_{\varphi}/\tau) \gg 1$. In particular, for the field B_{\pm} determined by the SO curvature (19), we require $l \ll l_{so} = \phi_0/2\lambda$, when λ is uniform. Note that for a fixed and finite SO interaction, Eq. (18) corresponds to a dip of the total conductivity as a function of the physical magnetic field B_m at zero field, since $(\phi_+^2 + \phi_-^2)/2 = \phi^2 + \phi_m^2$, while Eq. (20) has a peak at $B_m = 0$, since $|\phi_+\phi_-| = |\phi^2 - \phi_m^2|$.

It is now appropriate to discuss the legitimacy of the approximation (8). As mentioned earlier, this approximation requires that $r \ll l_{\rm so}$. Otherwise, the neglected non-Abelian corrections to the Stoke's theorem due to spin precession become appreciable. To study this, let us turn off magnetic field ($\phi_m = 0$) and consider two extreme cases: Homogeneous λ , such that the curvature field $\hat{\mathcal{F}}_{12}$ is determined by the second term in Eq. (19), and large constant gradient $\nabla \lambda$, such that the curvature is dominated by the first term. In the former case, the flux through the area $l_{\rm so}^2$ is $\phi \sim \lambda^2 l_{\rm so}^2/\phi_0 \sim \phi_0$, which means the non-Abelian corrections to the preceding weak-localization analysis based on Eq. (8) can be

appreciable if $l_{\rm so} \ll l_{\varphi}$. In fact, we in general cannot reduce the problem to uncorrelated propagation of two spin projections, since only the triplet subspace of the Cooperon precesses on the scale of $l_{\rm so}$, while the singlet channel is unaffected by SO coupling [13, 15]. Eq. (20) corresponding to $l_{\rm so} \ll l_{\varphi}$, in the case of a homogeneous Rashba SO with $\phi_m = 0$, thus should only be valid for the triplet channels, for which $l_{\rm so}$ sets the cutoff lengthscale for trajectories included in the Cooperon. Separating the antilocalization singlet contribution, we thus immediately find

$$\sigma_Q/2 \approx \ln(l_\varphi/l) - 3\ln(l_{\rm so}/2\pi l) \quad (l_\varphi \gg l_{\rm so}), \qquad (21)$$

in agreement with Ref. [15]. The factor of 3 accounts for the triplet degeneracy. In particular, note the crossover from localization to antilocalization, as $l_{\rm so}$ is made shorter. In the other extreme of a large spin-orbit gradient, we should generally have a similar concern of non-Abelian corrections beyond Eq. (8): Although the larger gradient increases the curvature field and shrinks the relevant lengthscale corresponding to $\phi \sim \phi_0$, it is easy to show the spin precession will remain significant. We will not pursue this problem in detail here, but it is worthwhile to remark that it can be overcome, e.g., by tuning the Rashba parameter λ in combination with the linear Dresselhaus SO interaction, to reduce the non-Abelian contribution to the curvature field [5].

It is also instructive to compare the limit of weak intrinsic SO coupling, Eq. (18), to the delocalizing correction in the presence of a weak extrinsic SO interaction due to quenched disorder [13]:

$$\sigma_Q/2 \approx -\ln(\tau_\varphi/\tau) + 2\tau_\varphi/\tau_{\rm so}\,,\tag{22}$$

where $\tau_{\rm so}^{-1}$ is the extrinsic SO scattering rate due to impurities. In the case of a homogeneous Rashba SO coupling (10), the D'yakonov-Perel (DP) spin-relaxation rate [1] is $\tau_{\rm DP}^{-1} \propto \lambda^2 \tau$, in the relevant regime of $l \ll l_{\rm so}$. The leading SO correction to the localization (18) is thus proportional to $\tau_{\rm DP}^{-2}$, i.e., the square of the DP spin-relaxation rate, since the curvature defining the effective flux ϕ_{\pm} is proportional to λ^2 , for a uniform λ .

We can easily understand the different localization dependence on the spin-relaxation rates, $\tau_{\rm so}^{-1}$ and $\tau_{\rm DP}^{-1}$, in the two cases, by examining how the SU(2) phase prefactor in the propagator (2) scales with the strength of the SO coupling for closed loops. The uncorrelated quenched SO disorder leads to a memoryless (Markovian) spin precession, which is similar for both open and closed trajectories. However, the intrinsic SO coupling (10) combined with ordinary momentum scattering results in a DP spinprecessional random walk, which is qualitatively different for open and closed paths. In the case of open trajectories responsible for DP spin relaxation, this random walk is Markovian. However, the net spin precession is reduced by closing a trajectory, since $\langle \hat{\mathbf{A}} \cdot d\mathbf{r}/dt \rangle_{\rm loop} = 0$, hence the higher-order scaling of the delocalizing correction with the spin-relaxation rate. Note that this vanishing of the average field driving spin precession along the closed loops is exact only for a homogeneous λ . Making λ inhomogeneous would enhance the delocalizing corrections, since now $\langle \hat{\mathbf{A}} \cdot d\mathbf{r}/dt \rangle_{\text{loop}} \neq 0$. For strongly inhomogeneous λ , the curvature (19) scales linearly with the SO strength λ , and the delocalizing correction linearly with the DP spin-relaxation rate $\tau_{\rm DP}^{-1}$. This is analogous to the effect of SO interaction on weak localization and quantum conductance fluctuations in a chaotic quantum dot [16]: The relevant SO scattering rate is substantially reduced due to the geometrical constraint on particle trajectories through a chaotic quantum dot, in the case of a uniform SO vector potential (10). By introducing SO nonuniformity, however, the rate of SO scattering can be augmented to a level comparable to that in the bulk [16].

Let us finally discuss some potential practical consequences of controlling the Yang-Mills curvature field (19) by tuning the appropriate SO coupling profile $\lambda(\mathbf{r})$. The local SO strength λ determines the curvature component $\propto \lambda^2$ with the spin-quantization axis out of plane, while the gradient $\nabla \lambda$ controls the in-plane curvature component. The combination of the two determines the net spin-quantization axis **n**, while the eigenvalues $\pm B$ of the curvature matrix $\hat{\mathcal{F}}_{12}$, which are opposite for the two spins, combined with the ordinary magnetic field B_m , which is the same for the two spins, can in principle result in any desirable spin-dependent field B_{\pm} . One interesting possibility would be to tune the field B_{\pm} deep in the localized regime such that it vanishes for one spin species only, say $B_{-} = 0$. In this case, only the spin \uparrow would be delocalized by a large enough field B_+ , along a desirable spin-quantization axis. This could pave the way for a gate- or strain-tunable spin filter in two dimensions, without the use of ferromagnetic materials, which may become useful for developing semiconductor-based spintronic applications.

We thank Arne Brataas and Xun Jia for numerous helpful discussions.

- I. Žutić, J. Fabian, and S. Das Sarma, Rev. Mod. Phys. 76, 323 (2004).
- [2] H.-A. Engel, E. I. Rashba, and B. I. Halperin, in Handbook of Magnetism and Advanced Magnetic Materials, edited by H. Kronmüller and S. S. P. Parkin (John Wiley & Sons, Chichester, UK, 2007), vol. 5, pp. 2858–2877; M. König, H. Buhmann, L. W. Molenkamp, T. L. Hughes, C.-X. Liu, X.-L. Qi, and S.-C. Zhang, arXiv:0801.0901.
- [3] Y. Tserkovnyak, B. I. Halperin, A. A. Kovalev, and A. Brataas, Phys. Rev. B 76, 085319 (2007).
- [4] Y. Tserkovnyak and A. Brataas, Phys. Rev. B 76, 155326 (2007).
- [5] B. A. Bernevig, J. Orenstein, and S.-C. Zhang, Phys.

Rev. Lett. 97, 236601 (2006).

- [6] P.-Q. Jin, Y.-Q. Li, and F.-C. Zhang, J. Phys. A: Math. Gen. **39**, 7115 (2006); P.-Q. Jin and Y.-Q. Li, condmat/0608303; N. Hatano, R. Shirasaki, and H. Nakamura, quant-ph/0701076.
- K. Osterloh, M. Baig, L. Santos, P. Zoller, and M. Lewenstein, Phys. Rev. Lett. 95, 010403 (2005); I. I. Satija, D. C. Dakin, J. Y. Vaishnav, and C. W. Clark, arXiv:0711.2433.
- [8] T. Koga, J. Nitta, T. Akazaki, and H. Takayanagi, Phys. Rev. Lett. **89**, 046801 (2002); J. B. Miller, D. M. Zumbühl, C. M. Marcus, Y. B. Lyanda-Geller, D. Goldhaber-Gordon, K. Campman, and A. C. Gossard, Phys. Rev. Lett. **90**, 076807 (2003); C. Schierholz, T. Matsuyama, U. Merkt, and G. Meier, Phys. Rev. B **70**, 233311 (2004); F. E. Meijer, A. F. Morpurgo, T. M. Klapwijk, T. Koga, and J. Nitta, Phys. Rev. B **70**, 201307(R) (2004); V. A. Guzenko, T. Schäpers, and H. Hardtdegen, arXiv:0709.1326; B. Grbić, R. Leturcq, T. Ihn, K. En-

sslin, D. Reuter, and A. D. Wieck, arXiv:0711.0492.

- [9] W. Knap, C. Skierbiszewski, A. Zduniak, E. Litwin-Staszewska, D. Bertho, F. Kobbi, J. L. Robert, G. E. Pikus, F. G. Pikus, S. V. Iordanskii, et al., Phys. Rev. B 53, 3912 (1996); V. M. Edelstein, J. Phys.: Condens. Matter. 7, 1 (1995); A. Punnoose, Appl. Phys. Lett. 88, 252113 (2006).
- [10] O. Chalaev and D. Loss, Phys. Rev. B 71, 245318 (2005).
- [11] B. Broda, math-ph/0012035.
- [12] I. Adagideli and G. E. W. Bauer, Phys. Rev. Lett. 95, 256602 (2005).
- [13] S. Chakravarty and A. Schmid, Phys. Rep. 140, 193 (1986), and references therein.
- [14] S. Maekawa and H. Fukuyama, J. Phys. Soc. Jpn. 50, 2516-2524 (1981).
- [15] M. A. Skvortsov, JETP Lett. 67, 133 (1998).
- [16] P. W. Brouwer, J. N. H. J. Cremers, and B. I. Halperin, Phys. Rev. B 65, 081302(R) (2002).