Hidden SDW order and effective low-energy theory for FeAs superconductors

Zheng-Yu Weng¹

¹Center for Advanced Study, Tsinghua University, Beijing, 100084, China

(Dated: September 6, 2021)

We propose a simple effective model to describe FeAs superconductors. This model is based on the assumption of a *local* spin-density-wave (SDW) order, with its magnetization direction allowed to fluctuate. It is shown that the long-range order with momentum $\mathbf{Q} = (\pi, \pi)$ is generally unstable in competing with the kinetic energy of the charge carriers. A true weak SDW order is formed in the undoped case with an additional momentum shift $\mathbf{Q}_s = (\pi, 0)$ due to the peculiar Fermi surface nesting. In the doped case, the fluctuating long-range order driven by kinetic energy can naturally result in a d-wave superconducting condensation. Such low-energy physics is protected by the presence of the local SDW which sustains some kind of "Mott gaps" for the multiband *d*-electrons near the Fermi energy.

PACS numbers: 71.10.Hf,74.20.Mn,71.27+a,75.20.Hr

Introduction. The recent discovery of the iron-based superconductors[1, 2, 3, 4, 5] has stimulated a lot of interest concerning underlying mechanism for superconductivity in this new superconductor family. It has been established by various measurements [6, 7, 8] that there exists an SDW order in the undoped LaOFeAs compound below $T \sim 150$ K, which quickly disappears with the electrons doped into the system, where the superconducting phase starts to set in with transition temperatures being raised beyond 50 K.[5, 9] While the general phase diagram[1, 2, 3, 4, 9] reminds us some interesting similarities with the cuprate superconductors, the delectrons on Fe seem much more itinerant with multiorbitals crossing the Fermi level as indicated by the band structure calculations, [10, 11, 12, 13, 14, 15, 16] compared to the isolated $Cu_{3d_{x^2-y^2}}$ - $O_{2p_{x,y}}$ antibonding orbitals in the cuprates. Many theoretical proposals are based on itinerant approaches [17, 18, 19, 20] with the emphasis on the important role played by various magnetic fluctuations, while some conjectures are also made from the side of large-spin Mott insulators.[21]

The LDA calculations[14, 15] have found an energetically robust SDW state at the antiferromagnetic (AF) momentum $\mathbf{Q} = (\pi, \pi)$ with a large Fe moment ~ 2.3 $\mu_{\rm B}$ per site, but experimentally only a weak SDW ordering with a different magnetic momentum $\mathbf{Q}_s = (\pi, 0)$, which further doubles the unit cell of the former SDW state composed of two Fe per cell, has been identified[7, 8] in the undoped case. The latter SDW (called stripe type below) was predicted by the first principle band structure calculation[6] due to the nesting Fermi surfaces of the hole and electron pockets, which is much more "fragile" and easily destroyed as the doped electrons fill up the small hole pockets at small doping.

In this paper, we will make a very simple proposal by assuming that the SDW state with an AF momentum \mathbf{Q} remains strong *locally* in both the undoped and small doped regime. The corresponding profile of the electron density of states is schematically illustrated in Fig. 1.



FIG. 1: (color online) Schematic profile of the density of states near the Fermi energy in the presence of the local SDW meanfields M_a and M_b , which decide the "charge gaps" for the *a*orbital and *b*-orbital bands according to H_0 [(9)]. The lower half of each band (the so-called α -band, see text) is assumed below the Fermi energy E_F and is filled, while the upper half (the β -band) is only partially filled. Here *a*-bands refer to the bands passing the Fermi level, which are more d_{xy} , d_{xz} , d_{yz} like, and *b*-bands are more of $d_{x^2-y^2}$ character according to the LDA result.[14, 15]

Based on this minimal model, we can show that the longrange part of this SDW is actually generically unstable, by coupling to the charge carriers near the Fermi level. Such instability can result in a weak stripe-type SDW order in the undoped case where the nesting Fermi surfaces are present, and naturally a superconducting condensation when the former is destroyed at small doping. In this whole regime, however, the local order of such an SDW remains robust to "protect" the low-energy physics, which resembles the role of the "Mott gap" in the cuprate superconductors.[22]

Minimal Model. In our model Hamiltonian $H_{\text{eff}} = H_{\text{band}} + H_{\text{I}}$, the first term is a tight-binding model

$$H_{\text{band}} = -\sum_{ij,a,b,\alpha} t_{ij}^{a,b} c_{ia\sigma}^{\dagger} c_{jb\sigma}$$
(1)

which describes the electron effective hoppings between



FIG. 2: (color online) (a) The iron atoms form a square lattice with the dashed diamond denoting the unit cell of the SDW order state with magnetic momentum \mathbf{Q} . (b) The magnetic Brillouin zone (BZ) is illustrated by the dashed diamond shape. Two circles at the Γ and M points are hole and electron pockets connected by a momentum \mathbf{Q}_s . The smaller square is a reduced BZ corresponding to an enlarged unit cell indicated by the largest square in (a).

the d-orbitals of the Fe ions on square lattice (Fig. 2(a)) at the nearest neighboring (NN) and next nearest neighboring (NNN) sites, including intra- and inter-orbital hoppings with the superscripts a, b specifying the orbitals. There have been several proposals[14, 16, 19, 20] for H_{band} based on the LDA calculations in order to capture the relevant bands near the Fermi energy.

The second term $H_{\rm I}$ reflects the effective influence of the Coulomb interaction on the *d*-electrons, which includes the on-site and NN repulsions as well as the Hund's rule ferromagnetic coupling, and has the following "mean-field" look

$$H_{\rm I} = -\sum_{i,a} \mathbf{M}_{ia} \cdot \mathbf{S}_{ia} \tag{2}$$

where \mathbf{M}_{ia} denotes an effective SDW mean-field felt by the electron spin \mathbf{S}_{ia} of the orbital *a* at site *i*. In an SDW ordered state, one would have $\mathbf{M}_{ia} \propto \sum_{b(\neq a)} J_H^{a,b} \langle \mathbf{S}_{ib} \rangle + U_a \langle \mathbf{S}_{ia} \rangle$, where $J_H^{a,b}$ is the Hund's rule coupling constant and U_a is the on-site repulsion. We shall make the following ansatz

$$\mathbf{M}_{ia} = M_a (-1)^i \mathbf{n}_i \tag{3}$$

where a single unit vector $(-1)^i \mathbf{n}_i$ will describe the true polarization direction of \mathbf{M}_{ia} . Here the staggered factor $(-1)^i$ is introduced such that $\mathbf{n}_i = \hat{\mathbf{n}}$ will correspond to a true AF order, but in a general case \mathbf{n}_i will not be fixed around a particular direction as only the relative change $\Delta_{\alpha} \mathbf{n}_i \equiv \mathbf{n}_{i+\hat{\alpha}} - \mathbf{n}_i \ (\hat{\alpha} = \hat{x}, \hat{y})$ will enter the Hamiltonian as shown below.

The local SDW field M_a will be assumed large according to the LDA calculation, [14, 15] with the magnetization at different orbitals tightly aligned together by the Hund's coupling. Under this assumption, the longwavelength, low-energy fluctuations of \mathbf{n}_i may be treated as an independent degree of freedom. \mathbf{n}_i will be selfconsistently determined by coupling to the electrons near the Fermi level in H_{eff} .

Effective theory. One may redefine \mathbf{n}_i as the new $\hat{\mathbf{z}}$ -axis for the spin index of the electron spinor operator:

$$\hat{c}_{ia} = U_i \hat{a}_{ia} \tag{4}$$

with $\hat{a}_{ia}^{\dagger}=(\hat{a}_{ia,\uparrow}^{\dagger},\hat{a}_{ia,\downarrow}^{\dagger})$ by an SU(2) rotation

$$U_i^{\dagger} \mathbf{n}_i \cdot \hat{\sigma} U_i = \hat{\sigma}_z. \tag{5}$$

Then $H_{\rm I}$ simply reduces to

$$H_{\rm I} = -\sum_{i,a} M_a (-1)^i S_{ia}^z \tag{6}$$

while H_{band} becomes

$$H_{\text{band}} = -\sum_{ij,a,b} t^{a,b}_{ij} \hat{a}^{\dagger}_{ia} \left(U^{\dagger}_{i} U_{j} \right) \hat{a}_{jb} \tag{7}$$

We may further rewrite

$$H_{\rm eff} = H_0 + H_1 \tag{8}$$

where $H_0 \equiv H_{\rm I} + H_{\rm band}[U_i^{\dagger}U_j = 1]$ is simply an SDW mean-field Hamiltonian for the multibands, which can be diagonalized as

$$H_{0} = \sum_{\mathbf{k},a} \xi_{\mathbf{k}}^{a+} \left(\hat{\alpha}_{\mathbf{k}a}^{\dagger} \hat{\alpha}_{\mathbf{k}a} + \hat{\beta}_{\mathbf{k}a}^{\dagger} \hat{\beta}_{\mathbf{k}a} \right) - \sum_{\mathbf{k},a} E_{\mathbf{k}}^{a} \left(\hat{\alpha}_{\mathbf{k}a}^{\dagger} \hat{\alpha}_{\mathbf{k}a} - \hat{\beta}_{\mathbf{k}a}^{\dagger} \hat{\beta}_{\mathbf{k}a} \right) + \text{const.} \quad (9)$$

by a canonical transformation $\hat{a}_{\mathbf{k}a} = u_{\mathbf{k}}^{a}\hat{\alpha}_{\mathbf{k}a} - v_{\mathbf{k}}^{a}\hat{\sigma}_{z}\hat{\beta}_{\mathbf{k}a}, \quad \hat{a}_{\mathbf{k}+\mathbf{Q}a} = v_{\mathbf{k}}^{a}\hat{\sigma}_{z}\hat{\alpha}_{\mathbf{k}a} + u_{\mathbf{k}}^{a}\hat{\beta}_{\mathbf{k}a} \text{ with } u_{\mathbf{k}}^{a} = \left[(1 - \xi_{\mathbf{k}}^{a-}/E_{\mathbf{k}}^{a})/2\right]^{1/2}, \quad v_{\mathbf{k}}^{a} = \left[(1 + \xi_{\mathbf{k}}^{a-}/E_{\mathbf{k}}^{a})/2\right]^{1/2}.$ Here $\xi_{\mathbf{k}}^{a\pm} \equiv \left(\varepsilon_{\mathbf{k}}^{a} \pm \varepsilon_{\mathbf{k}+\mathbf{Q}}^{a}\right)/2$ and

$$E_{\mathbf{k}}^{a} = \sqrt{\left(\xi_{\mathbf{k}}^{a-}\right)^{2} + \left(M_{a}/2\right)^{2}}$$
(10)

where $\varepsilon_{\mathbf{k}}^{a}$ denotes the bare spectrum determined by (1) (setting the chemical potential $\mu = 0$). Note that the band label *a* here can be different from the original orbital label in (1) because of the mixture of orbitals, and in obtaining (10), the same M_{a} is assumed for the mixed orbitals. Here **k** is defined in the magnetic Brillouin zone (BZ) with the magnetic momentum $\mathbf{Q} = (\pi, \pi)$, which coincides with the BZ of two irons per unit cell (Fig. 2(b)).

Now let us consider the term with $U_i^{\dagger}U_i \neq 1$:

$$H_{1} = -\sum_{ij,a,b} t_{ij}^{a,b} \hat{a}_{ia}^{\dagger} \left(U_{i}^{\dagger} U_{j} - 1 \right) \hat{a}_{jb}$$
(11)

We shall focus on the case in which the α -bands are all filled up by the electrons, and the Fermi energy is located in some of β -bands, which corresponds to both the undoped and electron-doped (or slightly hole-doped) situations as illustrated by Fig. 1. After integrating out the α -bands and by assuming $\Delta_{\alpha} \mathbf{n}_i$ is small, H_1 may be simplified[23] (at large M_a) to

$$H_{1} \simeq \frac{1}{2} \sum_{\mathbf{k},\mathbf{q},\sigma} \nabla \varepsilon_{\mathbf{k}} \cdot \left(\mathbf{D}_{\mathbf{q}}\right)_{\sigma,-\sigma} \left(-\theta_{\mathbf{k}+\mathbf{q}} + \sigma \bar{\theta}_{\mathbf{k}+\mathbf{q}}\right) \beta_{\mathbf{k}+\mathbf{q},\sigma}^{\dagger} \beta_{\mathbf{k},-\sigma} + \text{H.c.} + \frac{J_{\text{eff}}}{8} \sum_{i} \left(\Delta \mathbf{n}_{i}\right)^{2}$$
(12)

where $\mathbf{D}_{\mathbf{q}}$ is a Fourier transformation of

$$\mathbf{D}_{i} = \frac{1}{2} (\mathbf{\Delta} \mathbf{n}_{i} \times \mathbf{n}_{i}) \cdot U_{i}^{\dagger} \hat{\sigma} U_{i}$$
(13)

and $\theta_{\mathbf{k}+\mathbf{q}}(\bar{\theta}_{\mathbf{k}+\mathbf{q}})$ is the step function restricting $\mathbf{k} + \mathbf{q}$ within (outside) the magnetic BZ. Note that for simplicity we have omitted the band indices in (12), where the spiral field $\mathbf{D}_{\mathbf{q}}$ will couple to the electrons from all β -bands near the Fermi level. It shows that the twist of \mathbf{n}_i is kinetic energy driven. Such a spiral twist is balanced by the "superexchange" term in H_1 , where $J_{\text{eff}} \simeq 1/N \sum_{\mathbf{k}a} (1 - n_{\mathbf{k}a}^{\beta}) (\xi_{\mathbf{k}}^{a-})^2 / E_{\mathbf{k}}^{a}$ provides the spin stiffness against the twist of the SDW order and is mainly contributed by the filled α -bands and is reduced with the increasing filling $n_{\mathbf{k}}^{\beta}$ in the β -bands.

Some technical remarks with regarding the derivation of (12) are in order. Strictly speaking, the dynamic field \mathbf{n}_i should be introduced in a path-integral formalism, [23, 24] where the temporal term $U_i^{\dagger} \partial_t U_i$ will also enter the Lagrangian. As shown previously for the one-band Hubbard model, [23, 25] the terms like $U_i^{\dagger} \partial_t U_i$ will play an important role in determining the dynamics of \mathbf{n}_i at half-filling where an SDW long-range order is present. In this case, by expanding \mathbf{n}_i around $\mathbf{\hat{z}}, \ \mathbf{D_q} \simeq i \mathbf{q} (\mathbf{n_q} \times \mathbf{\hat{z}}) \cdot \hat{\sigma}, \ \text{the propagator of } \mathbf{D_q} \ \text{reduces}$ to $\mathcal{D}^{s}(\mathbf{q},t) \sim \mathbf{q}\mathbf{q} \langle \mathbf{n}_{\mathbf{q}}^{\perp}(t) \cdot \mathbf{n}_{\mathbf{q}}^{\perp}(0) \rangle$ which is proportional to the spin-wave propagator with the coupling to doped particles vanishing as $\mathbf{q} \to 0$. However, once a finite density of the holes or electrons are doped into the system, the dynamic spiral fluctuation of \mathbf{D}_q which couples to the doped charge carriers in (12) will become dominant over the other fluctuation terms like $U_i^{\dagger} \partial_t U_i$, leading to intrinsic instabilities [25] of the system (see below).

The propagator $\mathcal{D}(\mathbf{q}, t) = -i \langle T_t \operatorname{Tr} [\mathbf{D}_{\mathbf{q}}(t) \mathbf{D}_{-\mathbf{q}}(0)] \rangle$ can be obtained by integrating out the β -band

$$\mathcal{D}(\mathbf{q},\omega) \simeq -1/\left[\mathbf{\Pi}_{\beta}(\mathbf{q},\omega) + J_{\text{eff}}\right]$$
 (14)

where $\mathbf{\Pi}_{\beta}(\mathbf{q},\omega)$ is the "bubble" diagram contribution of the β -band electrons, with the "bare" $\mathbf{\Pi}_{\beta}^{0}(\mathbf{0},0) \sim$ $-v_{F}^{2}N_{F} \sim -t$ where v_{F} and N_{F} are the Fermi velocity and density of states based on H_{0} , which is expected to dominate over J_{eff} on general grounds,[25] reflecting the fact that the hopping energy is always dominant over the superexchange energy locally, leading to the instability of the AF long-range order.

"Stripe" instability in the undoped case. According to the band structure calculations, [14, 15] in the undoped case, there are hole-pockets around the Γ point and electron pockets around the M point around the Fermi level, which are connected by a particular nesting momentum \mathbf{Q}_s , and can be described by H_0 with the proper choice of $\varepsilon_{\mathbf{k}}^a$. Then, to take advantage of the enhanced response function at \mathbf{Q}_s in H_0 , the interaction term H_1 will naturally induce

$$\nabla \varepsilon_{\mathbf{k}} \cdot \langle \mathbf{D}_{\mathbf{Q}_s} \rangle \neq 0 \tag{15}$$

and result in a mean-field $\left< \beta^{\dagger}_{\mathbf{k}+\mathbf{Q}_{s},\sigma}\beta_{\mathbf{k},-\sigma} \right> \neq 0$ with a $\sqrt{2} \times \sqrt{2}$ folding of the BZ as shown in Fig. 2(b). Consequently a gap will be opened up at the Fermi energy to stabilize $\mathcal{D}(\mathbf{q},\omega)$ in (14). In contrast to the unstable SDW with the AF momentum \mathbf{Q} , the true magnetic ordering (15) is realized with an additional spiral twist at momentum \mathbf{Q}_{s} , which, known as a stripe-type SDW, was first predicted in Ref. [6] and has been recently confirmed experimentally by neutron scattering.[7, 8]

Superconducting instability. With introducing a small amount of doped electrons, the Fermi energy will move up such that the small hole pockets around the Γ point get filled up and the Fermi surface nesting disappears. Then the static "stripe" SDW order (15) vanishes, and $\mathbf{D}_{\mathbf{q}}$ will become unstable again.

Rewriting

$$U_{i}^{\dagger}U_{j} = U_{i}^{\dagger}U_{i_{1}}U_{i_{1}}^{\dagger}U_{i_{2}}U_{i_{2}}^{\dagger}...U_{i_{N_{i_{j}}-1}}^{\dagger}U_{j}$$
(16)

with $\{i_s\} = i_0, i_1, ..., i_{N_{ij}}$ a sequence of lattice sites connecting $i \equiv i_0$ and $j \equiv i_{N_{ij}}$ and using $U_{i_s}^{\dagger}U_{i_{s+1}} \simeq 1 + i\mathbf{D}_{i_s} \cdot \hat{\eta}_{i_{s+1},i_s} \simeq e^{i\mathbf{D}_{i_s} \cdot \hat{\eta}_{i_s+1,i_s}} (\hat{\eta}_{i_s+1,i_s} = \mathbf{r}_{i_s+1} - \mathbf{r}_{i_s})$ (neglecting the phase associated with the solid angle spanned by $\mathbf{n}_{i_s}, \mathbf{n}_{i_{s+1}}, \hat{\mathbf{z}}$), one finds

$$\left\langle U_i^{\dagger} U_j \right\rangle \sim \left\langle \exp i \int_i^j d\mathbf{r} \cdot \mathbf{D} \right\rangle$$

$$\simeq \exp \left[-\frac{1}{2} \int \int_i^j d\mathbf{r} \cdot \langle \mathbf{D} \mathbf{D} \rangle \cdot d\mathbf{r}' \right]$$

$$\simeq \exp \left[-\frac{|i-j|^2}{2\xi_s^2} \right]$$
(17)

where the spin correlation length $\xi_s \sim 1/\left[-\int d\omega Im \mathcal{D}(0,\omega)\right]^{1/2}$. Unless $Im \mathcal{D}(\mathbf{q},\omega) \to 0$ at $\mathbf{q} \to 0$ as in the spin wave case, the spiral fluctuations in (14) will generally lead to a spin disordered state with a finite ξ_s , but its precise nature has to be self-consistently determined in view of the divergent $\mathcal{D}(\mathbf{q},\omega)$ in the small \mathbf{q} and ω regime.

The dynamic spiral fluctuations of the magnetization directions are kinetic energy driven by the electrons near the Fermi level. However, this will also make the electrons lose their long-wavelength coherence. Indeed, the single-particle propagator for the electrons to leading order of approximation can be written as

$$G^{e}(i,j;t) \simeq \left\langle U_{i}^{\dagger}(t)U_{j}(0) \right\rangle G^{a}$$
 (18)

where G^a is the propagator for the *a*-particles whose leading term is coherent governed by the mean-field Hamiltonian H_0 in (9). Thus the dynamic spiral fluctuation will quickly damp the coherent motion of the quasiparticle beyond the spin correlation length ξ_s via $\langle U_i^{\dagger}(t)U_j(0) \rangle$.

By contrast, after averaging over the \mathbf{n}_i -field, the electron singlet pair operator $\hat{\Delta}_{ij}^{SC} \equiv \sum_{\sigma} \sigma c_{i\sigma}^{\dagger} c_{j-\sigma}^{\dagger}$ can be expressed by

$$\Delta_{ij}^{SC} \simeq \left\langle U_i^{\dagger} U_j \right\rangle \sum_{\sigma} \sigma a_{i\sigma}^{\dagger} a_{j-\sigma}^{\dagger} \\ \approx \sum_{\sigma} \sigma a_{i\sigma}^{\dagger} a_{j-\sigma}^{\dagger}$$
(19)

at $|i-j| \ll \xi_s$. It clearly shows that in such a spin disordered state, the singlet pairs of electrons can still propagate coherently. The above contrast between the single electron and singlet pairs of electrons indicates an instability of the system towards superconductivity with $\sum_{\sigma} \sigma \left\langle a_{i\sigma}^{\dagger} a_{j-\sigma}^{\dagger} \right\rangle \neq 0$. The latter can be indeed realized by exchanging the dynamic spiral fluctuations, $\mathcal{D}(\mathbf{q},\omega)$, between the β -electrons based on (12), which in turn stabilizes $\mathcal{D}(\mathbf{q},\omega)$ through a renormalized $\mathbf{\Pi}_{\beta}(\mathbf{q},\omega)$ as the β -electrons form Cooper pairs. It is noted that the stabilized $\operatorname{Re}\mathcal{D}(\mathbf{q},\omega) < 0$ at small \mathbf{q} and ω will lead to an attractive (repulsive) interaction between β -electrons if $\mathbf{k} + \mathbf{q}$ is within (outside) the magnetic BZ according to (12). It means that the β -electrons will form the dominantly $d_{x^2-y^2}$ -wave pairing at four M points in Fig. 2(b), similar to the case in the cuprates. Physically this kinetic-energy-driven pairing can be understood as that two electrons sitting at, say, NN sites, gain enhanced hopping energies by sharing the spiral twist of \mathbf{n}_i 's between the two sites.

Discussions. It is interesting to point out that T_c is upper bounded in the BCS theory because, whereas the softening phonon can enhance the attractive force, it also leads to the structural instability of the solids. Here the superconductivity is caused by strong dynamic spiral fluctuations which are in a "melting" SDW regime. T_c is expected to fall when such dynamic spiral fluctuations get reduced at higher electron doping concentration (the local magnetization M_a should be eventually destroyed when the Fermi level reaches beyond the highest β -band shown in Fig. 1, presumably the one with the dominant $d_{x^2-y^2}$ characters according to the LDA calculation[15]).

Although the d-electrons in Fe-based compounds are believed quite itinerant, our proposal suggests that the underlying physics is still far from the conventional itinerant magnetic metals in the following sense. Due to the presence of a large *local* SDW mean-field M_a , each band near the Fermi level are still split into the lower and upper Hubbard bands, with the lower one filled by the electrons in both undoped and electron-doped cases, which are responsible for the origin of large M_a 's. In contrast to the single-band Hubbard model relevant to the cuprate superconductors, [22] however, here some of the upper Hubbard bands are already at the Fermi level even in the undoped case. A search for the depleted density of states below the Fermi level by, say, photoemission, [26] may provide useful information concerning the correctness of the present model. The presence of a sizable local moment, which is distinguished from the itinerant approach, may be also investigated via various magnetic measurement above T_c , including the magnetic susceptibility. [27]

Finally we caution that in the present model the effect of the unit cell doubling with two irons per cell due to the crystal field reason (As ions are displaced above and below the Fe plane alternatively) has not been considered, which may drastically affect our results if it becomes sufficiently strong because in that case the α - and β -band splitting due to the local SDW may no longer be complete and α -bands can become partially filled.

In conclusion, we have proposed a simple effective model to describe the low-energy physics in FeAs superconductors. This model is based on an assumption that there exists a robust local SDW order, but its magnetization direction is intrinsically unstable against forming a true long range order due to the competition with the kinetic energy of the charge carriers near the Fermi energy. In the undoped case, a weak SDW order of stripe type is formed due to the peculiar nesting structure of the Fermi surfaces. In the doped case, the dynamic melting of the SDW order will result in the d-wave superconducting pairing of the doped electrons near the Fermi energy.

Acknowledgments: Stimulating discussions with Z.Y. Lu, Z. Fang, K. Wu, J.W. Mei, T. Li are acknowledged. We also thank N.L. Wang, H.H. Wen, and especially X.H. Chen for generously sharing their experimental results before publication. This work is supported by NSFC grant no. 10688401.

- Y. Kamihara, et al., J. Am. Chem. Sco. 128, 10012 (2006); Y. Kamihara, et al., J. Am. Chem. Sco. 130, 3296 (2008).
- [2] H.-H. Wen, et al., Europhys. Lett. 82, 17009 (2008).
- [3] X. H. Chen, et al., arXiv:0803.3603 (2008).
- [4] G. F. Chen, et al., arXiv:0803.3790 (2008).

- [5] Z. A. Ren, et al., arXiv:0803.4283 (2008);
 arXiv:0804.2053 (2008);
- [6] J. Dong, et al., arXiv:0803.3426 (2008).
- [7] C. Cruz, et al., arXiv:0804.0795(2008).
- [8] M. A. McGuire, et al., arXiv:0804.0796 (2008).
- [9] R. H. Liu, et al., arXiv:0804.2105 (2008).
- [10] S. Lebègue, Phys. Rev. B 75, 035110 (2008).
- [11] D. Singh and M. Du, arXiv:0803.0429 (2008); I. Mazin, et al., arXiv:0803.2740 (2008).
- [12] K. Haule, et al., arXiv:0803.1279 (2008).
- [13] G. Xu, et al., arXiv:0803.1282 (2008).
- [14] C. Cao, et al., arXiv:0803.3236(2008).
- [15] F. Ma and Z. Y. Lu arXiv:0803.3286(2008).
- [16] K. Kuroki, et al., arXiv:0803.3325 (2008).

- [17] X. Dai, et al., arXiv:0803.3982 (2008).
- [18] Q. Han, et al., Europhys. Lett. 82, 37007 (2008).
- [19] S. Raghu, et al., arXiv:0804.1113 (2008).
- [20] P. A. Lee and X. G. Wen, arXiv:0804.1113 (2008).
- T. Li, arXiv:0803.1279 (2008); G. Baskaran, arXiv:0803.1341 (2008); Q. M. Si and E. Abrahams, arXiv:0804.2480 (2008).
- [22] P. W. Anderson, Science 235, 1196 (1987).
- [23] Z. Y. Weng, et al., Phys. Rev. B43,3790 (1990).
- [24] H. J. Schulz, Phys. Rev. Lett. 65, 2462 (1990).
- [25] Z. Y. Weng, Phys. Rev. lett. 66, 2156 (1991).
- [26] H.W. Ou, et al., arXiv:0803.4328 (2008).
- [27] X. H. Chen, unpublished.