On Estimation of Finite Population Proportion *

Xinjia Chen

November 20, 2018

Abstract

In this paper, we study the classical problem of estimating the proportion of a finite population. First, we consider a fixed sample size method and derive an explicit sample size formula which ensures a mixed criterion of absolute and relative errors. Second, we consider an inverse sampling scheme such that the sampling is continue until the number of units having a certain attribute reaches a threshold value or the whole population is examined. We have established a simple method to determine the threshold so that a prescribed relative precision is guaranteed. Finally, we develop a multistage sampling scheme for constructing fixed-width confidence interval for the proportion of a finite population. Powerful computational techniques are introduced to make it possible that the fixed-width confidence interval ensures prescribed level of coverage probability.

1 Fixed Sample Size Method

The estimation of the proportion of a finite population is a basic and very important problem in probability and statistics [6, 8]. Such problem finds applications spanning many areas of sciences and engineering. The problem is formulated as follows.

Consider a finite population of N units, among which there are M units having a certain attribute. The objective is to estimate the proportion $p = \frac{M}{N}$ based on sampling without replacement.

One popular method of sampling is to draw n units without replacement from the population and count the number, \mathbf{k} , of units having the attribute. Then, the estimate of the proportion is taken as $\hat{p} = \frac{\mathbf{k}}{n}$. In this process, the sample size n is fixed.

Clearly, the random variable **k** possesses a hypergeometric distribution. The reliability of the estimator $\hat{p} = \frac{\mathbf{k}}{n}$ depends on n. For error control purpose, we are interested in a crucial question as follows:

^{*}The author is currently with Department of Electrical Engineering, Louisiana State University at Baton Rouge, LA 70803, USA, and Department of Electrical Engineering, Southern University and A&M College, Baton Rouge, LA 70813, USA; Email: chenxinjia@gmail.com

For prescribed margin of absolute error $\varepsilon_a \in (0,1)$, margin of relative error $\varepsilon_r \in (0,1)$, and confidence parameter $\delta \in (0,1)$, how large the sample size n should be to guarantee

$$\Pr\left\{\left|\widehat{\boldsymbol{p}}-\boldsymbol{p}\right| < \varepsilon_a \text{ or } \left|\frac{\widehat{\boldsymbol{p}}-\boldsymbol{p}}{\boldsymbol{p}}\right| < \varepsilon_r\right\} > 1 - \delta?$$
(1)

In this regard, we have

Theorem 1 Let $\varepsilon_a \in (0,1)$ and $\varepsilon_r \in (0,1)$ be real numbers such that $\frac{\varepsilon_a}{\varepsilon_r} + \varepsilon_a \leq \frac{1}{2}$. Then, (1) is guaranteed provided that

$$n > \frac{\varepsilon_r \ln \frac{2}{\delta}}{(\varepsilon_a + \varepsilon_a \varepsilon_r) \ln(1 + \varepsilon_r) + (\varepsilon_r - \varepsilon_a - \varepsilon_a \varepsilon_r) \ln \left(1 - \frac{\varepsilon_a \varepsilon_r}{\varepsilon_r - \varepsilon_a}\right)}.$$
(2)

The proof of Theorem 1 is given in Appendix A. It should be noted that conventional methods for determining sample sizes are based on normal approximation, see [6] and the references therein. In contrast, Theorem 1 offers a rigorous method for determining sample sizes. To reduce conservativeness, a numerical approach has been developed by Chen [4] which permits exact computation of the minimum sample size.

2 Inverse Sampling of Finite Population

To estimate the proportion p, a frequently-used sampling method is the *inverse sampling* scheme described as follows:

Continuing sampling from the population (without replacement) until r units found to carry the attribute or the number of sample size n reaches the population size N. The estimator of the proportion p is taken as the ratio $\tilde{p} = \frac{k}{n}$, where k is the number of units having the attribute among the n units.

Clearly, the reliability of the estimator \tilde{p} depends on the threshold value r. Hence, we are interested in a crucial question as follows:

For prescribed margin of relative error $\varepsilon \in (0,1)$ and confidence parameter $\delta \in (0,1)$, how large the threshold r should be to guarantee

$$\Pr\left\{|\widetilde{\boldsymbol{p}} - p| < \varepsilon p\right\} > 1 - \delta?$$

For this purpose, we have

Theorem 2 For any $\varepsilon \in (0, 1)$,

$$\Pr\left\{|\widetilde{\boldsymbol{p}} - p| \ge \varepsilon p\right\} \le \mathscr{Q}(\varepsilon, r)$$

where

$$\mathscr{Q}(\varepsilon, r) = (1+\varepsilon)^{-r} \exp\left(\frac{\varepsilon r}{1+\varepsilon}\right) + (1-\varepsilon)^{-r} \exp\left(-\frac{\varepsilon r}{1-\varepsilon}\right),$$

which is monotonically decreasing with respect to r. Moreover, for any $\delta \in (0,1)$, there exists a unique number r^* such that $\mathscr{Q}(\varepsilon, r^*) = \delta$ and

$$\max\left\{\frac{(1+\varepsilon)\ln\frac{1}{\delta}}{(1+\varepsilon)\ln(1+\varepsilon)-\varepsilon}, \ \frac{(1-\varepsilon)\ln\frac{2}{\delta}}{(1-\varepsilon)\ln(1-\varepsilon)+\varepsilon}\right\} < r^* < \frac{(1+\varepsilon)\ln\frac{2}{\delta}}{(1+\varepsilon)\ln(1+\varepsilon)-\varepsilon}.$$

The proof of Theorem 2 is given in Appendix B. As an immediate consequence of Theorem 2, we have

Corollary 1 Let ε , $\delta \in (0,1)$. Then, $\Pr\{|\widetilde{p} - p| < \varepsilon p\} > 1 - \delta$ provided that

$$r > \frac{(1+\varepsilon)\ln\frac{2}{\delta}}{(1+\varepsilon)\ln(1+\varepsilon) - \varepsilon}$$
(3)

3 Multistage Fixed-width Confidence Intervals

So far we have only considered point estimation for the proportion p. Interval estimation is also an important method for estimating p. Motivated by the fact that a confidence interval must be sufficiently narrow to be useful, we shall develop a multistage sampling scheme for constructing a fixed-width confidence interval for the proportion, p, of the finite population discussed in previous sections.

Note that the procedure of sampling without replacement can be precisely described as follows:

Each time a single unit is drawn without replacement from the remaining population so that every unit of the remaining population has equal chance of being selected.

Such a sampling process can be exactly characterized by random variables X_1, \dots, X_N defined in a probability space $(\Omega, \mathscr{F}, \Pr)$ such that X_i denotes the characteristics of the *i*-th sample in the sense that $X_i = 1$ if the *i*-th sample has the attribute and $X_i = 0$ otherwise. By the nature of the sampling procedure, it can be shown that

$$\Pr\{X_i = x_i, \ i = 1, \cdots, n\} = \binom{M}{\sum_{i=1}^n x_i} \binom{N-M}{n-\sum_{i=1}^n x_i} \Big/ \left[\binom{n}{\sum_{i=1}^n x_i} \binom{N}{n} \right]$$

for any $n \in \{1, \dots, N\}$ and any $x_i \in \{0, 1\}$, $i = 1, \dots, n$. Based on random variables X_1, \dots, X_N , we can define a multistage sampling scheme of the following basic structure. The sampling process is divided into s stages with sample sizes $n_1 < n_2 < \dots < n_s$. The continuation or termination of sampling is determined by decision variables. For each stage with index ℓ , a decision variable $D_{\ell} = \mathcal{D}_{\ell}(X_1, \dots, X_{n_{\ell}})$ is defined based on random variables $X_1, \dots, X_{n_{\ell}}$. The decision variable D_{ℓ} assumes only two possible values 0, 1 with the notion that the sampling is continued until $D_{\ell} = 1$ for some $\ell \in \{1, \dots, s\}$. Since the sampling must be terminated at or before the s-th stage, it is required that $D_s = 1$. For simplicity of notations, we also define $D_{\ell} = 0$ for $\ell = 0$. Our goal is to construct a fixed-width confidence interval $(\boldsymbol{L}, \boldsymbol{U})$ such that $\boldsymbol{U} - \boldsymbol{L} \leq 2\varepsilon$ and that $\Pr\{\boldsymbol{L} 1 - \delta$ for any $p \in \{\frac{i}{N} : 0 \leq i \leq N\}$ with prescribed $\varepsilon \in (0, \frac{1}{2})$ and $\delta \in (0, 1)$. Toward this goal, we need to define some multivariate functions as follows.

For $\alpha \in (0,1)$ and integers $0 \leq k \leq n \leq N$, let $\mathcal{L}(N,n,k,\alpha)$ be the smallest integer M_l such that $\sum_{i=k}^{n} {M_l \choose i} {N-M_l \choose n-i} / {N \choose n} > \frac{\alpha}{2}$. Let $\mathcal{U}(N,n,k,\alpha)$ be the largest integer M_u such that $\sum_{i=0}^{k} {M_u \choose i} {N-M_u \choose n-i} / {N \choose n} > \frac{\alpha}{2}$. Let $n_{\max}(N,\alpha)$ be the smallest number n such that $\mathcal{U}(N,n,k,\alpha) - \mathcal{L}(N,n,k,\alpha) \leq 2\varepsilon N$ for $0 \leq k \leq n$. Let $n_{\min}(N,\alpha)$ be the largest number n such that $\mathcal{U}(N,n,k,\alpha) - \mathcal{L}(N,n,k,\alpha) > 2\varepsilon N$ for $0 \leq k \leq n$.

Theorem 3 Let $\zeta > 0$ and $\rho > 0$. Let $n_1 < n_2 < \cdots < n_s$ be the ascending arrangement of all distinct elements of $\left\{ \left[\left[\frac{n_{\max}(N,\zeta\delta)}{n_{\min}(N,\zeta\delta)} \right]^{\frac{1}{\tau}} n_{\min}(N,\zeta\delta) \right] : i = 0, 1, \cdots, \tau \right\}$ with $\tau = \left[\frac{1}{\ln(1+\rho)} \ln \frac{n_{\max}(N,\zeta\delta)}{n_{\min}(N,\zeta\delta)} \right]$. For $\ell = 1, \cdots, s$, define $K_{\ell} = \sum_{i=1}^{n_{\ell}} X_i$ and D_{ℓ} such that $D_{\ell} = 1$ if $\mathcal{U}(N, n_{\ell}, K_{\ell}, \zeta\delta) - \mathcal{L}(N, n_{\ell}, K_{\ell}, \zeta\delta) \leq 2\varepsilon N$; and $D_{\ell} = 0$ otherwise. Suppose the stopping rule is that sampling is continued until $D_{\ell} = 1$ for some $\ell \in \{1, \cdots, s\}$. Define $\mathbf{L} = \frac{1}{N} \times \mathcal{L}(N, \mathbf{n}, \sum_{i=1}^{\mathbf{n}} X_i, \zeta\delta)$ and $\mathbf{U} = \frac{1}{N} \times \mathcal{U}(N, \mathbf{n}, \sum_{i=1}^{\mathbf{n}} X_i, \zeta\delta)$, where \mathbf{n} is the sample size when the sampling is terminated. Then, a sufficient condition to guarantee $\Pr\{\mathbf{L} 1 - \delta$ for any $p \in \{\frac{i}{N} : 0 \leq i \leq N\}$ is that

$$\sum_{\ell=1}^{s} [\Pr\{\mathcal{L}(N, n_{\ell}, K_{\ell}, \zeta\delta) \ge M, \ \boldsymbol{D}_{\ell-1} = 0, \ \boldsymbol{D}_{\ell} = 1 \mid M\} + \Pr\{\mathcal{U}(N, n_{\ell}, K_{\ell}, \zeta\delta) \le M, \ \boldsymbol{D}_{\ell-1} = 0, \ \boldsymbol{D}_{\ell} = 1 \mid M\}] < \delta$$
(4)

for all $M \in \{0, 1, \dots, N\}$, where (4) is satisfied if $\zeta > 0$ is sufficiently small.

It should be noted that Theorem 3 has employed the double-decision-variable method recently proposed by Chen in [1]. To further reduce computational complexity, the techniques of bisection confidence tuning and domain truncation developed in [1, 2] can be very useful.

A Proof of Theorem 1

To prove the theorem, we shall introduce function

$$g(\varepsilon, p) = (p + \varepsilon) \ln \frac{p}{p + \varepsilon} + (1 - p - \varepsilon) \ln \frac{1 - p}{1 - p - \varepsilon}$$

where $0 < \varepsilon < 1 - p$. We need some preliminary results.

The following lemma is due to Hoeffding [7].

Lemma 1

$$\Pr\{\widehat{\boldsymbol{p}} \ge p + \varepsilon\} \le \exp(n \ g(\varepsilon, p)) \quad for \quad 0 < \varepsilon < 1 - p < 1,$$
$$\Pr\{\widehat{\boldsymbol{p}} \le p - \varepsilon\} \le \exp(n \ g(-\varepsilon, p)) \quad for \quad 0 < \varepsilon < p < 1.$$

The following Lemmas 2–4 have been established in [3].

Lemma 2 Let $0 < \varepsilon < \frac{1}{2}$. Then, $g(\varepsilon, p)$ is monotonically increasing with respective to $p \in (0, \frac{1}{2} - \varepsilon)$ and monotonically decreasing with respective to $p \in (\frac{1}{2}, 1 - \varepsilon)$. Similarly, $g(-\varepsilon, p)$ is monotonically increasing with respective to $p \in (\varepsilon, \frac{1}{2})$ and monotonically decreasing with respective to $p \in (\varepsilon, \frac{1}{2})$ and monotonically decreasing with respective to $p \in (\frac{1}{2} + \varepsilon, 1)$.

Lemma 3 Let $0 < \varepsilon < \frac{1}{2}$. Then,

$$g(\varepsilon, p) > g(-\varepsilon, p) \qquad \forall p \in \left(\varepsilon, \frac{1}{2}\right],$$
$$g(\varepsilon, p) < g(-\varepsilon, p) \qquad \forall p \in \left(\frac{1}{2}, 1 - \varepsilon\right).$$

Lemma 4 Let $0 < \varepsilon < 1$. Then, $g(\varepsilon p, p)$ is monotonically decreasing with respect to $p \in \left(0, \frac{1}{1+\varepsilon}\right)$. Similarly, $g(-\varepsilon p, p)$ is monotonically decreasing with respect to $p \in (0, 1)$.

Lemma 5 Suppose $0 < \varepsilon_r < 1$ and $0 < \frac{\varepsilon_a}{\varepsilon_r} + \varepsilon_a \leq \frac{1}{2}$. Then,

$$\Pr\{\widehat{\boldsymbol{p}} \le p - \varepsilon_a\} \le \exp\left(n \ g\left(-\varepsilon_a, \frac{\varepsilon_a}{\varepsilon_r}\right)\right)$$
(5)

for 0 .

Proof. We shall show (5) by investigating three cases as follows. In the case of $p < \varepsilon_a$, it is clear that

$$\Pr\{\widehat{\boldsymbol{p}} \le p - \varepsilon_a\} = 0 < \exp\left(n \ g\left(-\varepsilon_a, \frac{\varepsilon_a}{\varepsilon_r}\right)\right)$$

In the case of $p = \varepsilon_a$, we have

$$\Pr\{\widehat{\boldsymbol{p}} \le p - \varepsilon_a\} = \Pr\{\widehat{\boldsymbol{p}} = 0\} = \Pr\{\mathbf{k} = 0\}$$
$$= \frac{\binom{N-M}{n}}{\binom{N}{n}} \le \left(\frac{N-M}{N}\right)^n$$
$$= (1-p)^n = (1-\varepsilon_a)^n$$
$$= \lim_{p \to \varepsilon_a} \exp(n \ g(-\varepsilon_a, p))$$
$$< \exp\left(n \ g\left(-\varepsilon_a, \frac{\varepsilon_a}{\varepsilon_r}\right)\right),$$

where the last inequality follows from Lemma 2 and the fact that $\varepsilon_a < \frac{\varepsilon_a}{\varepsilon_r} \leq \frac{1}{2} - \varepsilon_a$.

In the case of $\varepsilon_a , we have$

$$\Pr\{\widehat{\boldsymbol{p}} \le p - \varepsilon_a\} \le \exp(n \ g(-\varepsilon_a, p)) < \exp\left(n \ g\left(-\varepsilon_a, \frac{\varepsilon_a}{\varepsilon_r}\right)\right),$$

where the first inequality follows from Lemma 1 and the second inequality follows from Lemma 2 and the fact that $\varepsilon_a < \frac{\varepsilon_a}{\varepsilon_r} \leq \frac{1}{2} - \varepsilon_a$. So, (5) is established. \Box

Lemma 6 Suppose $0 < \varepsilon_r < 1$ and $0 < \frac{\varepsilon_a}{\varepsilon_r} + \varepsilon_a \leq \frac{1}{2}$. Then,

$$\Pr\{\widehat{\boldsymbol{p}} \ge (1+\varepsilon_r)p\} \le \exp\left(n \ g\left(\varepsilon_a, \frac{\varepsilon_a}{\varepsilon_r}\right)\right)$$
(6)

for $\frac{\varepsilon_a}{\varepsilon_r} .$

Proof. We shall show (6) by investigating three cases as follows. In the case of $p > \frac{1}{1+\varepsilon_r}$, it is clear that

$$\Pr\{\widehat{\boldsymbol{p}} \ge (1+\varepsilon_r)p\} = 0 < \exp\left(n \ g\left(\varepsilon_a, \frac{\varepsilon_a}{\varepsilon_r}\right)\right)$$

In the case of $p = \frac{1}{1+\varepsilon_r}$, we have

$$\Pr\{\widehat{p} \ge (1+\varepsilon_r)p\} = \Pr\{\widehat{p}=1\} = \Pr\{\mathbf{k}=n\}$$
$$= \frac{\binom{M}{n}}{\binom{N}{n}} \le \left(\frac{M}{N}\right)^n$$
$$= p^n = \left(\frac{1}{1+\varepsilon_r}\right)^n$$
$$= \lim_{p \to \frac{1}{1+\varepsilon_r}} \exp(n \ g(\varepsilon_r p, p))$$
$$< \exp\left(n \ g\left(\varepsilon_a, \frac{\varepsilon_a}{\varepsilon_r}\right)\right),$$

where the last inequality follows from Lemma 4 and the fact that $\frac{\varepsilon_a}{\varepsilon_r} \leq \frac{1}{2} \frac{1}{1+\varepsilon_r} < \frac{1}{1+\varepsilon_r}$ as a result of $0 < \frac{\varepsilon_a}{\varepsilon_r} + \varepsilon_a \leq \frac{1}{2}$. In the case of $\frac{\varepsilon_a}{\varepsilon_r} , we have$

$$\Pr\{\widehat{\boldsymbol{p}} \le (1+\varepsilon_r)p\} \le \exp(n \ g(\varepsilon_r p, p)) < \exp\left(n \ g\left(\varepsilon_a, \frac{\varepsilon_a}{\varepsilon_r}\right)\right),$$

where the first inequality follows from Lemma 1 and the second inequality follows from Lemma 4. So, (6) is established.

We are now in a position to prove the theorem. We shall assume (2) is satisfied and show that (1) is true. It suffices to show that

$$\Pr\{|\widehat{\boldsymbol{p}} - p| \ge \varepsilon_a, |\widehat{\boldsymbol{p}} - p| \ge \varepsilon_r p\} < \delta.$$

For 0 , we have

$$\Pr\{|\widehat{\boldsymbol{p}} - p| \ge \varepsilon_a, \ |\widehat{\boldsymbol{p}} - p| \ge \varepsilon_r p\} = \Pr\{|\widehat{\boldsymbol{p}} - p| \ge \varepsilon_a\}$$
$$= \Pr\{\widehat{\boldsymbol{p}} \ge p + \varepsilon_a\} + \Pr\{\widehat{\boldsymbol{p}} \le p - \varepsilon_a\}.$$
(7)

Noting that 0 , we have

$$\Pr\{\widehat{\boldsymbol{p}} \ge p + \varepsilon_a\} \le \exp(n \ g(\varepsilon_a, p)) \le \exp\left(n \ g\left(\varepsilon_a, \frac{\varepsilon_a}{\varepsilon_r}\right)\right),$$

where the first inequality follows from Lemma 1 and the second inequality follows from Lemma 2. It can be checked that (2) is equivalent to

$$\exp\left(n \ g\left(\varepsilon_a, \frac{\varepsilon_a}{\varepsilon_r}\right)\right) < \frac{\delta}{2}$$

Therefore,

$$\Pr{\{\widehat{p} \ge p + \varepsilon_a\}} < \frac{\delta}{2}$$

for 0 .

On the other hand, since $\varepsilon_a < \frac{\varepsilon_a}{\varepsilon_r} < \frac{1}{2}$, by Lemma 5 and Lemma 3, we have

$$\Pr\{\widehat{\boldsymbol{p}} \le p - \varepsilon_a\} \le \exp\left(n \ g\left(-\varepsilon_a, \frac{\varepsilon_a}{\varepsilon_r}\right)\right) \le \exp\left(n \ g\left(\varepsilon_a, \frac{\varepsilon_a}{\varepsilon_r}\right)\right) < \frac{\delta}{2}$$

for 0 . Hence, by (7),

$$\Pr\{|\widehat{\boldsymbol{p}} - p| \ge \varepsilon_a, |\widehat{\boldsymbol{p}} - p| \ge \varepsilon_r p\} < \frac{\delta}{2} + \frac{\delta}{2} = \delta.$$

This proves (1) for 0 .

For $\frac{\varepsilon_a}{\varepsilon_r} , we have$

$$\Pr\{|\widehat{\boldsymbol{p}} - p| \ge \varepsilon_a, \ |\widehat{\boldsymbol{p}} - p| \ge \varepsilon_r p\} = \Pr\{|\widehat{\boldsymbol{p}} - p| \ge \varepsilon_r p\} \\ = \Pr\{\widehat{\boldsymbol{p}} \ge p + \varepsilon_r p\} + \Pr\{\widehat{\boldsymbol{p}} \le p - \varepsilon_r p\}.$$

Invoking Lemma 6, we have

$$\Pr\{\widehat{\boldsymbol{p}} \ge p + \varepsilon_r p\} \le \exp\left(n \ g\left(\varepsilon_a, \frac{\varepsilon_a}{\varepsilon_r}\right)\right).$$

On the other hand,

$$\Pr\{\widehat{\boldsymbol{p}} \le p - \varepsilon_r p\} \le \exp(n \ g(-\varepsilon_r p, p)) \le \exp\left(n \ g\left(-\varepsilon_a, \frac{\varepsilon_a}{\varepsilon_r}\right)\right) \le \exp\left(n \ g\left(\varepsilon_a, \frac{\varepsilon_a}{\varepsilon_r}\right)\right)$$

where the first inequality follows from Lemma 1, the second inequality follows from Lemma 4, and the last inequality follows from Lemma 3. Hence,

$$\Pr\{|\widehat{\boldsymbol{p}} - p| \ge \varepsilon_a, \ |\widehat{\boldsymbol{p}} - p| \ge \varepsilon_r p\} \le 2 \exp\left(n \ g\left(\varepsilon_a, \frac{\varepsilon_a}{\varepsilon_r}\right)\right) < \delta.$$

This proves (1) for $\frac{\varepsilon_a}{\varepsilon_r} . The proof of Theorem 1 is thus completed.$

B Proof Theorem 2

We need some preliminary results. We shall introduce functions

$$\mathcal{M}(z,p) = \ln\left(\frac{p}{z}\right) + \left(\frac{1}{z} - 1\right)\ln\left(\frac{1-p}{1-z}\right)$$

and

$$\mathscr{H}(z,p) = z \ \mathscr{M}(z,p)$$

for 0 < z < 1 and 0 .

Lemma 7 Suppose $1 \le r \le M < N$. Then,

$$\Pr\left\{\frac{r}{n} \le (1-\varepsilon)p\right\} \le (1-\varepsilon)^{-r} \exp\left(-\frac{\varepsilon r}{1-\varepsilon}\right).$$

Proof. Clearly,

$$\Pr\left\{\frac{r}{n} \le (1-\varepsilon)p\right\} = \Pr\left\{n \ge \frac{r}{(1-\varepsilon)p}\right\}$$
$$= \Pr\{n \ge m\}$$

where

$$m = \left\lceil \frac{r}{(1-\varepsilon)p} \right\rceil$$

It can be seen that there exists a real number $\varepsilon^* \in (0,1)$ such that $\varepsilon^* \geq \varepsilon$ and

$$\frac{r}{(1-\varepsilon^*)p} = \left\lceil \frac{r}{(1-\varepsilon)p} \right\rceil.$$

Now let K_m be the number of units having a certain attribute among m units drawn by a sampling without replacement from a finite population of size N with M units having the attribute. Then,

$$\Pr\{\boldsymbol{n} \ge m\} = \Pr\{K_m \le r\}$$
$$= \Pr\left\{\frac{K_m}{m} \le \frac{r}{m}\right\}$$
$$= \Pr\left\{\frac{K_m}{m} \le (1 - \varepsilon^*)p\right\}.$$

Applying the well-known Hoeffding inequality [7] for the case of finite population, we have

$$\Pr\left\{\frac{K_m}{m} \le (1 - \varepsilon^*)p\right\} \le \exp\left(m\mathcal{H}(p - \varepsilon^*p, p)\right)$$
$$= \exp\left(\frac{r}{(1 - \varepsilon^*)p}\mathcal{H}(p - \varepsilon^*p, p)\right)$$
$$= \exp\left(r\mathcal{H}(p - \varepsilon^*p, p)\right)$$
$$\le \exp\left(r\mathcal{M}(p - \varepsilon p, p)\right)$$

where the last inequality follows from $\varepsilon^* \geq \varepsilon$ and the monotone property of $\mathcal{M}(p - \varepsilon p, p)$ with respect to ε , which has been established as Lemma 5 in [5].

From the proof of Lemma 6 of [5], we know that $\mathcal{M}(p - \varepsilon p, p)$ is monotonically decreasing with respect to $p \in (0, 1)$. Hence,

$$\Pr\left\{\frac{r}{n} \le (1-\varepsilon)p\right\} \le \exp\left(r\mathcal{M}(p-\varepsilon p,p)\right) \le \lim_{p \to 0} \exp\left(r\mathcal{M}(p-\varepsilon p,p)\right) = (1-\varepsilon)^{-r} \exp\left(-\frac{\varepsilon r}{1-\varepsilon}\right).$$

The proof of the lemma is thus completed.

Lemma 8 Suppose $1 \le r \le M < N$ and $p + \varepsilon p < 1$. Then,

$$\Pr\left\{\frac{r}{n} \ge (1+\varepsilon)p\right\} \le (1+\varepsilon)^{-r} \exp\left(\frac{\varepsilon r}{1+\varepsilon}\right).$$

Proof. It is clear that

$$\Pr\left\{\frac{r}{n} \ge (1+\varepsilon)p\right\} = \Pr\left\{n \le \frac{r}{(1+\varepsilon)p}\right\}$$
$$= \Pr\{n \le m\}$$

where

$$m = \left\lfloor \frac{r}{(1+\varepsilon)p} \right\rfloor.$$

It can be seen that there exists a real number $\varepsilon^* \in (0,1)$ such that $\varepsilon^* \ge \varepsilon$ and

$$\frac{r}{(1+\varepsilon^*)p} = \left\lfloor \frac{r}{(1+\varepsilon)p} \right\rfloor.$$

Now let K_m be the number of units having a certain attribute among m units drawn by a sampling without replacement from a finite population of size N with M units having the attribute. Then,

$$\Pr\{\boldsymbol{n} \le m\} = \Pr\{K_m \ge r\}$$
$$= \Pr\left\{\frac{K_m}{m} \ge \frac{r}{m}\right\}$$
$$= \Pr\left\{\frac{K_m}{m} \ge (1 + \varepsilon^*)p\right\}.$$

Applying the well-known Hoeffding inequality [7] for the case of finite population, we have

$$\Pr\left\{\frac{K_m}{m} \ge (1+\varepsilon^*)p\right\} \le \exp\left(m\mathcal{H}(\varepsilon^*p,p)\right)$$
$$= \exp\left(\frac{r}{(1+\varepsilon^*)p}\mathcal{H}(\varepsilon^*p,p)\right)$$
$$= \exp\left(r\mathcal{M}(p+\varepsilon^*p,p)\right)$$
$$\le \exp\left(r\mathcal{M}(p+\varepsilon p,p)\right)$$

where the last inequality follows from $\varepsilon^* \geq \varepsilon$ and the monotone property of $\mathcal{M}(p + \varepsilon p, p)$ with respect to ε , which has been established as Lemma 5 in [5].

From the proof of Lemma 6 of [5], we know that $\mathcal{M}(p + \varepsilon p, p)$ is monotonically decreasing with respect to $p \in \left(0, \frac{1}{1+\varepsilon}\right)$. Hence,

$$\Pr\left\{\frac{r}{n} \ge (1+\varepsilon)p\right\} \le \exp\left(r\mathcal{M}(p+\varepsilon p,p)\right) \le \lim_{p \to 0} \exp\left(r\mathcal{M}(p+\varepsilon p,p)\right) = (1+\varepsilon)^{-r} \exp\left(\frac{\varepsilon r}{1+\varepsilon}\right).$$

The proof of the lemma is thus completed.

Now we are in a position to prove Theorem 2. We shall consider the following cases: Case (i): M < r;

 $\begin{array}{ll} \text{Case (ii):} & M = N;\\ \text{Case (iii):} & r = N;\\ \text{Case (iv):} & 1 \leq r \leq M < N \text{ and } p < \frac{1}{1+\varepsilon};\\ \text{Case (v):} & 1 \leq r \leq M < N \text{ and } p = \frac{1}{1+\varepsilon};\\ \text{Case (vi):} & 1 \leq r \leq M < N \text{ and } p > \frac{1}{1+\varepsilon}. \end{array}$

In Case (i), we have $\boldsymbol{n} = N$ and $\boldsymbol{k} = M$. Hence, $\widetilde{\boldsymbol{p}} = p$ and $\Pr\{|\widetilde{\boldsymbol{p}} - p| \ge \varepsilon p\} = 0 \le \mathscr{Q}(\varepsilon, r)$.

In Case (ii), we have $\widetilde{\boldsymbol{p}} = p$ and $\Pr\{|\widetilde{\boldsymbol{p}} - p| \ge \varepsilon p\} = 0 \le \mathscr{Q}(\varepsilon, r)$.

In Case (iii), we have $\widetilde{\boldsymbol{p}} = p$ and $\Pr\{|\widetilde{\boldsymbol{p}} - p| \ge \varepsilon p\} = 0 \le \mathscr{Q}(\varepsilon, r)$.

In Case (iv), we have $\mathbf{k} = r$ and, by Lemma 7 and Lemma 8,

$$\Pr\left\{ |\tilde{\boldsymbol{p}} - \boldsymbol{p}| \ge \varepsilon \boldsymbol{p} \right\} = \Pr\left\{ \frac{r}{\boldsymbol{n}} \le (1 - \varepsilon)\boldsymbol{p} \right\} + \Pr\left\{ \frac{r}{\boldsymbol{n}} \ge (1 + \varepsilon)\boldsymbol{p} \right\}$$
$$\le (1 - \varepsilon)^{-r} \exp\left(-\frac{\varepsilon r}{1 - \varepsilon}\right) + (1 + \varepsilon)^{-r} \exp\left(\frac{\varepsilon r}{1 + \varepsilon}\right)$$
$$= \mathscr{Q}(\varepsilon, r).$$

In Case (v), we have $\boldsymbol{k} = r$ and

$$\Pr\left\{ |\tilde{\boldsymbol{p}} - p| \ge \varepsilon p \right\} = \Pr\left\{ \frac{r}{\boldsymbol{n}} \le (1 - \varepsilon)p \right\} + \Pr\left\{ \frac{r}{\boldsymbol{n}} \ge (1 + \varepsilon)p \right\}$$
$$= \Pr\left\{ \frac{r}{\boldsymbol{n}} \le (1 - \varepsilon)p \right\} + \Pr\left\{ \boldsymbol{k} = \boldsymbol{n} = r \right\}.$$

Notice that

$$\Pr\left\{\boldsymbol{k}=\boldsymbol{n}=r\right\} = \frac{\binom{M}{r}}{\binom{N}{r}} < \left(\frac{M}{N}\right)^r = p^r = \left(\frac{1}{1+\varepsilon}\right)^r < (1+\varepsilon)^{-r} \exp\left(\frac{\varepsilon r}{1+\varepsilon}\right)$$

as a result of $M \leq N$. Therefore, by Lemma 7,

$$\Pr\left\{|\widetilde{\boldsymbol{p}}-\boldsymbol{p}| \ge \varepsilon \boldsymbol{p}\right\} \le (1-\varepsilon)^{-r} \exp\left(-\frac{\varepsilon r}{1-\varepsilon}\right) + (1+\varepsilon)^{-r} \exp\left(\frac{\varepsilon r}{1+\varepsilon}\right) = \mathscr{Q}(\varepsilon,r).$$

In Case (vi), we have $\mathbf{k} = r$, $\Pr\left\{\frac{r}{\mathbf{n}} \ge (1 + \varepsilon)p\right\} = 0$ and, by Lemma 7,

$$\Pr\left\{ \left| \tilde{p} - p \right| \ge \varepsilon p \right\} = \Pr\left\{ \frac{r}{n} \le (1 - \varepsilon)p \right\} + \Pr\left\{ \frac{r}{n} \ge (1 + \varepsilon)p \right\}$$
$$= \Pr\left\{ \frac{r}{n} \le (1 - \varepsilon)p \right\}$$
$$\le (1 - \varepsilon)^{-r} \exp\left(-\frac{\varepsilon r}{1 - \varepsilon}\right) < \mathcal{Q}(\varepsilon, r).$$

So, we have shown $\Pr\{|\tilde{p} - p| \ge \varepsilon p\} \le \mathscr{Q}(\varepsilon, r)$. The other statements of Theorem 2 have been established in [5].

This concludes the proof of Theorem 2.

References

- CHEN, X., "A new framework of multistage estimation," arXiv:0809.1241v5 [math.ST], January 2009.
- [2] X. Chen, "A truncation approach for fast computation of distribution functions," arXiv:0802.3455 [math.ST], February 2008.
- [3] X. Chen, "On Estimation and Optimization of Probability," arXiv:0804.1399, April 2008.
- [4] X. Chen, "Exact computation of minimum sample size for estimating proportion of finite population," arXiv:0707.2115, July 2007.
- [5] X. Chen, "Inverse sampling for nonasymptotic sequential estimation of bounded variable means," arXiv:0711.2801, November 2007.
- [6] M. M. Desu and D. Raghavarao, Sample Size Methodology, Academic Press, 1990.
- [7] W. Hoeffding, "Probability inequalities for sums of bounded variables," J. Amer. Statist. Assoc., vol. 58, 13–29, 1963.
- [8] S. K. Thompson, *Sampling*, Wiley, 2002.