

**REMARK ON THE BOUNDEDNESS OF THE CAUCHY
SINGULAR INTEGRAL OPERATOR ON VARIABLE LEBESGUE
SPACES WITH RADIAL OSCILLATING WEIGHTS**

ALEXEI YU. KARLOVICH

To Professor Kokilashvili on his seventieth birthday

ABSTRACT. Recently V. Kokilashvili, N. Samko, and S. Samko have proved a sufficient condition for the boundedness of the Cauchy singular integral operator on variable Lebesgue spaces with radial oscillating weights over Carleson curves. This condition is formulated in terms of Matuszewska-Orlicz indices of weights. We prove a partial converse of their result.

1. INTRODUCTION AND MAIN RESULT

Let Γ be a rectifiable curve in the complex plane. We equip Γ with Lebesgue length measure $|d\tau|$. We say that a curve Γ is simple if it does not have self-intersections. In other words, Γ is said to be simple if it is homeomorphic either to a line segment or to a circle. In the latter situation we will say that Γ is a Jordan curve. The *Cauchy singular integral* of $f \in L^1(\Gamma)$ is defined by

$$(Sf)(t) := \frac{1}{\pi i} \int_{\Gamma} \frac{f(\tau)}{\tau - t} d\tau \quad (t \in \Gamma).$$

This integral is understood in the principal value sense, that is,

$$\int_{\Gamma} \frac{f(\tau)}{\tau - t} d\tau := \lim_{R \rightarrow 0} \int_{\Gamma \setminus \Gamma(t, R)} \frac{f(\tau)}{\tau - t} d\tau,$$

where $\Gamma(t, R) := \{\tau \in \Gamma : |\tau - t| < R\}$ for $R > 0$. David [4] (see also [3, Theorem 4.17]) proved that the Cauchy singular integral generates the bounded operator S on the Lebesgue space $L^p(\Gamma)$, $1 < p < \infty$, if and only if Γ is a *Carleson (Ahlfors-David regular) curve*, that is,

$$\sup_{t \in \Gamma} \sup_{R > 0} \frac{|\Gamma(t, R)|}{R} < \infty,$$

where for any measurable set $\Omega \subset \Gamma$ the symbol $|\Omega|$ denotes its measure. To have a better idea about Carleson curves, consider the following example. Let $\alpha > 0$ and

$$\Gamma := \{0\} \cup \{\tau \in \mathbb{C} : \tau = x + ix^\alpha \sin(1/x), 0 < x \leq 1\}.$$

Date: October 24, 2018.

2000 Mathematics Subject Classification. Primary 42B20; Secondary 47B38.

Key words and phrases. Variable Lebesgue space, Carleson curve, variable exponent, radial oscillating weight, Matuszewska-Orlicz indices, submultiplicative function.

The author is partially supported by F.C.T. (Portugal) grant FCT/FEDER/POCTI/MAT/59972/2004.

One can show (see [3, Example 1.3]) that Γ is not rectifiable for $0 < \alpha \leq 1$, Γ is rectifiable but not Carleson for $1 < \alpha < 2$, and Γ is a Carleson curve for $\alpha \geq 2$.

A measurable function $w : \Gamma \rightarrow [0, \infty]$ is referred to as a *weight function* or simply a *weight* if $0 < w(\tau) < \infty$ for almost all $\tau \in \Gamma$. Suppose $p : \Gamma \rightarrow [1, \infty]$ is a measurable a.e. finite function. Denote by $L^{p(\cdot)}(\Gamma, w)$ the set of all measurable complex-valued functions f on Γ such that

$$\int_{\Gamma} |f(\tau)w(\tau)/\lambda|^{p(\tau)} |d\tau| < \infty$$

for some $\lambda = \lambda(f) > 0$. This set becomes a Banach space when equipped with the Luxemburg-Nakano norm

$$\|f\|_{p(\cdot), w} := \inf \left\{ \lambda > 0 : \int_{\Gamma} |f(\tau)w(\tau)/\lambda|^{p(\tau)} |d\tau| \leq 1 \right\}.$$

If p is constant, then $L^{p(\cdot)}(\Gamma, w)$ is nothing else but the weighted Lebesgue space. Therefore, it is natural to refer to $L^{p(\cdot)}(\Gamma, w)$ as a *weighted generalized Lebesgue space with variable exponent* or simply as a *weighted variable Lebesgue space*. This is a special case of Musielak-Orlicz spaces [19] (see also [13]). Nakano [20] considered these spaces (without weights) as examples of so-called modular spaces, and sometimes the spaces $L^{p(\cdot)}(\Gamma, w)$ are referred to as weighted Nakano spaces.

Following [12, Section 2.3], denote by \mathbb{W} the class of all continuous functions $\varrho : [0, |\Gamma|] \rightarrow [0, \infty)$ such that $\varrho(0) = 0$, $\varrho(x) > 0$ if $0 < x \leq |\Gamma|$, and ϱ is almost increasing, that is, there is a universal constant $C > 0$ such that $\varrho(x) \leq C\varrho(y)$ whenever $x \leq y$. Further, let \mathbb{W} be the set of all functions $\varrho : [0, |\Gamma|] \rightarrow [0, \infty)$ such that $x^\alpha \varrho(x) \in \mathbb{W}$ and $x^\beta / \varrho(x) \in \mathbb{W}$ for some $\alpha, \beta \in \mathbb{R}$. Clearly, the functions $\varrho(x) = x^\gamma$ belong to \mathbb{W} for all $\gamma \in \mathbb{R}$. For $\varrho \in \mathbb{W}$, put

$$\Phi_{\varrho}^0(x) := \limsup_{y \rightarrow 0} \frac{\varrho(xy)}{\varrho(y)}, \quad x \in (0, \infty).$$

Since $\varrho \in \mathbb{W}$, one can show that the limits

$$m(\varrho) := \lim_{x \rightarrow 0} \frac{\log \Phi_{\varrho}^0(x)}{\log x}, \quad M(\varrho) := \lim_{x \rightarrow \infty} \frac{\log \Phi_{\varrho}^0(x)}{\log x}$$

exist and $-\infty < m(\varrho) \leq M(\varrho) < +\infty$. These numbers were defined by Matuszewska and Orlicz [17, 18] (see also [15] and [16, Chapter 11]). We refer to $m(\varrho)$ (resp. $M(\varrho)$) as the *lower* (resp. *upper*) *Matuszewska-Orlicz index* of ϱ . For $\varrho(x) = x^\gamma$ one has $m(\varrho) = M(\varrho) = \gamma$. Examples of functions $\varrho \in \mathbb{W}$ with $m(\varrho) < M(\varrho)$ can be found, for instance, in [1], [16, p. 93], [21, Section 2].

Fix pairwise distinct points $t_1, \dots, t_n \in \Gamma$ and functions $w_1, \dots, w_n \in \mathbb{W}$. Consider the following weight

$$(1.1) \quad w(t) := \prod_{k=1}^n w_k(|t - t_k|), \quad t \in \Gamma.$$

Each function $w_k(|t - t_k|)$ is a radial oscillating weight. The weight (1.1) is a continuous function on $\Gamma \setminus \{t_1, \dots, t_n\}$. This is a natural generalization of so-called Khvedelidze weights $w(t) = \prod_{k=1}^n |t - t_k|^{\lambda_k}$, where $\lambda_k \in \mathbb{R}$ (see, e.g., [3, Section 2.2], [9], [10]). Recently V. Kokilashvili, N. Samko, and S. Samko have proved the following (see [12, Theorem 4.3] and also [11] for similar results for maximal functions).

Theorem 1.1 (V. Kokilashvili, S. Samko, N. Samko [12, Theorem 4.3]). *Suppose Γ is a simple Carleson curve and $p : \Gamma \rightarrow (1, \infty)$ is a continuous function satisfying*

$$(1.2) \quad |p(\tau) - p(t)| \leq -A_\Gamma / \log |\tau - t| \quad \text{whenever} \quad |\tau - t| \leq 1/2,$$

where A_Γ is a positive constant depending only on Γ . Let $w_1, \dots, w_n \in \mathbb{W}$ and the weight w be given by (1.1). If

$$(1.3) \quad 0 < 1/p(t_k) + m(w_k), \quad 1/p(t_k) + M(w_k) < 1 \quad \text{for all} \quad k \in \{1, \dots, n\},$$

then the Cauchy singular integral operator S is bounded on $L^{p(\cdot)}(\Gamma, w)$.

For the weight $w(t) = \prod_{k=1}^n |t - t_k|^{\lambda_k}$, (1.3) reads as $0 < 1/p(t_k) + \lambda_k < 1$ for all $k \in \{1, \dots, n\}$. This condition is also necessary for the boundedness of S on the variable Lebesgue space $L^{p(\cdot)}(\Gamma, w)$ with the Khvedelidze weight w (see [10]).

The author have proved in [8] that for Jordan curves condition (1.3) is necessary for the boundedness of the operator S .

Theorem 1.2 ([8, Corollary 4.3]). *Suppose Γ is a rectifiable Jordan curve and $p : \Gamma \rightarrow (1, \infty)$ is a continuous function satisfying (1.2). Let $w_1, \dots, w_n \in \mathbb{W}$ and the weight w be given by (1.1). If the Cauchy singular integral operator S is bounded on $L^{p(\cdot)}(\Gamma, w)$, then Γ is a Carleson curve and (1.3) is fulfilled.*

The proof of this result given in [8] essentially uses that Γ is closed. In this paper we embark on the situation of non-closed curves. Our main result is a partial converse of Theorem 1.1. It follows from our results [6, 8] based on further development of ideas from [3, Chap. 1–3].

Theorem 1.3 (Main result). *Let Γ be a rectifiable curve homeomorphic to a line segment and $p : \Gamma \rightarrow (1, \infty)$ be a continuous function satisfying (1.2). Suppose $w_1, \dots, w_n \in \mathbb{W}$ and the weight w is given by (1.1). If the Cauchy singular integral operator S is bounded on $L^{p(\cdot)}(\Gamma, w)$, then Γ is a Carleson curve and*

$$0 \leq 1/p(t_k) + m(w_k), \quad 1/p(t_k) + M(w_k) \leq 1 \quad \text{for all} \quad k \in \{1, \dots, n\}.$$

Moreover, if there exists an $\varepsilon_0 > 0$ such that the Cauchy singular integral operator S is bounded on $L^{p(\cdot)}(\Gamma, w^{1+\varepsilon})$ for all $\varepsilon \in (-\varepsilon_0, \varepsilon_0)$, then

$$0 < 1/p(t_k) + m(w_k), \quad 1/p(t_k) + M(w_k) < 1 \quad \text{for all} \quad k \in \{1, \dots, n\}.$$

For standard Lebesgue spaces, the boundedness of the operator S on $L^p(\Gamma, w)$, $1 < p < \infty$, implies that S is also bounded on $L^p(\Gamma, w^{1+\varepsilon})$ for all ε in a sufficiently small neighborhood of zero (see [3, Theorems 2.31 and 4.15]). Hence if $1 < p < \infty$, Γ is a simple Carleson curve, $w_1, \dots, w_n \in \mathbb{W}$, and the weight w is given by (1.1), then S is bounded on the standard Lebesgue space $L^p(\Gamma, w)$, $1 < p < \infty$, if and only if

$$0 < 1/p + m(w_k), \quad 1/p + M(w_k) < 1 \quad \text{for all} \quad k \in \{1, \dots, n\}.$$

We believe that all weighted variable Lebesgue spaces have this stability property.

Conjecture 1.4. *Let Γ be a simple rectifiable curve, $p : \Gamma \rightarrow [1, \infty]$ be a measurable a.e. finite function, and $w : \Gamma \rightarrow [0, \infty]$ be a weight such that the Cauchy singular integral operator S is bounded on $L^{p(\cdot)}(\Gamma, w)$. Then there is a number $\varepsilon_0 > 0$ such that S is bounded on $L^{p(\cdot)}(\Gamma, w^{1+\varepsilon})$ for all $\varepsilon \in (-\varepsilon_0, \varepsilon_0)$.*

If this conjecture would be true, we were able to prove the complete converse of Theorem 1.1 for non-closed curves, too.

2. PROOF

In this section we formulate several results from [3, 6, 8] and show that Theorem 1.3 easily follows from them.

2.1. Muckenhoupt type condition. Suppose Γ is a simple rectifiable curve and $p : \Gamma \rightarrow (1, \infty)$ is a continuous function. Since Γ is compact, one has

$$1 < \min_{\tau \in \Gamma} p(\tau), \quad \max_{\tau \in \Gamma} p(\tau) < \infty$$

and the conjugate exponent

$$q(\tau) := p(\tau)/(p(\tau) - 1) \quad (\tau \in \Gamma)$$

is well defined and also bounded and bounded away from zero. We say that a weight $w : \Gamma \rightarrow [0, \infty]$ belongs to $A_{p(\cdot)}(\Gamma)$ if

$$\sup_{t \in \Gamma} \sup_{R > 0} \frac{1}{R} \|w \chi_{\Gamma(t,R)}\|_{p(\cdot)} \|w^{-1} \chi_{\Gamma(t,R)}\|_{q(\cdot)} < \infty.$$

If $p = \text{const} \in (1, \infty)$, then this class coincides with the well known Muckenhoupt class. From the Hölder inequality for $L^{p(\cdot)}(\Gamma)$ (see e.g. [19, Theorems 13.12 and 13.13] for Musielak-Orlicz spaces over arbitrary measure spaces and also [13, Theorem 2.1] for variable Lebesgue spaces over domains in \mathbb{R}^n) it follows that if $w \in A_{p(\cdot)}(\Gamma)$, then Γ is a Carleson curve.

Since $L^{p(\cdot)}(\Gamma, w)$ is a Banach function space in the sense of [2, Definition 1.1], the next result follows from [6, Theorem 6.1] (stated in [6] for Jordan curves, however its proof remains the same for curves homeomorphic to line segments, see also [7, Theorem 3.2]).

Theorem 2.1. *Let Γ be a simple rectifiable curve and let $p : \Gamma \rightarrow (1, \infty)$ be a continuous function. If $w : \Gamma \rightarrow [0, \infty]$ is an arbitrary weight such that the operator S is bounded on $L^{p(\cdot)}(\Gamma, w)$, then $w \in A_{p(\cdot)}(\Gamma)$.*

If $p = \text{const} \in (1, \infty)$, then $w \in A_p(\Gamma)$ is also sufficient for the boundedness of S on the weighted Lebesgue space $L^p(\Gamma, w)$ (see e.g. [3, Theorem 4.15]).

2.2. Submultiplicative functions. Following [3, Section 1.4], we say a function $\Phi : (0, \infty) \rightarrow (0, \infty]$ is *regular* if it is bounded in an open neighborhood of 1. A function $\Phi : (0, \infty) \rightarrow (0, \infty]$ is said to be *submultiplicative* if

$$\Phi(xy) \leq \Phi(x)\Phi(y) \quad \text{for all } x, y \in (0, \infty).$$

It is easy to show that if Φ is regular and submultiplicative, then Φ is bounded away from zero in some open neighborhood of 1. Moreover, in this case $\Phi(x)$ is finite for all $x \in (0, \infty)$. Given a regular and submultiplicative function $\Phi : (0, \infty) \rightarrow (0, \infty)$, one defines

$$\alpha(\Phi) := \sup_{x \in (0,1)} \frac{\log \Phi(x)}{\log x}, \quad \beta(\Phi) := \inf_{x \in (1,\infty)} \frac{\log \Phi(x)}{\log x}.$$

Clearly, $-\infty < \alpha(\Phi)$ and $\beta(\Phi) < \infty$.

Theorem 2.2 (see [3, Theorem 1.13] or [14, Chap. 2, Theorem 1.3]). *If a function $\Phi : (0, \infty) \rightarrow (0, \infty)$ is regular and submultiplicative, then*

$$\alpha(\Phi) = \lim_{x \rightarrow 0} \frac{\log \Phi(x)}{\log x}, \quad \beta(\Phi) = \lim_{x \rightarrow \infty} \frac{\log \Phi(x)}{\log x}$$

and $-\infty < \alpha(\Phi) \leq \beta(\Phi) < +\infty$.

The quantities $\alpha(\Phi)$ and $\beta(\Phi)$ are called the *lower* and *upper indices of the regular and submultiplicative function* Φ , respectively.

2.3. Indices of powerlikeness. Fix $t \in \Gamma$ and put $d_t := \max_{\tau \in \Gamma} |\tau - t|$. Suppose $w : \Gamma \rightarrow [0, \infty]$ is a weight such that $\log w \in L^1(\Gamma(t, R))$ for every $R \in (0, d_t]$. Put

$$H_{w,t}(R_1, R_2) := \frac{\exp\left(\frac{1}{|\Gamma(t, R_1)|} \int_{\Gamma(t, R_1)} \log w(\tau) |d\tau|\right)}{\exp\left(\frac{1}{|\Gamma(t, R_2)|} \int_{\Gamma(t, R_2)} \log w(\tau) |d\tau|\right)}, \quad R_1, R_2 \in (0, d_t].$$

Consider the function

$$(V_t^0 w)(x) := \limsup_{R \rightarrow 0} H_{w,t}(xR, R), \quad x \in (0, \infty).$$

Combining Lemmas 4.8–4.9 and Theorem 5.9 of [6] with Theorem 3.4, Lemma 3.5 of [3], we arrive at the following.

Theorem 2.3. *Let Γ be a simple rectifiable curve, $p : \Gamma \rightarrow (1, \infty)$ be a continuous function satisfying (1.2), and $w : \Gamma \rightarrow [0, \infty]$ be a weight such that $w \in A_{p(\cdot)}(\Gamma)$. Then, for every $t \in \Gamma$, the function $V_t^0 w$ is regular and submultiplicative and*

$$0 \leq 1/p(t) + \alpha(V_t^0 w), \quad 1/p(t) + \beta(V_t^0 w) \leq 1.$$

The numbers $\alpha(V_t^0 w)$ and $\beta(V_t^0 w)$ are called the *lower* and *upper indices of powerlikeness* of w at $t \in \Gamma$, respectively (see [3, Chap. 3]). This terminology can be explained by the simple fact that for the power weight $w(\tau) := |\tau - t|^\lambda$ its indices of powerlikeness coincide and are equal to λ .

2.4. Matuszewska-Orlicz indices as indices of powerlikeness. If $\varrho \in \mathbb{W}$, then Φ_ϱ^0 is a regular and submultiplicative function and its indices are nothing else but the Matuszewska-Orlicz indices $m(\varrho)$ and $M(\varrho)$. The next result shows that for radial oscillating weights indices of powerlikeness and Matuszewska-Orlicz indices coincide.

Theorem 2.4 (see [8, Theorem 2.8]). *Suppose Γ is a simple Carleson curve. If $w_1, \dots, w_n \in \mathbb{W}$ and $w(\tau) = \prod_{k=1}^n w_k(|\tau - t_k|)$, then for every $t \in \Gamma$ the function $V_t^0 w$ is regular and submultiplicative and*

$$\begin{aligned} \alpha(V_{t_k}^0 w) &= m(w_k), & \beta(V_{t_k}^0 w) &= M(w_k) & \text{for } k \in \{1, \dots, n\}, \\ \alpha(V_t^0 w) &= 0, & \beta(V_t^0 w) &= 0 & \text{for } t \in \Gamma \setminus \{t_1, \dots, t_n\}. \end{aligned}$$

Note that in [8], Theorem 2.4 is proved for Jordan curves. But the proof does not use the assumption that Γ is closed. It works also for non-closed curves considered in this paper.

2.5. Proof of Theorem 1.3.

Proof. Suppose S is bounded on $L^{p(\cdot)}(\Gamma, w)$. From Theorem 2.1 it follows that $w \in A_{p(\cdot)}(\Gamma)$. By Hölder's inequality this implies that Γ is a Carleson curve. Fix an arbitrary $t \in \Gamma$. Then, in view of Theorems 2.2 and 2.3 the function $V_t^0 w$ is regular

and submultiplicative, so its indices are well defined and satisfy $0 \leq 1/p(t) + \alpha(V_t^0 w)$ and $1/p(t) + \beta(V_t w) \leq 1$. From these inequalities and Theorem 2.4 it follows that

$$(2.1) \quad 0 \leq 1/p(t_k) + m(w_k), \quad 1/p(t_k) + M(w_k) \leq 1$$

for all $k \in \{1, \dots, n\}$.

If S is bounded on all spaces $L^{p(\cdot)}(\Gamma, w^{1+\varepsilon})$ for all ε in a neighborhood of zero, then as before

$$0 \leq 1/p(t_k) + m(w_k^{1+\varepsilon}), \quad 1/p(t_k) + M(w_k^{1+\varepsilon}) \leq 1$$

for every $k \in \{1, \dots, n\}$. It is easy to see that $m(w_k^{1+\varepsilon}) = (1 + \varepsilon)m(w_k)$ and $M(w_k^{1+\varepsilon}) = (1 + \varepsilon)M(w_k)$. Therefore

$$0 \leq 1/p(t_k) + (1 + \varepsilon)m(w_k), \quad 1/p(t_k) + (1 + \varepsilon)M(w_k) \leq 1$$

for all ε in a neighborhood of zero and for all $k \in \{1, \dots, n\}$. These inequalities immediately imply that $0 < 1/p(t_k) + m(w_k)$ and $1/p(t_k) + M(w_k) < 1$ for all k . \square

Remark 2.5. The presented proof involves the notion of indices of powerlikeness, which were invented to treat general Muckenhoupt weights (see [3]). Weights considered in the present paper are continuous except for a finite number of points. So, it would be rather interesting to find a direct proof of the fact that $w \in A_{p(\cdot)}(\Gamma)$ implies (2.1), which does not involve the indices of powerlikeness $\alpha(V_t^0 w)$ and $\beta(V_t^0 w)$.

2.6. Final remarks. In connection with Conjecture 1.4, we would like to note that for standard Lebesgue spaces $L^p(\Gamma, w)$ there are two different proofs of the stability of the boundedness of S on $L^p(\Gamma, w^{1+\varepsilon})$ for small ε . Simonenko's proof [22] is based on the stability of the Fredholm property of some singular integral operators related to the Riemann boundary value problem. Another proof is based on the self-improving property of Muckenhoupt weights (see e.g. [3, Theorem 2.31]). One may ask whether does $w \in A_{p(\cdot)}(\Gamma)$ imply $w^{1+\varepsilon} \in A_{p(\cdot)}(\Gamma)$ for all $\varepsilon \in (-\varepsilon_0, \varepsilon_0)$ with some fixed $\varepsilon > 0$? The positive answer would give a proof of the complete converse of (1.3). The author does not know any stability result for the boundedness of S or a self-improving property for $w \in A_{p(\cdot)}(\Gamma)$.

After this paper had been submitted, P. Hästö and L. Diening [5] have found a necessary and sufficient condition for the boundedness of the classical Hardy-Littlewood maximal function on weighted variable Lebesgue spaces in the setting of \mathbb{R}^n . Note that they write a weight as a measure (outside of $|\cdot|^{p(\tau)}$). Their condition is another generalization of the classical Muckenhoupt condition. In the setting of Carleson curves (and the weight written inside of $|\cdot|^{p(\tau)}$), the Hästö-Diening condition takes the form

$$(2.2) \quad \sup_{t \in \Gamma} \sup_{R > 0} \left(\frac{1}{R^{p_{\Gamma}(t, R)}} \int_{\Gamma(t, R)} w(\tau)^{p(\tau)} |d\tau| \right) \|w(\cdot)^{-p(\cdot)} \chi_{\Gamma(t, R)}(\cdot)\|_{q(\cdot)/p(\cdot)} < \infty,$$

where

$$p_{\Gamma}(t, R) := \left(\frac{1}{|\Gamma(t, R)|} \int_{\Gamma(t, R)} \frac{1}{p(\tau)} |d\tau| \right)^{-1}.$$

Let $HD_{p(\cdot)}(\Gamma)$ denote the class of weights $w : \Gamma \rightarrow [0, \infty]$ satisfying (2.2). Following the arguments contained in [5, Remark 3.10], one can show that

$$A_{L^{p(\cdot)}}(\Gamma) \supset HD_{p(\cdot)}(\Gamma)$$

whenever $p : \Gamma \rightarrow (1, \infty)$ satisfies the Dini-Lipschitz condition (1.2). We conjecture that the Hästö-Diening characterization remains true also for the operator S in the setting of Carleson curves.

Conjecture 2.6. *Let Γ be a simple Carleson curve, $w : \Gamma \rightarrow [0, \infty]$ be a weight, and $p : \Gamma \rightarrow (1, \infty)$ be a continuous function satisfying the Dini-Lipschitz condition (1.2). The operator S is bounded on $L^{p(\cdot)}(\Gamma, w)$ if and only if $w \in HD_{p(\cdot)}(\Gamma)$.*

Acknowledgement. The author would like to thank the anonymous referee for several useful remarks.

REFERENCES

- [1] V. D. Aslanov and Yu. I. Karlovich, *One-sided invertibility of functional operators in reflexive Orlicz spaces*. Akad. Nauk Azerbaidzhan. SSR Dokl. **45** (1989), no. 11–12, 3–7 (in Russian).
- [2] C. Bennett and R. Sharpley, *Interpolation of Operators*. Academic Press, Boston, 1988.
- [3] A. Böttcher and Yu. I. Karlovich, *Carleson Curves, Muckenhoupt Weights, and Toeplitz Operators*. Birkhäuser, Basel, 1997.
- [4] G. David, *Opérateurs intégraux singuliers sur certaines courbes du plan complexe*. Ann. Sci. École Norm. Super. **17** (1984), 157–189.
- [5] P. Hästö and L. Diening, Muckenhoupt weights in variable exponent spaces, (December 15, 2008), preprint is available at http://www.helsinki.fi/~hasto/pp/p75_submit.pdf.
- [6] A. Yu. Karlovich, *Fredholmness of singular integral operators with piecewise continuous coefficients on weighted Banach function spaces*. J. Integr. Equat. Appl. **15** (2003), 263–320.
- [7] A. Yu. Karlovich, *Singular integral operators with piecewise continuous coefficients in reflexive rearrangement-invariant spaces*. Integral Equations Operator Theory **32** (1998), 436–481.
- [8] A. Yu. Karlovich, *Singular integral operators on variable Lebesgue spaces with radial oscillating weights*. Proceedings of IWOTA 2007, Operator Theory: Advances and Applications, Birkhäuser, to appear. Preprint is available at <http://arxiv.org/abs/0708.0778>.
- [9] B. V. Khvedelidze, *Linear discontinuous boundary problems in the theory of functions, singular integral equations and some of their applications*. Trudy Tbiliss. Mat. Inst. Razmadze **23** (1956), 3–158 (in Russian).
- [10] V. Kokilashvili, V. Paatashvili, and S. Samko, *Boundedness in Lebesgue spaces with variable exponent of the Cauchy singular operator on Carleson curves*. In: “Modern Operator Theory and Applications. The Igor Borisovich Simonenko Anniversary Volume”. Operator Theory: Advances and Applications **170** (2006), 167–186.
- [11] V. Kokilashvili, N. Samko, and S. Samko, *The maximal operator in weighted variable spaces $L^{p(\cdot)}$* . J. Func. Spaces Appl. **5** (2007), 299–317.
- [12] V. Kokilashvili, N. Samko, and S. Samko, *Singular operators in variable spaces $L^{p(\cdot)}(\Omega, \rho)$ with oscillating weights*. Math. Nachr. **280** (2007), 1145–1156.
- [13] O. Kováčik and J. Rákosník, *On spaces $L^{p(x)}$ and $W^{k,p(x)}$* . Czechoslovak Math. J. **41(116)** (1991), 592–618.
- [14] S. G. Krein, Ju. I. Petunin, and E. M. Semenov, *Interpolation of Linear Operators*. AMS Translations of Mathematical Monographs **54**, Providence, RI, 1982.
- [15] L. Maligranda, *Indices and interpolation*. Dissert. Math. **234** (1985), 1–49.
- [16] L. Maligranda, *Orlicz Spaces and Interpolation*. Sem. Math. 5, Dep. Mat., Universidade Estadual de Campinas, Campinas SP, Brazil, 1989.
- [17] W. Matuszewska and W. Orlicz, *On certain properties of φ -functions*. Bull. Acad. Polon. Sci. Ser. Sci. Math. Astronom. Phys. **8** (1960), 439–443. Reprinted in: W. Orlicz, *Collected Papers*, PWN, Warszawa, 1988, 1112–1116.
- [18] W. Matuszewska and W. Orlicz, *On some classes of functions with regard to their orders of growth*. Studia Math. **26** (1965), 11–24. Reprinted in: W. Orlicz, *Collected Papers*, PWN, Warszawa, 1988, 1217–1230.
- [19] J. Musielak, *Orlicz Spaces and Modular Spaces*. Lecture Notes in Mathematics **1034**. Springer-Verlag, Berlin, 1983.
- [20] H. Nakano, *Modulated Semi-Ordered Linear Spaces*. Maruzen Co., Ltd., Tokyo, 1950.

- [21] N. Samko, *Singular integral operators in weighted spaces of continuous functions with oscillating continuity moduli and oscillating weights*. In: “The Extended Field of Operator Theory”. Operator Theory: Advances and Applications **171** (2006), 323–347.
- [22] I. B. Simonenko, *Stability of the weight properties of functions with respect to a singular Cauchy integral*, Math Notes **33** (1983), 208–212.

DEPARTAMENTO DE MATEMÁTICA, FACULDADE DE CIÊNCIAS E TECNOLOGIA, UNIVERSIDADE
NOVA DE LISBOA, QUINTA DA TORRE, 2829-516 CAPARICA, PORTUGAL
E-mail address: oyk@fct.unl.pt