Fourier-Mukai Partners of a K3 Surface and the Cusps of its Kahler Moduli

Shouhei Ma

Graduate School of Mathematical Sciences, University of Tokyo 3-8-1 Komaba Meguro-ku, Tokyo 153-8914, Japan E-mail: sma@ms.u-tokyo.ac.jp

Abstract

Using lattice theory, we establish a one-to-one correspondence between the set of Fourier-Mukai partners of a projective K3 surface and the set of 0-dimensional standard cusps of its Kahler moduli. We also study the relation between twisted Fourier-Mukai partners and general 0-dimensional cusps, and the relation between Fourier-Mukai partners with elliptic fibrations and certain 1-dimensional cusps.

1 Introduction

The excellent work of Mukai ([9], [10]) and Orlov ([12]) opened a way to study derived equivalence for K3 surfaces via their periods. One of their conclusions is that two projective K3 surfaces are derived equivalent if and only if there exists an isometry between their Mukai lattices preserving the periods. This theorem can be viewed as a generalization of the global Torelli theorem for K3surfaces. Let FM(S) be the set of isomorphism classes of Fourier-Mukai (FM) partners of a projective K3 surface S, i.e., K3 surfaces derived equivalent to S. As an application of Mukai-Orlov's theorem, Hosono-Lian-Oguiso-Yau ([5]) gave a counting formula for #FM(S).

In this paper, we construct a bijection between $\operatorname{FM}(S)$ and the set of embeddings of the hyperbolic plane U into the lattice $\widetilde{NS}(S) := NS(S) \oplus U$ up to the action of a certain finite-index subgroup $\Gamma_S \subset O(\widetilde{NS}(S))$. Then, by considering the isotropic vector $(0,1) \in U$, we obtain 0-dimensional standard cusps of the modular variety $\Gamma_S^+ \setminus \Omega_{\widetilde{NS}(S)}^+$ associated with Γ_S^+ and $\widetilde{NS}(S)$ ([1], [13]). Let $\Gamma_S^+ \setminus I^1(\widetilde{NS}(S))$ be the set of 0-dimensional standard cusps of $\Gamma_S^+ \setminus \Omega_{\widetilde{NS}(S)}^+$. We shall prove the following.

Theorem 1.1 (Theorem 3.6). There exists a bijective map

$$\mu_0: \mathrm{FM}(S) \to \Gamma_S^+ \backslash I^1(NS(S)).$$

In particular,

$$\# FM(S) = \# \{ \text{ 0-dimensional standard cusps of } \Gamma_S^+ \setminus \Omega_{\widetilde{NS}(S)}^+ \}$$

Of course, not every 0-dimensional cusp of $\Gamma_S^+ \setminus \Omega_{\widetilde{NS}(S)}^+$ is standard. It turns out that the existence of non-standard cusps corresponds to the existence of *twisted* FM-partners of S. Let $\mathrm{FM}^d(S)$ be the set of isomorphism classes of twisted FM-partners (S', α') of S with $\mathrm{ord}(\alpha') = d$, and let $\Gamma_S^+ \setminus I^d(\widetilde{NS}(S))$ be the set of 0-dimensional cusps [l] of $\Gamma_S^+ \setminus \Omega_{\widetilde{NS}(S)}^+$ with $\mathrm{div}(l) = d$. Introducing certain quotient sets $\mathcal{FM}^d(S)$, $r(\Gamma_S^+) \setminus I^d(A_{\widetilde{NS}(S)})$ of $\mathrm{FM}^d(S)$, $\Gamma_S^+ \setminus I^d(\widetilde{NS}(S))$ respectively (see Sect.3.2 for the definitions), and using Huybrechts-Stellari's solution of Căldăraru conjecture ([8]), we shall prove the following.

Theorem 1.2 (Theorem 3.10). There exist a map

$$\nu_0: \Gamma_S^+ \setminus I^d(NS(S)) \longrightarrow \mathrm{FM}^d(S)$$

and a bijective map

$$\xi_0: r(\Gamma_S^+) \setminus I^d(A_{\widetilde{NS}(S)}) \xrightarrow{\simeq} \mathcal{FM}^d(S)$$

which fit in the following commutative diagram:

$$\begin{array}{ccc} \Gamma_{S}^{+} \backslash I^{d}(\widetilde{NS}(S)) & \stackrel{\nu_{0}}{\longrightarrow} & \mathrm{FM}^{d}(S) \\ & p \\ & & \downarrow \pi \\ r(\Gamma_{S}^{+}) \backslash I^{d}(A_{\widetilde{NS}(S)}) & \stackrel{-}{\longrightarrow} & \mathcal{FM}^{d}(S). \end{array}$$

Via Theorem 1.1 and 1.2, we can obtain informations about the abstract set $\mathrm{FM}^d(S)$ by studying the 0-dimensional cusps of the modular variety $\Gamma^+_S \setminus \Omega^+_{\widetilde{NS}(S)}$.

Besides 0-dimensional cusps, $\Gamma_S^+ \setminus \Omega_{\widetilde{NS}(S)}^+$ has also 1-dimensional cusps. Let $\operatorname{FM}_{ell}(S)$ be the set of isomorphism classes of pairs (S', L'), where $S' \in \operatorname{FM}(S)$ and $L' \in \operatorname{Pic}(S')$ is the line bundle associated to a smooth elliptic curve on S'. In the same manner as in the case of 0-dimensional standard cusps, we relate $\operatorname{FM}_{ell}(S)$ to the set $\Gamma_S^+ \setminus I_2^{st}(\widetilde{NS}(S))$ of those 1-dimensional cusps of $\Gamma_S^+ \setminus \Omega_{\widetilde{NS}(S)}^+$ whose closure contains a 0-dimensional standard cusp. The map μ_1 from $\operatorname{FM}_{ell}(S)$ to the set of such 1-dimensional cusps is surjective but not injective in general. For example, for generic K3 surface S with NS(S) = U(r) (r > 2), we will calculate μ_1 explicitly to conclude that μ_1 is far from being injective (Example 4.13). On the other hand, if det NS(S) is square-free, then μ_1 is bijective (Corollary 4.11). It turns out that $\operatorname{FM}_{ell}(S)$ carries informations about the compactifications of 1-dimensional cusps which belong to $\Gamma_S^+ \setminus I_2^{st}(\widetilde{NS}(S))$. We shall observe that if two elliptic K3 surfaces (S, L) and (S', L') give the same 1-dimensional cusp, there exists a coherent sheaf on $S \times S'$ via which L

gives rise to L'. (For precise statement see Proposition 4.12 and the remark after it.)

According to Bridgeland's results ([2], [3]), $\Gamma_S^+ \setminus \Omega_{\widetilde{NS}(S)}^+$ is biholomorphic to a natural quotient of the space of certain stability conditions on $D^b(S)$, after removing some divisors on $\Omega_{\widetilde{NS}(S)}^+$. That is, we have a canonical isomorphism (Proposition 5.1)

$$Aut^{\dagger}(D^{b}(S))\backslash Stab^{\dagger}(S)/\widetilde{GL}_{2}^{+}(\mathbb{R}) \simeq \Gamma_{S}^{+} \backslash \Big(\Omega_{\widetilde{NS}(S)}^{+} - \bigcup_{\delta \in \Delta(S)} \delta^{\perp} \Big).$$

Hence Theorem 1.1 asserts that $\operatorname{FM}(S)$ is identified with the 0-dimensional standard cusps of the Baily-Borel compactification of $\operatorname{Aut}^{\dagger}(D^b(S)) \setminus \operatorname{Stab}^{\dagger}(S)/\widetilde{GL}_2^+(\mathbb{R})$. However, the lattice theoretic approach employed in this paper does not explain any intrinsic reason for the correspondence between (twisted) FM-partners and 0-dimensional cusps. It may be interesting to understand the correspondence in more derived-categorical way.

This paper is organized as follows. In Sect.2.1, we recall some facts about even lattices and their discriminant forms, following Nikulin ([11]). Sect.2.2 is devoted to the study of isotropic elements of an even lattice. In Sect.2.3, we recall the Baily-Borel compactification of an arithmetic quotient of a type IV symmetric domain, following [1], [13], [15]. In Sect.3.1, we prove Theorem 1.1. In Sect.3.2, we prove Theorem 1.2. In Sect.4.1, we define $\operatorname{FM}_{ell}(S)$ and relate it to primitive embeddings of $U \oplus \mathbb{Z}l$ into $\widetilde{NS}(S)$. In Sect.4.2, we study the relation between $\operatorname{FM}_{ell}(S)$ and certain 1-dimensional cusps of $\Gamma_S^+ \setminus \Omega_{\widetilde{NS}(S)}^+$. In Sect.5, we observe the action of Γ_S^+ on $\Omega_{\widetilde{NS}(S)}^+$, and we also describe $\Gamma_S^+ \setminus \Omega_{\widetilde{NS}(S)}^+$ in terms of the space of stability conditions.

Notation 1.3. By an *even lattice*, we mean a free \mathbb{Z} -module L of finite rank equipped with a non-degenerate symmetric bilinear form $(,): L \times L \to \mathbb{Z}$ satisfying $(l, l) \in 2\mathbb{Z}$ for all $l \in L$. Denote by $\operatorname{rk}(L)$ and $\operatorname{sign}(L)$ the rank and the signature of L, respectively. For a lattice L and a field \mathbb{K} , $L_{\mathbb{K}}$ denotes the K-vector space $L \otimes K$. For two lattices L and M, $L \oplus M$ is the lattice defined as the orthogonal direct sum of L and M, while L + M denotes the direct sum of the \mathbb{Z} -modules underlying L and M. The projection from $L \oplus M$ to L is denoted by $pr_L : L \oplus M \to L$. The group of isometries of L is denoted by O(L). For an element $l \in L$, we define the positive integer $\operatorname{div}(l)$ to be the generator of the ideal $(x,L) \subseteq \mathbb{Z}$. A sublattice $M \subset L$ is called *primitive* if L/M is a free \mathbb{Z} -module. For (possibly degenerate) lattices L and M, we denote by $\operatorname{Emb}(L, M)$ the set of primitive embeddings of L into M. A sublattice $M \subseteq L$ is called *isotropic* if (x, y) = 0 for all $x, y \in M$. We denote by $I_r(L)$ the set of primitive isotropic sublattices of L of rk = r. A non-zero element $l \in L$ is called isotropic (resp. primitive) if $\mathbb{Z}l$ is isotropic (resp. primitive). We denote $I^{d}(L) := \{l \in L | l \text{ is primitive }, (l, l) = 0, \operatorname{div}(l) = d \}.$

By a K3 surface, we mean a projective K3 surface over \mathbb{C} . For a K3 surface S, we denote by NS(S) (resp. T(S)) the Neron-Severi (resp. transcendental)

lattice of S. Set $\widetilde{NS}(S) := H^0(S, \mathbb{Z}) + NS(S) + H^4(S, \mathbb{Z}), \Lambda_{K3} := U^3 \oplus E_8^2$, and $\widetilde{\Lambda}_{K3} := U^4 \oplus E_8^2$.

2 Preliminaries from lattice theory

2.1 Even lattice and its discriminant form

For an even lattice L, we can associate a finite Abelian group $A_L := L^{\vee}/L$ equipped with a quadratic form $q_L : A_L \to \mathbb{Q}/2\mathbb{Z}$, $q_L(x+L) = (x,x) + 2\mathbb{Z}$. The corresponding bilinear form on A_L is $b_L : A_L \times A_L \to \mathbb{Q}/\mathbb{Z}$, $b_L(x+L,y+L) =$ $(x,y) + \mathbb{Z}$. We call (A_L,q_L) or (A_L,b_L) the discriminant form of L. There is a natural homomorphism $r_L : O(L) \to O(A_L,q_L)$, whose kernel is denoted by $O(L)_0 \subset O(L)$. We often write just A_L or (A_L,q) instead of (A_L,q_L) , and write r instead of r_L . The following well-known facts due to Nikulin will be used frequently in this paper :

Proposition 2.1 ([11]). Let M be a primitive sublattice of an even unimodular lattice L with the orthogonal complement M^{\perp} . Then

(1) There is an isometry $\lambda_L : (A_M, q_M) \xrightarrow{\simeq} (A_{M^{\perp}}, -q_{M^{\perp}}).$

(2) For isometries $\gamma_M \in O(M)$ and $\gamma_{M^{\perp}} \in O(M^{\perp})$, there is an isometry $\gamma_L \in O(L)$ such that $\gamma_L|_{M \oplus M^{\perp}} = \gamma_M \oplus \gamma_{M^{\perp}}$ if and only if γ_M and $\gamma_{M^{\perp}}$ are compatible on the discriminant group, i.e, $\lambda_L \circ r_M(\gamma_M) = r_{M^{\perp}}(\gamma_{M^{\perp}}) \circ \lambda_L$.

Two even lattices L and M are said to be *isogenus* if $L \otimes \mathbb{Z}_p \simeq M \otimes \mathbb{Z}_p$ for every prime number p and $\operatorname{sign}(L) = \operatorname{sign}(M)$. By [11], this is equivalent to the condition that $(A_L, q_L) \simeq (A_M, q_M)$ and $\operatorname{sign}(L) = \operatorname{sign}(M)$. The set of isometry classes of lattices isogenus to L is denoted by $\mathcal{G}(L)$.

Proposition 2.2 ([11]). Let $l(A_L)$ be the minimal number of the generators of the Abelian group A_L . If L is indefinite and $\operatorname{rk}(L) \geq l(A_L) + 2$, then $\mathcal{G}(L) = \{L\}$ and the homomorphism r_L is surjective.

By Proposition 2.2, L and M are isogenus if and only if $L \oplus U \simeq M \oplus U$, where U is the even indefinite unimodular lattice $\mathbb{Z}e + \mathbb{Z}f$, (e, f) = 1, (e, e) = (f, f) = 0. Note that if an even unimodular lattice L is embedded in another even lattice M, L must be an orthogonal summand of M.

Lemma 2.3. Let L and M be even lattices. If there is an isometry $\varphi : L \oplus U \simeq M \oplus U$ with $\varphi(f) = f$, then the composition $pr_M \circ (\varphi|_L) : L \hookrightarrow \varphi(f)^{\perp} = M \oplus \mathbb{Z}f \twoheadrightarrow M$ is an isometry.

Proof. Since $L \oplus \mathbb{Z}f = f^{\perp} \simeq \varphi(f)^{\perp} = M \oplus \mathbb{Z}f$, we can write $\varphi(l) = m + \alpha f$ for each $l \in L$, where $m \in M$ and $\alpha \in \mathbb{Z}$. Then $(l, l) = (\varphi(l), \varphi(l)) = (m, m)$. Thus the correspondence $\varphi(l) \mapsto m$ gives an embedding of lattice $L \hookrightarrow M$. By the inclusions $L \subseteq M \subseteq M^{\vee} \subseteq L^{\vee}$, we have $|A_M| = |M/M^{\vee}| \leq |L/L^{\vee}| = |A_L|$. Here the equality holds if and only if $L \simeq M$ by $pr_M \circ (\varphi|_L)$. Since $A_L \simeq$ $A_{L \oplus U} \simeq A_{M \oplus U} \simeq A_M$, we get $|A_M| = |A_L|$, so that $L \simeq M$.

2.2 Primitive isotropic vectors

Let L be an even lattice possessing a primitive isotropic vector l. We shall study some properties of L related to l.

Proposition 2.4. Let $l \in I^1(L)$ be an arbitrary element. Then there is a canonical bijection

$$O(L) \setminus I^1(L) \simeq \mathcal{G}(l^\perp / \mathbb{Z}l).$$

Proof. For $l \in I^1(L)$, there exists $m' \in L$ with (l,m') = 1. Setting $m := m' - \frac{(m',m')}{2}l$, we have (l,l) = (m,m) = 0 and (l,m) = 1. Hence there is an embedding $\varphi: U \hookrightarrow L$ with $\varphi(f) = l$. Since the lattice U is unimodular, we have

$$L = \varphi(U) \oplus \varphi(U)^{\perp} \simeq \varphi(U) \oplus (l^{\perp}/\mathbb{Z}l)$$

so that $A_L \simeq A_{l^{\perp}/\mathbb{Z}l}$ for each $l \in I^1(L)$. In particular, $\mathcal{G}(l^{\perp}/\mathbb{Z}l)$ is independent of the choices of $l \in I^1(L)$.

We define the map $\mu : O(L) \setminus I^1(L) \to \mathcal{G}(l^{\perp}/\mathbb{Z}l)$ by $\mu(l') := (l')^{\perp}/\mathbb{Z}l'$. If there is an isometry $\gamma : l_1^{\perp}/\mathbb{Z}l_1 \simeq l_2^{\perp}/\mathbb{Z}l_2$, we can extend γ to the isometry $\tilde{\gamma} \in O(L)$ with $\tilde{\gamma}(l_1) = l_2$. Therefore l_1 and l_2 are O(L)-equivalent so that μ is injective. Given a lattice $K \in \mathcal{G}(l^{\perp}/\mathbb{Z}l)$, we have an isometry $\Psi : K \oplus U \xrightarrow{\simeq} L$ by Proposition 2.2, which gives $\Psi(f) \in I^1(L)$. Since $\mu(\Psi(f)) = K$, μ is surjective.

Proposition 2.5. For a primitive isotropic vector $l \in I^d(L)$, we have the equality $d^2 \cdot \#(D_{l^{\perp}/\mathbb{Z}l}) = \#(D_L)$.

Proof. The free \mathbb{Z} -module $\widetilde{L} := \langle L, \frac{l}{d} \rangle \subset L^{\vee}$ is an even overlattice of L. Since $\frac{l}{d} \in I^1(\widetilde{L})$, we have $\widetilde{L} \simeq U \oplus (l^{\perp}/\mathbb{Z}l)$ so that $D_{\widetilde{L}} \simeq D_{l^{\perp}/\mathbb{Z}l}$. If we set $H := \widetilde{L}/L \subset D_L$, then H is an isotropic cyclic group of order d satisfying $D_{\widetilde{L}} \simeq H^{\perp}/H$. Hence $\#(D_L) = d^2 \cdot \#(D_{\widetilde{L}}) = d^2 \cdot \#(D_{l^{\perp}/\mathbb{Z}l})$.

Corollary 2.6. If det(L) is square-free, every primitive isotropic vector $l \in L$ satisfies div(l) = 1.

Let $l \in I^1(L)$. Choose an element $m \in L$ so that (m,m) = 0 and (l,m) = 1. Then we define $L_0 := \mathbb{Z}l + \mathbb{Z}m \simeq U$ and $L_1 := L_0^{\perp} \simeq l^{\perp}/\mathbb{Z}l$. We can describe the group

$$O(L)^{l} := \{ \gamma \in O(L) \, | \, \gamma(l) = l \}.$$

in terms of the free Abelian group L_1 and the group $O(L_1)$ as follows. For an element $v \in L_1$, define the isometry $T_v \in O(L)^l$ by

$$T_{v}(l) = l,$$

$$T_{v}(m) = m + v - \frac{(v, v)}{2}l,$$

$$T_{v}(v') = v' - (v', v)l, \quad v' \in L_{1}.$$
(1)

Proposition 2.7. For $l \in I^1(L)$, we regard $L_1 \subset O(L)^l$ by the correspondence $v \mapsto T_v$ and regard $O(L_1) \subset O(L)^l$ by the correspondence $g \mapsto \operatorname{id}_{L_0} \oplus g$. Then

$$O(L)^l \simeq O(L_1) \ltimes L_1.$$

Moreover, the subgroup $L_1 \subset O(L)^l$ acts trivially on $D_L \simeq D_{L_1}$.

Proof. Take an isometry $\gamma \in O(L)^l$. Since $\gamma(l^{\perp}) = l^{\perp}$ and $l^{\perp} = \mathbb{Z}l \oplus L_1$, γ induces the isometry $pr_{L_1} \circ (\gamma|_{L_1}) \in O(L_1)$. The correspondence $\gamma \mapsto pr_{L_1} \circ (\gamma|_{L_1})$ induces the homomorphism $\pi : O(L)^l \to O(L_1)$. Then π is surjective because $\pi(\operatorname{id}_{L_0} \oplus g) = g$ for $g \in O(L_1)$. The correspondence $g \mapsto \operatorname{id}_{L_0} \oplus g$ is a section of π .

We prove that $\operatorname{Ker}(\pi) = L_1 \subset O(L)^l$. The inclusion $\operatorname{Ker}(\pi) \supset L_1$ is apparent. For $\gamma \in \operatorname{Ker}(\pi)$, we can write $\gamma(m) = m + v + \alpha l$ for some integer $\alpha \in \mathbb{Z}$ and some vector $v \in L_1$. Since $(\gamma(m), \gamma(m)) = (m, m) = 0$, we can determine α as $\alpha = -\frac{(v,v)}{2}$. On the other hand, if we take $v' \in L_1$, we can write $\gamma(v') = v' + \beta(v')l$ for some integer $\beta(v') \in \mathbb{Z}$. Since $(\gamma(v'), \gamma(m)) = (v', m) = 0$, we have $\beta(v') = -(v, v')$. Hence we have $\gamma = T_v$. The claim that $r_L(L_1) = \{\operatorname{id}\}$ is obvious.

2.3 The Baily-Borel compactification

In this subsection, we recall the Baily-Borel compactification of an arithmetic quotient of a type IV symmetric domain ([1], [13]), following [15]. Although the Baily-Borel compactifications are defined for general arithmetic groups, we restrict ourselves to finite-index subgroups of $O(L)^+$ containing $\{\pm id\}$.

Let L be an even lattice of $sign(L) = (2, b^{-})$. Set

$$\Omega_L := \{ \mathbb{C}\omega \in \mathbb{P}(L_{\mathbb{C}}) \, | \, (\omega, \omega) = 0, \, (\omega, \bar{\omega}) > 0 \},\$$

which has two connected components Ω_L^+ and Ω_L^- exchanged by the complex conjugation on Ω_L . Denote by $O(L)^+ \subset O(L)$ the subgroup of index at most 2 which consists of isometries preserving Ω_L^+ . By associating the oriented twoplane $\mathbb{R}\text{Re}\omega \oplus \mathbb{R}\text{Im}\omega$ to $\mathbb{C}\omega \in \Omega_L$, we have the isomorphism

$$\Omega_L \simeq \left\{ \left(E, \tau \right) \right| \qquad E \subset L_{\mathbb{R}} \text{ is a positive-definite two-plane,}$$

$$\tau \text{ is an orientation of } E \right\}.$$
(2)

The right hand side of (2) is an open subset of the oriented Grassmannian. Choosing a connected component of Ω_L , we can attach the orientation to every positive-definite two-plane in $L_{\mathbb{R}}$. A choice of a component of Ω_L is sometimes called an *orientation* of L.

Regarding Ω_L^+ as an open subset of $Q := \{\mathbb{C}\omega \in \mathbb{P}(L_{\mathbb{C}}) \mid (\omega, \omega) = 0\}$, we can decompose its boundary in Q as follows:

$$\partial \Omega_L^+ = \bigsqcup_I B_I,$$

where I is an isotropic subspace of $L_{\mathbb{R}}$ and B_I is the interior of $\mathbb{P}(I_{\mathbb{C}}) \cap \overline{\Omega_L^+}$. Each B_I is biholomorphic to the upper half-plane \mathbb{H} or a point. The set B_I is called a *rational boundary component* if $I = E_{\mathbb{R}}$ for some isotropic sublattice E of L. Thus there is a canonical identification

$$I_{i+1}(L) = \left\{ i \text{-dimensional rational boundary components of } \Omega_L^+ \right\}$$

for i = 0, 1, and Ω_L^+ has no higher dimensional rational boundary components.

Now assume that we are given a finite-index subgroup $\Gamma \subset O(L)^+$ with $\{\pm id\} \subset \Gamma$. The Baily-Borel compactification of $\Gamma \setminus \Omega_L^+$ is set-theoretically the set

$$\overline{\Gamma \backslash \Omega_L^+} = \Gamma \backslash \left(\Omega_L^+ \sqcup \bigcup_{I_1(L)} B_{\mathbb{Z}l} \sqcup \bigcup_{I_2(L)} B_E \right) = \left(\Gamma \backslash \Omega_L^+ \right) \sqcup \bigcup_{\Gamma \backslash I_1(L)} [\mathbb{C}l] \sqcup \bigcup_{\Gamma \backslash I_2(L)} \Gamma^E \backslash B_E,$$

where $\Gamma^E = \{\gamma \in \Gamma \mid \gamma(E) = E\}$ for $E \in I_2(L)$. It turns out that $\overline{\Gamma \setminus \Omega_L^+}$ is a normal projective variety. The boundary components of $\Gamma \setminus \Omega_L^+$ are often called *cusps*. We have the canonical identification

$$\Gamma \setminus I_{i+1}(L) = \left\{ i \text{-dimensional cusps of } \Gamma \setminus \Omega_L^+ \right\}.$$

Definition 2.8. A 0-dimensional cusp corresponding to $[\mathbb{Z}l]$ with $\operatorname{div}(l) = 1$ is called a standard cusp.

By our assumption that $\Gamma \subset O(L)^+$, the notion of standardness is welldefined. Since $\{\pm id\} \subset \Gamma$, the set of 0-dimensional standard cusps of $\Gamma \setminus \Omega_L^+$ is identified with $\Gamma \setminus I^1(L)$.

For a rank 2 primitive isotropic sublattice $E \in I_2(L)$, the compact curve

$$\left(\Gamma^{E} \backslash B_{E}\right) \sqcup \left(\Gamma \backslash \bigcup_{\gamma \in \Gamma} \bigcup_{\mathbb{Z}l \in \gamma(E)} [\mathbb{C}l]\right)$$
(3)

is obtained from the curve

$$\left(\Gamma^{E} \backslash B_{E}\right) \sqcup \left(\Gamma^{E} \backslash \bigcup_{\mathbb{Z}l \in E} [\mathbb{C}l]\right)$$

$$\tag{4}$$

by identifying Γ -equivalent points in $\bigcup_{\mathbb{Z}l\in E} [\mathbb{C}l]$. The curve (4) is the canonical compactification of $\Gamma^E \setminus B_E \simeq \Gamma^E \setminus \mathbb{H}$ itself.

When there exists an embedding $\varphi: U \hookrightarrow L$, we can write $L = \varphi(U) \oplus L_{\varphi}$ for the lattice $L_{\varphi} := \varphi(U)^{\perp} \cap L$ of $\operatorname{sign}(L_{\varphi}) = (1, b^{-} - 1)$. Assume that we have chosen an orientation, say Ω_{L}^{+} , of L. Then we can choose the connected component L_{φ}^{+} of the open set $\{v \in (L_{\varphi})_{\mathbb{R}}, (v, v) > 0\}$ so that for each vector $v \in L_{\varphi}^{+}$ the oriented two-plane $\mathbb{R}\varphi(e+f) \oplus \mathbb{R}v$ belongs to Ω_{L}^{+} . If there is another embedding $\varphi': U \hookrightarrow L$ with $\varphi(f) = \varphi'(f)$, the projection $\gamma_{\varphi,\varphi'} := (pr_{L_{\varphi'}})|_{L_{\varphi}}:$ $L_{\varphi} \to L_{\varphi'}$ is an isometry (Lemma 2.3). Then it can be immediately checked that $\gamma_{\varphi,\varphi'}$ maps the cone L_{φ}^{+} to the cone $L_{\varphi'}^{+}$.

3 FM(S), FM^d(S) and the 0-dimensional cusps of $\Gamma^+_S \setminus \Omega^+_{\widetilde{NS}(S)}$

3.1 0-dimensional standard cusps and FM-partners

Let S be a K3 surface. We denote by $D^b(S)$ the bounded derived category of S. A K3 surface S' is called a Fourier-Mukai (FM) partner of S if $D^b(S) \simeq D^b(S')$ as triangulated categories. Let FM(S) be the set of isomorphism classes of FM-partners of S. By the results of Mukai and Orlov ([9], [10], [12]), FM(S) can be studied via the period of S. We denote by $\tilde{H}(S,\mathbb{Z})$ the total cohomology group $H^*(S,\mathbb{Z})$ equipped with the Mukai pairing

$$((r_1, l_1, s_1), (r_2, l_2, s_2)) := (l_1, l_2) - (r_1, s_2) - (s_1, r_2)$$

where $r_i \in H^0(S,\mathbb{Z}), l_i \in H^2(S,\mathbb{Z}), s_i \in H^4(S,\mathbb{Z})$. Since $H^2(S,\mathbb{Z}) \simeq \Lambda_{K3}$ as a lattice, $\widetilde{H}(S,\mathbb{Z}) \simeq H^2(S,\mathbb{Z}) \oplus U \simeq \widetilde{\Lambda}_{K3}$ as a lattice. We fix an isometry $H^0(S,\mathbb{Z}) + H^4(S,\mathbb{Z}) \simeq U$ by identifying (1,0,0) with e, and (0,0,-1) with f. The lattice $\widetilde{H}(S,\mathbb{Z})$ inherits the weight-two Hodge structure from $H^2(S,\mathbb{Z})$. If we denote by $\omega_S \in H^2(S,\mathbb{C})$ the period of S, then

$$\omega_S^{\perp} \cap \widetilde{H}(S, \mathbb{Z}) = NS(S) \simeq NS(S) \oplus U.$$

It is obvious that $T(S) = \widetilde{NS}(S)^{\perp} \cap \widetilde{H}(S, \mathbb{Z})$. Let $NS(S)^{+}$ be the positive cone, i.e., the component of $\{v \in NS(S)_{\mathbb{R}}, (v, v) > 0\}$ containing ample classes. We choose an orientation, say $\Omega^{+}_{\widetilde{NS}(S)}$, of $\widetilde{NS}(S)$ so that for a vector $v \in NS(S)^{+}$ the oriented positive-definite two-plane $\mathbb{R}(1, 0, -1) \oplus \mathbb{R}(0, v, 0)$ belongs to $\Omega^{+}_{\widetilde{NS}(S)}$.

Theorem 3.1 ([10], [12]). Let S and S' be K3 surfaces. Then $D^b(S) \simeq D^b(S')$ if and only if there exists a Hodge isometry $\widetilde{H}(S,\mathbb{Z}) \simeq \widetilde{H}(S',\mathbb{Z})$.

Thanks to this theorem, we are able to obtain every member of FM(S) by an embedding of U into $\widetilde{NS}(S)$.

Lemma 3.2. For an embedding $\varphi \in \operatorname{Emb}(U, NS(S))$, there exists a unique (up to isomorphism) K3 surface $S_{\varphi} \in \operatorname{FM}(S)$ such that $H^2(S_{\varphi}, \mathbb{Z})$ is Hodge isometric to $\Lambda_{\varphi} := \varphi(U)^{\perp} \cap \widetilde{H}(S, \mathbb{Z})$. The correspondence $\varphi \mapsto S_{\varphi}$ gives a surjection

$$\operatorname{Emb}(U, NS(S)) \twoheadrightarrow \operatorname{FM}(S).$$

Proof. The lattice Λ_{φ} is an even unimodular lattice of signature (3, 19), hence is isometric to the K3 lattice Λ_{K3} . Since $T(S) \subset \Lambda_{\varphi}$, Λ_{φ} inherits the period from that of T(S). By the surjectivity of period map ([16], [17]), there is a K3 surface S_{φ} such that $H^2(S_{\varphi}, \mathbb{Z})$ is Hodge isometric to Λ_{φ} . By the Torelli theorem ([14], [17]), S_{φ} is unique up to isomorphism. Since

$$\widetilde{H}(S_{\varphi},\mathbb{Z})\simeq U\oplus\Lambda_{\varphi}\simeq\varphi(U)\oplus\Lambda_{\varphi}=\widetilde{H}(S,\mathbb{Z})$$

is an isometry of lattices preserving the periods, $S_{\varphi} \in FM(S)$ by Theorem 3.1.

Conversely, if $S' \in FM(S)$, there is a Hodge isometry $\Phi : \tilde{H}(S', \mathbb{Z}) \xrightarrow{\simeq} \tilde{H}(S, \mathbb{Z})$ which induces a Hodge isometry

$$H^2(S',\mathbb{Z})\simeq \widetilde{\Phi}((H^0(S',\mathbb{Z})+H^4(S',\mathbb{Z}))^{\perp}\cap \widetilde{H}(S,\mathbb{Z}).$$

It follows that $S' \simeq S_{\tilde{\Phi}((H^0(S',\mathbb{Z})+H^4(S',\mathbb{Z})))}$ by the Torelli theorem.

By the construction of S_{φ} , we can identify $T(S_{\varphi}) = T(S)$, and $NS(S_{\varphi}) = \varphi(U)^{\perp} \cap \widetilde{NS}(S)$. We shall introduce an equivalence relation on $\operatorname{Emb}(U, \widetilde{NS}(S))$ to make the above correspondence bijective.

Definition 3.3. Set

$$\Gamma_S := r_{\widetilde{NS}(S)}^{-1}(\lambda \circ r_{T(S)}(O_{Hodge}(T(S)))),$$

where $O_{Hodge}(T(S))$ is the group of Hodge isometries of T(S), $r_{\widetilde{NS}(S)} : O(\widetilde{NS}(S)) \to O(A_{\widetilde{NS}(S)})$ and $r_{T(S)} : O(T(S)) \to O(A_{T(S)})$ are the natural homomorphisms, and the isomorphism $\lambda : O(A_{T(S)}) \simeq O(A_{\widetilde{NS}(S)})$ is induced from the isometry $(A_{T(S)}, -q) \simeq (A_{\widetilde{NS}(S)}, q)$ (cf. Proposition 2.1).

By the identifications $NS(S) = NS(S_{\varphi})$ and $T(S) = T(S_{\varphi})$, We have $\Gamma_S = \Gamma_{S_{\varphi}}$. If we denote by $O_{Hodge}(\tilde{H}(S,\mathbb{Z}))$ the group of Hodge isometries of $\tilde{H}(S,\mathbb{Z})$, then Γ_S is the image of the natural homomorphism

$$O_{Hodge}(\tilde{H}(S,\mathbb{Z})) \to O(NS(S))$$

by Proposition 2.1.

In generic case, the period ω_S is not contained in any eigenspace of any $\varphi \in O(T(S)) - \{\pm id\}$ so that $O_{Hodge}(T(S)) = \{\pm id\}$. By the inclusions

$$O(NS(S))_0 \times \{\pm \mathrm{id}\} \subseteq \Gamma_S \subseteq O(NS(S)),$$

 Γ_S is a finite-index subgroup of $O(\widetilde{NS}(S))$. Then Γ_S acts on $\operatorname{Emb}(U, \widetilde{NS}(S))$ from left as $\gamma(\varphi) := \gamma \circ \varphi$ for $\gamma \in \Gamma_S$ and $\varphi \in \operatorname{Emb}(U, \widetilde{NS}(S))$.

Proposition 3.4. The correspondence $\varphi \mapsto S_{\varphi}$ of Lemma 3.2 induces the bijection

$$\Gamma_S \setminus \operatorname{Emb}(U, NS(S)) \simeq \operatorname{FM}(S).$$

Proof. It suffices to show that $S_{\varphi_1} \simeq S_{\varphi_2}$ if and only if there exists an isometry $\gamma \in \Gamma_S$ such that $\varphi_1 = \gamma \circ \varphi_2$. By the Torelli theorem, the former condition is equivalent to the existence of a Hodge isometry $\Phi : H^2(S_{\varphi_2}, \mathbb{Z}) \xrightarrow{\simeq} H^2(S_{\varphi_1}, \mathbb{Z})$.

Given such Φ , we get a Hodge isometry $\widetilde{\Phi} : \widetilde{H}(S_{\varphi_2}, \mathbb{Z}) \xrightarrow{\simeq} \widetilde{H}(S_{\varphi_1}, \mathbb{Z})$ by adding $\varphi_1 \circ \varphi_2^{-1}|_{\varphi_2(U)} : \varphi_2(U) \xrightarrow{\simeq} \varphi_1(U)$. Then, by Proposition 2.1 (2), the Hodge isometry $\widetilde{\Phi}|_{T(S)} : T(S) \xrightarrow{\simeq} T(S)$ and the isometry $\widetilde{\Phi}|_{\widetilde{NS}(S)} : NS(S_{\varphi_2}) \oplus$ $\varphi_2(U) \xrightarrow{\simeq} NS(S_{\varphi_1}) \oplus \varphi_1(U)$ are compatible on the discriminant form. It follows that $\widetilde{\Phi}|_{\widetilde{NS}(S)} \in \Gamma_S$. By the construction $\varphi_1 = (\widetilde{\Phi}|_{\widetilde{NS}(S)}) \circ \varphi_2$.

Conversely, if there exists $\gamma \in \Gamma_S$ such that $\varphi_1 = \gamma \circ \varphi_2$, we can find a Hodge isometry $g: T(S) \xrightarrow{\simeq} T(S)$ which is compatible with γ on the discriminant group. Then, again by Proposition 2.1 (2), $\gamma \oplus g$ extends to $\widetilde{\Phi} \in O_{Hodge}(\widetilde{H}(S,\mathbb{Z}))$, which gives a Hodge isometry between $H^2(S_{\varphi_2},\mathbb{Z}) = \varphi_2(U)^{\perp} \cap \widetilde{H}(S,\mathbb{Z})$ and $H^2(S_{\varphi_1},\mathbb{Z}) = \varphi_1(U)^{\perp} \cap \widetilde{H}(S,\mathbb{Z})$.

Now we turn to the 0-dimensional standard cusps. Let $\Omega^+_{\widetilde{NS}(S)}$ be the bounded symmetric domain associated with the lattice $\widetilde{NS}(S)$ (cf. Section 2.3). Denote by Γ^+_S the arithmetic group $\Gamma_S \cap O(\widetilde{NS}(S))^+$. Since the isometry $\mathrm{id}_{NS(S)} \oplus -\mathrm{id}_U \in O(\widetilde{NS}(S))_0$ exchanges $\Omega^+_{\widetilde{NS}(S)}$ and $\Omega^-_{\widetilde{NS}(S)}$, we know that Γ^+_S is of index 2 in Γ_S . Recall from Section 2.3 the canonical identification

$$\Gamma_{S}^{+} \setminus I^{1}(\widetilde{NS}(S)) = \Big\{ \text{ 0-dimensional standard cusps of } \Gamma_{S}^{+} \setminus \Omega_{\widetilde{NS}(S)}^{+} \Big\}.$$

Lemma 3.5. The natural surjection

$$\Gamma_S^+ \setminus I^1(\widetilde{NS}(S)) \longrightarrow \Gamma_S \setminus I^1(\widetilde{NS}(S))$$

is bijective.

Proof. For $l \in I^1(\widetilde{NS}(S))$, there exist an embedding $\varphi \in \operatorname{Emb}(U, \widetilde{NS}(S))$ such that $l \in \varphi(U)$. Then the isometry $\operatorname{id}_{\varphi(U)^{\perp}} \oplus -\operatorname{id}_{\varphi(U)} \in O(\widetilde{NS}(S))_0$ preserves $\mathbb{Z}l$ and interchanges $\Omega^+_{\widetilde{NS}(S)}$ with $\Omega^-_{\widetilde{NS}(S)}$. Therefore $\Gamma^+_S \cdot l = \Gamma_S \cdot l$.

Theorem 3.6. Let $\{e, f\}$ be the canonical basis of the lattice U. The map

$$\mu_0: \mathrm{FM}(S) \ni [S_{\varphi}] \longrightarrow [\varphi(f)] \in \Gamma_S \setminus I^1(NS(S)) .$$

is bijective.

Proof. The map μ_0 is well-defined by Proposition 3.4. Since $l \in I^1(NS(S))$ induces an embedding $\varphi \in \operatorname{Emb}(U, \widetilde{NS}(S))$ with $\varphi(f) = l, \mu_0$ is surjective. If $[\varphi_1(f)] = [\varphi_2(f)]$, then by identifying $\varphi_i(\mathbb{Z}f) = H^4(S_{\varphi_i}, \mathbb{Z})$, the isometry $id_{\widetilde{H}(S,\mathbb{Z})}$ induces the Hodge isometry $\widetilde{\Phi} : \widetilde{H}(S_{\varphi_1}, \mathbb{Z}) \simeq \widetilde{H}(S_{\varphi_2}, \mathbb{Z})$ with $\widetilde{\Phi}(H^4(S_{\varphi_1}, \mathbb{Z})) = H^4(S_{\varphi_2}, \mathbb{Z})$. Now Lemma 2.3 implies that $pr_{H^2(S_{\varphi_2}, \mathbb{Z})} \circ (\widetilde{\Phi}|_{H^2(S_{\varphi_1}, \mathbb{Z})})$ gives a Hodge isometry between $H^2(S_{\varphi_1}, \mathbb{Z})$ and $H^2(S_{\varphi_2}, \mathbb{Z})$, since $pr_{H^2(S_{\varphi_2}, \mathbb{Z})}$ is identity on $T(S_{\varphi_2}) = \widetilde{\Phi}(T(S_{\varphi_1}))$.

In this way, we have

$$\# FM(S) = \# \{ \text{ 0-dimensional standard cusps of } \Gamma^+_S \setminus \Omega^+_{\widetilde{NS}(S)} \}.$$

3.2 General 0-dimensional cusps and twisted FM-partners

In this subsection, using Huybrechts-Stellari's solution of Căldăraru conjecture ([8]), we study the relation between general 0-dimensional cusps of $\Gamma_S^+ \setminus \Omega_{\widetilde{NS}(S)}^+$ and twisted Fourier-Mukai partners of S. For twisted K3 surfaces, see [7].

Let (S', α') be a twisted K3 surface. By the natural isomorphism

 $\operatorname{Br}(S') \simeq \operatorname{Hom}(T(S'), \mathbb{Q}/\mathbb{Z}),$

we identify the twisting $\alpha' \in \operatorname{Br}(S')$ with a surjective homomorphism $\alpha' : T(S') \to \mathbb{Z}/d\mathbb{Z}$ where $d = \operatorname{ord}(\alpha')$. Then $\operatorname{Ker}(\alpha')$ is denoted by $T(S', \alpha')$. A twisted K3 surface (S', α') is called a *twisted Fourier-Mukai partner* of S if there is an equivalence $D^b(S', \alpha') \simeq D^b(S)$. By the result of Canonaco-Stellari ([4]), the equivalence is of Fourier-Mukai type. We define

$$\mathrm{FM}^{d}(S) := \left\{ (S', \alpha') : \mathrm{twisted} K3 \mathrm{surface}, D^{b}(S', \alpha') \simeq D^{b}(S), \mathrm{ord}(\alpha') = d \right\} / \simeq$$

It is obvious that $FM^1(S) = FM(S)$.

Proposition 3.7. There exists a map

$$\nu_0: \Gamma_S^+ \setminus I^d(\widetilde{NS}(S)) \longrightarrow \mathrm{FM}^d(S).$$

If d = 1, we have $\nu_0 = \mu_0^{-1}$ on $\Gamma_S^+ \setminus I^1(\widetilde{NS}(S))$.

Proof. For $l \in I^d(\widetilde{NS}(S))$, we define

$$\widetilde{M}_l := \left\langle \frac{l}{d}, \widetilde{NS}(S) \right\rangle \subset \widetilde{NS}(S)^{\vee},$$

which is an even overlattice of NS(S). Via the isometry

$$\lambda := \lambda_{\widetilde{H}(S,\mathbb{Z})} : (A_{\widetilde{NS}(S)}, q) \simeq (A_{T(S)}, -q),$$

we have an isotropic element $\lambda(\frac{l}{d}) \in A_{T(S)}$ of order = d. Set

$$T_l := \left\langle \lambda(\frac{l}{d}), T(S) \right\rangle \subset T(S)^{\vee}.$$

We have a surjective homomorphism

$$\alpha_l: T_l \to \mathbb{Z}/d\mathbb{Z}$$
 with $\operatorname{Ker}(\alpha_l) = T(S), \ \alpha_l(\lambda(\frac{l}{d})) = \overline{1}.$

Since the isometry λ induces the isometry $\overline{\lambda} : (A_{\widetilde{M}_l}, q) \simeq (A_{T_l}, -q)$, we have an embedding $\widetilde{M}_l \oplus T_l \hookrightarrow \widetilde{\Lambda}_{K3}$ with both \widetilde{M}_l and T_l embedded primitively. Since $\frac{l}{d} \in I^1(\widetilde{M}_l)$, there exists an embedding $\varphi : U \hookrightarrow \widetilde{M}_l$ with $\varphi(f) = \frac{l}{d}$. The orthogonal complement $\Lambda_{\varphi} := \varphi(U)^{\perp} \cap \widetilde{\Lambda}_{K3}$ is isometric to the K3 lattice Λ_{K3} and has the period induced from T_l . Moreover, via the orientation of \widetilde{M}_l (induced from $\widetilde{NS}(S)$), we can choose the positive cone M_{φ}^+ of the lattice $M_{\varphi} := \varphi(U)^{\perp} \cap \widetilde{M}_l$ (see Section 2.3). By the surjectivity of the period map, there exist a K3 surface S_{φ} and a Hodge isometry $\Phi : H^2(S_{\varphi}, \mathbb{Z}) \xrightarrow{\cong} \Lambda_{\varphi}$ such that $\Phi(NS(S_{\varphi})^+) = M_{\varphi}^+$. Pulling back the homomorphism $\alpha_l : T_l \to \mathbb{Z}/d\mathbb{Z}$ by $\Phi|_{T(S_{\varphi})}$, we obtain a surjective homomorphism $\alpha_{\Phi} : T(S_{\varphi}) \to \mathbb{Z}/d\mathbb{Z}$. Thus we constructed a twisted K3 surface $(S_{\varphi}, \alpha_{\Phi})$.

If there are another K3 surface S'_{φ} and a Hodge isometry $\Phi': H^2(S'_{\varphi}, \mathbb{Z}) \xrightarrow{\simeq} \Lambda_{\varphi}$ such that $\Phi'(NS(S'_{\varphi})^+) = M_{\varphi}^+$, we have a Hodge isometry $\Phi'' := (\Phi')^{-1} \circ \Phi : H^2(S_{\varphi}, \mathbb{Z}) \xrightarrow{\simeq} H^2(S'_{\varphi}, \mathbb{Z})$ with $\Phi''(NS(S_{\varphi})^+) = NS(S'_{\varphi})^+$ and $(\Phi''|_{T(S_{\varphi})})^* \alpha_{\Phi'} = \alpha_{\Phi}$. Composing Φ'' with an element of the Weyl group $W(S'_{\varphi})$, we may assume that Φ'' is effective so that we have $(S_{\varphi}, \alpha_{\Phi}) \simeq (S'_{\varphi}, \alpha_{\Phi'})$ by the Torelli theorem. Thus we constructed a twisted K3 surface $(S_{\varphi}, \alpha_{\varphi})$ from an embedding φ .

If we take another embedding $\varphi' : U \hookrightarrow M_l$ with $\varphi'(f) = \frac{l}{d}$, it follows from Lemma 2.3 and Section 2.3 that the projection defines an isometry $\Lambda_{\varphi} \xrightarrow{\simeq} \Lambda_{\varphi'}$ which is identity on T_l and maps the cone M_{φ}^+ to the cone $M_{\varphi'}^+$. So we have a Hodge isometry $\Phi : H^2(S_{\varphi}, \mathbb{Z}) \xrightarrow{\simeq} H^2(S_{\varphi'}, \mathbb{Z})$ such that $\Phi(NS(S_{\varphi})^+) =$ $NS(S_{\varphi'})^+$ and $(\Phi|_{T(S_{\varphi})})^* \alpha_{\varphi'} = \alpha_{\varphi}$. As above, we obtain an isomorphism $(S_{\varphi}, \alpha_{\varphi}) \simeq (S_{\varphi'}, \alpha_{\varphi'})$. Hence we constructed a twisted K3 surface (S_l, α_l) from $l \in I^d(\widetilde{NS}(S))$.

Next we consider the action of the group Γ_S^+ . For an isometry $\gamma \in \Gamma_S^+$, set $l' := \gamma(l)$. The isometry γ extends to the isometry $\tilde{\gamma} : \widetilde{M}_l \xrightarrow{\rightarrow} \widetilde{M}_{l'}$. Take an embedding $\varphi : U \hookrightarrow \widetilde{M}_l$ with $\varphi(f) = \frac{l}{d}$ and set $\varphi' := \tilde{\gamma} \circ \varphi : U \hookrightarrow \widetilde{M}_{l'}$, which satisfies $\varphi'(f) = \frac{l'}{d}$. On the other hand, there exists a Hodge isometry $g \in O_{Hodge}(T(S))$ such that $\lambda \circ r(\gamma) = r(g) \circ \lambda$. Since $r(g)(\lambda(\frac{l}{d})) = \lambda(\frac{l'}{d})$, gextends to the Hodge isometry $\tilde{g} : T_l \xrightarrow{\rightarrow} T_{l'}$ with $\tilde{g}^* \alpha_{l'} = \alpha_l$. Because we have $\bar{\lambda} \circ r(\tilde{\gamma}) = r(\tilde{g}) \circ \bar{\lambda}$, the isometry $\tilde{\gamma} \oplus \tilde{g}$ extends to the Hodge isometry

$$(\varphi' \circ \varphi^{-1}) \oplus \Phi : \varphi(U) \oplus \Lambda_{\varphi} \xrightarrow{\simeq} \varphi'(U) \oplus \Lambda_{\varphi'} \text{ with } (\Phi|_{T_l})^* \alpha_{l'} = \alpha_l.$$

Since γ preserves the orientation, we have $\Phi(M_{\varphi}^+) = M_{\varphi'}^+$. In this way we obtain a Hodge isometry $\Phi : H^2(S_l, \mathbb{Z}) \xrightarrow{\simeq} H^2(S_{l'}, \mathbb{Z})$ with $\Phi(NS(S_l)^+) = NS(S_{l'})^+$ and $(\Phi|_{T(S_l)})^* \alpha_{l'} = \alpha_l$. As above we have $(S_l, \alpha_l) \simeq (S_{l'}, \alpha_{l'})$.

Finally, we see that $D^b(S_l, \alpha_l) \simeq D^b(S)$. Let $B_l \in H^2(S_l, \mathbb{Q})$ be a B-field lift of $\alpha_l \in Br(S_l)$. Since we have a Hodge isometry $T(S) \simeq T(S_l, B_l) \subset \widetilde{H}(S_l, B_l, \mathbb{Z})$, the lattice $\widetilde{NS}(S_l, B_l) := T(S_l, B_l)^{\perp} \cap \widetilde{H}(S_l, B_l, \mathbb{Z})$ is isogenus to $\widetilde{NS}(S)$, so we have an isometry $\widetilde{NS}(S) \simeq \widetilde{NS}(S_l, B_l)$ by Proposition 2.2. Hence there is a Hodge embedding

$$\gamma \oplus g : NS(S) \oplus T(S) \hookrightarrow H(S_l, B_l, \mathbb{Z}).$$

Since the natural homomorphism $r: O(\widetilde{NS}(S)) \to O(A_{\widetilde{NS}(S)})$ is surjective by Proposition 2.2, after composing γ with an element of $O(\widetilde{NS}(S))$ (and with $\mathrm{id}_{NS(S)} \oplus -\mathrm{id}_U \in O(NS(S))_0$ if necessary), $\gamma \oplus g$ extends to an orientationpreserving Hodge isometry $\widetilde{H}(S,\mathbb{Z}) \simeq \widetilde{H}(S_l, B_l, \mathbb{Z})$. Then our claim follows from Huybrechts-Stellari's theorem ([8]).

The proof of the assertion that $\nu_0 = \mu_0^{-1}$ for d = 1 can be left to the reader.

Definition 3.8. Set

$$\mathcal{FM}^d(S) := \mathrm{FM}^d(S) / \sim$$

where $(S_1, \alpha_1) \sim (S_2, \alpha_2)$ if there exists a Hodge isometry $g: T(S_1) \xrightarrow{\simeq} T(S_2)$ with $g^* \alpha_2 = \alpha_1$. Since $(S_1, \alpha_1) \simeq (S_2, \alpha_2)$ implies $(S_1, \alpha_1) \sim (S_2, \alpha_2)$, the set $\mathcal{FM}^d(S)$ is well-defined. Denote by $\pi: \mathrm{FM}^d(S) \to \mathcal{FM}^d(S)$ the quotient map.

Definition 3.9. Set

$$I^{d}(A_{\widetilde{NS}(S)}) := \Big\{ x \in A_{\widetilde{NS}(S)} \ \Big| \ q_{\widetilde{NS}(S)}(x) = 0 \ \in \mathbb{Q}/2\mathbb{Z}, \ \mathrm{ord}(x) = d \Big\}.$$

Since $NS(S) = NS(S) \oplus U$, the map

$$p: I^d(\widetilde{NS}(S)) \longrightarrow I^d(A_{\widetilde{NS}(S)}), \quad l \mapsto \frac{l}{d}$$

is surjective by Proposition 4.1.1 of [15].

Theorem 3.10. There exists a bijective map $\xi_0 : r(\Gamma_S^+) \setminus I^d(A_{\widetilde{NS}(S)}) \to \mathcal{FM}^d(S)$ which fits in the following commutative diagram.

$$\begin{array}{ccc} \Gamma_{S}^{+} \backslash I^{d}(\widetilde{NS}(S)) & \stackrel{\nu_{0}}{\longrightarrow} & \mathrm{FM}^{d}(S) \\ & p \\ & & \downarrow \pi \\ r(\Gamma_{S}^{+}) \backslash I^{d}(A_{\widetilde{NS}(S)}) & \stackrel{-}{\longrightarrow} & \mathcal{FM}^{d}(S). \end{array}$$

Proof. We define ξ_0 by $\pi \circ \nu_0 \circ p^{-1}$. For $x = \frac{l}{d} \in I^d(A_{\widetilde{NS}(S)})$ we have an isotropic cyclic subgroup $\lambda(\langle x \rangle) \subset A_{T(S)}$ of order d and its generator $\lambda(x) \in A_{T(S)}$, which define an overlattice $T_x \supset T(S)$ and a surjective homomorphism $\alpha_x : T_x \to \mathbb{Z}/d\mathbb{Z}$ with $\operatorname{Ker}(\alpha_x) = T(S)$, $\alpha_x(\lambda(x)) = \overline{1} \in \mathbb{Z}/d\mathbb{Z}$. Here we write λ for $\lambda_{\widetilde{H}(S,\mathbb{Z})} : (A_{\widetilde{NS}(S)}, q) \xrightarrow{\cong} (A_{T(S)}, -q)$. The isotropic subgroup $\langle x \rangle \subset A_{\widetilde{NS}(S)}$ defines an overlattice $\widetilde{M}_x = \left\langle \frac{l}{d}, \widetilde{NS}(S) \right\rangle \supset \widetilde{NS}(S)$. Similarly as the construction of ν_0 in Proposition 3.7, λ induces an isometry $\overline{\lambda} : (A_{\widetilde{M}_x}, q) \simeq (A_{T_x}, -q)$, which gives an embedding $\widetilde{M}_x \oplus T_x \hookrightarrow \widetilde{\Lambda}_{K3}$. Then the isotropic vector $\frac{l}{d} \in I^1(\widetilde{M}_x)$ defines the twisted K3 surface $(S_l, \alpha_x) \in \operatorname{FM}^d(S)$.

Since the definitions of T_x and α_x depend only on $x \in I^d(A_{\widetilde{NS}(S)})$, the equivalence class $[(S_l, \alpha_x)] \in \mathcal{FM}^d(S)$ is independent of the choices of $l \in I^d(\widetilde{NS}(S))$ with $x = \frac{l}{d}$.

Take an isometry $\gamma \in \Gamma_S^+$ and set $x' := r(\gamma)(x)$. If we choose an isotropic vector $l \in I^d(\widetilde{NS}(S))$ with $\underline{l}_d = x$, then $\gamma(\underline{l}_d) = x' \in A_{\widetilde{NS}(S)}$. Now we have $(S_l, \alpha_l) \simeq (S_{\gamma(l)}, \alpha_{\gamma(l)})$ so that $[(S_l, \alpha_l)] = [(S_{\gamma(l)}, \alpha_{\gamma(l)})]$. Hence the map ξ_0 is well-defined.

We prove the surjectivity of ξ_0 . For a twisted FM partner $(S', \alpha') \in FM^d(S)$, we have a Hodge isometry $T(S) \simeq T(S', \alpha')$ and an inclusion $T(S', \alpha') \subset T(S')$ with $\operatorname{Ker}(\alpha') = T(S', \alpha')$. Set $y := (\alpha')^{-1}(\overline{1}) \in A_{T(S)} \simeq A_{T(S', \alpha')}$, where $\overline{1} \in \mathbb{Z}/d\mathbb{Z}$ is the generator. Then we have an isotropic element $x := \lambda^{-1}(y) \in \mathbb{Z}$ $I^d(A_{\widetilde{NS}(S)})$. By the construction of T_x and α_x , we have a Hodge isometry $(T_x, \alpha_x) \simeq (T(S'), \alpha')$ so that $[(S_l, \alpha_x)] = [(S', \alpha')] \in \mathcal{FM}^d(S)$, where $l \in$ $I^d(NS(S))$ is such that $\frac{l}{d} = x$.

Finally, we prove the injectivity of ξ_0 . Suppose we have $[(S_l, \alpha_l)] = [(S_{l'}, \alpha_{l'})] \in$ $\mathcal{FM}^d(S)$. Since there exists a Hodge isometry $\widetilde{g}: T(S_l) \xrightarrow{\simeq} T(S_{l'})$ with $\widetilde{g}^* \alpha_{l'} =$ $\alpha_l, \ \widetilde{g} \text{ maps } T(S) = \operatorname{Ker}(\alpha_l) \subset T(S_l) \text{ to } T(S) = \operatorname{Ker}(\alpha_{l'}) \subset T(S_{l'}) \text{ isometri-}$ cally. Setting $g := \widetilde{g}|_{T(S)} \in O_{Hodge}(T(S))$, we have $r(g)\left(\lambda(\frac{l}{d})\right) = \lambda\left(\frac{l'}{d}\right)$. Hence $\lambda^{-1} \circ r(g) \circ \lambda \in r(\Gamma_S) = r(\Gamma_S^+)$ maps $\frac{l}{d}$ to $\frac{l'}{d}$.

The set $r(\Gamma_S^+) \setminus I^d(A_{\widetilde{NS}(S)})$ is relatively easy to calculate.

Corollary 3.11. If there is an embedding $U \hookrightarrow NS(S)$, then $\nu_0 : \Gamma_S^+ \setminus I^d(\widetilde{NS}(S)) \to$ $FM^{d}(S)$ is bijective. In particular, we have the formula

 $\# \mathrm{FM}^{d}(S) = \# \{ \text{ 0-dimensional cusp } [l] \text{ of } \Gamma^{+}_{S} \backslash \Omega^{+}_{\widetilde{NS}(S)} \text{ with } \mathrm{div}(l) = d \}.$

Proof. Since there is an embedding $U \oplus U \hookrightarrow \widetilde{NS}(S)$, p is bijective by Proposition 4.1.3 of [15]. On the other hand, we have an embedding $U \oplus U \hookrightarrow NS(S) \subset$ M_x so that Proposition 2.2 implies that π is bijective.

In this way, we can obtain informations about $FM^d(S)$ by studying 0dimensional cusps of $\Gamma_S^+ \setminus \Omega_{\widetilde{NS}(S)}^+$ and vice versa.

$\operatorname{FM}_{ell}(S)$ and certain 1-dimensional cusps of $\Gamma_S^+ \setminus \Omega_{\widetilde{NS}(S)}^+$ 4

Now we study the relation between FM-partners with elliptic fibrations and 1-dimensional cusps containing 0-dimensional standard cusps in their closures.

FM-partners with elliptic fibrations 4.1

Definition 4.1. Define

 $\mathrm{FM}_{ell}(S) := \left\{ (S', \mathcal{O}_{S'}(C')) \, \middle| \, S' \in \mathrm{FM}(S), C' \text{ is a smooth elliptic curve on } S' \right\} / \simeq$ where $(S_1, \mathcal{O}_{S_1}(C_1)) \simeq (S_2, \mathcal{O}_{S_2}(C_2))$ if there exists an isomorphism $\varphi : S_1 \simeq S_2$ such that $\varphi^* \mathcal{O}_{S_2}(C_2) \simeq \mathcal{O}_{S_1}(C_1)$. In other words, $(S_1, \mathcal{O}_{S_1}(C_1)) \simeq (S_2, \mathcal{O}_{S_2}(C_2))$

if they are isomorphic as elliptic K3 surfaces.

For an even lattice L, let $I(L) = \bigcup_d I^d(L)$ be the set of primitive isotropic vectors in L.

Lemma 4.2. By associating to $\mathcal{O}_{S'}(C')$ its class in NS(S'), there exists a one-to-one correspondence between $FM_{ell}(S)$ and the set

$$\left\{ (S', l') \left| S' \in FM(S), l' \in I(NS(S')) \right\} \right\} \simeq .$$
(5)

In (5), $(S_1, l_1) \simeq (S_2, l_2)$ if there exists a Hodge isometry $\Phi : H^2(S_1, \mathbb{Z}) \simeq H^2(S_2, \mathbb{Z})$ with $\Phi(l_1) = l_2$.

Proof. It is known (Corollary 6.1 of [14]) that every primitive isotropic element of NS(S) can be transformed by the action of $\{\pm id\} \times W(S)$ to the class of a smooth elliptic curve. Thus, each element of the set (5) is represented by the divisor class of a smooth elliptic curve. Let $l_1 \in NS(S_1), l_2 \in NS(S_2)$ be the classes of smooth elliptic curves. By the Torelli theorem, it suffices to show that, if there is a Hodge isometry $\Phi: H^2(S_1,\mathbb{Z}) \xrightarrow{\simeq} H^2(S_2,\mathbb{Z})$ with $\Phi(l_1) = l_2$, then there is an effective Hodge isometry $\Phi' : H^2(S_1, \mathbb{Z}) \xrightarrow{\simeq} H^2(S_2, \mathbb{Z})$ with $\Phi'(l_1) = l_2$. Let $\mathcal{A}(S)_{\mathbb{R}}$ be the cone in $NS(S)_{\mathbb{R}}$ generated by the ample classes of S over $\mathbb{R}_{>0}$. Since l_i is contained in the boundary of the positive cone $NS(S_i)^+$, we have $\Phi(\mathcal{A}(S_1)_{\mathbb{R}}) \subset NS(S_2)^+$. Then we have two chambers $\Phi(\mathcal{A}(S_1)_{\mathbb{R}})$ and $\mathcal{A}(S_2)_{\mathbb{R}}$ in $NS(S_2)^+$, and $\Phi(l_1) = l_2$ is contained in the closures of both chambers. If we take $x_1 \in \Phi(\mathcal{A}(S_1)_{\mathbb{R}})$ and $x_2 \in \mathcal{A}(S_2)_{\mathbb{R}}$, only finite number of walls $\{W_i\}_1^N$ intersect with the segment $\overline{x_1x_2}$, and each W_i must passes through l_2 . Writing $W_i \cap \overline{x_1 x_2} = t_i x_2 + (1 - t_i) x_1$, we may assume that $0 < t_1 < \cdots < t_N < 1$. If we denote by s_{W_i} the reflection with respect to W_i , then $s_{W_N} \circ \ldots \circ s_{W_1}$ maps $\Phi(\mathcal{A}(S_1)_{\mathbb{R}})$ to $\mathcal{A}(S_2)_{\mathbb{R}}$ and fixes l_2 . Hence $s_{W_N} \circ \ldots \circ s_{W_1} \circ \Phi$ gives the desired effective Hodge isometry Φ' .

In what follows, we identify the two sets in Lemma 4.2. Let $\mathbb{Z}l$ be a degenerate lattice of rank 1, and consider the degenerate lattice $U \oplus \mathbb{Z}l = \mathbb{Z}e + \mathbb{Z}f + \mathbb{Z}l$ of rank 3, where (e, f) = 1, (e, l) = (f, l) = (l, l) = (e, e) = (f, f) = 0. The proof of the following theorem is parallel to that of Lemma 3.2 and Proposition 3.4.

Theorem 4.3. For an embedding $\varphi \in \operatorname{Emb}(U \oplus \mathbb{Z}l, \widetilde{NS}(S))$ we have $\varphi(l) \in \varphi(U)^{\perp} \cap \widetilde{NS}(S) = NS(S_{\varphi})$ so that we get $[(S_{\varphi}, \varphi(l))] \in \operatorname{FM}_{ell}(S)$. Then the correspondence

$$\Gamma_S \setminus \operatorname{Emb}(U \oplus \mathbb{Z}l, NS(S)) \ni [\varphi] \mapsto [(S_{\varphi}, \varphi(l))] \in \operatorname{FM}_{ell}(S)$$

is a bijection.

Corollary 4.4. The following counting formula for $FM_{ell}(S)$ holds:

$$\#(\mathrm{FM}_{ell}(S)) = \#\left(\bigsqcup_{[M]}\bigsqcup_{[k]} r_{T(S)}(O_{Hodge}(T(S))) \backslash O(A_M) / r_M(O(M)^k)\right)$$

where [M] runs over $\mathcal{G}(NS(S))$ and [k] runs over $O(M)\setminus I(M) = O(M)\setminus I_1(M)$.

Proof. We have

$$\Gamma_{S} \setminus \operatorname{Emb}(U \oplus \mathbb{Z}l, \widetilde{NS}(S))$$

$$= \bigsqcup_{[M]} \bigsqcup_{[k]} \Gamma_{S} \setminus \left\{ \varphi \in \operatorname{Emb}(U \oplus \mathbb{Z}l, \widetilde{NS}(S)) \mid \varphi(U)^{\perp} \simeq M, [\varphi(l)] = [k] \right\}$$

Since $O(M \oplus U)$ acts transitively on each set $\{\varphi \in \operatorname{Emb}(U \oplus \mathbb{Z}l, \widetilde{NS}(S)) | \varphi(U)^{\perp} \simeq M, [\varphi(l)] = [k]\}$ with stabilizer $O(M)^k$, the last line can be written as

$$= \bigsqcup_{[M]} \bigsqcup_{[k]} \Gamma_S \setminus O(M \oplus U) / O(M)^k$$
$$= \bigsqcup_{[M]} \bigsqcup_{[k]} r_{T(S)}(O_{Hodge}(T(S))) \setminus O(A_M) / r_M(O(M)^k),$$

where we used Proposition 2.2 and the inclusion $O(M \oplus U)_0 \subset \Gamma_S$ in the last equality.

Definition 4.5. Define

$$\operatorname{FM}_{ell,sec}(S) := \Big\{ \left[(S', \mathcal{O}_{S'}(C')) \right] \in \operatorname{FM}_{ell}(S) \ \Big| \ [C'] \in NS(S'), \operatorname{div}([C']) = 1 \Big\}.$$

Note that $\operatorname{div}([C']) = 1$ if and only if the elliptic fibration associated to $\mathcal{O}_{S'}(C')$ has a section. If S is an elliptic K3 surface admitting a section, then $\operatorname{FM}(S) = \{S\}$ by Corollary 2.7 of [5]. Hence $\operatorname{FM}_{ell,sec}(S)$ (if not empty) is exactly the set of isomorphism classes of elliptic fibrations on S admitting a section.

Corollary 4.6. The following equality holds:

$$\#(\mathrm{FM}_{ell,sec}(S)) = \#\left(\bigsqcup_{L \in \mathcal{G}(l^{\perp}/\mathbb{Z}l)} r_{T(S)}(O_{Hodge}(T(S))) \setminus O(A_L)/r_L(O(L))\right),$$

where l is an arbitrary standard isotropic vector in NS(S), and $A_{NS(S)}$ is identified with A_L by the isometry $NS(S) \simeq U \oplus L$.

Proof. By Proposition 2.4, $O(NS(S)) \setminus I^1(NS(S))$ is identified with $\mathcal{G}(l^{\perp}/\mathbb{Z}l)$. On the other hand, $r_{NS(S)}(O(NS(S))^l) \simeq r_L(O(l^{\perp}/\mathbb{Z}l))$ by Proposition 2.7. Now the assertion follows from Corollary 4.4.

4.2 $FM_{ell}(S)$ and certain 1-dimensional cusps

Definition 4.7. Set

$$I_2^{st}(\widetilde{NS}(S)) := \left\{ E \in I_2(\widetilde{NS}(S)) \mid \text{there is } e \in \widetilde{NS}(S) \text{ with}(e, E) = \mathbb{Z} \right\}.$$
(6)

If we take $f, l \in E$ so that (e, f) = 1 and $\operatorname{Ker}(e, \cdot)|_E = \mathbb{Z}l$, then $E + \mathbb{Z}e = \mathbb{Z}l + \mathbb{Z}f + \mathbb{Z}e \simeq U \oplus \mathbb{Z}l$. We may assume that e in (6) is taken to be isotropic.

Lemma 4.8. Let $E \in I_2^{st}(\widetilde{NS}(S))$.

(1) If $f, f' \in E$ satisfy the equalities (e, f) = (e, f') = 1, then

$$(S_{\mathbb{Z}e+\mathbb{Z}f},l)\simeq (S_{\mathbb{Z}e+\mathbb{Z}f'},l)$$

(2) If $e, e' \in NS(S)$ satisfy the relations $(e, E) = (e', E) = \mathbb{Z}$ and $\operatorname{Ker}(e, \cdot)|_E = \operatorname{Ker}(e', \cdot)|_E = \mathbb{Z}l$, then

$$(S_{\mathbb{Z}e+\mathbb{Z}f}, l) \simeq (S_{\mathbb{Z}e'+\mathbb{Z}f'}, l).$$

Here $f' \in E$ is an arbitrary element satisfying (e', f') = 1. (3) If $e, e' \in \widetilde{NS}(S)$ satisfy (e, f) = (e', f) = 1 for some $f \in E$, then

$$(S_{\mathbb{Z}e+\mathbb{Z}f}, l) \simeq (S_{\mathbb{Z}e'+\mathbb{Z}f}, l')$$

Here l (resp. l') is a generator of $\operatorname{Ker}(e, \cdot)|_E$ (resp. $\operatorname{Ker}(e', \cdot)|_E$).

Proof. (1) Since $l \in (\mathbb{Z}e + \mathbb{Z}f)^{\perp} \cap (\mathbb{Z}e + \mathbb{Z}f')^{\perp}$, Lemma 2.3 tells that the projection from $(\mathbb{Z}e + \mathbb{Z}f)^{\perp}$ to $(\mathbb{Z}e + \mathbb{Z}f')^{\perp}$ gives a Hodge isometry $H^2(S_{\mathbb{Z}e + \mathbb{Z}f'}, \mathbb{Z}) \simeq H^2(S_{\mathbb{Z}e + \mathbb{Z}f'}, \mathbb{Z})$, which is identity on $\mathbb{Z}l$.

(2) Both $\{l, f\}$ and $\{l, f'\}$ are basis of E, so that $f' = \pm f + \alpha l$ for some integer $\alpha \in \mathbb{Z}$. Hence $(e, f) = (e, \pm f') = 1$, and $(S_{\mathbb{Z}e+\mathbb{Z}f}, l) \simeq (S_{\mathbb{Z}e+\mathbb{Z}(\pm f')}, l) \simeq (S_{\mathbb{Z}(\pm e)+\mathbb{Z}f'}, l)$ by (1). On the other hand, $(S_{\mathbb{Z}(\pm e)+\mathbb{Z}f'}, l) \simeq (S_{\mathbb{Z}e'+\mathbb{Z}f'}, l)$ by Lemma 2.3.

(3) As in the proof of (2), $l' = \pm l + \beta f$ for some integer $\beta \in \mathbb{Z}$. If we project $l' \in (\mathbb{Z}e' + \mathbb{Z}f)^{\perp}$ to $(\mathbb{Z}e + \mathbb{Z}f)^{\perp}$, the image of l' is given by $\pm l$. Now the claim follows from Lemma 2.3.

Now we relate $\operatorname{FM}_{ell}(S)$ to $\Gamma_S^+ \setminus I_2^{st}(\widetilde{NS}(S))$, the set of 1-dimensional cusps of $\Gamma_S^+ \setminus \Omega_{\widetilde{NS}(S)}^+$ whose closures contain 0-dimensional standard cusps, via Theorem 4.3. Similarly as Lemma 3.5, the projection

$$\Gamma_{S}^{+} \backslash I_{2}^{st}(\widetilde{NS}(S)) \longrightarrow \Gamma_{S} \backslash I_{2}^{st}(\widetilde{NS}(S))$$

is bijective.

Definition 4.9. Define the map $\mu_1 : \operatorname{FM}_{ell}(S) \longrightarrow \Gamma_S \setminus I_2^{st}(\widetilde{NS}(S))$ by

$$\mu_1((S_{\varphi},\varphi(l))) := [\mathbb{Z}\varphi(f) \oplus \mathbb{Z}\varphi(l)].$$

By the definition of $I_2^{st}(\widetilde{NS}(S))$, μ_1 is surjective. It follows that

$$\# \Big(\mathrm{FM}_{ell}(S) \Big) \geq \# \Big\{ 1 \text{-dimensional cusp of } \Gamma^+_S \backslash \Omega^+_{\widetilde{NS}(S)} \text{ whose} \\ \text{closure contains a 0-dimensional standard cusp } \Big\}.$$

Proposition 4.10. The map

$$\mu_1: \mu_1^{-1}\Big(\mu_1(\mathrm{FM}_{ell,sec}(S))\Big) \longrightarrow \mu_1(\mathrm{FM}_{ell,sec}(S))$$

is bijective. In particular, we have $\mu_1^{-1}\Big(\mu_1(\mathrm{FM}_{ell,sec}(S))\Big) = \mathrm{FM}_{ell,sec}(S).$

Proof. Take $[E] = \mu_1((S_{\mathbb{Z}e+\mathbb{Z}f}, l)) \in \mu_1(\mathrm{FM}_{ell,sec}(S))$. We can find an element $m \in (\mathbb{Z}e + \mathbb{Z}f)^{\perp} \cap \widetilde{NS}(S)$ with (m, m) = 0, (l, m) = 1. Write $U_1 := \mathbb{Z}e + \mathbb{Z}f$ and $U_2 := \mathbb{Z}l + \mathbb{Z}m$. Let $e', f', l' \in \widetilde{NS}(S)$ be such that $[E] = \mu_1((S_{\mathbb{Z}e'+\mathbb{Z}f'}, l'))$. We may assume that $E = \mathbb{Z}f + \mathbb{Z}l = \mathbb{Z}f' + \mathbb{Z}l'$. Then we can write $f' = \alpha f + \beta l, l' = \gamma f + \delta l$ with $\alpha \delta - \beta \gamma = \pm 1$. Since $(\pm (\delta e - \gamma m), f') = 1$ and $(\pm (\delta e - \gamma m), l') = 0$, we get $(S_{\mathbb{Z}(\pm \delta e \mp \gamma m) + \mathbb{Z}f', l') \simeq (S_{\mathbb{Z}e'+\mathbb{Z}f'}, l')$ by Lemma 4.8 (2). If we set $e'' := \pm (\delta e - \gamma m)$ and $m' := \pm (-\beta e + \alpha m)$, then the correspondence $e \mapsto e'', f \mapsto f', l \mapsto l', m \mapsto m'$ gives an isometry $\varphi \in O(U_1 \oplus U_2)$. The isometry $\widetilde{\varphi} := \varphi \oplus \mathrm{id}_{(U_1 \oplus U_2)^{\perp}} \in O(\widetilde{NS}(S))$ is identity on $A_{\widetilde{NS}(S)} \simeq A_{(U_1 \oplus U_2)^{\perp}}$, so that $\widetilde{\varphi} \in \Gamma_S$. It follows that

$$(S_{\mathbb{Z}e+\mathbb{Z}f},l) \stackrel{\varphi}{\simeq} (S_{\mathbb{Z}e''+\mathbb{Z}f'},l') \simeq (S_{\mathbb{Z}e'+\mathbb{Z}f'},l').$$

Corollary 4.11. If det NS(S) is square-free, then every 0-dimensional cusp of $\Gamma_S^+ \setminus \Omega_{\widetilde{NS}(S)}^+$ is standard. Every elliptic fibration on $S' \in FM(S)$ (if exists) has a section. Therefore, μ_1 is bijective in this case. If $rk(NS(S)) \ge 3$ in addition, $\Gamma_S^+ \setminus \Omega_{\widetilde{NS}(S)}^+$ has exactly one 0-dimensional cusp corresponding to S.

Proof. The assertions except the last one are consequences of Corollary 2.6 and Proposition 4.10, while the last assertion is Corollary 2.7 of [5]. \Box

However, the map μ_1 is not injective in general. By Lemma 4.8 (3), it is easily seen that

$$\mu_1^{-1}\left([E]\right) \simeq \Gamma_S^E \setminus \left\{ f \in E \mid \operatorname{div}(f) = 1 \right\}$$
(7)

for $E \in I_2^{st}(\widetilde{NS}(S))$, where $\Gamma_S^E = \{\gamma \in \Gamma_S \mid \gamma(E) = E\}$. The right hand side of (7) is the set of standard cusps appearing in the canonical compactification of the curve $\Gamma_S^E \setminus B_E \simeq \Gamma_S^E \setminus \mathbb{H}$ (cf. (4)). From the identification (7), we observe the following two facts. Firstly, for two elliptic K3 surfaces (S_1, l_1) and (S_2, l_2) with $S_1 \not\simeq S_2$, $\mu_1((S_1, l_1)) = \mu_1((S_2, l_2))$ if and only if the two distinct standard cusps $\mu_0(S_1)$ and $\mu_0(S_2)$ are connected by the 1-dimensional cusp $\mu_1((S_1, l_1))$. Secondly, fixing an elliptic K3 surface (S, l), we have

$$#\left\{ \left[(S,l') \right] \in \mathrm{FM}_{ell}(S) \ \Big| \ \mu_1((S,l')) = \mu_1((S,l)) \right\} = \mathrm{mult}_{\mu_0(S)}\left(\overline{\mu_1((S,l))}\right), \ (8)$$

where $\overline{\mu_1((S,l))}$ is the closure of the 1-dimensional cusp $\mu_1((S,l))$ in $\Gamma_S^+ \setminus \Omega_{\overline{NS}(S)}^+$, and $\operatorname{mult}_x(C)$ is the multiplicity of the curve C at the point x. Hence $\operatorname{FM}_{ell}(S)$ carries informations about canonical compactifications of certain 1-dimensional cusps themselves.

We remark that, by Proposition 3.7 and Theorem 4.3, an elliptic fibration on an FM-partner $S' \in FM(S)$ gives rise to a twisted FM-partner of S. If

the elliptic fibration has a section, then $S' \simeq S$, and the resulting twisted FMpartner is also isomorphic to S. In particular, untwisted FM-partner except Sitself can never be obtained in this way.

When two elliptic K3 surfaces are connected by a 1-dimensional cusp, there certainly exists a geometric (but not so direct) relation between them. Let $L \in$ Pic(S) be the line bundle associated to an elliptic fibration, and denote by $l \in$ NS(S) the class of L. Define the isotropic lattice $E := \mathbb{Z}l \oplus H^4(S, \mathbb{Z})$. Assume we are given a hyperbolic plane $\mathbb{Z}e' + \mathbb{Z}f' \subset \widetilde{NS}(S)$ such that $f' \in E$ and $(S, l) \not\simeq$ $(S_{\mathbb{Z}e'+\mathbb{Z}f'}, l')$. Here $l' \in E$ is a generator of $\operatorname{Ker}(\cdot, e')|_E$. For brevity, we write S' instead of $S_{\mathbb{Z}e'+\mathbb{Z}f'}$. The situation is that, two non-isomorphic elliptic K3 surfaces (S, l) and (S', l') are connected by the boundary curve corresponding to E. Let $\pi_1 : S \times S' \to S$ and $\pi_2 : S \times S' \to S'$ be the projections. For a coherent sheaf \mathcal{E} on $S \times S'$, we can associate the Mukai vector $v_{\mathcal{E}} := ch(\mathcal{E})\sqrt{td(S \times S')} \in$ $H^*(S \times S', \mathbb{Z})$. The following proposition gives a way to obtain $l \in NS(S)$ from $l' \in NS(S')$.

Proposition 4.12. (1) There is a coherent sheaf \mathcal{E} on $S \times S'$ such that the cohomological FM transform

$$\Phi_{\mathcal{E}}^{H}: \widetilde{H}(S',\mathbb{Z}) \to \widetilde{H}(S,\mathbb{Z}), \quad a \mapsto \pi_{1*}(v_{\mathcal{E}} \wedge \pi_{2}^{*}a)$$

is a Hodge isometry with $\Phi_{\mathcal{E}}^{H}(H^{4}(S',\mathbb{Z})) = \mathbb{Z}e', \ \Phi_{\mathcal{E}}^{H}(H^{0}(S',\mathbb{Z})) = \mathbb{Z}f'.$ By the construction of S', we may assume that $\Phi_{\mathcal{E}}^{H}$ gives the identification $H^{2}(S',\mathbb{Z}) \simeq (\mathbb{Z}e' + \mathbb{Z}f')^{\perp} \cap \widetilde{H}(S,\mathbb{Z}).$ (In fact, \mathcal{E} induces an equivalence $D^{b}(S') \simeq D^{b}(S).$) (2) Let $L' \in Pic(S')$ be the line bundle representing $l' \in NS(S')$. Then the

line bundle

$$\det(R\pi_{1*}(\mathcal{E}\otimes\pi_2^*L'))\otimes\det(R\pi_{1*}\mathcal{E})^{-1}\in Pic(S)$$

is a multiple of L.

Proof. (1) Firstly, we claim that the $H^0(S, \mathbb{Z})$ component of e' is not 0. Assume e' = (0, m, s). Since $f' \in E = \mathbb{Z}l \oplus H^4(S, \mathbb{Z})$, we can write f' degreewise as $f' = (0, \alpha l, t)$ for some integer α. Then we have $1 = (e', f') = \alpha(m, l)$. It follows that div(l) = 1, so we have $(S, l) \simeq (S', l')$ by Proposition 4.10. This contradicts the assumption. Since (e', e') = 0 and (e', f') = 1, then according to [10], [12] (or Proposition 10.10 of [6]), there exists a coherent sheaf \mathcal{E}' on $S \times S'$ such that the cohomological FM transform $\Phi^H_{\mathcal{E}'}$: $\widetilde{H}(S', \mathbb{Z}) \to \widetilde{H}(S, \mathbb{Z})$ is a Hodge isometry with $\Phi^H_{\mathcal{E}'}(H^4(S', \mathbb{Z})) = \mathbb{Z}e'$. Replacing $\{e', f'\}$ by $\{-e', -f'\}$ if necessary, we may assume that the positive generator v' of $H^4(S', \mathbb{Z})$ satisfies $\Phi^H_{\mathcal{E}'}(v') = e'$. Since (e', f') = 1, the $H^0(S', \mathbb{Z})$ component of $(\Phi^H_{\mathcal{E}'})^{-1}(f')$ is equal to -1. Then, as $(\Phi^H_{\mathcal{E}'})^{-1}(f')$ is isotropic, $(\Phi^H_{\mathcal{E}'})^{-1}(f') = -(1, [M], \frac{[M]^2}{2}) = -ch(M)$ for some line bundle $M \in Pic(S')$. Setting $\mathcal{E} := \mathcal{E}' \otimes \pi_2^* M$, we have $\Phi^H_{\mathcal{E}}(v') = \Phi^H_{\mathcal{E}'}(v' \wedge ch(M)) = \Phi^H_{\mathcal{E}'}(v') = e'$ and $\Phi^H_{\mathcal{E}}(1) = \Phi^H_{\mathcal{E}'}(ch(M)) = -f'$.

(2) We compute

$$c_1\left(\det(R\pi_{1*}(\mathcal{E}\otimes\pi_2^*L'))\right)$$

= $ch_1\left(R\pi_{1*}(\mathcal{E}\otimes\pi_2^*L)\right)$
= the $H^2(S,\mathbb{Z})$ component of $ch\left(R\pi_{1*}(\mathcal{E}\otimes\pi_2^*L')\right)\wedge\sqrt{td(S)}$
= the $H^2(S,\mathbb{Z})$ component of $\Phi_{\mathcal{E}}^H\left(ch(L')\wedge\sqrt{td(S')}\right)$.

The last equation is a consequence of the Grothendieck-Riemann-Roch formula (cf. Corollary 5.29 of [6]). Decomposing $ch(L') \in H(S', \mathbb{Z})$ degreewise, we can write ch(L') = (1, l', 0). Since $\sqrt{td(S')} = (1, 0, 1)$, we have $ch(L') \wedge \sqrt{td(S')} =$ (1, l', 1). Similarly,

$$c_1\left(\det(R\pi_{1*}\mathcal{E})\right) = \text{ the } H^2(S,\mathbb{Z}) \text{ component of } \Phi^H_{\mathcal{E}}\left((1,0,1)\right).$$

Hence,

$$c_1\left(\det(R\pi_{1*}(\mathcal{E}\otimes\pi_2^*L'))\otimes\det(R\pi_{1*}\mathcal{E})^{-1}\right) = \text{ the } H^2(S,\mathbb{Z}) \text{ component of } \Phi_{\mathcal{E}}^H(l').$$

Since the identification $H^2(S',\mathbb{Z}) \simeq (\mathbb{Z}e' + \mathbb{Z}f')^{\perp} \cap \widetilde{H}(S,\mathbb{Z})$ is given by $\Phi_{\mathcal{E}}^H$, $\Phi^H_{\mathcal{E}}(l')$ belongs to $E = \mathbb{Z}l \oplus H^4(S, \mathbb{Z})$. Hence the $H^2(S, \mathbb{Z})$ component of $\Phi^H_{\mathcal{E}}(l')$ lies in $\mathbb{Z}l$.

Note that the statement of Proposition 4.12 is symmetric with respect to (S, l) and (S', l'). In this way, we can construct L' (resp. L) from L (resp. L') via a certain sheaf on $S \times S'$. (Of course, even if a sheaf \mathcal{E} on $S \times S'$ induces a Hodge isometry $\widetilde{H}(S,\mathbb{Z}) \simeq \widetilde{H}(S',\mathbb{Z})$, it is not true in general that the first Chern class of the line bundle $\det(R\pi_{2*}(\mathcal{E}\otimes\pi_1^*L))\otimes\det(R\pi_{2*}\mathcal{E})^{-1}$ is isotropic.) To obtain an elliptic curve on S', we first eliminate the fixed components of the linear system $\left|\det(R\pi_{2*}(\mathcal{E}\otimes\pi_1^*L))^{\pm 1}\otimes\det(R\pi_{2*}\mathcal{E})^{\mp 1}\right|$ and then we take a connected component of generic member of the moving part.

We give an example for which μ_1 is not injective.

Example 4.13. Let S be a K3 surface with $NS(S) \simeq U(r), r > 2$, and denote by $r = \prod_{i=1}^{\tau(r)} p_i^{e_i}$ the prime decomposition of r. Assume that S is generic so that $O_{Hodge}(T(S)) = \{\pm id\}$. Then the followings hold :

- (1) $\mathcal{G}(U(r)) = \{U(r)\}.$
- (2) $\#(\mathrm{FM}(S)) = 2^{\tau(r)-2}\varphi(r)$, where φ is the Euler function.

(3) S has two non-isomorphic elliptic fibrations, whose image by μ_1 are different 1-dimensional cusps.

- (4) $\#\left(\operatorname{FM}_{ell}(S)\right) = 2^{\tau(r)-1}\varphi(r).$ (5) For $E \in I_2^{st}(\widetilde{NS}(S)))$, $\#\mu_1^{-1}(E) = \frac{\varphi(r)}{2}.$ (6) $\#\left(\Gamma_S \setminus I_2^{st}(\widetilde{NS}(S))\right) = 2^{\tau(r)}.$

Proof. Write $U(r) = \mathbb{Z}l + \mathbb{Z}m$ with (l, l) = (m, m) = 0, (l, m) = r. Then $A_{U(r)} = \langle \frac{l}{r} \rangle + \langle \frac{m}{r} \rangle \simeq (\mathbb{Z}/r\mathbb{Z}) \oplus (\mathbb{Z}/r\mathbb{Z})$ with

$$(\frac{l}{r},\frac{l}{r}) \equiv (\frac{m}{r},\frac{m}{r}) \equiv 0 \mod 2\mathbb{Z}, \quad (\frac{l}{r},\frac{m}{r}) \equiv \frac{1}{r} \mod \mathbb{Z}$$

We have $O(U(r)) = \{ \mathrm{id}, -\mathrm{id}, \iota, -\iota \} \simeq (\mathbb{Z}/2\mathbb{Z}) \oplus (\mathbb{Z}/2\mathbb{Z}), \text{ where } \iota(l) = m, \iota(m) = l.$ Therefore $O(U(r))^l = O(U(r))^m = \{ \mathrm{id} \}.$

(1) Let L be an even lattice with sign(L) = (1, 1), A_L ≃ ⟨l/r⟩ + ⟨m/r⟩. Given a basis {l̃, m̃} of L[∨], we have rl̃, rm̃ ∈ L. The inclusions Zrl̃ + Zrm̃ ⊂ L ⊂ L[∨] = Zl̃ + Zm̃ imply that Zrl̃ + Zrm̃ = L. Thus the Gram matrix of L with respect to the basis {rl̃, rm̃} is divisible by r. Then |det(L(1/r))| = #(A_L)/r² = 1, sign(L(1/r)) = (1, 1), and L(1/r) is even so that L(1/r) must be isometric to U.
(2) To calculate #FM(S), we shall calculate O(A_{U(r)}).

Claim 4.14. $O(A_{U(r)}) \simeq \prod_{i=1}^{\tau(r)} O(A_{U(p_i^{e_i})})$. Thus $\#O(A_{U(r)}) = 2^{\tau(r)}\varphi(r)$.

With respect to the basis $\{\frac{l}{r}, \frac{m}{r}\}$ of $(\mathbb{Z}/r\mathbb{Z}) \oplus (\mathbb{Z}/r\mathbb{Z})$,

$$O(A_{U(r)}) = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in GL_2(\mathbb{Z}/r\mathbb{Z}) \mid ad + bc \equiv 1 \mod r, \ ab \equiv cd \equiv 0 \mod r \right\}$$
$$= \prod_{i=1}^{\tau(r)} \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in GL_2(\mathbb{Z}/p_i^{e_i}\mathbb{Z}) \mid ad + bc - 1 \equiv ab \equiv cd \equiv 0 \mod p_i^{e_i} \right\}$$
$$= \prod_{i=1}^{\tau(r)} O(A_{U(p_i^{e_i})}),$$

where the second equality follows from the Chinese Remainder theorem. Direct calculations show that $O(A_{U(p_i^{e_i})}) = \left\{ \begin{pmatrix} a & 0 \\ 0 & a^{-1} \end{pmatrix}; a \in (\mathbb{Z}/p_i^{e_i}\mathbb{Z})^{\times} \right\} \cup \left\{ \begin{pmatrix} 0 & b \\ b^{-1} & 0 \end{pmatrix}; b \in (\mathbb{Z}/p_i^{e_i}\mathbb{Z})^{\times} \right\}$, so that $\#O(A_{U(p_i^{e_i})}) = 2\varphi(p_i^{e_i})$. Since $\varphi(r) = \prod \varphi(p_i^{e_i})$, the second assertion follows.

Now (2) follows from (1), Claim 4.2, and the counting formula for #FM(S) ([5]).

(3) NS(S) has exactly four primitive isotropic elements $\{\pm l, \pm m\}$. We may assume that l, m are classes of effective divisors. Since $W(S) = \{\text{id}\}$, both of l, m are classes of elliptic fibrations by the first part of the proof of Lemma 4.2. The only isometry of NS(S) mapping l to m is ι , which is not compatible with any element of $O_{Hodge}(T(S)) = \{\pm \text{id}\}$ on the discriminant group. Therefore (S, l) and (S, m) are not isomorphic. If there exists $\gamma \in \Gamma_S$ which sends $\mu_1((S, l)) = H^4(S, \mathbb{Z}) \oplus \mathbb{Z}l$ to $\mu_1((S, m)) = H^4(S, \mathbb{Z}) \oplus \mathbb{Z}m$, γ maps $\langle \frac{l}{r} \rangle$ to $\langle \frac{m}{r} \rangle$ on $A_{U(r) \oplus U} \simeq A_{U(r)}$, which does not coincide with $\{\pm \text{id}\}$.

The assertion (4) follows from (2) and (3).

(5) By (1), for each $E \in I_2^{st}(U(r) \oplus U)$, we can find a basis $\{l_1, m_1, e_1, f_1\}$ of $U(r) \oplus U$ such that $E = \mathbb{Z}f_1 + \mathbb{Z}l_1$ and $(l_1, l_1) = (m_1, m_1) = (e_1, e_1) = (f_1, f_1) = (l_1, e_1) = (m_1, e_1) = (m_1, f_1) = 0, (l_1, m_1) = r, (e_1, f_1) = 1.$ Take integers $\alpha, \beta \in \mathbb{Z}$ so that β and $r\alpha$ are coprime to each other. Then there exist $\gamma, \delta \in \mathbb{Z}$ such that $\beta\delta + r\alpha\gamma = 1$. Set $f_2 := \alpha l_1 + \beta f_1 \in E, e_2 := \gamma m_1 + \delta e_1, l_2 := \delta l_1 - r\gamma f_1 \in E, m_2 := \beta m_1 - r\alpha e_1$. We have an isometry $\varphi_{(\alpha,\beta,\gamma,\delta)} \in O(U \oplus U(r))^E$ which maps $e_1(\text{resp. } f_1, l_1, m_1)$ to $e_2(\text{resp. } f_2, l_2, m_2)$. On the discriminant group $A_{U(r)\oplus U}, \varphi_{(\alpha,\beta,\gamma,\delta)}(\frac{m_1}{r}) = \beta \frac{m_1}{r}, \varphi_{(\alpha,\beta,\gamma,\delta)}(\frac{l_1}{r}) = \delta \frac{l_1}{r}$. Note that, the image of δ in $\mathbb{Z}/r\mathbb{Z}$ is uniquely determined by α, β as above. Conversely, given $f_2 = \alpha l_1 + \beta f_1 \in E$ with $\operatorname{div}(f_2) = 1$, we can find $e_2 = \gamma m_1 + \delta e_1 + \epsilon f_1 + \zeta l_1$ with $(f_2, e_2) = 1, (e_2, e_2) = 0$. We have $\beta\delta + r\alpha\gamma = 1$. Then Lemma 4.8 (3) assures that we are allowed to take $e'_2 := \gamma m_1 + \delta e_1$ instead of e_2 . In this case, $l_2 = \delta l_1 - r\gamma f_1$. Setting $m_2 := \beta m_1 - r\alpha e_1$, we get $\varphi_{(\alpha,\beta,\gamma,\delta)} \in O(\widetilde{NS}(S))$ which maps $f_1(\text{resp. } e_1, l_1, m_1)$ to $f_2(\text{resp. } e'_2, l_2, m_2)$. Therefore, each $\mathbb{Z}e_2 + \mathbb{Z}f_2 + \mathbb{Z}l_2 \in \mu_1^{-1}(E)$ can be written as $\varphi_{(\alpha,\beta,\gamma,\delta)}(\mathbb{Z}e_1 + \mathbb{Z}f_1 + \mathbb{Z}l_1)$ for some $\alpha, \beta, \gamma, \delta \in \mathbb{Z}$ with $\beta\delta + r\alpha\gamma = 1$. Recall that $\varphi_{(\alpha,\beta,\gamma,\delta)} \in \Gamma_S \cdot \varphi_{(\alpha',\beta',\gamma',\delta')}$ if and only if $r_{U(r)\oplus U}(\varphi_{(\alpha,\beta,\gamma,\delta)}) = \pm r_{U(r)\oplus U}(\varphi_{(\alpha',\beta',\gamma',\delta')})$. Since $r_{U(r)\oplus U}(\varphi_{(\alpha,\beta,\gamma,\delta)})$ is expressed as the matrix $\binom{\beta \ 0}{0 \ \delta}$, we see that $\Gamma_S \varphi_{(\alpha,\beta,\gamma,\delta)} = \Gamma_S \varphi_{(\alpha',\beta',\gamma',\delta')}$ if and only if $\beta \equiv \pm \beta', \delta \equiv \pm \delta'$ mod $r\mathbb{Z}$. It follows that

$$\mu_1^{-1}(E) \simeq \left\{ [\beta], [\gamma] \in \mathbb{Z}/r\mathbb{Z} \mid [\beta][\gamma] \equiv 1 \in \mathbb{Z}/r\mathbb{Z} \right\} / \{\pm \mathrm{id}\}$$
$$\simeq (\mathbb{Z}/r\mathbb{Z})^{\times} / \{\pm \mathrm{id}\}.$$

The assertion (6) follows from (4) and (5).

5 Remarks on
$$\Gamma^+_S \setminus \Omega^+_{\widetilde{NS}(S)}$$

In this section, we mention two remarks on the modular variety $\Gamma_S^+ \setminus \Omega_{\widetilde{NS}(S)}^+$.

For $\varphi \in \operatorname{Emb}(U, \widetilde{NS}(S))$, the splitting $\widetilde{NS}(S) = NS(S_{\varphi}) \oplus \varphi(U)$ induces the realization of $\Omega^+_{\widetilde{NS}(S)}$ as a tube domain

$$\iota_{\varphi}: NS(S_{\varphi})_{\mathbb{R}} + \sqrt{-1}NS(S_{\varphi})^{+} \xrightarrow{\simeq} \Omega^{+}_{\widetilde{NS}(S)}, \quad y \mapsto \mathbb{C}\Big(\varphi(e) + y - \frac{(y,y)}{2}\varphi(f)\Big).$$

Let us observe briefly the action of $O(NS(S))^{\varphi(f)} \cap \Gamma_S^+$ on the tube domain $NS(S_{\varphi})_{\mathbb{R}} + \sqrt{-1}NS(S_{\varphi})^+$. By Proposition 2.7 we have

$$O(NS(S))^{\varphi(f)} \simeq O(NS(S_{\varphi})) \ltimes NS(S_{\varphi}),$$

and the action of $NS(S_{\varphi})$ is identity on the discriminant group. Take an element $\alpha + \sqrt{-1}\omega \in NS(S_{\varphi})_{\mathbb{R}} + \sqrt{-1}NS(S_{\varphi})^+$. According to the equations (1), $m \in NS(S_{\varphi})$ and $g \in O(NS(S_{\varphi}))$ act as

$$\begin{array}{rcl} m(\alpha+\sqrt{-1}\omega) &=& \alpha+m+\sqrt{-1}\omega,\\ g(\alpha+\sqrt{-1}\omega) &=& g(\alpha)+\sqrt{-1}g(\omega). \end{array} \end{array}$$

There is a normal subgroup $W(S_{\varphi}) \ltimes NS(S_{\varphi}) \subset O(\widetilde{NS}(S))^{\varphi(f)} \cap \Gamma_{S}^{+}$, where $W(S_{\varphi})$ is the Weyl group of $NS(S_{\varphi})$. Since $\overline{\mathcal{A}(S_{\varphi})_{\mathbb{R}}}$ is the fundamental domain for the action of $W(S_{\varphi})$ on $NS(S_{\varphi})^{+}$,

$$\left(NS(S_{\varphi})_{\mathbb{R}}/NS(S_{\varphi})\right) + \sqrt{-1}\,\overline{\mathcal{A}(S_{\varphi})_{\mathbb{R}}} \tag{9}$$

is the fundamental domain for the action of $W(S_{\varphi}) \ltimes NS(S_{\varphi})$ on $NS(S_{\varphi})_{\mathbb{R}} + \sqrt{-1}NS(S_{\varphi})^+$. Hence the fundamental domain for the action of $O(\widetilde{NS}(S))^{\varphi(f)} \cap \Gamma_S^+$ on $NS(S_{\varphi})_{\mathbb{R}} + \sqrt{-1}NS(S_{\varphi})^+$ is a quotient of (9). Next, we explain $\Gamma_S^+ \setminus \Omega_{\widetilde{NS}(S)}^+$ in connection with Stab(S), the space of nu-

Next, we explain $\Gamma_S^+ \setminus \Omega_{\widetilde{NS}(S)}^+$ in connection with Stab(S), the space of numerical locally finite stability conditions on $D^b(S)$, following Bridgeland ([2], [3]). The space Stab(S) admits a right action of $\widetilde{GL}_2^+(\mathbb{R})$, the universal covering of $GL_2^+(\mathbb{R})$, and a left action of $Aut(D^b(S))$. These two actions commute. By definition, the central charge of $\sigma \in Stab(S)$ takes the form

$$Z: E \mapsto \left(\pi(\sigma), ch(E)\sqrt{Td(S)}\right), \ E \in K(D^b(S))$$

for some vector $\pi(\sigma) \in \widehat{NS}(S)_{\mathbb{C}}$. The correspondence $\sigma \mapsto \pi(\sigma)$ defines a continuous map $\pi : Stab(S) \to \widetilde{NS}(S)_{\mathbb{C}}$. Set

$$P(S) := \left\{ \omega \in \widetilde{NS}(S)_{\mathbb{C}} \mid \mathbb{R}\operatorname{Re}\omega + \mathbb{R}\operatorname{Im}\omega \text{ is a positive-definite two-plane} \right\}.$$

Via the isomorphism (2), P(S) is a principal $GL_2^+(\mathbb{R})$ -bundle over $\Omega_{\widetilde{NS}(S)}$ by the projection

$$p: P(S) \to \Omega_{\widetilde{NS}(S)}, \ \omega \mapsto \mathbb{R}\mathrm{Re}\omega + \mathbb{R}\mathrm{Im}\omega.$$

Here the right action of $GL_2^+(\mathbb{R})$ is given by

$$\omega \cdot g := (a \operatorname{Re}\omega + b \operatorname{Im}\omega) + \sqrt{-1}(c \operatorname{Re}\omega + d \operatorname{Im}\omega),$$

where $\omega \in P(S)$ and $g = \begin{pmatrix} a & b \\ c & d \end{pmatrix}^{-1} \in GL_2^+(\mathbb{R})$. Each tube domain realization ι_{φ} induces a section $\Omega_{\widetilde{NS}(S)} \to P(S)$. Denote by $P^+(S) \subset P(S)$ the component lying over $\Omega^+_{\widetilde{NS}(S)}$, and set

$$\Delta(S) := \{ \delta \in NS(S) \, | \, (\delta, \delta) = -2 \}.$$

As an immediate corollary of Bridgeland's results, we have the following description of the modular variety $\Gamma_S^+ \setminus \Omega_{\widetilde{NS}(S)}^+$.

Proposition 5.1. There is a connected component $Stab^{\dagger}(S) \subset Stab(S)$ and a subgroup $Aut^{\dagger}(D^{b}(S)) \subset Aut(D^{b}(S))$ such that $p \circ \pi$ induces the isomorphism

$$Aut^{\dagger}(D^{b}(S))\backslash Stab^{\dagger}(S)/\widetilde{GL}_{2}^{+}(\mathbb{R}) \simeq \Gamma_{S}^{+} \backslash \Big(\Omega_{\widetilde{NS}(S)}^{+} - \bigcup_{\delta \in \Delta(S)} \delta^{\perp}\Big).$$

Proof. According to [2] and [3], π induces the isomorphism

$$Aut^{\dagger}(D^{b}(S))\backslash Stab^{\dagger}(S) \simeq \Gamma_{S}^{+} \backslash \Big(P^{+}(S) - \bigcup_{\delta \in \Delta(S)} \delta^{\perp} \Big).$$

Since π is $\widetilde{GL}_2^+(\mathbb{R})$ -equivariant, we have

$$\begin{aligned} Aut^{\dagger}(D^{b}(S)) \backslash Stab^{\dagger}(S) / \widetilde{GL}_{2}^{+}(\mathbb{R}) &\simeq \Gamma_{S}^{+} \backslash \Big(P^{+}(S) - \bigcup_{\delta \in \Delta(S)} \delta^{\perp} \Big) / GL_{2}^{+}(\mathbb{R}) \\ &\simeq \Gamma_{S}^{+} \backslash \Big(\Omega_{\widetilde{NS}(S)}^{+} - \bigcup_{\delta \in \Delta(S)} \delta^{\perp} \Big). \end{aligned}$$

Acknowledgments

The author would like to express his gratitude to Professor K.-I. Yoshikawa for many helpful advises, discussions, and teaching. The relation between hyperbolic planes and standard cusps was taught by him. The author is grateful to Professor S.Hosono for valuable comments. The author would also like to thank Professors T.Bridgeland and D.Kaledin and Doctors S.Iida and S.Ookubo for answering warmly to his questions.

References

- Baily, W. L., Jr.; Borel, A. Compactification of arithmetic quotients of bounded symmetric domains. Ann. of Math. (2) 84 1966 442–528.
- [2] Bridgeland, T. Stability conditions on K3 surfaces. Duke Math. J. 141 (2008), No.2, 241–291.
- [3] Bridgeland, T. Spaces of stability conditions. math.AG/0611510v1
- [4] Canonaco, A.; Stellari, P. Twisted Fourier-Mukai functors. Adv. Math. 212 (2007), no. 2, 484–503.
- [5] Hosono, S.; Lian, B. H.; Oguiso, K.; Yau, S.-T. Fourier-Mukai number of a K3 surface. Algebraic structures and moduli spaces, 177–192, CRM Proc. Lecture Notes, 38, Amer. Math. Soc., Providence, RI, 2004
- [6] Huybrechts, D. Fourier-Mukai transforms in algebraic geometry. The Clarendon Press, Oxford University Press, Oxford, 2006
- [7] Huybrechts, D.; Stellari, P. Equivalences of twisted K3 surfaces. Math. Ann. 332 (2005), no. 4, 901–936.

- [8] Huybrechts, D.; Stellari, P. Proof of Căldăraru's conjecture. Appendix: "Moduli spaces of twisted sheaves on a projective variety" by K. Yoshioka. Moduli spaces and arithmetic geometry, 31–42, Adv. Stud. Pure Math., 45, Math. Soc. Japan, Tokyo, 2006.
- [9] Mukai, S. Symplectic structure of the moduli space of sheaves on an abelian or K3 surface. Invent. Math. 77 (1984), no. 1, 101–116.
- [10] Mukai, S. On the moduli space of bundles on K3 surfaces. I. Vector bundles on algebraic varieties (Bombay, 1984), 341–413, Tata Inst. Fund. Res. Stud. Math., 11, Tata Inst. Fund. Res., Bombay, 1987.
- [11] Nikulin, V. V. Integral symmetric bilinear forms and some of their geometric applications. Izv. Akad. Nauk SSSR Ser. Mat. 43 (1979), no. 1, 111–177
- [12] Orlov, D. O. Equivalences of derived categories and K3 surfaces. J. Math. Sci. 84 (1997), no. 5, 1361–1381
- [13] Pyateskii-Shapiro, I. I. Automorphic functions and the geometry of classical domains. Gordon and Breach Science Publishers, 1969
- [14] Pyateskii-Shapiro, I. I.; Shafarevich, I. R. Torelli's theorem for algebraic surfaces of type K3. Izv. Akad. Nauk SSSR Ser. Mat. 35 1971 530–572.
- [15] Scattone, F. On the compactification of moduli spaces for algebraic K3 surfaces. Mem. Amer. Math. Soc. 70 (1987), no. 374
- [16] Todorov, A. N. Applications of the Kahler-Einstein-Calabi-Yau metric to moduli of K3 surfaces. Invent. Math. 61 (1980), no. 3, 251–265
- [17] Geometrie des surfaces K3: modules et periodes. Asterisque No. 126 (1985).