Quantum states for perfectly secure secret sharing

Dong Pyo Chi,¹ Jeong Woon Choi,¹ Jeong San Kim,² Taewan Kim,¹ and Soojoon Lee³

 1 Department of Mathematical Sciences, Seoul National University, Seoul 151-742, Korea

² Institute for Quantum Information Science, University of Calgary, Alberta T2N 1N4, Canada

 3 Department of Mathematics and Research Institute for Basic Sciences, Kyung Hee University, Seoul 130-701, Korea

(Dated: November 3, 2018)

In this work, we investigate what kinds of quantum states are feasible to perform perfectly secure secret sharing, and present its necessary and sufficient conditions. We also show that the states are bipartite distillable for all bipartite splits, and hence the states could be distillable into the Greenberger-Horne-Zeilinger state. We finally exhibit a class of secret-sharing states, which have an arbitrarily small amount of bipartite distillable entanglement for a certain split.

PACS numbers: 03.67.Hk, 03.65.Ud, 03.67.Mn

Introduction.— Entanglement has been considered as one of the most crucial resources for quantum communication, which has been shown to be perfectly secure against any interior/exterior eavesdropper. The perfect security seems to be due to the pure entanglement. However, in the case of the quantum key distribution, appropriately defining its perfect security as in Refs. [\[1](#page-3-0), [2\]](#page-3-1), we can see that only pure entanglement does not guarantee the perfect security.

We here focus on another quantum communication protocol, the quantum secret sharing of classical information, originally presented by Hillery *et al.* [\[3](#page-3-2)]. Our question is what kinds of quantum states are feasible to perform the perfectly secure secret sharing (PSSS). In order to answer this question, first of all, it is required to present the conditions for the PSSS.

One of the most important problems in secret sharing of classical information is how to share random bits securely between one dealer, Alice and other players, Bob and Charlie. If each participant would securely share one

TABLE I: Random bits for secret sharing

| | | Alice Bob Charlie |
|--------------|-----|-------------------|
| $\mathbf{0}$ | ' ' | 0 |
| | | |
| | 0 | |
| | | |

of the random bit sequences as in TABLE I, then Alice could secretly make Bob and Charlie share her secret bit.

Thus, for the PSSS, the two following conditions must be satisfied: (i) Probability distributions of all participants' secret bits should be unbiased and perfectly correlated, that is, if we let p_{ijk} be the probability that Alice, Bob, and Charlie get the random bits i, j , and k , respectively, then $p_{000} = p_{011} = p_{101} = p_{110} = 1/4$ and $p_{ijk} = 0$ for other i, j, k . (ii) Eavesdropper should not be able to obtain any information about participants' secret bits.

In this work, according to the two above conditions, we show that $\rho_{ABCA'B'C'}$ is a quantum state for the PSSS if and only if it is of the form

$$
\frac{1}{4} \sum_{\substack{i+j+k \equiv 0 \pmod{2} \\ i'+j'+k' \equiv 0 \pmod{2}}} |ijk\rangle_{ABC} \langle i'j'k'|\otimes U_{ijk}\rho_{A'B'C'}U^{\dagger}_{i'j'k'},\tag{1}
$$

where $\rho_{A'B'C'}$ is a state on subsystem $A'B'C'$, and U_{ijk} 's are unitary operators. We call this form of states in [\(1\)](#page-0-0) the *secret-sharing states*.

We also show that the states are bipartite distillable for all bipartite splits. From the results of Dür *et al.* [\[4](#page-3-3)] we can readily derive the fact that if any n -qubit state has negative partial transposition for all bipartite splits then it is distillable into the Greenberger-Horne-Zeilinger (GHZ) [\[5\]](#page-3-4) state. Hence, the secret-sharing states could be also distillable into the GHZ state.

Furthermore, we show that our results can be generalized into multipartite cases, that is, ρ_n is an *n*-qubit state for the PSSS consisting of one dealer and $n-1$ players if and only if it is of the form

$$
\frac{1}{2^{n-1}} \sum_{\substack{I,J \in \mathbb{Z}_2^n \\ \text{even parity}}} |I\rangle_{A_1 A_2 \cdots A_n} \langle J| \otimes U_I \rho_{A'_1 A'_2 \cdots A'_n} U_J^{\dagger}, \quad (2)
$$

where $\rho_{A'_1A'_2\cdots A'_n}$ is a state on subsystem $A'_1A'_2\cdots A'_n$, and U_I 's are unitary operators, and that ρ_n is bipartite distillable for its all bipartite splits. Hence, as in the three-party case, ρ_n could be distillable into the *n*-qubit GHZ state.

Secret-Sharing States.— We first provide necessary and sufficient conditions for a state to perform the PSSS. Let A , B , and C be qubit systems, and A' , B' , and C' be of arbitrary dimensions. Here AA' , BB' , and CC' are Alice's, Bob's, and Charlie's systems, respectively. Then we obtain the following theorem.

Theorem 1. *Any state is a quantum state for the 3 party PSSS if and only if it is a secret-sharing state of the form in [\(1\)](#page-0-0).*

Proof. We first assume that $\rho_{ABCA'B'C'}$ is a quantum state for the PSSS, and let $|\Psi\rangle_{ABCA'B'C'E}$ be its purifi-

$$
|\Psi\rangle_{ABCA'B'C'E} = \sum_{i,j,k} \sqrt{p_{ijk}} |ijk\rangle_{ABC} |\Psi_{ijk}\rangle_{A'B'C'E},
$$
\n(3)

where E is the system of the eavesdropper, Eve. Since probability distributions of all participants' secret bits are unbiased and perfectly correlated, it is clear that $p_{000} = p_{011} = p_{101} = p_{110} = 1/4$ and $p_{ijk} = 0$ for other i, j, k. Thus, the state $|\Psi\rangle$ becomes

$$
\frac{1}{2} \sum_{i+j+k \equiv 0 \pmod{2}} |ijk\rangle_{ABC} |\Psi_{ijk}\rangle_{A'B'C'E}. \tag{4}
$$

For each of all participants' measurement result ijk , let ρ_{ijk}^E be Eve's state after the measurement. Then $\rho_{ijk}^E = \text{tr}_{A'B'C'}(|\Psi_{ijk}\rangle\langle\Psi_{ijk}|)$. Since Eve cannot obtain any information about participants' secret bit at all, we have $\rho_{000}^E = \rho_{011}^E = \rho_{101}^E = \rho_{110}^E$. For each *i*, *j*, *k*, let $\rho_{ijk}^E = \sum_l \lambda_l |\phi_l\rangle\langle\phi_l|$ be its spectral decomposition. Then it follows from Gisin-Hughston-Jozsa-Wootters (GHJW) theorem [\[6\]](#page-3-5) that for each i, j, k , there are unitary operators U_{ijk} on the system $A'B'C'$ such that

$$
|\Psi_{ijk}\rangle = \sum_{l} \sqrt{\lambda_{l}} U_{ijk} |\psi_{l}\rangle_{A'B'C'} |\phi_{l}\rangle_{E}, \tag{5}
$$

where $|\psi_l\rangle$ forms an orthonormal set for the system $A'B'C'$. Hence, by Eqs. [\(4\)](#page-1-0) and [\(5\)](#page-1-1), $\rho_{ABCA'B'C'}$ is of the form

$$
\frac{1}{4} \sum_{\substack{i+j+k \equiv 0 \pmod{2} \\ i'+j'+k' \equiv 0 \pmod{2}}} |ijk\rangle_{ABC} \langle i'j'k'|\otimes U_{ijk}\rho_{A'B'C'}U_{i'j'k'}^{\dagger},\tag{6}
$$

where $\rho_{A'B'C'} = \sum_l \lambda_l |\psi_l\rangle \langle \psi_l|.$

Conversely, we now assume that a given state $\rho_{ABCA'B'C'}$ is of the form [\(1\)](#page-0-0). Then since for the probabilities p_{ijk} that participants get the bits ijk p_{000} = $p_{011} = p_{101} = p_{110} = 1/4$ and $p_{ijk} = 0$ for other i, j, k, probability distributions for the secret bits are clearly unbiased and perfectly correlated. Thus, it suffices to show that Eve cannot any information about the secret bits, that is, $\rho_{000}^E = \rho_{011}^E = \rho_{101}^E = \rho_{110}^E$, where ρ_{ijk}^E is Eve's state when participants' measurement result is ijk . For convenience, we consider the following block matrix form of $\rho_{ABCA'B'C'}$:

ù.

$$
\frac{1}{4} \begin{pmatrix} X_{000,000} & 0 & 0 & X_{000,011} & 0 & X_{000,101} & X_{000,110} & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ X_{011,000} & 0 & 0 & X_{011,011} & 0 & X_{011,101} & X_{011,110} & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ X_{101,000} & 0 & 0 & X_{101,011} & 0 & X_{101,101} & X_{101,110} & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix},
$$
\n
$$
(7)
$$

where $X_{ijk,i'j'k'} = U_{ijk} \rho_{A'B'C'} U_{i'j'k'}^{\dagger}$. Then we can readily obtain that the trace norm of $X_{ijk,i'j'k'}$ is one, that is, $||X_{ijk,i'j'k'}||_1 = 1$. Let the state in Eq. [\(4\)](#page-1-0) be the purification of $\rho_{ABCA'B'C'}$. Then we have $X_{ijk,i'j'k'} =$ $\text{tr}_E\left(|\Psi_{ijk}\rangle\langle\Psi_{i'j'k'}| \right)$. It follows from straightforward calculations that

$$
||X_{ijk,i'j'k'}||_1 = \text{tr}\left|\sqrt{\rho_{ijk}^E}\sqrt{\rho_{i'j'k'}^E}\right| = F(\rho_{ijk}^E, \rho_{i'j'k'}^E),
$$
\n
$$
\text{where } F \text{ is the fidelity. Since } F(\rho_{ijk}^E, \rho_{i'j'k'}^E) = 1 \text{ for every}
$$
\n
$$
(8)
$$

i ′j $(v_{k'}) = 1$ for every i, j, k , the proof is completed.

We remark that, as seen in the proof of Theorem [1,](#page-0-1) in order to prove its converse, it is sufficient to use that the trace norms of three well-chosen off-diagonal blocks are one, for example, $||X_{000,011}||_1 = ||X_{011,101}||_1 =$ $||X_{101,110}||_1 = 1$. Moreover, any block matrix of the form in [\(7\)](#page-1-2) whose three well-chosen off-diagonal blocks have trace norm 1/4 forms a secret state as follows.

Theorem 2. $\sigma_{ABCA'B'C'}$ *is a state which can be expressed as the following block-matrix form:*

$$
\sigma_{ABCA'B'C'} = \begin{pmatrix}\n\star & 0 & 0 & X & 0 & \star & \star & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\
\star & 0 & 0 & \star & 0 & Y & \star & 0 \\
\hline\n0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\
\star & 0 & 0 & \star & 0 & \star & Z & 0 \\
\star & 0 & 0 & \star & 0 & \star & \star & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 & 0\n\end{pmatrix}, \quad (9)
$$

where $||X||_1 = ||Y||_1 = ||Z||_1 = 1/4$ *if and only if* σABCA′B′C′ *is a secret-sharing state.*

Proof. Since any secret-sharing state is of the form [\(9\)](#page-1-3), it suffices to show that $\sigma_{ABCA'B'C'}$ in [\(9\)](#page-1-3) is a secret-sharing state. For each ijk , let p_{ijk} be the trace of (ijk, ijk) block entry of $\sigma_{ABCA'B'C'}$. Then

$$
|\Psi\rangle = \sqrt{p_{000}} |000\rangle_{ABC} |\Psi_{000}\rangle_{A'B'C'E} + \sqrt{p_{011}} |011\rangle_{ABC} |\Psi_{011}\rangle_{A'B'C'E} + \sqrt{p_{101}} |101\rangle_{ABC} |\Psi_{101}\rangle_{A'B'C'E} + \sqrt{p_{110}} |110\rangle_{ABC} |\Psi_{110}\rangle_{A'B'C'E}
$$
(10)

is its purification, and hence we obtain

$$
X = \sqrt{p_{000}p_{011}} \text{tr}_E (|\Psi_{000}\rangle \langle \Psi_{011}|),
$$

\n
$$
Y = \sqrt{p_{011}p_{101}} \text{tr}_E (|\Psi_{011}\rangle \langle \Psi_{101}|),
$$

\n
$$
Z = \sqrt{p_{101}p_{110}} \text{tr}_E (|\Psi_{101}\rangle \langle \Psi_{110}|).
$$
 (11)

As in the proof of Theorem [1,](#page-0-1) we have

$$
||X||_1 = \sqrt{p_{000}p_{011}}F(\rho_{000}^E, \rho_{011}^E),
$$

\n
$$
||Y||_1 = \sqrt{p_{011}p_{101}}F(\rho_{011}^E, \rho_{101}^E),
$$

\n
$$
||Z||_1 = \sqrt{p_{101}p_{110}}F(\rho_{101}^E, \rho_{110}^E),
$$
\n(12)

where $\rho_{ijk}^E = \text{tr}_{A'B'C'}(|\Psi_{ijk}\rangle\langle\Psi_{ijk}|)$. Since $||X||_1 =$ $||Y||_1 = ||Z||_1 = 1/4$, we have the following inequalities:

$$
\frac{p_{000} + p_{011}}{2} \ge \sqrt{p_{000}p_{011}} \ge \frac{1}{4},
$$

\n
$$
\frac{p_{011} + p_{101}}{2} \ge \sqrt{p_{011}p_{101}} \ge \frac{1}{4},
$$

\n
$$
\frac{p_{101} + p_{110}}{2} \ge \sqrt{p_{101}p_{110}} \ge \frac{1}{4}.
$$
 (13)

It follows from the fact $p_{000} + p_{011} + p_{101} + p_{110} = 1$ that $p_{000} = p_{011} = p_{101} = p_{110} = 1/4$ and $F(\rho_{000}^E, \rho_{011}^E) =$ $F(\rho_{011}^E, \rho_{101}^E) = F(\rho_{101}^E, \rho_{110}^E) = 1$. This implies that $\sigma_{ABCA'B'C'}$ is a state for the PSSS. Therefore, it is a secret-sharing state by Theorem [1.](#page-0-1) □

We note that every private state is distillable [\[7\]](#page-3-6). By employing this note, we now show that every secretsharing state is bipartite distillable for its all bipartite splits.

Theorem 3. *Let* ρ *be a secret-sharing state for the 3 party secret sharing. Then* ρ *is bipartite distillable for its all bipartite splits of the 3 parties.*

Proof. By Theorem [1,](#page-0-1) ρ can be expressed as the form of [\(1\)](#page-0-0) for some state $\rho_{A'B'C'}$ and unitary operators U_{ijk} . Let $CNOT_{ij}$ be the controlled-NOT operation such that i and j represent its control system and target system, respectively. Then applying CNOT_{BC} to ρ and performing the projective measurement on system B with respect to the standard basis $\{|0\rangle, |1\rangle\}$, when the measurement result is r , the resulting state becomes a private state,

$$
\frac{1}{2} \sum_{i,j=0}^{1} |ii\rangle_{AC} \langle jj| \otimes U_{iri} \rho_{A'B'C'} U^{\dagger}_{jrj}, \qquad (14)
$$

which is distillable [\[7\]](#page-3-6). Thus, a given ρ is bipartite distillable for the split \AA ^{\AA}- $BB^{\prime}CC^{\prime}$. Similarly, we can show that ρ is bipartite distillable for the splits BB' -CC'AA' and CC' - $AA'BB'$. П

Generalization into multipartite cases.— We now generalize our results into multipartite cases.

Theorem 4. ρⁿ *is a quantum state for the* n*-party PSSS consisting of one dealer and* n − 1 *players if and only if it is a secret-sharing state of the form in [\(2\)](#page-0-2).*

Proof. Let $|\Psi\rangle$ be a purification of ρ_n as follows:

$$
|\Psi\rangle = \sum_{I \in \mathbb{Z}_2^n} \sqrt{p_I} |I\rangle_{A_1 A_2 \cdots A_n} |\Psi_I\rangle_{A'_1 A'_2 \cdots A'_n E}.
$$
 (15)

As in the case of the 3-party case, it is clear that $p_I = 1/2^{n-1}$ for all I with even parity and $p_J = 0$ for other J. For each of all participants' measurement result I, Eve's state after the measurement, ρ_I^E becomes $\rho_I^E = \text{tr}_{A'_1 A'_2 \cdots A'_n} (|\Psi_I\rangle \langle \Psi_I|).$ Since Eve cannot obtain any information about participants' secret bit at all, all ρ_I^E 's are the same. Thus, by GHJW theorem there are a state $\rho_{A'_1 A'_2 \cdots A'_n}$ and unitary operators U_I on the system $A'_1A'_2 \cdots A'_n$ such that ρ_n is of the form in [\(2\)](#page-0-2) and $\rho_{A'_1 A'_2 \cdots A'_n}$ has the same spectrum as ρ_I^E .

Conversely, assuming that a given state ρ_n is of the form [\(2\)](#page-0-2), it can be readily shown that ρ_n is a state for nparty PSSS, by the same way as the proof of Theorem [1.](#page-0-1) 口

We call the state in (2) the *n*-party secret-sharing state. Remark that, as in Theorem [1](#page-0-1) and Theorem [2,](#page-1-4) any quantum state of the form of $2^n \times 2^n$ block matrix, whose block entries vanish if they are in the rows or columns of odd parity, has well-chosen $2^{n-1} - 1$ off-diagonal block entries of the trace norm $1/2^{n-1}$ if and only if the state is an n-party secret-sharing state.

We now consider the bipartite distillability of the n party secret-sharing states.

Theorem 5. *Any n*-party secret-sharing state ρ_n is bi*partite distillable for all bipartite splits of the* n *parties.*

Proof. We use the mathematical induction on $n \geq 3$. Then if $n = 3$ then this theorem is true by Theorem [3.](#page-2-0) We assume that this theorem is true for $(n - 1)$ -party secret-sharing states. Let P be an arbitrary bipartite split $I_0 - I_1$ of the *n* parties, $\{A_1 A'_1, A_2 A'_2, \ldots, A_n A'_n\}$. Then for $A_j A'_j, A_k A'_k \in I_0$ applying $\text{CNOT}_{A_j A_k}$ to ρ_n and performing the projective measurement on system A_i with respect to the standard basis $\{|0\rangle, |1\rangle\}$, the resulting state becomes an $(n - 1)$ -party secret-sharing state. By the induction hypothesis, the state is bipartite distillable for the split, $I_0 - \{A_j A'_j\}$ - I_1 , and hence ρ_n is also bipartite distillable for I_0 - I_1 . This completes the proof. □

Example.— We construct a class of secret-sharing states in a similar way to one presented in [\[1\]](#page-3-0). Consider the following state:

$$
\rho = a_0 |\psi_0\rangle \langle \psi_0| \otimes \sigma_0 + a_1 |\psi_1\rangle \langle \psi_1| \otimes \sigma_1 + a_2 |\psi_2\rangle \langle \psi_2| \otimes \sigma_2 + a_3 |\psi_3\rangle \langle \psi_3| \otimes \sigma_3, \quad (16)
$$

where

$$
|\psi_0\rangle = \frac{1}{2}(|000\rangle + |011\rangle + |101\rangle + |110\rangle),
$$

\n
$$
|\psi_1\rangle = \frac{1}{2}(|000\rangle + |011\rangle - |101\rangle - |110\rangle),
$$

\n
$$
|\psi_2\rangle = \frac{1}{2}(|000\rangle - |011\rangle + |101\rangle - |110\rangle),
$$

\n
$$
|\psi_3\rangle = \frac{1}{2}(|000\rangle - |011\rangle - |101\rangle + |110\rangle),
$$
 (17)

and the states σ_j have support on orthogonal subspaces. Then one can readily verify that ρ is a 3-party secretsharing state, since $\|a_0\sigma_0 \pm a_1\sigma_1 \pm a_2\sigma_2 \pm a_3\sigma_3\|_1 = 1.$

As in Ref. [\[1](#page-3-0)], one can find a secret-sharing state which can have an arbitrarily small amount of bipartite distillable entanglement for a certain split. In order to find such a state, take $a_0 = a_1$ and $a_2 = a_3$ such that $a_1 + a_2 = 1/2$, and

$$
\sigma_0 = \rho_s \otimes |0\rangle\langle 0|,
$$

\n
$$
\sigma_1 = \rho_a \otimes |0\rangle\langle 0|,
$$

\n
$$
\sigma_2 = \rho_a \otimes |1\rangle\langle 1|,
$$

\n
$$
\sigma_3 = \rho_s \otimes |1\rangle\langle 1|,
$$
\n(18)

where ρ_s and ρ_a and two extreme $d \otimes d$ Werner states

$$
\rho_s = \frac{2}{d^2 + d} P_{sym} = \frac{\mathcal{I} + \mathcal{F}}{d^2 + d},
$$

$$
\rho_a = \frac{2}{d^2 - d} P_{as} = \frac{\mathcal{I} - \mathcal{F}}{d^2 - d} \tag{19}
$$

with the identity operator $\mathcal I$ on the $d \otimes d$ system and the flip operator $\mathcal{F} = \sum_{i,j=0}^{d-1} |ij\rangle\langle ji|$. Then we have $\|\rho^{T_{AA'}}\|_1 = (d+2)/d$. Therefore, since the lognegativity is an upper bound of the distillable entanglement [\[8](#page-3-7)], the bipartite distillable entanglement for the

- [1] K. Horodecki, M. Horodecki, P. Horodecki, and J. Oppenheim, Phys. Rev. Lett. 94, 160502 (2005).
- [2] K. Horodecki, M. Horodecki, P. Horodecki, and J. Oppenheim, [quant-ph/0506189.](http://arxiv.org/abs/quant-ph/0506189)
- [3] M. Hillery, V. Bužek, and A. Berthiaume, Phys. Rev. A 59, 1829 (1999).
- [4] W. Dür, J. I. Cirac, and R. Tarrach, Phys. Rev. Lett. 83, 3562 (1999); W. Dür, and J. I. Cirac, Phys. Rev. A 61, 042314 (2000).
- [5] D.M. Greenberger, M.A. Horne, and A. Zeilinger, in Bell's Theorem, Quantum Theory, and Conceptions of the Uni-

split AA' - $BB'CC'$ can be arbitrarily small by increasing d. Nevertheless, the state is always a secret-sharing state for any d.

In conclusion, we have presented necessary and sufficient conditions for secret-sharing states, and have also shown that any secret-sharing state is bipartite distillable for its all bipartite splits, and hence the states could be distillable into the GHZ state. We have furthermore generalized our results into multipartite cases, and have exhibited a class of secret-sharing states, which have an arbitrarily small amount of bipartite distillable entanglement for a certain split.

D.P.C. was supported by the Korea Science and Engineering Foundation (KOSEF) grant funded by the Korea government (MOST) (No. R01-2006-000-10698-0), J.S.K was supported by Alberta's informatics Circle of Research Excellence (iCORE), and S.L. was supported by the Korea Research Foundation Grant funded by the Korean Government (MOEHRD, Basic Research Promotion Fund) (KRF-2007-331-C00049).

verse edited by M. Kafatos (Kluwer, Dordrecht) p. 69 (1989).

- [6] N. Gisin, Helv. Phys. Acta 62, 363 (1989); L.P. Hughston, R. Jozsa, and W.K. Wootters, Phys. Lett. A 183, 14 (1993).
- [7] P. Horodecki and R. Augusiak, Phys. Rev. A 74, 010302 (2006).
- [8] G. Vidal and R.F. Werner, Phys. Rev. A 65, 032314 (2002).