Decoupling of Eulerian and Lagrangian variables in Lagrangian velocity correlations.

Moshe Schwartz

Raymond and Beverly School of Physics and Astronomy

Tel Aviv University, Tel Aviv 69978, Israel

bricki@netvision.net.il

Abstract: The motion of a particle carried by a liquid is described by the differential equation $\dot{\mathbf{r}} = \mathbf{v}(\mathbf{r},t)$, where \mathbf{v} is the Eulerian velocity field of the liquid .Assuming the velocity field to be random with a stationary, isotropic , translational invariant, zero mean distribution, the Lagrangian velocity correlation of the particle can be expressed in terms of the Eulerian correlations and the function $f(q,t) = \langle \exp i \mathbf{q} \cdot \mathbf{r}(t) \rangle$, where the particle is taken to be at the origin at time t = 0 and $\langle \cdots \rangle$ denotes averaging over the Eulerian velocity distribution. This is a result of a decoupling, which is exact to leading order in the volume of the system.

Consider an incompressible liquid confined in a large cubical volume Ω with periodic boundary conditions. Let the Fourier coefficient of the velocity field, $\mathbf{v}(\mathbf{r},t)$, be given by

$$\mathbf{v}(\mathbf{q},t) = \frac{1}{\sqrt{\Omega}} \int d\mathbf{r} \, \mathbf{v}(\mathbf{r},t) \exp(-i\mathbf{q} \cdot \mathbf{r}) \,. \tag{1}$$

The velocity Fourier components of the velocity field be bounded and random with zero average and correlations,

$$\langle \mathbf{v}_{i}(\mathbf{q},t)\mathbf{v}_{j}(\mathbf{p},t')\rangle = [\delta_{ij} - \frac{\mathbf{q}_{i}\mathbf{q}_{j}}{\mathbf{q}^{2}}]\delta_{\mathbf{p},-\mathbf{q}}\Phi(\mathbf{q},t-t').$$
 (2)

Consider next a particle carried by the velocity field that obeys the equation

$$\dot{\mathbf{r}} = \mathbf{v}(\mathbf{r}, t) \,. \tag{3}$$

The problem is to obtain the Lagrangian velocity correlation of the particle,

$$\Lambda(t-t') = \left\langle \dot{\mathbf{r}}(t) \cdot \dot{\mathbf{r}}(t') \right\rangle,\tag{4}$$

where the average is over realizations of the velocity field.

It follows from equation (3) that

$$\Lambda(t-t') = \frac{1}{\Omega} \sum_{\mathbf{q},\mathbf{p}} \langle \mathbf{v}(\mathbf{q},t) \cdot \mathbf{v}(\mathbf{p},t') \exp[i(\mathbf{q} \cdot \mathbf{r}(t) + \mathbf{p} \cdot \mathbf{r}(t'))] \rangle.$$
(5)

Clearly, by the translational invariance of the Eulerian velocity distribution, the right hand side is unchanged if $\mathbf{r}(t)$ and $\mathbf{r}(t')$ are replaced by $\mathbf{r}(t) + \mathbf{d}$ and $\mathbf{r}(t') + \mathbf{d}$. Therefore, performing a spatial average by translating the \mathbf{r} 's by \mathbf{d} , then integrating over \mathbf{d} and dividing by the volume, Ω , still leaves the right hand side unchanged. Consequently,

$$\Lambda(t-t') = \frac{1}{\Omega} \sum_{qp} \langle \mathbf{v}(\mathbf{q},t) \cdot \mathbf{v}(\mathbf{p},t') \exp[i(\mathbf{q} \cdot \mathbf{r}(t) + \mathbf{p} \cdot \mathbf{r}(t'))] \rangle \delta_{\mathbf{p},-\mathbf{q}}, \qquad (6)$$

where δ is the Kronecker δ . Therefore,

$$\Lambda(t-t') = \frac{1}{\Omega} \sum_{\mathbf{q}} \langle \mathbf{v}(\mathbf{q},t) \cdot \mathbf{v}(-\mathbf{q},t') \exp[i(\mathbf{q} \cdot (\mathbf{r}(t) - \mathbf{r}(t')))] \rangle.$$
(7)

Due to invariance under time translations and taking , without loss of generality, ${\bm r}(0) = 0\,,$ finally

$$\Lambda(t) = \frac{1}{\Omega} \sum_{\mathbf{q}} \langle \mathbf{v}(\mathbf{q}, t) \cdot \mathbf{v}(-\mathbf{q}, 0) \exp[i(\mathbf{q} \cdot (\mathbf{r}(t)))] \rangle.$$
(8)

The claim now concerns the average in the sum on the right hand side of equation (8),

$$\langle \mathbf{v}(\mathbf{q},t) \cdot \mathbf{v}(-\mathbf{q},0) \exp[i\mathbf{q} \cdot \mathbf{r}(t)] \rangle = \langle \mathbf{v}(\mathbf{q},t) \mathbf{v}(-\mathbf{q},0) \rangle \langle \exp[i\mathbf{q} \cdot \mathbf{r}(t)] \rangle$$
 (9)

to leading order in Ω .

If the claim is true then the Lagrangian correlations can be written in the limit of the infinite system as

$$\Lambda(t-t') = \frac{1}{(2\pi)^{d}} \int d\mathbf{q} \, [\delta_{ij} - \frac{\mathbf{q}_{i} \mathbf{q}_{j}}{\mathbf{q}^{2}}] \Phi(q,t-t') f(q,t-t') \,. \tag{10}$$

The decoupling given by equation (9) and the resulting equation (10) have been used successfully in the study of a number of problems: The derivation of a Langevin equation [1] and the generalized Langevin equation [2,3] from a Lagrangian describing the interaction of a particle with longitudinal phonons. The diffusion of a deformable object in a random flow [4,5] and the trajectory of a particle swept by a turbulent liquid [6]. In the following I prove that decoupling.

To prove equation (9) consider the dependence of $\exp[i\mathbf{q}\cdot\mathbf{r}(t)]$ on the Fourier components of the velocity field of the liquid with momenta \mathbf{q} and $-\mathbf{q}$. Let us take first the situation, where those components are taken to be identically zero at all times. Let the corresponding solution of equation (3) with the initial condition $\mathbf{r}(0) = 0$ be denoted by $\overline{\mathbf{r}}(\mathbf{q},-\mathbf{q},t)$. Equation (3) will be expressed now in the form

$$\dot{\mathbf{r}} = \frac{1}{\sqrt{\Omega}} \sum_{\mathbf{p} \neq \mathbf{q}, -\mathbf{q}} \mathbf{v}(\mathbf{p}, t) \exp[i\mathbf{p} \cdot \mathbf{r}(t)] + \frac{1}{\sqrt{\Omega}} \mathbf{v}(\mathbf{q}, t) \exp[i\mathbf{q} \cdot \mathbf{r}(t)] + \mathbf{q} \rightarrow -\mathbf{q}.$$
(11)

The solution of the above may be written to order $\,\Omega^{^{-1/2}}\,$ as

$$\mathbf{r}(t) = \overline{\mathbf{r}}(\mathbf{q}, -\mathbf{q}, t) + \frac{\mathbf{a}(t)}{\sqrt{\Omega}},$$
(13)

where

$$\mathbf{a}(t) = \int_{0}^{t} dt' F_{+} \{ \mathbf{v}(\mathbf{p}, t''); \mathbf{q}, t', t \} \mathbf{v}(\mathbf{q}, t') + F_{-} \{ \mathbf{v}(\mathbf{p}, t''); -\mathbf{q}, t', t \} \mathbf{v}(-\mathbf{q}, t')$$
(14)

This is a linear functional of $\mathbf{v}(\mathbf{q},t)$ and $\mathbf{v}(-\mathbf{q},t)$ with coefficients F_+ and F_- which are (non-linear) functionals of all the functions $\mathbf{v}(\mathbf{p},t')$ with $\mathbf{p} \neq \mathbf{q},-\mathbf{q}$.

Consider now the average on the left hand side of equation (9) and perform it in two steps. The first step involves averaging only over $\mathbf{v}(\mathbf{q},t)$ and $\mathbf{v}(-\mathbf{q},t)$. Denote this average by $\langle \cdots \rangle_{\mathbf{q},-\mathbf{q}}$. Since $\overline{\mathbf{r}}(\mathbf{q},-\mathbf{q},t)$ does not depend on $\mathbf{v}(\mathbf{q},t)$ and $\mathbf{v}(-\mathbf{q},t)$,

$$\left\langle \mathbf{v}(\mathbf{q},t) \cdot \mathbf{v}(-\mathbf{q},0) \exp[i\mathbf{q} \cdot \mathbf{r}(t)] \right\rangle_{\mathbf{q},-\mathbf{q}} = \exp[i\mathbf{q} \cdot \overline{\mathbf{r}}(\mathbf{q},-\mathbf{q},t)] \times \left\langle \mathbf{v}(\mathbf{q},t) \cdot \mathbf{v}(-\mathbf{q},0) \exp\frac{1}{\sqrt{\Omega}} [i\mathbf{q} \cdot \mathbf{a}(t)] \right\rangle_{\mathbf{q},-\mathbf{q}}.$$
(15)

It is easy to verify that the average on the right hand side of equation (15) is given by

$$\left\langle \mathbf{v}(\mathbf{q},t) \cdot \mathbf{v}(-\mathbf{q},0) \exp \frac{1}{\sqrt{\Omega}} [i\mathbf{q} \cdot \mathbf{a}(t)] \right\rangle_{\mathbf{q},-\mathbf{q}} = \left\langle \mathbf{v}(\mathbf{q},t) \cdot \mathbf{v}(-\mathbf{q},0) \right\rangle + O(\frac{1}{\Omega}) .$$
 (16)

Consequently, after averaging over the remaining components of the velocity field

$$\langle \mathbf{v}(\mathbf{q},t) \cdot \mathbf{v}(-\mathbf{q},0) \exp[i\mathbf{q} \cdot \mathbf{r}(t)] \rangle = \langle \mathbf{v}(\mathbf{q},t) \cdot \mathbf{v}(-\mathbf{q},0) \rangle \langle \exp[i\mathbf{q} \cdot \overline{\mathbf{r}}(\mathbf{q},-\mathbf{q},t)] \rangle + O(\frac{1}{\Omega}).$$
 (17)

Replacing in all the above considerations $v(q,t) \cdot v(-q,0)$ by its bound, it is easy to see that

$$\langle \exp[i\mathbf{q}\cdot\mathbf{r}(t)]\rangle = \langle \exp[i\mathbf{q}\cdot\overline{\mathbf{r}}(\mathbf{q},-\mathbf{q},t)]\rangle + O(\frac{1}{\Omega}).$$
 (18)

Combining equations (17) and (18) the decoupling (9) is verified and its accuracy is of order $\frac{1}{\Omega}$.

The decoupling given by equation (9) is just a specific case of a more general principle of decoupling, which will be described in the following. Let a function of many

variables, $G_{\Omega}(x_1, x_2,...)$, be absolutely bounded and have the property that for large enough Ω

$$|G_{\Omega}(x_1, x_2, ...) - G_{\Omega}(0, x_2, ...)| < g(\Omega) f(x_1, x_2, ...),$$
(19)

where $g(\Omega)$ tends to zero as Ω tends to infinity and let the variables $\{x_i\}$ be random and independent, then,

$$\lim \Omega \to \infty \langle h(x_1) G_{\Omega}(x_1, x_2, \ldots) \rangle = \langle h(x_1) \rangle \lim \Omega \to \infty \langle G_{\Omega}(x_1, x_2, \ldots) \rangle,$$
(20)

which is very simple to understand.

References:

- (1) M. Schwartz and R. Brustein, Jour. Stat. Phys, 51 (1988) 585.
- (2) R. Brustein, S. Marianer and M. Schwartz, Physica A, 175 (1991) 47.
- (3) G. Frenkel and M. Schwartz , Europhys. Lett. , 50 , (2000) 628.
- (4) G. Frenkel and M. Schwartz , Physica A, 298 , (2001) 278.
- (5) M. Schwartz and G. Frenkel, Phys. Rev. E, 65, (2002) 041104.
- (6) M. Schwartz, G. Frenkel and S. F. Edwards, arXiv:0710.2844, 2007.