## Dynamics of nonlinear resonances in Hamiltonian systems

Miguel D. Bustamante<sup>†</sup> and Elena Kartashova<sup>\*</sup>

<sup>†</sup> Mathematics Institute, University of Warwick, Coventry CV4 7AL, UK

\* RISC, J.Kepler University, Linz 4040, Austria

It is well known that the dynamics of a Hamiltonian system depends crucially on whether or not it possesses nonlinear resonances. In the generic case, the set of nonlinear resonances consists of independent clusters of resonantly interacting modes, described by a few low-dimensional dynamical systems. We show that 1) most frequently met clusters are described by integrable dynamical systems, and 2) construction of clusters can be used as the base for the Clipping method, substantially more effective for these systems than the Galerkin method. The results can be used directly for system with cubic Hamiltonian.

PACS numbers: 47.10.Df, 47.10.Fg, 02.70.Dh

1. Introduction. A notion of resonance runs through all our life. Without resonance we wouldn't have radio, television, music, etc. The general properties of linear resonances are quite well-known; their nonlinear counterpart is substantially less studied though interest in understanding nonlinear resonances is enormous. Famous experiments of Tesla show how disastrous resonances can be: he studied experimentally vibrations of an iron column which ran downward into the foundation of the building, and caused sort of a small earthquake in Manhattan, with smashed windows and swayed buildings [1]. Another example is Tacoma Narrows Bridge which tore itself apart and collapsed (in 1940) under a wind of only 42 mph, though designed for winds of 120 mph.

Nonlinear resonances are ubiquitous in physics. Euler equations, regarded with various boundary conditions and specific values of some parameters, describe an enormous number of nonlinear dispersive wave systems (capillary waves, surface water waves, atmospheric planetary waves, drift waves in plasma, etc.) all possessing nonlinear resonances [2]. Nonlinear resonances appear in a great amount of typical mechanical systems such as an infinite straight bar, a circular ring, and a flat plate [3]. The so-called "nonlinear resonance jump", important for the analysis of a turbine governor positioning system of hydroelectric power plants, can cause severe damage to the mechanical, hydraulic and electrical systems [4]. It was recently established that nonlinear resonance is the dominant mechanism behind outer ionization and energy absorption in near infrared laser-driven rare-gas or metal clusters [5]. The characteristic resonant frequencies observed in accretion disks allow astronomers to determine whether the object is a black hole, a neutron star, or a quark star [6]. Thermally induced variations of the helium dielectric permittivity in superconductors are due to microwave nonlinear resonances [7]. Temporal processing in the central auditory nervous system analyzes sounds using networks of nonlinear neural resonators [8]. The non-linear resonant response of biological tissue to the action of an electromagnetic field is used to investigate cases of suspected disease or cancer [9].

The very special role of resonant solutions of nonlinear ordinary differential equations (ODEs) has been first investigated by Poincaré at the end of the 19th century. Poincaré proved that if a nonlinear ODE has no resonance solutions, then it can be linearized by an invertible change of variables (for details see [10] and refs. therein). This simplifies both analytical and numerical investigations of the original nonlinear equation, allows for the introduction of corresponding normal forms of ODEs, etc. In the middle of the 20th century, Poincaré's approach has been generalized to the case of nonlinear partial differential equations (PDEs) yielding what is nowadays known as KAM theory ([11]–[15] and others). This theory allows us to transform a nonlinear dispersive PDE into a Hamiltonian equation of motion in Fourier space [16],

$$i \dot{a}_{\mathbf{k}} = \partial \mathcal{H} / \partial a_{\mathbf{k}}^*,$$
 (1)

where  $a_{\mathbf{k}}$  is the amplitude of the Fourier mode corresponding to the wavevector  $\mathbf{k}$  and the Hamiltonian  $\mathcal{H}$  is represented as an expansion in powers  $\mathcal{H}_j$  which are proportional to the product of j amplitudes  $a_{\mathbf{k}}$ . In this Letter we are going to consider expansions of Hamiltonians up to third order in wave amplitude, i.e. a cubic Hamiltonian of the form

$$\mathcal{H}_3 = \sum_{\mathbf{k}_1, \mathbf{k}_2, \mathbf{k}_3} V_{23}^1 a_1^* a_2 a_3 \delta_{23}^1 + \text{ complex conj.},$$

where for brevity we introduced the notation  $a_j \equiv a_{\mathbf{k}_j}$ and  $\delta_{23}^1 \equiv \delta(\mathbf{k_1} - \mathbf{k_2} - \mathbf{k_3})$  is the Kronecker symbol. If  $\mathcal{H}_3 \neq 0$ , three-wave process is dominant and the main contribution to the nonlinear evolution comes from the waves satisfying the following resonance conditions:

$$\begin{cases} \omega(\mathbf{k}_1) + \omega(\mathbf{k}_2) - \omega(\mathbf{k}_3) = \Omega, \\ \mathbf{k}_1 + \mathbf{k}_2 - \mathbf{k}_3 = 0, \end{cases}$$

where  $\omega(\mathbf{k})$  is a dispersion relation for the linear wave frequency and  $\Omega \geq 0$  is called resonance width.

If  $\Omega > 0$ , the equation of motion (1) turns into

$$i\dot{A}_{\mathbf{k}} = \omega_{\mathbf{k}}A_{\mathbf{k}} + \sum_{\mathbf{k}_{1},\mathbf{k}_{2}} V_{12}^{\mathbf{k}}A_{1}A_{2}\delta_{12}^{\mathbf{k}} + 2V_{\mathbf{k}2}^{1*}A_{1}A_{2}^{*}\delta_{\mathbf{k}2}^{1}.$$
 (2)

If  $\Omega = 0$ , the equation of motion (1) turns into

$$i\dot{B}_{\mathbf{k}} = \sum_{\mathbf{k}_{1},\mathbf{k}_{2}} \left( V_{12}^{\mathbf{k}} B_{1} B_{2} \delta_{12}^{\mathbf{k}} \delta(\omega_{\mathbf{k}} - \omega_{1} - \omega_{2}) \right. \\ \left. + 2 V_{\mathbf{k}2}^{1*} B_{1} B_{2}^{*} \delta_{\mathbf{k}3}^{1} \delta(\omega_{1} - \omega_{\mathbf{k}} - \omega_{2}) \right).$$
(3)

The co-existence of these two substantially different types of wave interactions, described by Eqs.(2) and (3), has been observed in numerical simulations [17]and proven analytically in the frame of the kinematic two-layer model of laminated turbulence [18]. Dynamics of the layer (2) is described by wave kinetic equations and is well studied [16]. Dynamics of the layer (3)is practically not studied though a lot of preliminary results is already known. Namely, the layer (3) is described by a few independent wave clusters formed by the waves which are in exact nonlinear resonance [19]. The corresponding solutions of (2) can be computed by the specially developed q-class method, presented in [20] and implemented in [21]. The general form of dynamical systems describing resonant clusters can also be found algorithmically [22], as well as coefficients of dynamical systems [23]. Moreover, as it was demonstrated in [24] (numerically) and in [25] (analytically), these clusters "survive" for small enough but non-zero  $\Omega$ . The main goal of this Letter is to study the dynamics of the most frequently met resonant clusters.

2. Clusters. In this Letter we present some analytical and numerical results for the three most commonly met dynamical systems corresponding to non-isomorphic clusters of nonlinear resonances  $-a \ triad$ ,  $a \ kite$ , and  $a \ butterfly$  consisting of 3, 4 and 5 complex variables correspondingly.

The dynamical system for a triad has the form

$$\dot{B}_1 = ZB_2^*B_3, \quad \dot{B}_2 = ZB_1^*B_3, \quad \dot{B}_3 = -ZB_1B_2.$$
 (4)

A kite consists of two triads a and b, with wave amplitudes  $B_{ja}$ ,  $B_{jb}$ , j = 1, 2, 3, connected via two common modes. Analogously to [26], one can point out 4 types of kites according to the properties of connecting modes. For our considerations, this is not important: the general method to study integrability of kites will be the same. For the concreteness of presentation, in this Letter a kite with  $B_{1a} = B_{1b}$  and  $B_{2a} = B_{2b}$  has been chosen:

$$\begin{cases} \dot{B}_{1a} = B_{2a}^*(Z_a B_{3a} + Z_b B_{3b}), \\ \dot{B}_{2a} = B_{1a}^*(Z_a B_{3a} + Z_b B_{3b}), \\ \dot{B}_{3a} = -Z_a B_{1a} B_{2a}, \ \dot{B}_{3b} = -Z_b B_{1a} B_{2a}. \end{cases}$$
(5)

A butterfly consists of two triads a and b, with wave amplitudes  $B_{ja}$ ,  $B_{jb}$ , j = 1, 2, 3, connected via one common mode. As it was shown in [26], there exist 3 different types of butterflies, according to the choice of the connecting mode. Let us take, for instance,  $B_{1a} = B_{1b} (\equiv B_1)$ . The corresponding dynamical system is then as follows:

$$\begin{cases} \dot{B}_1 = Z_a B_{2a}^* B_{3a} + Z_b B_{2b}^* B_{3b} ,\\ \dot{B}_{2a} = Z_a B_1^* B_{3a} , \quad \dot{B}_{2b} = Z_b B_1^* B_{3b} ,\\ \dot{B}_{3a} = -Z_a B_1 B_{2a} , \quad \dot{B}_{3b} = -Z_b B_1 B_{2b} . \end{cases}$$
(6)

3. Integrability of resonance clusters. From here on, general notations and terminology will follow Olver's book [27]. We use hereafter Einstein convention on repeated indices and  $f_{,i} \equiv \partial f / \partial x^i$ . Consider a general *N*-dimensional system of autonomous evolution equations of the form:

$$\frac{dx^i}{dt}(t) = \Delta^i(x^j(t)), \quad i = 1, \dots, N.$$
(7)

Any scalar function  $f(x^i, t)$  that satisfies  $\frac{d}{dt}(f(x^i(t), t)) = \frac{\partial}{\partial t}f + \Delta^i f_{,i} = 0$  is called a conservation law in [27]. It is easy to see that this definition gives us two types of conservation laws. The first type is the standard notion used in classical physics: the conservation law is of the form  $f(x^i)$ , i.e. it does not depend explicitly on time. The second type looks more like a mathematical trick: it is of the form  $f(x^i, t)$ , where the time dependence is explicit. In this Letter we will be interested in both types of conservation law. We claim that they are both physically important and to show that we present an illustrative example from Mechanics. Let us regard the damped harmonic oscillator. The equations of motion in non-dimensional form can be written as:

$$\dot{q} = p, \quad \dot{p} = -q - \alpha p, \tag{8}$$

where  $\alpha \geq 0$  is the damping coefficient. If this coefficient is equal to zero,  $\alpha = 0$ , then the total energy of the system

$$E(q,p) = 1/2 \left( p^2 + q^2 \right)$$

is a conservation law of the first type. If  $\alpha > 0$ , then the system does not conserve the energy anymore but one can still define a conserved quantity which is a generalization of E for the case  $\alpha > 0$ . This new quantity

$$F(q, p, t) = \exp(\alpha t) \left(p^2 + q^2 + \alpha p q\right)$$

is a conservation law of the second type. This means that for an arbitrary solution q(t), p(t) of the system (8) we have

$$\frac{d}{dt} \left[ \exp(\alpha t) \left( p(t)^2 + q(t)^2 + \alpha p(t) q(t) \right) \right] = 0$$

These two types of conservation law are very different but complementary. While the first type, E(q, p) in the case  $\alpha = 0$ , defines where the motion takes place, the second type, F(q, p, t) in the case  $\alpha > 0$ , defines how the motion takes place. In other words, the first type defines orbits of the dynamical system (8) and the second type defines its motion within the orbit.

To keep in mind the difference between these two types of conservation laws, we will call the first type just a conservation law (CL), and we will call the second type a *dynamical invariant*.

We say that Sys.(7) is *integrable* if there are N functionally independent dynamical invariants. Obviously, if Sys.(7) possesses (N - 1) functionally independent CLs, then it is constrained to move along a 1-dimensional manifold, and the way it moves is dictated by 1 dynamical invariant. This dynamical invariant can be obtained from the knowledge of the (N - 1) CLs and the explicit form of the Sys.(7), i.e. Sys.(7) is integrable then. It follows from the Theorem below that in many cases the knowledge of only (N-2) CLs is enough for the integrability of the Sys.(7).

**Theorem.** Let us assume that the system (7) possesses a standard Liouville volume density

$$\rho(x^i): \ (\rho\Delta^i)_{,i} = 0,$$

and (N - 2) functionally independent CLs,  $H^1, \ldots, H^{N-2}$ . Then a new CL can be constructed, which is functionally independent of the original ones, and therefore the system is integrable.

The (lengthy) proof follows from the existence of a Poisson bracket for the original Sys.(7) and is an extension of the general approach used in [28] for three dimensional first order autonomous equations. The proof is constructive and allows us to find the explicit form of a new CL for dynamical systems of the form (4),(5),(6), etc.

In the examples below, we always need to eliminate socalled slave phases, which corresponds to the well-known order reduction in Hamiltonian systems [29]. The number N used below corresponds to the effective number of degrees of freedom after this reduction has been performed.

Integrability of *a triad*, dynamical system (4), is a well-known fact (e.g. [30]) and its two conservation laws are

$$I_{23} = |B_2|^2 + |B_3|^2, \ I_{13} = |B_1|^2 + |B_3|^2.$$

Sys.(4) has been used for a preliminary check of our method; in this case N = 4. The method can thus be applied and we obtain the following CL:

$$I_T = \operatorname{Im}(B_1 B_2 B_3^*)$$

together with the time-dependent dynamical invariant of the form:

$$S_0 = Z t - \frac{F\left(\arcsin\left(\left(\frac{R_3 - v}{R_3 - R_2}\right)^{1/2}\right), \left(\frac{R_3 - R_2}{R_3 - R_1}\right)^{1/2}\right)}{2^{1/2}(R_3 - R_1)^{1/2}(I_{13}^2 - I_{13}I_{23} + I_{23}^2)^{1/4}}.$$

Here F is the elliptic integral of the first kind,  $R_1 < R_2 < R_3$  are the three real roots of the polynomial

$$x^3 + x^2 = 2/27 - (27I_T^2 -$$

$$(I_{13}+I_{23})(I_{13}-2I_{23})(I_{23}-2I_{13}))/27(I_{13}^2-I_{13}I_{23}+I_{23}^2)^{3/2}$$

and

$$v = |B_1|^2 - (2I_{13} - I_{23} + (I_{13}^2 - I_{13}I_{23} + I_{23}^2)^{1/2})/3$$

is always within the interval  $[R_2, R_3] \ni 0$ .

A kite, dynamical system (5), is also an integrable system. Indeed, after reduction of slave variables the system corresponds to N = 6 and has 5 CLs (2 linear, 2 quadratic, 1 cubic):

$$\begin{cases} L_R = \operatorname{Re}(Z_b B_{3a} - Z_a B_{3b}), \ L_I = \operatorname{Im}(Z_b B_{3a} - Z_a B_{3b}), \\ I_{1ab} = |B_{1a}|^2 + |B_{3a}|^2 + |B_{3b}|^2, \\ I_{2ab} = |B_{2a}|^2 + |B_{3a}|^2 + |B_{3b}|^2, \\ I_K = \operatorname{Im}(B_{1a} B_{2a} (Z_a B_{3a}^* + Z_b B_{3b}^*)), \end{cases}$$

with a dynamical invariant that is essentially the same as for a triad,  $S_0$ , after replacing  $Z = Z_a + Z_b, I_T = I_K(Z_a^2 + Z_b^2)/Z^3, I_{13} = I_{1ab}(Z_a^2 + Z_b^2)/Z^2 - (L_R^2 + L_I^2)/Z^2, I_{23} = I_{2ab}(Z_a^2 + Z_b^2)/Z^2 - (L_R^2 + L_I^2)/Z^2.$ 

The dynamics of a *butterfly* is governed by Eqs.(6) and its 4 CLs (3 quadratic and 1 cubic) can easily be obtained:

$$\begin{cases} I_{23a} = |B_{2a}|^2 + |B_{3a}|^2, & I_{23b} = |B_{2b}|^2 + |B_{3b}|^2, \\ I_{ab} = |B_1|^2 + |B_{3a}|^2 + |B_{3b}|^2, \\ I_0 = \operatorname{Im}(Z_a B_1 B_{2a} B_{3a}^* + Z_b B_1 B_{2b} B_{3b}^*), \end{cases}$$
(9)

while a Liouville volume density is  $\rho = 1$ . Notice that all cubic CLs are canonical Hamiltonians for the respective *triad*, *kite* and *butterfly* systems. From now on we consider the *butterfly* case when no amplitude is identically zero; otherwise the system would become integrable.

The use of standard amplitude-phase representation  $B_j = C_j \exp(i\theta_j)$  of the complex amplitudes  $B_j$  in terms of real amplitudes  $C_j$  and phases  $\theta_j$  shows immediately that only two phase combinations are important:

$$\varphi_a = \theta_{1a} + \theta_{2a} - \theta_{3a}, \quad \varphi_b = \theta_{1b} + \theta_{2b} - \theta_{3b},$$

*a*- and *b*-triad phases (with the requirement  $\theta_{1a} = \theta_{1b}$  which correspond to the chosen resonance condition).



FIG. 1: Color online. To facilitate view, color hue of the plot is a linear function of time t, varying from 0 to 1 as t runs through one period. Left panel:  $Z_a = Z_b = 10/100 (integrablecase), I_{ab} \approx 2.1608$ . Middle panel:  $Z_a = 8/100, Z_b = 12/100, I_{ab} \approx 2.2088$ . Right panel:  $Z_a = 9/100, Z_b = 11/100, I_{ab} \approx 2.1846$ .

This reduces five complex equations (6) to only four real ones:

$$\frac{dC_{3a}}{dt} = -Z_a C_1 C_{2a} \cos \varphi_a , \qquad (10)$$

$$\frac{dC_{3b}}{dt} = -Z_b C_1 C_{2b} \cos \varphi_b , \qquad (11)$$

$$\frac{d\varphi_a}{dt} = Z_a C_1 \left( \frac{C_{2a}}{C_{3a}} - \frac{C_{3a}}{C_{2a}} \right) \sin \varphi_a - \frac{I_0}{(C_1)^2} \,, \ (12)$$

$$\frac{d\varphi_b}{dt} = Z_b C_1 \left( \frac{C_{2b}}{C_{3b}} - \frac{C_{3b}}{C_{2b}} \right) \sin \varphi_b - \frac{I_0}{(C_1)^2} \,. \tag{13}$$

The cubic CL reads

$$I_0 = C_1 \left( Z_a C_{2a} C_{3a} \sin \varphi_a + Z_b C_{2b} C_{3b} \sin \varphi_b \right)$$
(14)

in terms of the amplitudes and phases. This means that the dynamics of a butterfly cluster is, in the generic case, confined to a 3-dimensional manifold. Below we regard a few particular cases in which Sys.(6) is integrable.

**Example 1: Real amplitudes,**  $\varphi_a = \varphi_b = 0$ . In this case the Hamiltonian  $I_0$  becomes identically zero while Liouville density in coordinates  $C_{3a}, C_{3b}$  is  $\rho(C_{3a}, C_{3b}) = 1/C_1C_{2a}C_{2b}$ . So in this case the equations for the unknown CL  $H(C_{3a}, C_{3b})$  are:

$$\rho \Delta^{C_{3a}} = -\frac{Z_a}{C_{2b}} = \frac{\partial}{\partial C_{3b}} H, \quad \rho \Delta^{C_{3b}} = -\frac{Z_b}{C_{2a}} = -\frac{\partial}{\partial C_{3a}} H,$$

and from Eqs.(9) we readily obtain

$$H(C_{3a}, C_{3b}) = Z_b \arctan(C_{3a}/C_{2a}) - Z_a \arctan(C_{3b}/C_{2b}),$$

i.e. Sys.(6) is integrable in this case. Of course, this case is degenerate for  $I_0 \equiv 0$  yields no constraint on the remaining independent variables  $C_{3a}, C_{3b}$  satisfying equations (10), (11).

General change of coordinates. Going back to Eqs.(10)-(14), we can jump from this degenerate case to a more generic case by defining new coordinates which

are suggested by  $H(C_{3a}, C_{3b})$ . The new coordinates are to replace the amplitudes  $C_{3a}, C_{3b}$ :

$$\alpha_a = \arctan\left(C_{3a}/C_{2a}\right), \ \ \alpha_b = \arctan\left(C_{3b}/C_{2b}\right).$$

We choose the inverse transformation to be

$$\begin{cases} C_{2a} = \sqrt{I_{23a}} \cos(\alpha_a), & C_{3a} = \sqrt{I_{23a}} \sin(\alpha_a), \\ C_{2b} = \sqrt{I_{23b}} \cos(\alpha_b), & C_{3b} = \sqrt{I_{23b}} \sin(\alpha_b), \end{cases}$$
(15)

so that the domain for the new variables is  $0 < \alpha_a < \pi/2$ ,  $0 < \alpha_b < \pi/2$ . For these new coordinates, the evolution equations simplify enormously:

$$\begin{cases} \frac{d\alpha_a}{dt} = -Z_a C_1 \cos \varphi_a, & \frac{d\alpha_b}{dt} = -Z_b C_1 \cos \varphi_b, \\ \frac{d\varphi_a}{dt} = Z_a C_1 \left( \cot \alpha_a - \tan \alpha_a \right) \sin \varphi_a - \frac{I_0}{(C_1)^2}, \\ \frac{d\varphi_b}{dt} = Z_b C_1 \left( \cot \alpha_b - \tan \alpha_b \right) \sin \varphi_b - \frac{I_0}{(C_1)^2}, \end{cases}$$
(16)

where the amplitude  $C_1 > 0$  is obtained using eqs.(9):

$$C_1 = \sqrt{I_{ab} - I_{23a} \sin^2 \alpha_a - I_{23b} \sin^2 \alpha_b}$$
(17)

and the cubic CL is now

$$I_{0} = \frac{C_{1}}{2} \left( Z_{a} I_{23a} \sin(2\alpha_{a}) \sin(\varphi_{a}) + Z_{b} I_{23b} \sin(2\alpha_{b}) \sin(\varphi_{b}) \right).$$
(18)

Equations (16)-(18) represent the final form of our 3-dimensional general system.

**Example 2: Complex amplitudes,**  $I_0 = 0$ . Here, we just impose the condition  $I_0 = 0$  but the phases are otherwise arbitrary: this case is therefore not degenerate anymore and we have a 3-dimensional system which requires the existence of only 1 CL in order to be integrable: a CL is  $A_a = \sin(2\alpha_a)\sin(\varphi_a)$ , which can be deduced from Eqs.(16). Making use of the Theorem, one can find another CL for this case:  $H_{new}(C_{3a}, C_{3b}) = (1 + Z_b/Z_a) \arccos\left(\frac{\cos 2\alpha_a}{\sqrt{1-A_a^2}}\right) - (1 + Z_a/Z_b) \arccos\left(\frac{\cos 2\alpha_b}{\sqrt{1-A_b^2}}\right)$ .



FIG. 2: Color online. Chain of 4 triads, real amplitudes for all 9 modes. Time and amplitudes in non-dimensioned units.

Obviously  $A_a$  and  $H_{new}$  are functionally independent, i.e. the case  $I_0 = 0$  is integrable.

**Example 3: Complex amplitudes,**  $Z_a = Z_b$ . In this case a new CL has the form

$$\frac{I_0^2}{Z_a}E = C_{2a}^2 C_{3a}^2 + C_{2b}^2 C_{3b}^2 + 2C_{2a}C_{3a}C_{2b}C_{3b}\cos(\varphi_a - \varphi_b) - C_1^2 (C_1^2 - C_{2a}^2 + C_{3a}^2 - C_{2b}^2 + C_{3b}^2), \qquad (19)$$

which is functionally independent of the other known constants of motion. Therefore, according to the Theorem the case  $Z_b = Z_a$  is integrable.

The numerical scheme is programmed in *Mathematica* with *stiffness-switching* method in single precision. For arbitrary  $Z_a, Z_b$  the scheme has been checked by computing  $I_0$  from eq.(18) at all consequent time steps; there is no noticeable change of  $I_0$  up to machine precision.

**4**. Numerical simulations. To investigate the general behavior of a butterfly cluster with  $Z_a \neq Z_b$ , we integrated directly Eqs. (16), (17) with  $I_0$  computed from Eq.(18) evaluated at t = 0 and used to check numerical scheme afterwards. Some results of the simulations with Sys.(16) are presented in Fig.1. Initial conditions  $\alpha_a(0) = 78/100, \alpha_b(0) = 60/100, \varphi_a(0) =$  $147/100, \varphi_b(0) = 127/100$  and values of the constants of motion  $I_0 = 11/2000, I_{23a} = 4/100, I_{23b} = 4/100$  are the same for all three parts of Fig.1. 3D parametric plots are shown in space  $(\alpha_a, \varphi_a, \varphi_b)$  with color hue depending on time, so the plots are effectively 4D. The main goal of this series of numerical simulations was to study changes in the dynamics of a butterfly cluster according to the magnitude of the ratio  $\zeta = Z_a/Z_b$ . In Fig.1, left panel  $\zeta_l = 1$ , middle panel  $\zeta_m = 2/3$  and right panel  $\zeta_r = 9/11$ . As it was shown above, the case  $\zeta_l = 1$  is integrable, and one can see the closed trajectory with period  $T_l \approx 21.7$ . Quite unexpectedly, rational  $\zeta_m, \zeta_r$  produce what appear to be periodic motions and closed trajectories with periods  $T_m\approx 53$  and  $T_r\approx 215$ correspondingly. A few dozen of simulations made with different rational ratios  $\zeta = Z_a/Z_b$  show that periodicity depends - or is even defined by - the commensurability of the coefficients  $Z_a$  and  $Z_b$ . Fig.1 shows that  $\zeta_m = 2/3$ 

gives 2 spikes in one direction and 3 spikes in the perpendicular direction, while  $\zeta_r = 9/11$  gives 9 and 11 spikes correspondingly; and so on. This does not mean, of course, that in the general case  $Z_a \neq Z_b$  solutions of Sys.(6) are periodic. But the opposite can only be proven analytically for in numerical simulations any choice of the ratio  $\zeta$  necessarily turns into a rational number. Some preliminary series of simulations have been performed with the chains of triads with connection types as in (6), with maximum number of triads in a chain being 8 (corresponds to 17 modes). For our numerical simulations, resonant clusters of spherical Rossby waves were taken, with initial (non-dimensioned) energies of the order of measured atmospheric data, as in [31]. The dynamical system for computations was taken in the original variables  $B_i$ . The case of real amplitudes is shown in Fig.2: all resonant modes behave (almost) periodically. Non-dimensioned units for time and amplitudes were chosen to illustrate clearly the characteristic behavior of the amplitudes.

5. Galerkin method versus Clipping method. Speaking very generally, Galerkin method (GM) allows one to reduce infinite dimensional systems, described by PDEs, to low dimensional systems of ODEs. Three steps have to be performed: a) choice of ansatz functions (modes); b) choice of number of modes N; c) construction of the resulting reduced system. While in the case of a *linear* PDE, Fourier modes is the natural choice of the ansatz functions, the number of modes is mostly defined by the available computer facilities and the construction of the corresponding ODE is obvious, the case of a *nonlinear* PDE is much more involved [32].

The Clipping method (CM) has been introduced in [33] in order to deal with evolutionary dispersive nonlinear PDEs. The idea of the CM is very simple: under some physically relevant conditions, it is enough to regard the dynamics of the resonantly interacting modes. These resonant modes can be found systematically using the q-class method [20]. All other modes behave as linear and can be clipped out. Numerical evidence of the effectiveness of this approach was presented in [24], using a pseudo-spectral model of the barotropic vorticity equation. There, it was observed that the most energetically active modes were the resonant modes (221) modes were excited, among them 39 resonant). These resonant modes appear in clusters, i.e. a small number of modes, and the energy of each cluster is conserved (see [24], Fig.1). As for each of the non-resonant modes, their energies are approximately constant during many periods of energy exchange of the resonant modes ([24], Fig.3).

Implementation of the q-class method for various dispersion functions [21], together with explicit construction of the dynamical systems corresponding to the exact resonances [22], allows one to eliminate the arbitrariness of the choices at all three steps a), b) and c) of the GM. Instead of one big system of ODEs of order N, we have a few independent systems of ODEs, of order  $N \ll N$ , and these systems are often integrable. Another example can be found in [22]: ocean planetary motions, 128 resonant modes among 2500 Fourier harmonics in the chosen spectral domain, all clusters and their dynamical systems are written out explicitly. In dozens of studied 3-wave systems, the amount of resonant modes does not exceed 20% of all modes in the spectral domain.

6. Conclusions. Our analysis and general mathematical results [19] on resonant clusters are valid for arbitrary Hamiltonian  $\mathcal{H}_j, j \geq 3$ , though computation of clusters in the case j > 3 is more involved [21].

Clipping method has at least three advantages compared to Galerkin truncation: i) Numerical schemes in CM can be truncated at a substantially higher wavenumber than in GM, depending not on the computer facilities but on some physically relevant parameters (say, dissipation range of wavenumbers). ii) Most of the resulting dynamical systems corresponding to each resonant cluster is analytically integrable. iii) The solutions obtained from the integrable cases could be used to parameterize the numerical solutions of non-integrable systems found for bigger clusters. This work is in progress.

Last not least. Even for one specific PDE, it is a highly non-trivial task to prove that Galerkin truncation is a Hamiltonian system and to construct additional conserved quantity [34]. Clipping method combined with the constructive procedure based on the Theorem above (Section 3), allows us to produce physically relevant dynamical systems and to find additional conservation laws systematically, for a wide class of evolutionary dispersive nonlinear PDEs.

Acknowledgements. E.K. acknowledges the support of the Austrian Science Foundation (FWF) under project P20164-N18 "Discrete resonances in nonlinear wave systems". M.B. acknowledge the support of the Transnational Access Programme at RISC-Linz, funded by European Commission Framework 6 Programme for Integrated Infrastructures Initiatives under the project SCI-Ence (Contract No. 026133).

- [1] Cheney M. Tesla Man Out Of Time. NY, USA: Dorset Press (1989)
- [2] Zakharov, V.S., V.S. Lvov, and G. Falkovich. Kolmogorov Spectra of Turbulence. Springer-Verlag, Berlin (1992)
- [3] Kovriguine, D.A., and G.A. Maugin Mathematical Problems in Engineering (2006)
- [4] Horvat, K., M. Miskovic and O. Kuljaca. Industrial Technology 2: 881 (2003)

- [5] Kundu, M., and D. Bauer. Phys. Rev. Lett. 96: 123401 (2005)
- Kluzniak W. Acta Physica Polonica B 37 (4): 1361 [6](2006)
- Karuzskii, A.L., A.N. Lykov, A.V. Perestoronin and A.I. [7]Golovashkin. Physica C: Superconductivity **408-410**: 739
- Almonte, F., V.K. Jirsa, E.W. Large and B. Tuller. Phys-[8] ica D 212 (1-2): 137-159 (2005)
- Vedruccio, C., E. Mascia and V. Martines. International Review of the Armed Forces Medical Services (IRAFMS), 78 (2): 120 (2005)
- [10] Arnold V.I. Geometrical methods in the theory of ordinary differential equations. Grundleheren der mathematischen Wissenschaften 250 (A Series of Comprehensive Studies in Mathematics) New York Heidelberg Berlin: Springer-Verlag (1983)
- [11] Arnold V.I. Russian Math. Surveys 18: 9 (1963)
- [12]Arnold V.I. Soviet Mathematics 5 (3): 581 (1964)
- [13]Kolmogorov A.N. J. Fluid Mech. 13: 82 (1962)
- [14] Kolmogorov A.N. Dokl. Akad. Nauk SSSR 98: 527 (1954). English translation in 1979: Lecture notes in *Physics* **93**, Springer
- [15] Moser J. Nachr. Akad. Wiss. Goett., Math. Phys. Kl., 1-20(1962)
- [16] Zakharov, V.E., V.S. L'vov and G. Falkovich. Kolmogorov Spectra of Turbulence. Series in Nonlinear Dynamics, Springer (1992)
- [17] Zakharov, V.E., A.O. Korotkevich, A.N. Pushkarev and A.I. Dyachenko. JETP Lett. 82(8): 491 (2005)
- [18] Kartashova E.A. JETP Lett. 83(7): 341 (2006)
- [19] Kartashova E.A. AMS Transl. 2, pp. 95-129 (1998)
- [20] Kartashova, E. Low Temp. Phys. 145 (1-4): 286 (2006)
- [21] Kartashova, E., and A. Kartashov: Int. J. Mod. Phys. C **17**(11): 1579 (2006): Comm. Comp. Phys. **2**(4): 783: (2007); Physica A: Stat. Mech. Appl. 380: 66 (2007)
- [22] Kartashova, E., and G. Mayrhofer. Physica A: Stat. Mech. Appl. 385: 527 (2007)
- [23] Kartashova, E., C. Raab, Ch. Feurer, G. Mayrhofer and W. Schreiner. Extreme Ocean Waves, pp. 97-128. Eds: E. Pelinovsky and Ch. Harif, Springer (2008)
- [24] Kartashova, E. Phys. Rev. Lett. 72: 2013 (1994)
- [25]Kartashova, E. Phys. Rev. Lett. 98(21): 214502 (2007)
- [26] Kartashova, E., and V.S. L'vov. Europhys. Lett. 83: 50012 (2008)
- [27] Olver, P. J. Applications of Lie groups to differential equations. Graduated texts in Mathematics 107, Springer (1993)
- [28] Bustamante, M.D., and S.A. Hojman. J. Phys. A: Math. Gen. 360: 151 (2003)
- Marsden, J. E. and Weinstein, A. Rep. Math. Phys. 5:121 [29](1974)
- [30]Whittaker, E.T., and G.N. Watson. A Course in Mod-
- \* Electronic address: mig busta@yahoo.com, lena@risc.uni-linz.ac.aern Analysis, 4th ed. Cambridge, England: Cambridge University Press (1990)
  - [31] Kartashova, E., and V.S. L'vov. Phys. Rev. Lett. 98(19): 198501 (2007)
  - [32] S. Brenner, S., and R.L. Scott. The Mathematical Theory of Finite Element Methods, 2nd ed. Springer, Berlin Heidelberg New York, (2005)
  - [33]Kartashova, E. Theor. Math. Phys. 99: 675 (1994)
  - [34]Abramov, R., Kovac G., and Majda A.J. Comm. Pure App. Math. 56(1):1 (2003)