Gaussian wave packet solution of the Schrodinger equation in the presence of a time-dependent linear potential

M. Maamache and Y. Saadi

Laboratoire de Physique Quantique et Systèmes Dynamiques, Faculté des Sciences, Université Ferhat Abbas de Sétif, Sétif 19000, Algeria

Abstract

We argue that the way to get the general solution of a Schrodinger equation in the presence of a timedependent linear potential based on the Lewis-Riesenfeld framework is to use a Hermitian linear invariant operator. We demonstrate that the linear invariant proposed in p and q is an Hermitian operator which has the Gaussian wave packet as its eigenfunction.

PACS: 03.65.Ge, 03.65.Fd

The time evolution of a quantum system subject to a spatially uniform, time-dependent force has attracted considerable interest lately. The exact propagator for this system has long been known [1], as have a set of exact solutions (the Volkov solutions) [2-3]. Recent work [4–13] focuses further on exact solutions and their properties. First, Guedes [5] obtained, by means of the invariant operator method introduced by Lewis and Riesenfeld (L-R) [14], a special solution for the time-dependent linear potential. The idea is that any operator satisfying the quantum Liouville-von Neumann equation provides its eigenstate as a solution of the time-dependent Schrodinger equation up to a time dependent phase factor. Later on, Feng [6] followed a method based on spatiotemporal transformations of the Schrodinger equation to get the plane-wave-type and the Airy-packet solutions where the one in Ref. [5] constitutes a particular case which corresponds to the so-called "standing" particle case in a linear potential [6].

However, Bekkar et al. [8] pointed out that the Airy-packet solution is in fact only a superposition of the plane-wave-type solution. In his Comment [7], Bauer explained that the solution found by Guedes [5] is simply a special case of the Volkov solution, with a zero wave vector k, to the time dependent Schrodinger equation describing a nonrelativistic charged particle moving in an electromagnetic field. Dunkel and Trigger [10] considered the initial minimum-uncertainty Gaussian for a sinusoidally time-dependent linear potential. Bowman [11] investigated the time evolution of the general quantum state for the time-dependent linear system, which was shown to be that of a free-particle state, plus an overall motion arising from the classical force . Sang Pyo Kim [12] has shown that a charged free particle in a constant and/or oscillating electric field has a bounded Gaussian wave packet, which is a coherent state of a one-parameter and time-dependent ground state for the free-particle Hamiltonian. From the test-function method, Gengbiao Lu et al.[13] constructed an exact n wave-packet-train (n GWPT) solution , whose center moves acceleratively along the corresponding classical trajectory.

Very recently, Luan et al. [9] have reexamined the linear invariant proposed by Guedes [5] and indicated that besides the solutions described in Refs. [5-8], a Gaussian wave packet eigenfunction solution can naturally be derived from the LR method if a non-Hermitian linear invariant is used, and they argue that this solution was ruled out before, because the authors in Refs. [5-8] assumed in advance the linear L-R invariant I(t) = A(t)p + B(t)x + C(t) be a Hermitian operator.

The authors [9] stated that in Ref.[5, 8], the Gaussian wave packet eigenfunction solution was ruled out, because the linear LR invariant is supposed to be Hermitian operator, and the *B* parameter was set to zero for hermiticity reason. In fact this is not the reason for setting B = 0, in Ref. [8], it was meant to justify the B = 0 imposed in Ref. [5]. We will show and we will correct in the present paper, that if *B* is different from zero ($B \neq 0$), we still obtain Gaussian wave packet (GWP) solutions. In fact, we could find all the results using the Hermitian invariant operator linear in *p* and *q* with any conditions imposed to get physically acceptable solutions as was done in Luan et al. paper [9]. In the latter approach, the authors obtained solutions labelled by complex parameters and complex eigenvalues, this is ambiguous.

We recall that according to the theory of Lewis and Riesenfeld [14], an invariant is an operator that must necessarily satisfy three requirements: 1. It is Hermitian. 2. It satisfies the Von Neumann equation. 3. Its eigenvalues are real and time-independent (we signal that the Hermiticity of the invariant is one of the essential conditions which makes the eigenvalues of the invariant time-independent). Furthermore, any invariant satisfying these three requirements leads to a complete set of solutions of the corresponding Schrodinger equation. So, a conventional solution is constructed as a linear combination of these solutions.

The problem is to find the solutions of the Schrödinger equation,

$$i\hbar\frac{\partial}{\partial t}\psi(x,t) = H \ \psi(x,t) \tag{1}$$

for the Hamiltonian

$$H(x, p, t) = \frac{1}{2m}p^2 - F(t)x,$$
(2)

where F(t) is a time-dependent function.

According to the theory of Lewis and Riesenfeld [14], a solution of the Schro"dinger equation with a timedependent Hamiltonian is easily found if a nontrivial Hermitian operator I(t) exists and satisfies the invariant equation

$$\frac{dI}{dt} = \frac{\partial I}{\partial t} + \frac{1}{i\hbar} \left[I, H \right] = 0.$$
(3)

Indeed, this equation is equivalent to saying that if $\varphi_{\lambda}(x, t)$ is an eigenfunction of I(t) with a time-independent eigenvalue λ , we can find a solution of the Schrödinger equation in the form $\psi_{\lambda}(x, t) = \exp[i\alpha_{\lambda}(t)]\varphi_{\lambda}(x, t)$ where $\alpha_{\lambda}(t)$ satisfies the eigenvalue equation for the Schrödinger operator,

$$\hbar\dot{\alpha}_{\lambda}\left(t\right)\varphi_{\lambda} = i\left[\hbar\frac{\partial}{\partial t} - H\right]\varphi_{\lambda}.$$
(4)

It turns out that the time-dependent invariant operator takes the linear form [9]

$$I(t) = A(t)p + B(t)x + C(t).$$
 (5)

The invariant equation is satisfied if the time-dependent coefficients are such that

$$A(t) = A_0 - \frac{B_0}{m}t, \qquad B(t) = B_0,$$
(6)

$$C(t) = C_0 - A_0 \int_0^t F(\tau) d\tau + \frac{B_0}{m} \int_0^t F(\tau) \tau d\tau,$$
(7)

where A_0 , B_0 , C_0 are arbitrary real constants.

The eigenstates of I(t) corresponding to time-independent eigenvalues are the solutions of the equation

$$I(t)\varphi_{\lambda}(x,t) = \lambda \ \varphi_{\lambda}(x,t). \tag{8}$$

It is easy to see that the solutions of Eq. (8) are of the form

$$\varphi_{\lambda}(x,t) = \exp\left\{\frac{i}{\hbar} \left[\frac{2\left(\lambda - C\left(t\right)\right)x - B_0 x^2}{2A\left(t\right)}\right]\right\}.$$
(9)

Substituting Eq. (9) into Eq. (4) and accomplishing the integration, we obtain

$$\alpha_{\lambda}(t) = \alpha_{\lambda}(0) - \int_{0}^{t} \left\{ \frac{(\lambda - C(\tau))^{2}}{2m\hbar A(\tau)^{2}} + \frac{iB_{0}}{2mA(\tau)} \right\} d\tau$$
$$= \alpha_{\lambda}(0) - \int_{0}^{t} \left\{ \frac{(\lambda - C(\tau))^{2}}{2m\hbar A(\tau)^{2}} \right\} d\tau - i\operatorname{Ln}\left(\sqrt{\frac{A_{0}}{A(t)}}\right),$$
(10)

note that the logarithmic term goes downstairs as a time-dependent normalization factor in $\psi_{\lambda}(x,t)$.

Therefore the physical orthogonal wave functions $\psi_{\lambda}(x,t)$ solutions of the Schrödinger equation (1) are given by

$$\psi_{\lambda}(x,t) = \sqrt{\frac{A_0}{A(t)}} \exp\left\{-i \int_0^t \frac{(\lambda - C(t'))^2}{2\hbar m A(t')^2} dt' + \right\} \exp\left\{\frac{i}{\hbar} \left[\frac{2(\lambda - C(t))x - B_0 x^2}{2A(t)}\right]\right\}.$$
 (11)

It is easy to verify that

$$\langle \psi_{\lambda} \mid \psi_{\lambda'} \rangle = \delta(\lambda - \lambda'). \tag{12}$$

Furthermore, the evolution of the general Schrodinger state can be writen as

$$\Psi(x,t) = \int_{-\infty}^{+\infty} g(\lambda) \psi_{\lambda}(x,t) d\lambda, \qquad (13)$$

where $g(\lambda)$ is a weight function which determines the state of the system such that $\Psi(x,t)$ is square integrable i.e. $\int |\Psi(x,t)| dx$ is time-independent finite constant.

In Ref. [8], it has been shown that the choice

$$g\left(\lambda\right) = \exp\left(\frac{i\lambda^3}{3}\right) \tag{14}$$

leads to Airy function solutions.

Here we shall show that one obtains a general Gaussian wave-packet solution by choosing weight function as a gaussian too,

$$g(\lambda) = \sqrt{\frac{\sqrt{a}}{\hbar A_0 \pi \sqrt{2\pi}}} \exp\left(-a\lambda^2\right),\tag{15}$$

where a is a positive real constant.

Substituting Eq. (11) and (15) into Eq. (13) and accomplishing the integration, we obtain the normalized Gaussian solution as

$$\Psi(x,t) = \sqrt{\frac{\sqrt{a}}{\hbar A(t)\sqrt{2\pi}\left(a+i\int_{0}^{t}\frac{1}{2\hbar m A(t')^{2}}dt'\right)}} \exp\left\{-i\int_{0}^{t}\frac{C(t')^{2}}{2\hbar m A(t')^{2}}dt'\right\}$$
$$\exp\left\{\frac{i}{\hbar}\left[\frac{-2C(t)x-B_{0}x^{2}}{2A(t)}\right]\right\} \exp\left\{\frac{-\left[\int_{0}^{t}\frac{C(t')}{\hbar m A(t')^{2}}dt'+\frac{x}{\hbar A(t)}\right]^{2}}{\left(4a+i\int_{0}^{t}\frac{2}{\hbar m A(t')^{2}}dt'\right)}\right\}$$
(16)

Let us now evaluate the mean value of x, p and the quantum coordinate and momentum fluctuation in the state $\Psi(x,t)$. After some algebra we find that

$$\langle x \rangle = \langle \Psi(t) | x | \Psi(t) \rangle = -A(t) \int_0^t \frac{C(t')}{mA(t')^2} dt', \qquad (17)$$

which is nothing but the classical position $x_c(t)$, and

$$\langle p \rangle = \langle \Psi(t) | p | \Psi(t) \rangle = \frac{-C(t)}{A(t)} + \int_0^t \frac{B_0 C(t')}{m A(t')^2} dt'$$
 (18)

is a classical momentum $p_c(t)$. The position uncertainty

$$\Delta x = \sqrt{\langle x^2 \rangle - \langle x \rangle^2} = \hbar A(t) \sqrt{\frac{\left(a^2 + \left(\int_0^t \frac{1}{2\hbar m A(t')^2} dt'\right)^2\right)}{a}},\tag{19}$$

and the momentum uncertainty

$$\Delta p = \sqrt{\langle p^2 \rangle - \langle p \rangle^2} \\ = \frac{1}{\Delta x} \sqrt{\frac{\hbar^2}{4} + \left[\int_0^t \frac{1}{4a \ mA(t')^2} dt' - \frac{\hbar^2 B_0 A(t)}{a} \left(a^2 + \left(\int_0^t \frac{1}{2\hbar mA(t')^2} dt' \right)^2 \right) \right]^2}, \quad (20)$$

leads to the uncertainty relation

$$\Delta p \Delta x = \sqrt{\frac{\hbar^2}{4} + \left[\int_0^t \frac{1}{4a \ mA(t')^2} dt' - \frac{\hbar^2 B_0 A(t)}{a} \left(a^2 + \left(\int_0^t \frac{1}{2\hbar mA(t')^2} dt'\right)^2\right)\right]^2} \ge \frac{\hbar}{2}.$$
 (21)

We can rewrite (16) as follows

$$\Psi(x,t) = \sqrt{\frac{\hbar A(t) \left(a - i \int_{0}^{t} \frac{1}{2\hbar m A(t')^{2}} dt'\right)}{\sqrt{2\pi a} \Delta x^{2}}} \exp\left\{-i \int_{0}^{t} \frac{C(t')^{2}}{2\hbar m A(t')^{2}} dt'\right\} \\ \exp\left\{\frac{i}{\hbar} \left[\frac{-2C(t) x - B_{0} x^{2}}{2A(t)}\right]\right\} \exp\left\{\frac{-\left(a - i \int_{0}^{t} \frac{1}{2\hbar m A(t')^{2}} dt'\right) [x - \langle x \rangle]^{2}}{4a \Delta x^{2}}\right\}.$$
 (22)

Morever, the time-dependent probability density associated with this Gaussian wave packet is Gaussian for all times

$$\left|\Psi\left(x,t\right)\right|^{2} = \frac{1}{\sqrt{2\pi}\Delta x} \exp\left\{\frac{\left(x-\langle x\rangle\right)^{2}}{2\Delta x^{2}}\right\},\tag{23}$$

we see that Δx represents the width of the wave packet at time t. It is also readily verified tat the time-dependent probability density is conserved

$$\int_{-\infty}^{\infty} \left|\Psi\left(x,t\right)\right|^2 dx = 1 \tag{24}$$

Equations (22) and (23) describe a Gaussian wave packet that is centered at $x = \langle x \rangle$ whose width $\Delta x(t)$ varies with time. So, during time t, the packet's center has moved from x = 0 to $x = -A(t) \int_0^t \frac{C(t')}{mA(t')^2} dt'$ and its width has expanded from $\Delta x_0 = \hbar A_0 \sqrt{a}$ to $\Delta x(t) = \Delta x_0 \frac{A(t)}{A_0} \sqrt{1 + \left(\int_0^t \frac{1}{2\hbar maA(t')^2} dt'\right)^2}$. The wave packet therefore undergoes a distortion; although it remains Gaussian, its width broadens with time whereas its height, $\frac{1}{\sqrt{2\pi\Delta x}}$, decreases with time. Further, it should be noted that the width of the Gaussian packet does not depend on the external force F(t). Thus the shape of the wave packet is not changed by the external force. This means that the external force F(t) acts uniformly in the wave packet.

Acknowledgments:

We wish to thank Professor A. Layadi for his help. References

- [1] R. Feynman and A. Hibbs, Quantum Mechanics and Path Integrals (New York: McGraw-Hill), (1965).
- [2] W. Gordon, Z. Phys. 40, 117 (1926).
- [3] D. Volkov, Z. Phys. 94, 250 (1935).
- [4] A. Rau and K. Unnikrishnan, Phys. Lett. A 222, 304 (1999).
- [5] I. Guedes, Phys. Rev. A 63, 034102 (2001).
- [6] M. Feng, Phys. Rev. A 64, 034101 (2001).
- [7] J. Bauer, Phys. Rev. A 65, 036101 (2002).
- [8] H. Bekkar, F. Benamira, and M. Maamache, Phys. Rev. A 68, 016101 (2003).
- [9] Pi-Gang Luan and Chi-Shung Tang, Phys. Rev. A 71, 014101 (2005).
- [10] J. Dunkel and S. A. Trigger, Phys. Rev.A 71, 052102 (2005).
- [11] G.E. Bowman, J. Phys. A. 39, 157 (2006).
- [12] S. P. Kim, J. Korean Phys. Soc. 44, 464 (2006).
- [13] Gengbiao Lu, Wenhua Hai and Lihua Cai, Phys. Lett. A 357, 181 (2006).
- [14] H. R. Lewis, Jr. and W. B. Reisenfeld, J. Math. Phys. 10, 1458 (1969).