

MOTIVIC INTEGRATION ON TORIC STACKS

A. STAPLEDON

ABSTRACT. We present a decomposition of the space of twisted arcs of a toric stack. As a consequence, we give a combinatorial description of the motivic integral associated to a torus-invariant divisor of a toric stack.

1. INTRODUCTION

Let N be a lattice of rank d and set $N_{\mathbb{R}} = N \otimes_{\mathbb{Z}} \mathbb{R}$. Let Σ be a simplicial, rational, d -dimensional fan in $N_{\mathbb{R}}$, and assume that the support $|\Sigma|$ of Σ is convex. Denote the primitive integer generators of Σ by v_1, \dots, v_r and fix positive integers a_1, \dots, a_r . The data $\mathbf{\Sigma} = (N, \Sigma, \{a_i v_i\})$ is called a *stacky fan*. In [5], Borisov, Chen and Smith associated to $\mathbf{\Sigma}$ a smooth Deligne-Mumford stack $\mathcal{X} = \mathcal{X}(\mathbf{\Sigma})$ over \mathbb{C} , called a *toric stack*, with coarse moduli space the toric variety $X = X(\Sigma)$ (see also [13]).

To any complex variety Y , one can associate its corresponding arc space $J_{\infty}(Y) := \text{Hom}(\text{Spec } \mathbb{C}[[t]], Y)$. The geometry of $J_{\infty}(Y)$ encodes a lot of information about the birational geometry of Y and has been the subject of intensive study in recent times (see, for example, [11, 12, 23]). Kontsevich introduced the theory of *motivic integration* in [21], which assigns a measure to subsets of $J_{\infty}(Y)$ and allows one to compare invariants on birationally equivalent varieties. This theory has been developed over the last decade by many authors including Denef and Loeser [7, 8, 9, 10] and Batyrev [3, 4].

Yasuda extended the theory of arc spaces and motivic integration to Deligne-Mumford stacks in [33] and [34]. More specifically, to any Deligne-Mumford stack \mathcal{Y} , he associated the space $|\mathcal{J}_{\infty}\mathcal{Y}|$ of *twisted arcs* of \mathcal{Y} and defined an associated measure. The goal of this paper is to give an explicit description of the space of twisted arcs of the toric stack \mathcal{X} above. We will prove a decomposition of $|\mathcal{J}_{\infty}\mathcal{X}|$ which allows us to compute motivic integrals on \mathcal{X} and we will show how these motivic integrals relate to combinatorial invariants from the theory of lattice point enumeration of polyhedral complexes.

We briefly recall Ishii's decomposition of the arc space $J_{\infty}(X)$ of the toric variety X in [19] (for more details, see Section 4). If T denotes the torus of X with corresponding arc space $J_{\infty}(T)$, then $J_{\infty}(T)$ has the structure of a group and acts on $J_{\infty}(X)$. If D_1, \dots, D_r denote the T -invariant divisors of X , then the open subset $J_{\infty}(X)' = J_{\infty}(X) \setminus \cup_i J_{\infty}(D_i)$ is invariant under the action of $J_{\infty}(T)$. Ishii gave a decomposition of $J_{\infty}(X)'$ into $J_{\infty}(T)$ -orbits indexed by $|\Sigma| \cap N$ and described the orbit closures.

Our first main result is a stacky analogue of Ishii's decomposition. We describe an action of $J_{\infty}(T)$ on $|\mathcal{J}_{\infty}\mathcal{X}|$, which restricts to an action on $|\mathcal{J}_{\infty}\mathcal{X}'| = |\mathcal{J}_{\infty}\mathcal{X}| \setminus \cup_i |\mathcal{J}_{\infty}\mathcal{D}_i|$, where $\mathcal{D}_1, \dots, \mathcal{D}_r$ are the T -invariant prime divisors of \mathcal{X} (see Section 4). We define a twisted arc $\tilde{\gamma}_v$ for each element v in $|\Sigma| \cap N$, and describe the $J_{\infty}(T)$ -orbits of $|\mathcal{J}_{\infty}\mathcal{X}'|$ (Theorem 4.2), as follows.

Theorem. *We have a decomposition of $|\mathcal{J}_\infty \mathcal{X}'|$ into $J_\infty(T)$ -orbits*

$$|\mathcal{J}_\infty \mathcal{X}'| = \coprod_{v \in |\Sigma| \cap N} \tilde{\gamma}_v \cdot J_\infty(T).$$

Moreover, $\tilde{\gamma}_w \cdot J_\infty(T) \subseteq \overline{\tilde{\gamma}_v \cdot J_\infty(T)}$ if and only if $w - v = \sum_{\rho_i \subseteq \sigma} \lambda_i a_i v_i$ for some non-negative integers λ_i and some cone σ containing v and w .

We now turn our attention to motivic integrals on \mathcal{X} . If $\mathcal{E} = \sum_{i=1}^r b_i \mathcal{D}_i$ is a T -invariant \mathbb{Q} -divisor on \mathcal{X} , then the pair $(\mathcal{X}, \mathcal{E})$ is *Kawamata log terminal* if $b_i < 1$ for $i = 1, \dots, r$. To any Kawamata log terminal pair $(\mathcal{X}, \mathcal{E})$, Yasuda associated the motivic integral $\Gamma(\mathcal{X}, \mathcal{E})$ of \mathcal{E} on \mathcal{X} . In the case when each $a_i = 1$, Theorem 3.41 in [34] implies that $\Gamma(\mathcal{X}, \mathcal{E})$ is equal to Batyrev's stringy invariant of a corresponding Kawamata log terminal pair of varieties [3]. We may view $\Gamma(\mathcal{X}, \mathcal{E})$ either as an element of a Grothendieck ring of varieties [33] or as an element in a ring of convergent stacks [34]. For simplicity, we will specialise $\Gamma(\mathcal{X}, \mathcal{E})$ and view it as a convergent power series in $\mathbb{Z}[(uv)^{1/N}][[(uv)^{-1/N}]]$, for some positive integer N , and refer the reader to section 6 for the relevant details.

In order to motivate and state our results, we recall some notions from the theory of lattice point enumeration of polyhedral complexes (see, for example, [2]). Denote by $\psi : |\Sigma| \rightarrow \mathbb{R}$ the piecewise \mathbb{Q} -linear function satisfying $\psi(a_i v_i) = 1$ for $i = 1, \dots, r$, and consider the polyhedral complex $Q = \{v \in |\Sigma| \mid \psi(v) \leq 1\}$. If for each positive integer m , $f_Q(m)$ denotes the number of lattice points in mQ , then $f_Q(m)$ is a polynomial in m of degree d , called the *Ehrhart polynomial* of Q . The generating series of $f_Q(m)$ can be written in the form

$$1 + \sum_{m \geq 1} f_Q(m) t^m = \delta_Q(t) / (1 - t)^{d+1},$$

where $\delta_Q(t)$ is a polynomial of degree less than or equal to d , called the *Ehrhart δ -polynomial* of Q . More generally, if $\mu : |\Sigma| \rightarrow \mathbb{R}_{\geq 0}$ is a piecewise linear function, one can consider the power series

$$\delta_Q(t; \mu) := (1 - t)^{d+1} \left(1 + \sum_{m \geq 1} \sum_{v \in mQ \cap N} t^{\mu(v)+m} \right),$$

so that $\delta_Q(t; \mu) = \delta_Q(t)$ when μ is identically zero. The Ehrhart δ -polynomial of Q and its associated generalisations have been studied extensively over the last forty years by many authors including Stanley [25, 26, 27, 28] and Hibi [15, 16, 17, 18].

Motivated precisely by computations of motivic integrals on toric stacks, the following variant of the above definition was recently introduced in [30]. If $\lambda : |\Sigma| \rightarrow \mathbb{R}$ is a piecewise \mathbb{Q} -linear function satisfying $\lambda(a_i v_i) > -1$ for $i = 1, \dots, r$, then the *weighted δ -vector* $\delta^\lambda(t)$ is defined by

$$\delta^\lambda(t) = (1 - t)^{d+1} \left(1 + \sum_{m \geq 1} \sum_{v \in mQ \cap N} t^{\psi(v) - \lceil \psi(v) \rceil + \lambda(v) + m} \right).$$

Observe that if λ is a piecewise linear function, then the coefficient of t^i in $\delta_Q(t; \lambda)$ is equal to the sum of the coefficients of t^j in $\delta^\lambda(t)$ for $i - 1 < j \leq i$. The weighted δ -vector satisfies the following symmetry property.

Theorem ([30]). *With the notation above, the expression $\delta^\lambda(t)$ is a rational function in $\mathbb{Q}(t^{1/N})$, for some positive integer N . If Σ is a complete fan, then*

$$\delta^\lambda(t) = t^d \delta^\lambda(t^{-1}).$$

The following result (Theorem 6.5) may be viewed as a geometric realisation of weighted δ -vectors.

Theorem. *Consider a Kawamata log terminal pair $(\mathcal{X}, \mathcal{E})$, where \mathcal{E} is a T -invariant \mathbb{Q} -divisor on the toric stack \mathcal{X} . There exists a corresponding piecewise \mathbb{Q} -linear function $\lambda : |\Sigma| \rightarrow \mathbb{R}$ satisfying $\lambda(a_i v_i) > -1$ for $i = 1, \dots, r$, such that*

$$\Gamma(\mathcal{X}, \mathcal{E}) = (uv)^d \delta^\lambda((uv)^{-1}).$$

In particular, $\Gamma(\mathcal{X}, \mathcal{E})$ is a rational function in $\mathbb{Q}(t^{1/N})$, for some positive integer N . If Σ is a complete fan, then

$$\Gamma(\mathcal{X}, \mathcal{E})(u, v) = (uv)^d \Gamma(\mathcal{X}, \mathcal{E})(u^{-1}, v^{-1}) = \delta^\lambda(uv).$$

Moreover, every weighted δ -vector has the form $\delta^\lambda(uv) = (uv)^d \Gamma(\mathcal{X}, \mathcal{E})(u^{-1}, v^{-1})$, for such a pair $(\mathcal{X}, \mathcal{E})$.

In the case when $\mathcal{E} = 0$, the invariant $\Gamma(\mathcal{X}, 0)$ is a polynomial in $\mathbb{Z}[(uv)^{1/N}]$, for some positive integer N , and the coefficient of $(uv)^j$ is equal to the dimension of the $2j^{\text{th}}$ orbifold cohomology group of \mathcal{X} with compact support [33]. It follows from Poincaré duality for orbifold cohomology [6] and the above discussion that the coefficients of the Ehrhart δ -polynomial of Q are sums of dimensions of orbifold cohomology groups of \mathcal{X} . A detailed discussion of this result is provided in [30].

We conclude the introduction with an outline of the contents of the paper. In Section 2, we recall Borisov, Chen and Smith's notion of a toric stack and give an explicit description of the local construction. We review Yasuda's theory of twisted jets in Section 3 and establish a concrete description of these spaces in the toric case. We use these results in Section 4 to prove our decomposition of the space of twisted arcs of a toric stack. In Section 5, we compute the contact order of a twisted arc along a torus-invariant divisor and, in Section 6, we use the results of the previous sections to compute certain motivic integrals on toric stacks. Finally, in Section 7, we interpret Yasuda's change of variables formula in the toric case to give a geometric proof of a combinatorial result in [30].

The author would like to thank his advisor Mircea Mustață for all his help. He would also like to thank Bill Fulton, Sam Payne and Kevin Tucker for some useful discussions. The author was supported by Mircea Mustață's Packard Fellowship and by an Eleanor Sophia Wood travelling scholarship from the University of Sydney.

2. TORIC STACKS

We fix the following notation throughout the paper. Let N be a lattice of rank d and set $N_{\mathbb{R}} = N \otimes_{\mathbb{Z}} \mathbb{R}$. Let Σ be a simplicial, rational, d -dimensional fan in $N_{\mathbb{R}}$, and let $X = X(\Sigma)$ denote the corresponding toric variety with torus T . We assume that the support $|\Sigma|$ of Σ in $N_{\mathbb{R}}$ is convex. Let ρ_1, \dots, ρ_r denote the rays of Σ , with primitive integer generators v_1, \dots, v_r in N and corresponding T -invariant divisors D_1, \dots, D_r . Fix elements b_1, \dots, b_r in N such that $b_i = a_i v_i$ for some positive integer a_i . The data $\Sigma = (N, \Sigma, \{b_i\})$ is called a *stacky fan*. We denote by $\psi : |\Sigma| \rightarrow \mathbb{R}$ the function that is \mathbb{Q} -linear on each cone of Σ and satisfies $\psi(b_i) = 1$ for $i = 1, \dots, r$. For each non-zero cone τ of Σ , set

$$\text{Box}(\tau) = \{v \in N \mid v = \sum_{\rho_i \subseteq \tau} q_i b_i \text{ for some } 0 < q_i < 1\}.$$

We set $\text{Box}(\{\mathbf{0}\}) = \{0\}$ and $\text{Box}(\Sigma) = \cup_{\tau \in \Sigma} \text{Box}(\tau)$. We will always work over \mathbb{C} and will often identify schemes with their \mathbb{C} -valued points.

Associated to a stacky fan, there is a smooth Deligne-Mumford toric stack $\mathcal{X} = \mathcal{X}(\Sigma)$ over \mathbb{C} with coarse moduli space X [5]. Each cone σ in Σ corresponds to an open substack $\mathcal{X}(\sigma)$ of \mathcal{X} . We identify $\mathcal{X}(\{\mathbf{0}\})$ with the torus T of X . The open substacks $\{\mathcal{X}(\sigma) \mid \dim \sigma = d\}$ give an open covering of \mathcal{X} . We will give an explicit description of $\mathcal{X}(\sigma)$, for a fixed cone σ of dimension d .

If N_σ denotes the sublattice of N generated by $\{b_i \mid \rho_i \subseteq \sigma\}$, then $N(\sigma) = N/N_\sigma$ is a finite group with elements in bijective correspondence with $\coprod_{\tau \subseteq \sigma} \text{Box}(\tau)$. Let M_σ be the dual lattice of N_σ and let M be the dual lattice of N . If σ' denotes the cone in N_σ generated by $\{b_i \mid \rho_i \subseteq \sigma\}$, with corresponding dual cone $(\sigma')^\vee$ in M_σ , then we will make the identifications

$$\text{Spec } \mathbb{C}[(\sigma')^\vee \cap M_\sigma] \cong \mathbb{A}^d, \text{Hom}_{\mathbb{Z}}(M_\sigma, \mathbb{C}^*) \cong (\mathbb{C}^*)^d.$$

If we regard \mathbb{Q}/\mathbb{Z} as a subgroup of \mathbb{C}^* by sending p to $\exp(2\pi\sqrt{-1}p)$, then we have a natural isomorphism

$$(1) \quad N(\sigma) \cong \text{Hom}_{\mathbb{Z}}(\text{Hom}_{\mathbb{Z}}(N(\sigma), \mathbb{Q}/\mathbb{Z}), \mathbb{C}^*) = \text{Hom}_{\mathbb{Z}}(\text{Ext}_{\mathbb{Z}}^1(N(\sigma), \mathbb{Z}), \mathbb{C}^*).$$

We apply the functor $\text{Hom}_{\mathbb{Z}}(\cdot, \mathbb{Z})$ to the exact sequence

$$0 \rightarrow N_\sigma \rightarrow N \rightarrow N(\sigma) \rightarrow 0,$$

to get

$$0 \rightarrow M \rightarrow M_\sigma \rightarrow \text{Ext}_{\mathbb{Z}}^1(N(\sigma), \mathbb{Z}) \rightarrow 0.$$

Applying $\text{Hom}(\cdot, \mathbb{C}^*)$ and the natural isomorphism (1), gives an injection $N(\sigma) \rightarrow (\mathbb{C}^*)^d$. The natural action of $(\mathbb{C}^*)^d$ on \mathbb{A}^d induces an action of $N(\sigma)$ on \mathbb{A}^d . We identify $\mathcal{X}(\sigma)$ with the quotient stack $[\mathbb{A}^d/N(\sigma)]$, with corresponding coarse moduli space the open subscheme $U_\sigma = \mathbb{A}^d/N(\sigma)$ of X [5].

Consider an element g in $N(\sigma)$ of order l corresponding to v in $\text{Box}(\sigma(v))$, where $\sigma(v)$ denotes the cone containing v in its relative interior. We can write $v = \sum_{i=1}^d q_i b_i$, for some $0 \leq q_i < 1$. Note that $q_i \neq 0$ if and only if $\rho_i \subseteq \sigma(v)$. If x_1, \dots, x_d are the coordinates on $\mathbb{A}^d = \text{Spec } \mathbb{C}[(\sigma')^\vee \cap M_\sigma]$ and $\zeta_l = \exp(2\pi\sqrt{-1}/l)$, then one can verify that the action of $N(\sigma)$ on \mathbb{A}^d is given by

$$(2) \quad g \cdot (x_1, \dots, x_d) = (\zeta_l^{lq_1} x_1, \dots, \zeta_l^{lq_d} x_d),$$

and the *age* of g (see Subsection 7.1 [1]) is equal to

$$(3) \quad \text{age}(g) = (1/l) \sum_{i=1}^d lq_i = \psi(v).$$

3. TWISTED JETS OF TORIC STACKS

We use the discussion of twisted jets of Deligne-Mumford stacks in [34] to give an explicit description of the toric case. More specifically, for any non-negative integer n , we describe the stack of twisted n -jets $\mathcal{J}_n \mathcal{X}$ of the toric stack \mathcal{X} .

Fix an affine scheme $S = \text{Spec } R$ over \mathbb{C} and let $D_{n,S}$ denote the affine scheme $\text{Spec } R[t]/(t^{n+1})$. If we fix a positive integer l and consider the group μ_l of l th roots of unity with generator $\zeta_l = \exp(2\pi\sqrt{-1}/l)$, then μ_l acts on $D_{nl,S}$ via the morphism $p : D_{nl,S} \times \mu_l \rightarrow D_{nl,S}$, corresponding to the ring homomorphism

$R[t]/(t^{nl+1}) \rightarrow R[t]/(t^{nl+1}) \otimes \mathbb{C}[x]/(x^l - 1), t \mapsto t \otimes x$. That is, μ_l acts on $D_{nl,S}$ by scaling t . If $\mathcal{D}_{n,S}^l$ denotes the quotient stack $[D_{nl,S}/\mu_l]$, then we have morphisms

$$D_{nl,S} \xrightarrow{\pi} \mathcal{D}_{n,S}^l \rightarrow D_{n,S},$$

such that π is an atlas for $\mathcal{D}_{n,S}^l$ and $D_{n,S}$ is the coarse moduli space of $\mathcal{D}_{n,S}^l$. The composition of the two maps is the quotient of $D_{nl,S}$ by μ_l and corresponds to the ring homomorphism $R[t]/(t^{nl+1}) \rightarrow R[t]/(t^{n+1}), t \mapsto t^l$. The atlas π corresponds to the object α of $\mathcal{D}_{n,S}^l$ over $D_{nl,S}$

$$\begin{array}{ccc} D_{nl,S} \times \mu_l & \xrightarrow{p} & D_{nl,S} \\ \downarrow pr_1 & & \\ D_{nl,S} & & \end{array}$$

and every object in $\mathcal{D}_{n,S}^l$ is locally a pullback of α . Consider the automorphism

$$\theta = \zeta_l \times \zeta_l^{-1} : D_{nl,S} \times \mu_l \rightarrow D_{nl,S} \times \mu_l$$

over $\zeta_l : D_{nl,S} \rightarrow D_{nl,S}$. Every automorphism of α is a power of θ and hence every object and automorphism in $\mathcal{D}_{n,S}^l$ is locally determined by a pullback of α and a power of θ .

A *twisted n -jet of order l* of \mathcal{X} over S is a representable morphism $\mathcal{D}_{n,S}^l \rightarrow \mathcal{X}$. By the above discussion, a twisted jet is determined by the images of α and θ . Yasuda defined the stack $\mathcal{J}_n^l \mathcal{X}$ of twisted n -jets of order l of \mathcal{X} . An object of $\mathcal{J}_n^l \mathcal{X}$ over S is a twisted n -jet $\gamma : \mathcal{D}_{n,S}^l \rightarrow \mathcal{X}$ of order l . Suppose $\gamma' : \mathcal{D}_{n,T}^l \rightarrow \mathcal{X}$ is another twisted n -jet of order l . If $f : S \rightarrow T$ is a morphism, there is an induced morphism $f' : \mathcal{D}_{n,S}^l \rightarrow \mathcal{D}_{n,T}^l$. A morphism in $\mathcal{J}_n^l \mathcal{X}$ from γ to γ' over $f : S \rightarrow T$ is a 2-morphism from γ to $\gamma' \circ f'$. The stack $\mathcal{J}_n \mathcal{X}$ of twisted n -jets is the disjoint union of the stacks $\mathcal{J}_n^l \mathcal{X}$ as l varies over the positive integers. Both $\mathcal{J}_n \mathcal{X}$ and the $\mathcal{J}_n^l \mathcal{X}$ are smooth Deligne-Mumford stacks (Theorem 2.9 [34]). The stack $\mathcal{J}_0 \mathcal{X}$ is identified with the *inertia stack* $\mathcal{I}(\mathcal{X})$ of \mathcal{X} . That is, an object of $\mathcal{J}_0 \mathcal{X}$ over S is determined by a pair (x, α) , where x is an object of \mathcal{X} over S and α is an automorphism of x . We identify the stack $\mathcal{J}_0^1 \mathcal{X}$ with \mathcal{X} . For any $m \geq n$ and for each $l > 0$, the truncation map $R[t]/(t^{ml+1}) \rightarrow R[t]/(t^{nl+1})$ induces a morphism $\mathcal{D}_{n,S}^l \rightarrow \mathcal{D}_{m,S}^l$. Via composition, we get a natural affine morphism (Theorem 2.9 [34])

$$\pi_n^m : \mathcal{J}_m \mathcal{X} \rightarrow \mathcal{J}_n \mathcal{X}.$$

The projective system $\{\mathcal{J}_n \mathcal{X}\}_n$ has a projective limit with a projection morphism (p15 [34])

$$\mathcal{J}_\infty \mathcal{X} = \lim_{\leftarrow} \mathcal{J}_n \mathcal{X}.$$

$$\pi_n : \mathcal{J}_\infty \mathcal{X} \rightarrow \mathcal{J}_n \mathcal{X}.$$

Remark 3.1. The open covering $\{\mathcal{X}(\sigma) \mid \dim \sigma = d\}$ of \mathcal{X} induces an open covering $\{\mathcal{J}_n \mathcal{X}(\sigma) \mid \dim \sigma = d\}$ of $\mathcal{J}_n \mathcal{X}$, for $0 \leq n \leq \infty$.

Recall that if \mathcal{Y} is a stack over \mathbb{C} , then we can consider the set of *points* $|\mathcal{Y}|$ of \mathcal{Y} over \mathbb{C} [22]. Its elements are equivalence classes of morphisms from $\text{Spec } \mathbb{C}$ to \mathcal{Y} . Two morphisms $\psi, \psi' : \text{Spec } \mathbb{C} \rightarrow \mathcal{Y}$ are equivalent if there is a 2-morphism from ψ to ψ' . It has a Zariski topology; for every open substack \mathcal{Y}' of \mathcal{Y} , $|\mathcal{Y}'|$ is

an open subset of $|\mathcal{Y}|$. If \mathcal{Y} has a coarse moduli space Y , then $|\mathcal{Y}|$ is homeomorphic to $Y(\mathbb{C})$. We will often identify Y with $Y(\mathbb{C})$.

Let $D_{\infty, \mathbb{C}} = \text{Spec } \mathbb{C}[[t]]$ and $\mathcal{D}_{\infty, \mathbb{C}}^l = [D_{\infty, \mathbb{C}}/\mu_l]$. A *twisted arc* of order l of \mathcal{X} over \mathbb{C} is a representable morphism from $\mathcal{D}_{\infty, \mathbb{C}}^l$ to \mathcal{X} . Two twisted arcs $\alpha, \alpha' : \mathcal{D}_{\infty, \mathbb{C}}^l \rightarrow \mathcal{X}$ of order l are equivalent if there is a 2-morphism from α to α' . The set of equivalence classes of twisted arcs over \mathcal{X} is identified with $|\mathcal{J}_{\infty} \mathcal{X}|$ (p16 [34]), which we will call the *space of twisted arcs* of \mathcal{X} .

For $0 \leq n \leq \infty$ and a positive integer l , the scheme $D_{n, \mathbb{C}}$ has a unique closed point. A (not necessarily representable) morphism $\gamma : \mathcal{D}_{n, \mathbb{C}}^l \rightarrow \mathcal{X}$ induces an object $\bar{\gamma}$ of \mathcal{X} over \mathbb{C} ,

$$\bar{\gamma} : \text{Spec } \mathbb{C} \rightarrow D_{n, \mathbb{C}} \rightarrow \mathcal{D}_{n, \mathbb{C}}^l \rightarrow \mathcal{X}.$$

The automorphism group of the object in $\mathcal{D}_{n, \mathbb{C}}^l$ corresponding to the morphism $\text{Spec } \mathbb{C} \rightarrow \mathcal{D}_{n, \mathbb{C}}^l$ is μ_l . Hence we get a homomorphism $\phi : \mu_l \rightarrow \text{Aut}(\bar{\gamma})$. Since every automorphism in $\mathcal{D}_{n, \mathbb{C}}^l$ is locally induced by θ , ϕ is injective if and only if $\text{Aut}(\chi) \rightarrow \text{Aut}(\gamma(\chi))$ is injective for all objects χ in $\mathcal{D}_{n, \mathbb{C}}^l$. This holds if and only if γ is representable [22]. We conclude that γ is representable if and only if ϕ is injective.

By considering coarse moduli spaces we have a commutative diagram

$$\begin{array}{ccc} \mathcal{D}_{n, \mathbb{C}}^l & \xrightarrow{\gamma} & \mathcal{X} \\ \downarrow & & \downarrow \\ D_{n, \mathbb{C}} & \xrightarrow{\gamma'} & X. \end{array}$$

We will consider the n th *jet scheme* $J_n(X) = \text{Hom}(D_{n, \mathbb{C}}, X)$ of X and identify jet schemes with their \mathbb{C} -valued points. When $n = \infty$, $J_{\infty}(X)$ is called the *arc space* of X . We have a map

$$(4) \quad \tilde{\pi}_n : |\mathcal{J}_n \mathcal{X}| \rightarrow J_n(X)$$

$$\tilde{\pi}_n(\gamma) = \gamma'.$$

Fix a d -dimensional cone σ of Σ and consider the open substack $\mathcal{X}(\sigma) = [\mathbb{A}^d/N(\sigma)]$ of \mathcal{X} . A twisted n -jet $\mathcal{D}_{n, S}^l \rightarrow \mathcal{X}(\sigma)$ can be lifted to a morphism between atlases, $D_{n, S} \rightarrow \mathbb{A}^d$. Yasuda used these lifts to describe the stack $\mathcal{J}_n^l \mathcal{X}$ (Proposition 2.8 [34]). We will present his result and a sketch of the proof in our situation. We first fix some notation. The action of μ_l on $D_{n, \mathbb{C}}$ extends to an action on $J_{nl}(\mathbb{A}^d)$. The action of $N(\sigma)$ on \mathbb{A}^d also extends to an action on $J_{nl}(\mathbb{A}^d)$. If g is an element of $N(\sigma)$ of order l , let $J_{nl}^{(g)}(\mathbb{A}^d)$ be the closed subscheme of $J_{nl}(\mathbb{A}^d)$ on which the actions of ζ_l in μ_l and g in $N(\sigma)$ agree.

Lemma 3.2 ([34]). *For $0 \leq n \leq \infty$, we have a homeomorphism*

$$|\mathcal{J}_n \mathcal{X}(\sigma)| \cong \coprod_{g \in N(\sigma)} J_{nl}^{(g)}(\mathbb{A}^d)/N(\sigma).$$

Proof. We will only show that there is a bijection between the two sets. We have noted that a representable morphism $\gamma : \mathcal{D}_{n,\mathbb{C}}^l \rightarrow [\mathbb{A}^d/N(\sigma)]$ is determined by the images of α and θ . These images have the form

$$\begin{array}{ccc}
 & & \mathbb{A}^d \\
 & \nearrow^{(v,\lambda) \mapsto \nu(v) \cdot \lambda} & \\
 D_{nl,\mathbb{C}} \times N(\sigma) & \xrightarrow{\zeta_l \times g^{-1}} & D_{nl,\mathbb{C}} \times N(\sigma) \\
 \downarrow pr_1 & & \downarrow pr_1 \\
 D_{nl,\mathbb{C}} & \xrightarrow{\zeta_l} & D_{nl,\mathbb{C}}
 \end{array}$$

for some g in $N(\sigma)$ of order l and some nl -jet $\nu : D_{nl,\mathbb{C}} \rightarrow \mathbb{A}^d$. Conversely, given such a diagram, we can construct a representable morphism. The diagram is determined by any choice of g and ν satisfying

$$\begin{array}{ccc}
 D_{nl,\mathbb{C}} & \xrightarrow{\nu} & \mathbb{A}^d \\
 \downarrow \zeta_l & & \downarrow g \\
 D_{nl,\mathbb{C}} & \xrightarrow{\nu} & \mathbb{A}^d.
 \end{array}$$

That is, γ is determined by the pair (g, ν) , where g has order l and ν lies in $J_{nl}^{(g)}(\mathbb{A}^d)$. Suppose γ' is a twisted n -jet of order l determined by the pair (g', ν') . A 2-morphism from γ to γ' is determined by a morphism $\beta : \gamma(\alpha) \rightarrow \gamma'(\alpha)$ in $[\mathbb{A}^d/N(\sigma)]$ over the identity morphism on $D_{nl,\mathbb{C}}$, such that the following diagram commutes

$$\begin{array}{ccc}
 \gamma(\alpha) & \xrightarrow{\beta} & \gamma'(\alpha) \\
 \downarrow \gamma(\theta) & & \downarrow \gamma'(\theta) \\
 \gamma(\alpha) & \xrightarrow{\beta} & \gamma'(\alpha).
 \end{array}$$

One verifies that the morphism β is determined by an element h in $N(\sigma)$ such that $\nu = \nu' \cdot h$ and that the diagram above is commutative if and only if $g = g'$. Hence the equivalence class of γ in $|\mathcal{J}_n \mathcal{X}(\sigma)|$ corresponds to a point in $J_{nl}^{(g)}(\mathbb{A}^d)/N(\sigma)$. This gives our desired bijection. \square

Remark 3.3. Borisov, Chen and Smith gave a decomposition of $|\mathcal{I}\mathcal{X}|$ into connected components indexed by $\text{Box}(\Sigma)$ (Proposition 4.7 [5]). Given v in $\text{Box}(\Sigma)$, let $\sigma(v)$ be the cone containing v in its relative interior and let $\Sigma_{\sigma(v)}$ be the simplicial fan in $(N/N_{\sigma(v)})_{\mathbb{R}}$ with cones given by the projections of the cones in Σ containing $\sigma(v)$. The associated toric variety $X(\Sigma_{\sigma(v)})$ is a T -invariant closed subvariety of X . The connected component of $|\mathcal{I}\mathcal{X}|$ corresponding to v is homeomorphic to the simplicial toric variety $X(\Sigma_{\sigma(v)})$. By taking $n = 0$ in Lemma 3.2, we recover this decomposition for $|\mathcal{I}\mathcal{X}(\sigma)|$.

Consider an element g in $N(\sigma)$ of order l corresponding to v in $\text{Box}(\tau)$, for some cone τ contained in σ . We will give an explicit description of $J_{nl}^{(g)}(\mathbb{A}^d)$. If we write $v = \sum_{i=1}^d q_i b_i$, for some $0 \leq q_i < 1$, then recall from (2) that the action of g on \mathbb{A}^d is given by

$$g \cdot (x_1, \dots, x_d) = (\zeta_l^{lq_1} x_1, \dots, \zeta_l^{lq_d} x_d).$$

An element ν of $J_{nl}(\mathbb{A}^d)$ can be written in the form

$$\nu = \left(\sum_{j=0}^{nl} \alpha_{1,j} t^j, \dots, \sum_{j=0}^{nl} \alpha_{d,j} t^j \right),$$

for some $\alpha_{i,j}$ in \mathbb{C} , $1 \leq i \leq d$, $0 \leq j \leq nl$. We have

$$g \cdot \nu = \left(\zeta_l^{lq_1} \sum_{j=0}^{nl} \alpha_{1,j} t^j, \dots, \zeta_l^{lq_d} \sum_{j=0}^{nl} \alpha_{d,j} t^j \right)$$

$$\zeta_l \cdot \nu = \left(\sum_{j=0}^{nl} \alpha_{1,j} \zeta_l^j t^j, \dots, \sum_{j=0}^{nl} \alpha_{d,j} \zeta_l^j t^j \right).$$

Hence ν lies in $J_{nl}^{(g)}(\mathbb{A}^d)$ if and only if $\alpha_{i,j} = 0$ whenever $j \not\equiv lq_i \pmod{l}$. We conclude that

$$(5) \quad J_{nl}^{(g)}(\mathbb{A}^d) = \left\{ \left(\sum_{j=0}^{nl} \alpha_{1,j} t^j, \dots, \sum_{j=0}^{nl} \alpha_{d,j} t^j \right) \mid \alpha_{i,j} = 0 \text{ if } j \not\equiv lq_i \pmod{l} \right\}.$$

4. TWISTED ARCS OF TORIC STACKS

In this section we describe an action of $J_\infty(T)$ on an open, dense subset of $|\mathcal{J}_\infty \mathcal{X}|$ and compute the corresponding orbits and orbit closures.

We first recall Ishii's decomposition of the arc space of the toric variety X [19]. Recall that D_1, \dots, D_r denote the T -invariant prime divisors of X and let $J_\infty(X)' = J_\infty(X) \setminus \cup_i J_\infty(D_i)$. The action of T on X extends to an action of $J_\infty(T)$ on $J_\infty(X)'$ (Proposition 2.6 [19]). Given an arc γ in $J_\infty(X)'$, we can find a d -dimensional cone σ such that $\gamma : \text{Spec } \mathbb{C}[[t]] \rightarrow U_\sigma \subseteq X$ corresponds to the ring homomorphism $\gamma^\# : \mathbb{C}[\sigma^\vee \cap M] \rightarrow \mathbb{C}[[t]]$. We have an induced semigroup morphism $\sigma^\vee \cap M \rightarrow \mathbb{N}$, $u \mapsto \text{ord}_t \gamma^\#(\chi^u)$, which extends to homomorphism from M to \mathbb{Z} , necessarily of the form $\langle \cdot, v \rangle$, for some v in $\sigma \cap N$. Consider the arc

$$\gamma_v : \text{Spec } \mathbb{C}[[t]] \rightarrow U_\sigma \subseteq X$$

corresponding to the ring homomorphism

$$\gamma_v^\# : \mathbb{C}[\sigma^\vee \cap M] \rightarrow \mathbb{C}[[t]]$$

$$\chi^u \mapsto t^{\langle u, v \rangle}.$$

If ϕ denotes the arc in $J_\infty(T)$ corresponding to the ring homomorphism $\phi^\# : \mathbb{C}[\sigma^\vee \cap M] \rightarrow \mathbb{C}[[t]]$, $\chi^u \mapsto \gamma^\#(\chi^u)/t^{\langle u, v \rangle}$, then $\gamma = \gamma_v \cdot \phi$ and both v in $|\Sigma| \cap N$ and ϕ in $J_\infty(T)$ are uniquely determined. We have shown the first part of the following theorem.

Theorem 4.1 ([19]). *We have a decomposition of $J_\infty(X)'$ into $J_\infty(T)$ -orbits*

$$J_\infty(X)' = \coprod_{v \in |\Sigma| \cap N} \gamma_v \cdot J_\infty(T).$$

Moreover, $\gamma_w \cdot J_\infty(T) \subseteq \overline{\gamma_v \cdot J_\infty(T)}$ if and only if $w - v$ lies in some cone σ containing v and w .

We will give a similar decomposition of the space $|\mathcal{J}_\infty \mathcal{X}|$ of twisted arcs of \mathcal{X} . For $i = 1, \dots, r$, we first describe a closed substack \mathcal{D}_i of \mathcal{X} with coarse moduli space D_i [5]. Fix a maximal cone σ . If ρ_i is not in σ then \mathcal{D}_i is disjoint from $\mathcal{X}(\sigma)$. Suppose ρ_i is a ray of σ and consider the projection $p_i : N \rightarrow N(\rho_i) = N/N_{\rho_i}$, where N_{ρ_i} is the sublattice of N generated by b_i . If σ_i denotes the cone $p_i(\sigma)$ in $N(\rho_i)$, then $(N(\rho_i), \sigma_i, \{p_i(b_j)\}_{\rho_j \subseteq \sigma})$ is the stacky fan corresponding to $\mathcal{D}_i \cap \mathcal{X}(\sigma)^\perp$. Note that p_i induces an isomorphism between $N(\sigma) = N/N_\sigma$ and $N(\rho_i)(\sigma_i) = N(\rho_i)/N(\rho_i)_{\sigma_i}$. We have an inclusion of \mathbb{A}^{d-1} into \mathbb{A}^d by setting the coordinate corresponding to ρ_i to be zero. We conclude that $\mathcal{D}_i \cap \mathcal{X}(\sigma) \cong [\mathbb{A}^{d-1}/N(\sigma)]$ and the inclusion of \mathbb{A}^{d-1} into \mathbb{A}^d induces the inclusion of $\mathcal{D}_i \cap \mathcal{X}(\sigma)$ into $\mathcal{X}(\sigma)$. Moreover, by Lemma 3.2, if g is an element of $N(\sigma)$, then we have an induced inclusion

$$(6) \quad J_\infty^{(g)}(\mathbb{A}^{d-1})/N(\sigma) \hookrightarrow J_\infty^{(g)}(\mathbb{A}^d)/N(\sigma),$$

corresponding to the closed inclusion $|\mathcal{J}_\infty(\mathcal{D}_i \cap \mathcal{X}(\sigma))| \hookrightarrow |\mathcal{J}_\infty \mathcal{X}(\sigma)|$. We define $|\mathcal{J}_\infty \mathcal{X}'|$ to be the open subset

$$|\mathcal{J}_\infty \mathcal{X}'| = |\mathcal{J}_\infty \mathcal{X}| \setminus \cup_{i=1}^r |\mathcal{J}_\infty \mathcal{D}_i|.$$

Similarly, we consider $|\mathcal{J}_\infty \mathcal{X}(\sigma)'|$ and let $J_\infty(\mathbb{A}^d)'$ be the open locus of arcs of \mathbb{A}^d that are not contained in a coordinate hyperplane. Setting $J_\infty^{(g)}(\mathbb{A}^d)' = J_\infty^{(g)}(\mathbb{A}^d) \cap J_\infty(\mathbb{A}^d)'$, it follows from Lemma 3.2 and (6) that

$$(7) \quad |\mathcal{J}_\infty \mathcal{X}(\sigma)'| \cong \coprod_{g \in N(\sigma)} J_\infty^{(g)}(\mathbb{A}^d)'/N(\sigma).$$

We will often make this identification.

For a fixed positive integer l , the open embedding of T in $\mathcal{X}(\sigma)$ extends to an open embedding of $J_\infty(T) \cong \mathcal{J}_\infty^l T$ in $\mathcal{J}_\infty^l \mathcal{X}(\sigma)$. We obtain an action of $J_\infty(T)$ on $|\mathcal{J}_\infty^l \mathcal{X}(\sigma)|$, which restricts to an action on $|\mathcal{J}_\infty^l \mathcal{X}(\sigma)'|$. More specifically, we identify $J_\infty(T) \cong J_\infty^{(l)}(T)/N(\sigma)$, where $J_\infty^{(l)}(T)$ is the closed subscheme of $J_\infty(T)$ fixed by the action of μ_l . In coordinates,

$$J_\infty^{(l)}(T) = \left(\sum_{j=0}^{\infty} \beta_{1,j} t^{lj}, \dots, \sum_{j=0}^{\infty} \beta_{d,j} t^{lj} \mid \beta_{i,0} \neq 0 \text{ for } i = 1, \dots, d \right).$$

Fix an element g in $N(\sigma)$ of order l and recall from (5) that

$$J_\infty^{(g)}(\mathbb{A}^d) = \left\{ \left(\sum_{j=0}^{\infty} \alpha_{1,j} t^j, \dots, \sum_{j=0}^{\infty} \alpha_{d,j} t^j \right) \mid \alpha_{i,j} = 0 \text{ if } j \not\equiv lq_i \pmod{l} \right\}.$$

The elements in $J_\infty^{(g)}(\mathbb{A}^d)'$ also satisfy the property that for each $i = 1, \dots, d$, there exists a non-negative integer j such that $\alpha_{i,j} \neq 0$. Componentwise multiplication of power series gives an action,

$$J_\infty^{(l)}(T)/N(\sigma) \times J_\infty^{(g)}(\mathbb{A}^d)'/N(\sigma) \rightarrow J_\infty^{(g)}(\mathbb{A}^d)'/N(\sigma).$$

By (7), this induces an action of $J_\infty(T)$ on $|\mathcal{J}_\infty^l \mathcal{X}(\sigma)'|$. By varying l , we obtain an action of $J_\infty(T)$ on $|\mathcal{J}_\infty \mathcal{X}'|$. We would like to describe the $J_\infty(T)$ -orbits.

Let w be an element of $\sigma \cap N$. There is a unique decomposition $w = v + \sum_{i=1}^d \lambda_i b_i$, where v lies in $\text{Box}(\tau)$, for some $\tau \subseteq \sigma$, and the λ_i are non-negative integers. We will use the notation $\{w\} = v$. Suppose

¹Note that $N(\rho_i)$ may have torsion and so σ_i is a cone in the image of $N(\rho_i)$ in $N(\rho_i)_{\mathbb{Q}}$. There are no difficulties in generalising to this situation. See Section 8 for a discussion of this issue. This is the level of generality used in [5].

v corresponds to an element g in $N(\sigma)$ of order l . If we write $v = \sum_{i=1}^d q_i b_i$, for some $0 \leq q_i < 1$, then $w = \sum_{i=1}^d w_i b_i = \sum_{i=1}^d (\lambda_i + q_i) b_i$, where $w_i = \lambda_i + q_i$. We define $\tilde{\gamma}_w \in |\mathcal{J}_\infty^l \mathcal{X}(\sigma)|' \subseteq |\mathcal{J}_\infty \mathcal{X}'|$ to be the equivalence class of twisted arcs corresponding to the element

$$(8) \quad (t^{l(\lambda_1+q_1)}, \dots, t^{l(\lambda_d+q_d)}) = (t^{lw_1}, \dots, t^{lw_d})$$

of $J_\infty^{(g)}(\mathbb{A}^d)'$, under the isomorphism (7).

Theorem 4.2. *We have a decomposition of $|\mathcal{J}_\infty \mathcal{X}'|$ into $J_\infty(T)$ -orbits*

$$|\mathcal{J}_\infty \mathcal{X}'| = \coprod_{v \in |\Sigma| \cap N} \tilde{\gamma}_v \cdot J_\infty(T).$$

Moreover, $\tilde{\gamma}_w \cdot J_\infty(T) \subseteq \overline{\tilde{\gamma}_v \cdot J_\infty(T)}$ if and only if $w - v = \sum_{\rho_i \subseteq \sigma} \lambda_i b_i$ for some non-negative integers λ_i and some cone σ containing v and w . With the notation of Theorem 4.1 and (4),

$$\tilde{\pi}_\infty : |\mathcal{J}_\infty \mathcal{X}'| \rightarrow J_\infty(X)'$$

is a $J_\infty(T)$ -equivariant bijection satisfying

$$\tilde{\pi}_\infty(\tilde{\gamma}_v) = \gamma_v.$$

Proof. Let σ be a d -dimensional cone in Σ and let g be an element in $N(\sigma)$ of order l corresponding to an element v in $\text{Box}(\tau)$, for some $\tau \subseteq \sigma$. By (5), and with the notation of the previous discussion, we have a decomposition

$$J_\infty^{(g)}(\mathbb{A}^d)' = \coprod_{\substack{w \in \sigma \cap N \\ \{w\}=v}} (t^{lw_1}, \dots, t^{lw_d}) \cdot J_\infty^{(l)}(T),$$

and hence

$$J_\infty^{(g)}(\mathbb{A}^d)' / N(\sigma) \cong \coprod_{\substack{w \in \sigma \cap N \\ \{w\}=v}} \tilde{\gamma}_w \cdot J_\infty(T).$$

Also, $\tilde{\gamma}_w \cdot J_\infty(T) \subseteq \overline{\tilde{\gamma}_{w'} \cdot J_\infty(T)}$ in $J_\infty^{(g)}(\mathbb{A}^d)' / N(\sigma)$ if and only if $w_i \geq w'_i$ for $i = 1, \dots, d$. We conclude that

$$|\mathcal{J}_\infty \mathcal{X}(\sigma)|' = \coprod_{w \in \sigma \cap N} \tilde{\gamma}_w \cdot J_\infty(T),$$

and $\tilde{\gamma}_w \cdot J_\infty(T) \subseteq \overline{\tilde{\gamma}_{w'} \cdot J_\infty(T)}$ in $|\mathcal{J}_\infty \mathcal{X}(\sigma)|'$ if and only if $\{w\} = \{w'\}$ and $w_i \geq w'_i$ for $i = 1, \dots, d$. This is equivalent to $w - w' = \sum_{i=1}^d \lambda_i b_i$ for some non-negative integers λ_i . Since the open subsets $|\mathcal{J}_\infty \mathcal{X}(\sigma)|'$ cover $|\mathcal{J}_\infty \mathcal{X}'|$ as σ varies over all maximal cones, we obtain the desired decomposition and closure relations.

Consider the sublattice $N_\sigma \subseteq N$ and recall that σ' is the cone in N_σ generated by b_1, \dots, b_d . If H denotes the quotient of N by the sublattice generated by v_1, \dots, v_d , then we have a pairing $\langle \cdot, \cdot \rangle : N \times M_\sigma \rightarrow \mathbb{Q}$, $\langle v_i, u_j \rangle = \delta_{i,j} |H|$, where $\delta_{i,j} = 1$ if $i = j$ and 0 otherwise. If u_1, \dots, u_d are the primitive integer generators of σ^\vee in M , then $u_1/a_1 |H|, \dots, u_d/a_d |H|$ are the primitive integer generators of $(\sigma')^\vee$ in M_σ . We have made an identification throughout that $\text{Spec } \mathbb{C}[(\sigma')^\vee \cap M_\sigma] \cong \mathbb{A}^d$. Consider an element γ in $J_\infty^{(g)}(\mathbb{A}^d)'$, for some g

in $N(\sigma)$ of order l , corresponding to an element of $|\mathcal{J}_\infty \mathcal{X}(\sigma)|'$. Applying $\tilde{\pi}_\infty$ gives an arc in U_σ , which we denote by $\tilde{\pi}_\infty(\gamma)$. We have a commutative diagram

$$\begin{array}{ccc} \mathbb{C}[(\sigma')^\vee \cap M_\sigma] & \xrightarrow{\gamma^\#} & \mathbb{C}[[t]] \\ \uparrow & & \uparrow t \rightarrow t^l \\ \mathbb{C}[\sigma^\vee \cap M] & \xrightarrow{\tilde{\pi}_\infty(\gamma)^\#} & \mathbb{C}[[t]]. \end{array}$$

It follows that $\tilde{\pi}_\infty$ is $J_\infty(T)$ -equivariant. Let w be an element of $\sigma \cap N$ and consider the notation of (8). With a slight abuse of notation, $(\tilde{\gamma}_w)^\#(\chi^{u_i/a_i|H|}) = t^{lw_i}$, and we compute

$$\tilde{\pi}_\infty(\tilde{\gamma}_w)^\#(\chi^{u_i}) = ((t^{1/l})^{lw_i})^{a_i|H|} = t^{a_i w_i |H|} = t^{\langle u_i, w \rangle}.$$

It follows from the definition of γ_w that $\tilde{\pi}_\infty(\tilde{\gamma}_w) = \gamma_w$. \square

Remark 4.3. The morphism from a Deligne-Mumford stack to its coarse moduli space is proper [22]. Hence the fact that the map $\tilde{\pi}_\infty : |\mathcal{J}_\infty \mathcal{X}'| \rightarrow J_\infty(X)'$ is bijective follows from Proposition 3.37 of [34].

Remark 4.4. We give a geometric description of a well-known involution ι on $|\Sigma| \cap N$ (see, for example, Section 2 [30]). If τ is a cone in Σ and v is a lattice point in $\text{Box}(\tau)$, then v can be uniquely written in the form $v = \sum_{\rho_i \subseteq \tau} q_i b_i$, for some $0 < q_i < 1$. We define $\iota = \iota_\Sigma : \text{Box}(\tau) \rightarrow \text{Box}(\tau)$ by $\iota(v) = \sum_{\rho_i \subseteq \tau} (1 - q_i) b_i$. By Remark 3.3, the connected components of $|\mathcal{I}\mathcal{X}|$ are indexed by $\text{Box}(\Sigma)$. Recall that the elements of $|\mathcal{I}\mathcal{X}|$ are equivalence classes of pairs (x, α) , where x is an object of \mathcal{X} over \mathbb{C} and α is an automorphism of x . There is a natural involution on $|\mathcal{I}\mathcal{X}|$, taking a pair (x, α) to (x, α^{-1}) . Observe that I induces the involution ι on the connected components of $|\mathcal{I}\mathcal{X}|$.

Every lattice point w in $|\Sigma| \cap N$ can be uniquely written in the form $w = \{w\} + \tilde{w}$, where $\{w\}$ lies in $\text{Box}(\tau)$ for some $\tau \subseteq \sigma(w)$ and \tilde{w} lies in $N_{\sigma(w)}$. Here $\sigma(w)$ is the cone of Σ containing w in its relative interior. The map ι extends to an involution on $|\Sigma| \cap N$, taking w to $\iota(\{w\}) + \tilde{w}$. Recall that we have a projection morphism $\pi : \mathcal{J}_\infty \mathcal{X} \rightarrow \mathcal{I}\mathcal{X} = \mathcal{J}_0 \mathcal{X}$. We see that I extends to a $J_\infty(T)$ -equivariant involution $I : |\mathcal{J}_\infty \mathcal{X}'| \rightarrow |\mathcal{J}_\infty \mathcal{X}'|$ satisfying $I(\tilde{\gamma}_v) = \tilde{\gamma}_{\iota(v)}$.

Remark 4.5. For any non-negative integer n , we can consider $|\mathcal{J}_n \mathcal{X}'| = |\mathcal{J}_n \mathcal{X}| \setminus \cup_{i=1}^r |\mathcal{J}_n \mathcal{D}_i|$ and an action of $J_n(T)$ on $|\mathcal{J}_n \mathcal{X}'|$. Recall that we have projection morphisms $\pi_n : \mathcal{J}_\infty \mathcal{X} \rightarrow \mathcal{J}_n \mathcal{X}$. For each non-zero cone τ of Σ , let

$$\overline{\text{Box}}(\mathbf{n}\tau) = \{v \in N \mid v = \sum_{\rho_i \subseteq \tau} q_i b_i \text{ for some } 0 < q_i \leq n\}.$$

We set $\overline{\text{Box}}(\mathbf{n}\{0\}) = \{0\}$ and $\overline{\text{Box}}(\mathbf{n}\Sigma) = \cup_{\tau \in \Sigma} \overline{\text{Box}}(\mathbf{n}\tau)$. It can be shown that there is a decomposition of $|\mathcal{J}_n \mathcal{X}'|$ into $J_n(T)$ -orbits

$$|\mathcal{J}_n \mathcal{X}'| = \coprod_{v \in \overline{\text{Box}}(\mathbf{n}\Sigma)} \pi_n(\tilde{\gamma}_v) \cdot J_n(T).$$

5. CONTACT ORDER ALONG A DIVISOR

In this section, we compute the contact order of a twisted arc along a T -invariant divisor on \mathcal{X} . Recall that for each maximal cone σ in Σ , we have morphisms

$$\mathbb{A}^d \xrightarrow{p} [\mathbb{A}^d/N(\sigma)] \cong \mathcal{X}(\sigma) \xrightarrow{q} \mathbb{A}^d/N(\sigma) \cong U_\sigma,$$

where \mathbb{A}^d is the atlas of $\mathcal{X}(\sigma)$ via p and U_σ is the coarse moduli space of $\mathcal{X}(\sigma)$. For every ray ρ_i in Σ , there is a corresponding divisor \mathcal{D}_i of $\mathcal{X}(\Sigma)$ (see Section 4). If $\rho_i \not\subseteq \sigma$ then \mathcal{D}_i does not intersect $\mathcal{X}(\sigma)$. If $\rho_i \subseteq \sigma$, then \mathcal{D}_i corresponds to the divisor $\{x_i = 0\}$ on \mathbb{A}^d , with an appropriate choice of coordinates. We say that a \mathbb{Q} -divisor \mathcal{E} on \mathcal{X} is T -invariant if $\mathcal{E} = \sum_i \beta_i \mathcal{D}_i$, for some $\beta_i \in \mathbb{Q}$. There is a natural isomorphism of Picard groups over \mathbb{Q} (Example 6.7 [32])

$$q^* : \text{Pic}_{\mathbb{Q}}(X(\Sigma)) \rightarrow \text{Pic}_{\mathbb{Q}}(\mathcal{X}(\Sigma))$$

$$D_i \mapsto q^* D_i = a_i \mathcal{D}_i.$$

The inverse map is induced by the pushforward functor $q_* : \text{Pic}_{\mathbb{Q}}(\mathcal{X}(\Sigma)) \rightarrow \text{Pic}_{\mathbb{Q}}(X(\Sigma))$. The canonical divisor on \mathcal{X} is given by $K_{\mathcal{X}} = -\sum_i \mathcal{D}_i$, and so $q_* K_{\mathcal{X}} = -\sum_i a_i^{-1} D_i$. Recall that a T -invariant \mathbb{Q} -divisor $E = \sum_{\rho_i \in \Sigma} \alpha_i D_i$ on X corresponds to a real-valued piecewise \mathbb{Q} -linear function $\psi_E : |\Sigma| \rightarrow \mathbb{R}$ satisfying $\psi_E(v_i) = -\alpha_i$ [14]. Note that $\psi_{q_* K_{\mathcal{X}}}(b_i) = 1$ for $i = 1, \dots, r$, and hence, with the notation of Section 2, $\psi = \psi_{q_* K_{\mathcal{X}}}$. Let $\mathcal{E} = \sum_i \beta_i \mathcal{D}_i$ be a T -invariant \mathbb{Q} -divisor on \mathcal{X} . Following Yasuda (Definition 4.17 [34]), we say that the pair $(\mathcal{X}, \mathcal{E})$ is *Kawamata log terminal* if $\beta_i < 1$ for $i = 1, \dots, r$. Geometrically, this says that for each maximal cone σ , the representative of \mathcal{E} in the atlas \mathbb{A}^d of $\mathcal{X}(\sigma) = [\mathbb{A}^d/N(\sigma)]$ is supported on the coordinate axes with all coefficients less than 1. Equivalently, the condition says that $\psi_{q_* \mathcal{E}}(b_i) > -1$ for $i = 1, \dots, r$.

If $\mathcal{E} = \sum u_j \mathcal{E}_j$ is a \mathbb{Q} -divisor on \mathcal{X} , for some prime divisors \mathcal{E}_j and rational numbers u_j , then Yasuda [34] defined a function

$$\text{ord } \mathcal{E} : |\mathcal{J}_{\infty} \mathcal{X}| \setminus \cup_j |\mathcal{J}_{\infty} \mathcal{E}_j| \rightarrow \mathbb{Q},$$

$$\text{ord } \mathcal{E} = \sum_j u_j \text{ord } \mathcal{E}_j.$$

When \mathcal{E} is a prime divisor, the function $\text{ord } \mathcal{E}$ is defined as follows: if $\gamma : \mathcal{D}_{\infty, \mathbb{C}}^l \rightarrow \mathcal{X}$ is a representable morphism, then consider the composition of γ with the atlas $\text{Spec } \mathbb{C}[[t]] \rightarrow \mathcal{D}_{\infty, \mathbb{C}}^l$, and choose a lifting to an arc $\bar{\gamma}$ of an atlas M of \mathcal{X} . If m is the contact order of $\bar{\gamma}$ along the representative of \mathcal{E} in M , then $\text{ord } \mathcal{E}(\gamma) = m/l$.

The following lemma describes the function $\text{ord } \mathcal{E}$ in the toric case. We will use the notation of Theorem 4.2.

Lemma 5.1. *If \mathcal{E} is a T -invariant \mathbb{Q} -divisor on \mathcal{X} , then*

$$\text{ord } \mathcal{E}(\tilde{\gamma}_w \cdot J_{\infty}(T)) = -\psi_{q_* \mathcal{E}}(w).$$

In particular,

$$\text{ord } K_{\mathcal{X}}(\tilde{\gamma}_w \cdot J_{\infty}(T)) = -\psi(w).$$

Proof. It will be enough to prove the result for $\mathcal{E} = \mathcal{D}_i$ and twisted arcs of the form $\tilde{\gamma}_w$. Consider the representable morphism $\tilde{\gamma}_w : \mathcal{D}_{\infty, \mathbb{C}}^l \rightarrow \mathcal{X}(\sigma) \subseteq \mathcal{X}$, for some lattice point w in a maximal cone σ . With the notation of (8), the composition of $\tilde{\gamma}_w$ with $\text{Spec } \mathbb{C}[[t]] \rightarrow \mathcal{D}_{\infty, \mathbb{C}}^l$ lifts to an arc $(t^{lw_1}, \dots, t^{lw_d})$ in $J_{\infty}(\mathbb{A}^d)$. If $\rho_i \not\subseteq \sigma$ then $\text{ord } \mathcal{D}_i(\tilde{\gamma}_w) = 0 = -\psi_{a_i^{-1}D_i}(w)$. If $\rho_i \subseteq \sigma$ then the divisor \mathcal{D}_i is represented by the divisor $\{x_i = 0\}$ in the atlas \mathbb{A}^d and we conclude that $\text{ord } \mathcal{D}_i(\tilde{\gamma}_w) = w_i = -\psi_{a_i^{-1}D_i}(w)$. The second statement follows since $\psi = \psi_{q_*K_{\mathcal{X}}}$. \square

6. MOTIVIC INTEGRATION ON TORIC STACKS

We consider motivic integration on a Deligne-Mumford stack as developed by Yasuda in [34]. We will compute motivic integrals associated to T -invariant divisors on \mathcal{X} and show that they correspond to weighted δ -vectors of an associated polyhedral complex.

Recall that to any complex algebraic variety X of dimension r , we can associate its *Hodge polynomial* (see, for example, [24])

$$E_X(u, v) = \sum_{i, j=0}^r (-1)^{i+j} h_{i, j} u^i v^j \in \mathbb{Z}[u, v].$$

The Hodge polynomial is determined by the properties

- (1) $h_{i, j} = \dim H^j(X, \Omega_X^i)$ if X is smooth and projective,
- (2) $E_X(u, v) = E_U(u, v) + E_{X \setminus U}(u, v)$ if U is an open subvariety of X ,
- (3) $E_{X \times Y}(u, v) = E_X(u, v)E_Y(u, v)$.

The second property means we can consider the Hodge polynomial of a constructible subset of a complex variety. For example, $E_{\mathbb{A}^1}(u, v) = E_{\mathbb{P}^1}(u, v) - E_{\{\text{pt}\}}(u, v) = uv$ and hence $E_{\mathbb{A}^n}(u, v) = (uv)^n$. Similarly, $E_{(\mathbb{C}^*)^n}(u, v) = (uv - 1)^n$. More generally, we can compute the Hodge polynomial of any toric variety. Recall that if Δ is a fan in a lattice $N'_{\mathbb{R}}$, for some lattice N' , then its associated h -vector $h_{\Delta}(t)$ is defined by

$$h_{\Delta}(t) = \sum_{\tau \in \Delta} t^{\dim \tau} (1-t)^{\text{codim } \tau}.$$

Lemma 6.1. *If $X = X(\Delta)$ is an r -dimensional toric variety associated to a fan Δ , then*

$$E_X(u, v) = (uv)^r h_{\Delta}((uv)^{-1}).$$

Proof. We can write X as a disjoint union of torus orbits indexed by the cones of Δ (see, for example, [14]). The orbit corresponding to τ is isomorphic to $(\mathbb{C}^*)^{\text{codim } \tau}$. We compute, using the example above,

$$E_X(u, v) = \sum_{\tau \in \Delta} E_{(\mathbb{C}^*)^{\text{codim } \tau}}(u, v) = \sum_{\tau \in \Delta} (uv - 1)^{\text{codim } \tau} = (uv)^r h_{\Delta}((uv)^{-1}).$$

\square

Remark 6.2. If Δ is an r -dimensional, simplicial fan with convex support, then $X = X(\Delta)$ has no odd cohomology and the coefficient of $(uv)^i$ in $E_X(u, v)$ is equal to the dimension of the $2i^{\text{th}}$ cohomology group of X with compact support and rational coefficients. This follows from the above lemma, Lemma 4.1 of [30] and Poincaré duality.

Recall that we have projection morphisms $\pi_n : |\mathcal{J}_\infty \mathcal{X}| \rightarrow |\mathcal{J}_n \mathcal{X}|$, for each non-negative integer n . A subset $A \subseteq |\mathcal{J}_\infty \mathcal{X}|$ is a *cylinder* if $A = \pi_n^{-1} \pi_n(A)$ and $\pi_n(A)$ is a constructible subset, for some non-negative integer n . In this case, the measure $\mu_{\mathcal{X}}(A)$ of A is defined to be

$$\mu_{\mathcal{X}}(A) = E_{\pi_n(A)}(u, v)(uv)^{-nd} \in \mathbb{Z}[u, u^{-1}, v, v^{-1}].$$

By Lemma 3.18 of [34], the right hand side is independent of the choice of n . The collection of cylinders is closed under taking finite unions and finite intersections and $\mu_{\mathcal{X}}$ defines a finite measure on $|\mathcal{J}_\infty \mathcal{X}|$.

We will compute the measure of certain cylinders. Recall the decomposition of $|\mathcal{J}_\infty \mathcal{X}'|$ in Theorem 4.2. Every w in $|\Sigma| \cap N$ can be uniquely written in the form $w = \{w\} + \tilde{w}$, where $\{w\}$ lies in $\text{Box}(\tau)$ for some $\tau \subseteq \sigma(w)$ and \tilde{w} lies in $N_{\sigma(w)}$. Recall that $\sigma(w)$ is the cone of Σ containing w in its relative interior and that $\psi : |\Sigma| \rightarrow \mathbb{R}$ is the piecewise \mathbb{Q} -linear function satisfying $\psi(b_i) = 1$ for $i = 1, \dots, r$.

Lemma 6.3. *For any w in $|\Sigma| \cap N$, the orbit $\tilde{\gamma}_w \cdot J_\infty(T)$ is a cylinder in $|\mathcal{J}_\infty \mathcal{X}'|$ with measure*

$$\mu_{\mathcal{X}}(\tilde{\gamma}_w \cdot J_\infty(T)) = (uv - 1)^d (uv)^{-\psi(w) + \psi(\{w\}) - \dim \sigma(\{w\})}.$$

Proof. Fix a d -dimensional cone σ containing w and consider the notation of (8). The twisted arc $\tilde{\gamma}_w$ lies in $J_\infty^{(g)} \mathbb{A}^d / N(\sigma)$ and $\pi_n(\tilde{\gamma}_w)$ lies in $J_{nl}^{(g)} \mathbb{A}^d / N(\sigma)$. If we consider the natural projection $\pi_n^{(g)} : J_\infty^{(g)} \mathbb{A}^d \rightarrow J_{nl}^{(g)} \mathbb{A}^d$, then $\pi_n^{(g)}((t^{lw_1}, \dots, t^{lw_d}) \cdot J_\infty^{(l)}(T))$ is equal to

$$\left\{ \left(\sum_{k=\lambda_1}^{n_1} \alpha_{1,l(k+q_1)} t^{l(k+q_1)}, \dots, \sum_{k=\lambda_d}^{n_d} \alpha_{d,l(k+q_d)} t^{l(k+q_d)} \right) \mid \alpha_{i,w_i} \neq 0 \right\},$$

for $n \geq \max\{\lambda_1 + 1, \dots, \lambda_d + 1\}$. Here $n_i = n$ if $q_i = 0$ and $n_i = n - 1$ if $q_i \neq 0$. Note that $q_i \neq 0$ if and only if $\rho_i \subseteq \sigma(\{w\})$. Hence $(\pi_n^{(g)})^{-1} \pi_n^{(g)}((t^{lw_1}, \dots, t^{lw_d}) \cdot J_\infty^{(l)}(T)) = (t^{lw_1}, \dots, t^{lw_d}) \cdot J_\infty^{(l)}(T)$, and $\pi_n^{(g)}((t^{lw_1}, \dots, t^{lw_d}) \cdot J_\infty^{(l)}(T))$ is isomorphic to

$$(9) \quad (\mathbb{C}^*)^d \times \mathbb{A}^{nd - \sum_{i=1}^d \lambda_i - \dim \sigma(\{w\})}.$$

Note that (9) has a decomposition into a disjoint union of locally closed subspaces, each isomorphic to $(\mathbb{C}^*)^i$ for some i , which is preserved after quotienting by the induced action of the finite group $N(\sigma)$. The result follows by considering the corresponding spaces and maps after quotienting by $N(\sigma)$, and observing that this doesn't affect the Hodge polynomial of (9). \square

Let $F : |\mathcal{J}_\infty \mathcal{X}'| \rightarrow \mathbb{Q}$ be a $J_\infty(T)$ -invariant function on the space of twisted arcs of \mathcal{X} and let A be a $J_\infty(T)$ -invariant subset of $|\mathcal{J}_\infty \mathcal{X}'|$. By Theorem 4.2, we have a decomposition

$$A \cap |\mathcal{J}_\infty \mathcal{X}'| = \coprod_{\substack{w \in |\Sigma| \cap N \\ \tilde{\gamma}_w \in A}} \tilde{\gamma}_w \cdot J_\infty(T).$$

We define

$$\int_A (uv)^F d\mu_{\mathcal{X}} := \sum_{\substack{w \in |\Sigma| \cap N \\ \tilde{\gamma}_w \in A}} \mu_{\mathcal{X}}(\tilde{\gamma}_w \cdot J_\infty(T)) (uv)^{F(\tilde{\gamma}_w)},$$

when the right hand sum is a well-defined element in $\mathbb{Z}[(uv)^{1/N}][[(uv)^{-1/N}]]$, for some positive integer N . With the definitions of Yasuda, this is the *motivic integral* of F over A .²

Recall the involution $\iota : \text{Box}(\Sigma) \rightarrow \text{Box}(\Sigma)$ (Remark (4.4)). Following [34], we define the *shift function* $s_{\mathcal{X}} : |\mathcal{J}_{\infty}\mathcal{X}'| \rightarrow \mathbb{Q}$ by

$$s_{\mathcal{X}}(\tilde{\gamma}_w \cdot J_{\infty}(T)) = \dim \sigma(\{w\}) - \psi(\{w\}) = \psi(\iota(\{w\})).$$

The shift function factors as $|\mathcal{J}_{\infty}\mathcal{X}'| \xrightarrow{\pi} |\mathcal{J}_0\mathcal{X}| = |\mathcal{I}\mathcal{X}| \xrightarrow{\text{sft}} \mathbb{Q}$, where the function sft is constant on the connected components of $|\mathcal{I}\mathcal{X}|$. Borisov, Chen and Smith [5] showed that the connected components of $|\mathcal{I}\mathcal{X}|$ are indexed by $\text{Box}(\Sigma)$ (c.f. Remark 3.3). The value of sft on the component of $|\mathcal{I}\mathcal{X}|$ corresponding to g in $N(\sigma)$ is the age of g^{-1} (see (3) and Remark 4.4).

Consider a Kawamata log terminal pair $(\mathcal{X}, \mathcal{E})$, where \mathcal{E} is a T -invariant \mathbb{Q} -divisor on \mathcal{X} (see Section 5). Following Yasuda (Definition 4.1 [34]), we consider the invariant $\Gamma(\mathcal{X}, \mathcal{E})$ defined by

$$\Gamma(\mathcal{X}, \mathcal{E}) = \int_{|\mathcal{J}_{\infty}\mathcal{X}'|} (uv)^{s_{\mathcal{X}} + \text{ord } \mathcal{E}} d\mu_{\mathcal{X}}.$$

Yasuda showed that $\Gamma(\mathcal{X}, \mathcal{E})$ is a well-defined element in $\mathbb{Z}[(uv)^{1/N}][[(uv)^{-1/N}]]$, for some positive integer N [34, Proposition 4.15]. Our goal is to compute such invariants and give them a combinatorial interpretation. We first recall the notion of weighted δ -vector from [30]. Consider the polyhedral complex $Q = \{v \in |\Sigma| \mid \psi(v) \leq 1\}$ and let $\lambda : |\Sigma| \rightarrow \mathbb{R}$ be a piecewise \mathbb{Q} -linear function satisfying $\lambda(b_i) > -1$ for $i = 1, \dots, r$. The *weighted δ -vector* $\delta^{\lambda}(t)$ is defined by

$$\delta^{\lambda}(t) = (1-t)^{d+1} \left(1 + \sum_{m \geq 1} \sum_{v \in mQ \cap N} t^{\psi(v) - \lceil \psi(v) \rceil + \lambda(v) + m} \right).$$

The following symmetry property was established in Corollary 2.13 of [30].

Theorem 6.4 ([30]). *With the notation above, the expression $\delta^{\lambda}(t)$ is a rational function in $\mathbb{Q}(t^{1/N})$, for some positive integer N . If Σ is a complete fan, then*

$$\delta^{\lambda}(t) = t^d \delta^{\lambda}(t^{-1}).$$

For each cone τ in Σ , let N_{τ} be the sublattice of N generated by $\{b_i \mid \rho_i \subseteq \tau\}$ and let Σ_{τ} be the simplicial fan in $(N/N_{\tau})_{\mathbb{R}}$ with cones given by the projections of the cones in Σ containing τ . We define $h_{\tau}^{\lambda}(t)$ to be the expression

$$h_{\tau}^{\lambda}(t) = \sum_{\sigma \supseteq \tau} t^{\sum_{\rho_i \subseteq \sigma \setminus \tau} \lambda(b_i)} t^{\dim \sigma - \dim \tau} (1-t)^{\text{codim } \sigma} \prod_{\rho_i \subseteq \sigma \setminus \tau} (1-t)/(1-t^{\lambda(b_i)+1}),$$

so that when $\lambda \equiv 0$, $h_{\tau}^{\lambda}(t)$ is the h -vector of Σ_{τ} (see, for example, [29]). By expanding and collecting terms, we have

$$(10) \quad t^{\text{codim } \tau} h_{\tau}^{\lambda}(t^{-1}) = (t-1)^{\text{codim } \tau} \sum_{\sigma \supseteq \tau} \prod_{\rho_i \subseteq \sigma \setminus \tau} 1/(t^{\lambda(b_i)+1} - 1).$$

²To check the definitions agree, we can replace A by $A \cap |\mathcal{J}_{\infty}\mathcal{X}'|$, by Propositions 3.24 and 3.25 in [34]. Now apply the decomposition of Theorem 4.2.

The following expression for the weighted δ -vector was established in Proposition 2.7 of [30],

$$(11) \quad \delta^\lambda(t) = \sum_{\tau \in \Sigma} h_\tau^\lambda(t) \sum_{v \in \text{Box}(\boldsymbol{\tau})} t^{\psi(v) + \lambda(v)} \prod_{\rho_i \subseteq \tau} (t-1)/(t^{\lambda(b_i)+1} - 1).$$

After rearranging we see that

$$(12) \quad \delta^\lambda(t^{-1}) = \sum_{\tau \in \Sigma} h_\tau^\lambda(t^{-1}) \sum_{v \in \text{Box}(\boldsymbol{\tau})} t^{-\psi(v) - \lambda(v) + \sum_{\rho_i \subseteq \tau} \lambda(b_i)} \prod_{\rho_i \subseteq \tau} (t-1)/(t^{\lambda(b_i)+1} - 1).$$

Theorem 6.5. *If $(\mathcal{X}, \mathcal{E})$ is a Kawamata log terminal pair, where \mathcal{E} is a T -invariant \mathbb{Q} -divisor on \mathcal{X} , then the corresponding piecewise \mathbb{Q} -linear function $\lambda = \psi_{q^* \mathcal{E}}$ satisfies $\lambda(b_i) > -1$ for $i = 1, \dots, r$, and*

$$\Gamma(\mathcal{X}, \mathcal{E}) = (uv)^d \delta^\lambda((uv)^{-1}).$$

In particular, $\Gamma(\mathcal{X}, \mathcal{E})$ is a rational function in $\mathbb{Q}(t^{1/N})$, for some positive integer N . If Σ is a complete fan, then

$$\Gamma(\mathcal{X}, \mathcal{E})(u, v) = (uv)^d \Gamma(\mathcal{X}, \mathcal{E})(u^{-1}, v^{-1}) = \delta^\lambda(uv).$$

Moreover, every weighted δ -vector has the form $\delta^\lambda(uv) = (uv)^d \Gamma(\mathcal{X}, \mathcal{E})(u^{-1}, v^{-1})$, for some such pair $(\mathcal{X}, \mathcal{E})$.

Proof. We showed in Lemma 5.1 that $\text{ord } \mathcal{E}(\tilde{\gamma}_w \cdot J_\infty(T)) = -\lambda(w)$ and hence $(s_{\mathcal{X}} + \text{ord } \mathcal{E})(\tilde{\gamma}_w \cdot J_\infty(T)) = \dim \sigma(\{w\}) - \psi(\{w\}) - \lambda(w)$. Using Lemma 6.3, we compute

$$\Gamma(\mathcal{X}, \mathcal{E}) = (uv-1)^d \sum_{w \in |\Sigma| \cap N} (uv)^{-\psi(w) - \lambda(w)}.$$

For each w in $|\Sigma| \cap N$, we have a unique decomposition

$$w = \{w\} + w' + \sum_{\rho_i \subseteq \sigma(w) \setminus \sigma(\{w\})} b_i,$$

where w' is a non-negative linear combination of $\{b_i \mid \rho_i \subseteq \sigma(w)\}$. Here $\sigma(w)$ is the cone containing w in its relative interior. Using this decomposition, we compute the following expression for $\Gamma(\mathcal{X}, \mathcal{E})$,

$$\sum_{\substack{\tau \in \Sigma \\ v \in \text{Box}(\boldsymbol{\tau})}} (uv-1)^d (uv)^{-(\psi+\lambda)(v)} \sum_{\tau \subseteq \sigma} (uv)^{-\sum_{\rho_i \subseteq \sigma \setminus \tau} (\lambda(b_i)+1)} \prod_{\rho_i \subseteq \sigma} 1/(1 - (uv)^{-\lambda(b_i)-1}).$$

Rearranging and using (10) gives

$$\begin{aligned} \Gamma(\mathcal{X}, \mathcal{E}) &= \sum_{\tau \in \Sigma} (uv)^{\text{codim } \tau} h_\tau^\lambda((uv)^{-1}) \times \\ &\quad \sum_{v \in \text{Box}(\boldsymbol{\tau})} (uv)^{\sum_{\rho_i \subseteq \tau} \lambda(b_i) + \dim \tau - \psi(v) - \lambda(v)} \prod_{\rho_i \subseteq \tau} (uv-1)/((uv)^{\lambda(b_i)+1} - 1). \end{aligned}$$

Comparing with (12) gives $\Gamma(\mathcal{X}, \mathcal{E}) = (uv)^d \delta^\lambda((uv)^{-1})$. The second statement follows from Theorem 6.4.

If $\lambda : |\Sigma| \rightarrow \mathbb{R}$ is a piecewise \mathbb{Q} -linear function satisfying $\lambda(b_i) > -1$ for $i = 1, \dots, r$, then we may consider the corresponding T -invariant \mathbb{Q} -divisor E on X . The pair (\mathcal{X}, q^*E) is a Kawamata log terminal pair and, by the above argument, $\delta^\lambda(uv) = (uv)^d \Gamma(\mathcal{X}, q^*E)(u^{-1}, v^{-1})$. Hence every weighted δ -vector corresponds to a motivic integral on \mathcal{X} . \square

Remark 6.6. By Theorem 6.5 and (12), we obtain a formula for the invariant $\Gamma(\mathcal{X}, \mathcal{E})$. This formula is a stacky analogue of Batyrev's formula for the motivic integral of a simple normal crossing divisor on a smooth variety (see [31], [3]).

Remark 6.7. When $\mathcal{E} = 0$, $\Gamma(\mathcal{X}, 0)$ is a polynomial in uv of degree d and the coefficient of $(uv)^j$ is equal to the dimension of the $2j^{\text{th}}$ orbifold cohomology group of \mathcal{X} with compact support [33]. When Σ is complete, the symmetry $\Gamma(\mathcal{X}, 0)(u, v) = (uv)^d \Gamma(\mathcal{X}, 0)(u^{-1}, v^{-1})$ is a consequence of Poincaré duality for orbifold cohomology [6].

7. THE TRANSFORMATION RULE

A morphism $f : \mathcal{Y} \rightarrow \mathcal{X}$ of smooth Deligne-Mumford stacks is *birational* if there exist open, dense substacks \mathcal{Y}_0 of \mathcal{Y} and \mathcal{X}_0 of \mathcal{X} such that f induces an isomorphism $\mathcal{Y}_0 \cong \mathcal{X}_0$ (Definition 3.36 [34]). If f is proper and birational then Yasuda [34] proved a transformation rule relating motivic integrals on \mathcal{X} to motivic integrals on \mathcal{Y} , generalising a classic result of Kontsevich for smooth varieties (see, for example, [31]). The goal of this section is to interpret the transformation rule in our context in order to give a geometric proof of a combinatorial result in [30].

If $\Delta = (N, \Delta, \{\bar{b}_i\})$ and $\Sigma = (N, \Sigma, \{b_i\})$ are stacky fans, we say that Δ *refines* Σ if

- (1) The fan Δ refines Σ in $N_{\mathbb{R}}$,
- (2) For any ray $\bar{\rho}_i$ in Δ , \bar{b}_i is an integer combination of the lattice points $\{b_j \mid \rho_j \subseteq \sigma\}$, where σ is any cone of Σ containing $\bar{\rho}_i$.

If Δ *refines* Σ , we have an induced morphism of toric stacks $f : \mathcal{X}(\Delta) \rightarrow \mathcal{X}(\Sigma)$ (Remark 4.5 [5]) such that f restricts to the identity map on the torus and hence is birational. Note that the corresponding map of coarse moduli spaces is proper [14]. Since the morphism from a Deligne-Mumford stack to its coarse moduli space is proper [20], it follows that f is proper [22]. We will give a local description of f . Let τ be a maximal cone in Δ , contained in a maximal cone σ of Σ . By Property (2), we have an inclusion of lattices $N_{\tau} \hookrightarrow N_{\sigma}$, inducing a homomorphism of groups $j : N(\tau) \rightarrow N(\sigma)$. Let τ' be the cone generated by $\{\bar{b}_i \mid \bar{\rho}_i \subseteq \tau\}$ in N_{τ} and let σ' be the cone generated by $\{b_i \mid \rho_i \subseteq \sigma\}$ in N_{σ} . The inclusion $\tau' \cap N_{\tau} \hookrightarrow \sigma' \cap N_{\sigma}$ induces a j -equivariant map $\phi : \text{Spec } \mathbb{C}[\check{\tau}' \cap M_{\tau}] \rightarrow \text{Spec } \mathbb{C}[(\sigma')^{\vee} \cap M_{\sigma}]$. Taking stacky quotients of both sides yields the restriction of f to $\mathcal{X}(\tau)$, $f : \mathcal{X}(\tau) = [\text{Spec } \mathbb{C}[\check{\tau}' \cap M_{\tau}]/N(\tau)] \rightarrow \mathcal{X}(\sigma) = [\text{Spec } \mathbb{C}[(\sigma')^{\vee} \cap M_{\sigma}]/N(\sigma)]$.

Given a morphism $g : \mathcal{Y} \rightarrow \mathcal{Z}$ of Deligne-Mumford stacks, Yasuda described a natural morphism $g_{\infty} : |\mathcal{J}_{\infty} \mathcal{Y}| \rightarrow |\mathcal{J}_{\infty} \mathcal{Z}|$ (Proposition 2.14 [34]). We outline his construction. Given a representable morphism $\gamma : \mathcal{D}_{\infty, \mathbb{C}}^l \rightarrow \mathcal{Y}$, we can consider the composition $\gamma' : \mathcal{D}_{\infty, \mathbb{C}}^l \rightarrow \mathcal{Y} \rightarrow \mathcal{X}$. If l' is a positive integer dividing l , then we have a group homomorphism $\mu_l \rightarrow \mu_{l'}$, $\zeta_l \mapsto (\zeta_l)^{l/l'} = \zeta_{l'}$, and an equivariant map $\text{Spec } \mathbb{C}[[t]] \rightarrow \text{Spec } \mathbb{C}[[t]]$, $t \mapsto t^{l/l'}$. Taking stacky quotients of both sides gives a morphism $\mathcal{D}_{\infty, \mathbb{C}}^l \rightarrow \mathcal{D}_{\infty, \mathbb{C}}^{l'}$. By Lemma 2.15 of [34], there exists a unique positive integer l' dividing l so that γ' factors (up to 2-isomorphism) as $\mathcal{D}_{\infty, \mathbb{C}}^l \rightarrow \mathcal{D}_{\infty, \mathbb{C}}^{l'} \xrightarrow{\psi} \mathcal{X}$, where ψ is representable, and then $g_{\infty}(\gamma) = \psi$.

Let Y be the coarse moduli space of \mathcal{Y} and let Z be the coarse moduli space of \mathcal{Z} . Then $g : \mathcal{Y} \rightarrow \mathcal{Z}$ induces a morphism $g' : Y \rightarrow Z$. Note that, by considering coarse moduli spaces, the morphism $\mathcal{D}_{\infty, \mathbb{C}}^l \rightarrow \mathcal{D}_{\infty, \mathbb{C}}^{l'}$ yields

the identity morphism on $\text{Spec } \mathbb{C}[[t]]$. It follows that we have a commutative diagram

$$(13) \quad \begin{array}{ccc} |\mathcal{J}_\infty \mathcal{Y}| & \xrightarrow{g_\infty} & |\mathcal{J}_\infty \mathcal{Z}| \\ \downarrow \tilde{\pi}_\infty & & \downarrow \tilde{\pi}_\infty \\ J_\infty(Y) & \xrightarrow{g'_\infty} & J_\infty(Z). \end{array}$$

Consider the notation of Theorem 4.2.

Lemma 7.1. *If $\Delta = (N, \Delta, \{\bar{b}_i\})$ is a stacky fan refining $\Sigma = (N, \Sigma, \{b_i\})$, then the birational morphism $f : \mathcal{X}(\Delta) \rightarrow \mathcal{X}(\Sigma)$ induces a $J_\infty(T)$ -equivariant map*

$$f_\infty : |\mathcal{J}_\infty \mathcal{X}(\Delta)|' \rightarrow |\mathcal{J}_\infty \mathcal{X}(\Sigma)|'$$

$$f_\infty(\tilde{\gamma}_v) = \tilde{\gamma}_v.$$

Proof. By considering coarse moduli spaces, $f : \mathcal{X}(\Delta) \rightarrow \mathcal{X}(\Sigma)$ gives rise to the toric morphism $f' : X(\Delta) \rightarrow X(\Sigma)$, with induced map of arc spaces $f'_\infty : J_\infty(X(\Delta))' \rightarrow J_\infty(X(\Sigma))'$. Consider the commutative diagram (13). By Theorem 4.2, the maps $\tilde{\pi}_\infty$ are $J_\infty(T)$ -equivariant bijections satisfying $\tilde{\pi}_\infty(\tilde{\gamma}_v) = \gamma_v$. Hence we only need to show that f'_∞ is $J_\infty(T)$ -equivariant and satisfies $f'_\infty(\gamma_v) = \gamma_v$. This fact is well-known but we recall a proof for the convenience of the reader. Suppose v lies in a cone τ of Δ and let σ be a cone in Σ containing τ . The arc γ_v corresponds to the ring homomorphism $\mathbb{C}[\check{\tau} \cap M] \rightarrow \mathbb{C}[[t]]$, $\chi^u \mapsto t^{\langle u, v \rangle}$ and hence $f'_\infty(\gamma_v)$ corresponds to the ring homomorphism $\mathbb{C}[\sigma^\vee \cap M] \hookrightarrow \mathbb{C}[\check{\tau} \cap M] \rightarrow \mathbb{C}[[t]]$, $\chi^u \mapsto \chi^u \mapsto t^{\langle u, v \rangle}$. We see that f'_∞ is $J_\infty(T)$ -equivariant and $f'_\infty(\gamma_v) = \gamma_v$. \square

We now state Yasuda's transformation rule in the case of toric stacks.

Theorem 7.2 (The Transformation Rule, [34]). *Let $\Delta = (N, \Delta, \{\bar{b}_i\})$ be a stacky fan refining $\Sigma = (N, \Sigma, \{b_i\})$, with corresponding birational morphism $f : \mathcal{X}(\Delta) \rightarrow \mathcal{X}(\Sigma)$. If $F : |\mathcal{J}_\infty \mathcal{X}(\Sigma)|' \rightarrow \mathbb{Q}$ is a $J_\infty(T)$ -invariant function and A is a $J_\infty(T)$ -invariant subset of $|\mathcal{J}_\infty \mathcal{X}(\Sigma)|'$, then*

$$\int_A (uv)^{s_{\mathcal{X}(\Sigma)} + F} d\mu_{\mathcal{X}} = \int_{f_\infty^{-1}(A)} (uv)^{s_{\mathcal{X}(\Delta)} + F \circ f_\infty - \text{ord } K_{\mathcal{X}(\Delta)/\mathcal{X}(\Sigma)}} d\mu_{\mathcal{X}'},$$

where $K_{\mathcal{X}(\Delta)/\mathcal{X}(\Sigma)} = K_{\mathcal{X}(\Delta)} - f^* K_{\mathcal{X}(\Sigma)}$.

As a corollary, we deduce a geometric proof of the following result, which was proved using combinatorial methods in Proposition 2.13 of [30]. If $\Sigma = (N, \Sigma, \{b_i\})$ is a stacky fan and λ is a piecewise \mathbb{Q} -linear function on $|\Sigma|$, we fix the following notation. Let $\psi_\Sigma = \psi_{q_* K_{\mathcal{X}(\Sigma)}}$ be the piecewise \mathbb{Q} -linear function on $|\Sigma|$ satisfying $\psi_\Sigma(b_i) = 1$, and let $\delta_\Sigma^\lambda(t)$ denote the weighted δ -vector corresponding to λ .

Corollary 7.3. *Let N be a lattice of rank d and let $\Sigma = (N, \Sigma, \{b_i\})$ and $\Delta = (N, \Delta, \{b'_j\})$ be stacky fans such that $|\Sigma| = |\Delta|$. Let λ be a piecewise \mathbb{Q} -linear function with respect to Σ satisfying $\lambda(b_i) > -1$ for every b_i , and set $\lambda' = \lambda + \psi_\Sigma - \psi_\Delta$. If λ' is piecewise \mathbb{Q} -linear with respect to Δ and satisfies $\lambda'(b'_j) > -1$ for every b'_j , then $\delta_\Sigma^\lambda(t) = \delta_\Delta^{\lambda'}(t)$.*

Proof. Let $\tilde{\Sigma}$ be a common refinement of Σ and Δ . We can choose lattice points $\{\tilde{b}_i\}$ on the rays of $\tilde{\Sigma}$ so that $\tilde{\Sigma} = (N, \tilde{\Sigma}, \{\tilde{b}_i\})$ is a stacky fan refining Σ and Δ . Hence we can reduce to the case when Δ refines Σ . In this case, consider the corresponding birational morphism $f : \mathcal{X}(\Delta) \rightarrow \mathcal{X}(\Sigma)$. By Lemma 5.1, there is a Kawamata log terminal pair $(\mathcal{X}(\Sigma), \mathcal{E})$ such that $\text{ord } \mathcal{E}(\tilde{\gamma}_w \cdot J_\infty(T)) = -\lambda(w)$. It follows from Lemma 7.1 that $(\text{ord } \mathcal{E} \circ f_\infty)(\tilde{\gamma}_w \cdot J_\infty(T)) = -\lambda(w)$ and Lemma 5.1 and Lemma 7.1 imply that $\text{ord } K_{\mathcal{X}(\Delta)/\mathcal{X}(\Sigma)}(\tilde{\gamma}_w \cdot J_\infty(T)) = \psi_\Sigma(w) - \psi_\Delta(w)$. The result now follows from Theorem 6.5 and Theorem 7.2, with $F = \text{ord } \mathcal{E}$ and $A = |\mathcal{J}_\infty \mathcal{X}(\Sigma)|$. \square

8. REMARKS

More generally, we can replace N by a finitely generated abelian group of rank d . All the results go through with minor modifications. We mention the local construction of the toric stack [5]. Let \bar{N} be the lattice given by the image of N in $N_{\mathbb{R}}$ and for each v in N , let \bar{v} denote the image of v in \bar{N} . Let Σ be a complete, simplicial, rational fan in $N_{\mathbb{R}}$ and let $\bar{v}_1, \dots, \bar{v}_r$ be the primitive integer generators of Σ in \bar{N} . Fix elements b_1, \dots, b_r in N such that $\bar{b}_i = a_i \bar{v}_i$, for some positive integer a_i . The data $\Sigma = (N, \Sigma, \{b_i\})$ is a *stacky fan*. For each maximal cone σ of Σ , let N_σ denote the subgroup of N generated by $\{b_i \mid \rho_i \subseteq \sigma\}$. We obtain a homomorphism of finite groups $N(\sigma) = N/N_\sigma \rightarrow \bar{N}(\sigma) = \bar{N}/\bar{N}_\sigma$. Composing with the injection $\bar{N}(\sigma) \rightarrow (\mathbb{C}^*)^d$ from Section 2 gives a homomorphism $N(\sigma) \rightarrow \bar{N}(\sigma) \rightarrow (\mathbb{C}^*)^d$, and $\mathcal{X}(\sigma) = [\mathbb{A}^d/N(\sigma)]$.

Using this more general setup, one can apply Theorem 4.2 to the T -invariant closed substacks of \mathcal{X} to give a decomposition of $|\mathcal{J}_\infty \mathcal{X}|$ into $J_\infty(T)$ -orbits.

REFERENCES

1. Dan Abramovich, Tom Graber, and Angelo Vistoli, *Algebraic orbifold quantum products*, Orbifolds in mathematics and physics (Madison, WI, 2001), Contemp. Math., vol. 310, Amer. Math. Soc., Providence, RI, 2002, pp. 1–24. MR MR1950940 (2004c:14104)
2. Alexander Barvinok, *Lattice points, polyhedra, and complexity*, Geometric combinatorics, IAS/Park City Math. Ser., vol. 13, Amer. Math. Soc., Providence, RI, 2007, pp. 19–62. MR MR2383125
3. Victor V. Batyrev, *Non-Archimedean integrals and stringy Euler numbers of log-terminal pairs*, J. Eur. Math. Soc. (JEMS) **1** (1999), no. 1, 5–33. MR MR1677693 (2001j:14018)
4. Victor V. Batyrev and Dimitrios I. Dais, *Strong McKay correspondence, string-theoretic Hodge numbers and mirror symmetry*, Topology **35** (1996), no. 4, 901–929. MR MR1404917 (97e:14023)
5. Lev A. Borisov, Linda Chen, and Gregory G. Smith, *The orbifold Chow ring of toric Deligne-Mumford stacks*, J. Amer. Math. Soc. **18** (2005), no. 1, 193–215 (electronic). MR MR2114820 (2006a:14091)
6. Weimin Chen and Yongbin Ruan, *A new cohomology theory of orbifold*, Comm. Math. Phys. **248** (2004), no. 1, 1–31. MR MR2104605 (2005j:57036)
7. J. Denef and F. Loeser, *Motivic integration and the Grothendieck group of pseudo-finite fields*, Proceedings of the International Congress of Mathematicians, Vol. II (Beijing, 2002) (Beijing), Higher Ed. Press, 2002, pp. 13–23. MR MR1957016 (2004f:14040)
8. Jan Denef and François Loeser, *Germes of arcs on singular algebraic varieties and motivic integration*, Invent. Math. **135** (1999), no. 1, 201–232. MR MR1664700 (99k:14002)
9. ———, *Geometry on arc spaces of algebraic varieties*, European Congress of Mathematics, Vol. I (Barcelona, 2000), Progr. Math., vol. 201, Birkhäuser, Basel, 2001, pp. 327–348. MR MR1905328 (2004c:14037)
10. ———, *Motivic integration, quotient singularities and the McKay correspondence*, Compositio Math. **131** (2002), no. 3, 267–290. MR MR1905024 (2004e:14010)

11. Lawrence Ein, Robert Lazarsfeld, and Mircea Mustață, *Contact loci in arc spaces*, Compos. Math. **140** (2004), no. 5, 1229–1244. MR MR2081163 (2005f:14006)
12. Lawrence Ein and Mircea Mustață, *Inversion of adjunction for local complete intersection varieties*, Amer. J. Math. **126** (2004), no. 6, 1355–1365. MR MR2102399 (2005j:14020)
13. Barbara Fantechi, Etienne Mann, and Fabio Nironi, *Smooth toric dm stacks*, arXiv:0708.1254, 2007.
14. William Fulton, *Introduction to toric varieties*, Annals of Mathematics Studies, vol. 131, Princeton University Press, Princeton, NJ, 1993, , The William H. Roever Lectures in Geometry. MR MR1234037 (94g:14028)
15. Takayuki Hibi, *Some results on Ehrhart polynomials of convex polytopes*, Discrete Math. **83** (1990), no. 1, 119–121. MR MR1065691 (91g:52008)
16. ———, *Ehrhart polynomials of convex polytopes, h-vectors of simplicial complexes, and nonsingular projective toric varieties*, Discrete and computational geometry (New Brunswick, NJ, 1989/1990), DIMACS Ser. Discrete Math. Theoret. Comput. Sci., vol. 6, Amer. Math. Soc., Providence, RI, 1991, pp. 165–177. MR MR1143294 (92j:52018)
17. ———, *A lower bound theorem for Ehrhart polynomials of convex polytopes*, Adv. Math. **105** (1994), no. 2, 162–165. MR MR1275662 (95b:52018)
18. ———, *Star-shaped complexes and Ehrhart polynomials*, Proc. Amer. Math. Soc. **123** (1995), no. 3, 723–726. MR MR1249883 (95d:52012)
19. Shihoko Ishii, *The arc space of a toric variety*, J. Algebra **278** (2004), no. 2, 666–683. MR MR2071659 (2005f:14100)
20. Seán Keel and Shigefumi Mori, *Quotients by groupoids*, Ann. of Math. (2) **145** (1997), no. 1, 193–213. MR MR1432041 (97m:14014)
21. Maxim Kontsevich, *Motivic integration*, Lecture at Orsay, 1995.
22. Gérard Laumon and Laurent Moret-Bailly, *Champs algébriques*, Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics], vol. 39, Springer-Verlag, Berlin, 2000. MR MR1771927 (2001f:14006)
23. Mircea Mustață, *Jet schemes of locally complete intersection canonical singularities*, Invent. Math. **145** (2001), no. 3, 397–424, With an appendix by David Eisenbud and Edward Frenkel. MR MR1856396 (2002f:14005)
24. V. Srinivas, *The hodge characteristic*, Lecture in Jet Schemes Seminar, MSRI, 2002.
25. Richard P. Stanley, *Hilbert functions of graded algebras*, Advances in Math. **28** (1978), no. 1, 57–83. MR MR0485835 (58 #5637)
26. ———, *Decompositions of rational convex polytopes*, Ann. Discrete Math. **6** (1980), 333–342, Combinatorial mathematics, optimal designs and their applications (Proc. Sympos. Combin. Math. and Optimal Design, Colorado State Univ., Fort Collins, Colo., 1978). MR MR593545 (82a:52007)
27. ———, *On the Hilbert function of a graded Cohen-Macaulay domain*, J. Pure Appl. Algebra **73** (1991), no. 3, 307–314. MR MR1124790 (92f:13017)
28. ———, *A monotonicity property of h-vectors and h*-vectors*, European J. Combin. **14** (1993), no. 3, 251–258. MR MR1215335 (94f:52016)
29. ———, *Enumerative combinatorics. Vol. 1*, Cambridge Studies in Advanced Mathematics, vol. 49, Cambridge University Press, Cambridge, 1997, With a foreword by Gian-Carlo Rota, Corrected reprint of the 1986 original. MR MR1442260 (98a:05001)
30. Alan Stapledon, *Weighted ehrhart theory and orbifold cohomology*, arXiv:0711.4382, 2007, to appear in Adv. Math.
31. Willem Veys, *Arc spaces, motivic integration and stringy invariants*, Singularity theory and its applications, Adv. Stud. Pure Math., vol. 43, Math. Soc. Japan, Tokyo, 2006, pp. 529–572. MR MR2325153
32. Angelo Vistoli, *Intersection theory on algebraic stacks and on their moduli spaces*, Invent. Math. **97** (1989), no. 3, 613–670. MR MR1005008 (90k:14004)
33. Takehiko Yasuda, *Twisted jets, motivic measures and orbifold cohomology*, Compos. Math. **140** (2004), no. 2, 396–422. MR MR2027195 (2004m:14037)
34. ———, *Motivic integration over Deligne-Mumford stacks*, Adv. Math. **207** (2006), no. 2, 707–761. MR MR2271984 (2007j:14002)

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MICHIGAN, ANN ARBOR, MI 48109, USA

E-mail address: `astapl@umich.edu`