

A set of exactly solvable Ising models with half-odd-integer spin

Onofre Rojas* and S. M. de Souza

Departamento de Ciências Exatas, Universidade Federal de Lavras,

CP 3037, 37200-000, Lavras - MG, Brazil.

November 28, 2018

Abstract

We present a set of exactly solvable Ising models, with half-odd-integer spin- S on a square-type lattice including a quartic interaction term in the Hamiltonian. The particular properties of the mixed lattice, associated with mixed half-odd-integer spin- $(S, 1/2)$ and only nearest-neighbor interaction, allow us to map this system either onto a purely spin- $1/2$ lattice or onto a purely spin- S lattice. By imposing the condition that the mixed half-odd-integer spin- $(S, 1/2)$ lattice must have an exact solution, we found a set of exact solutions that satisfy the *free fermion* condition of the eight vertex model. The number of solutions for a general half-odd-integer spin- S is given by $S + 1/2$. Therefore we conclude that this transformation is equivalent to a simple spin transformation which is independent of the coordination number.

Keywords: Mathematical physics, two dimensional Ising model, exact results.

Two-dimensional lattice models are one of the most interesting subjects of statistical mechanics, both experimentally[1, 2] and theoretically. Several approximation methods are used to investigate these models on the lattice, such as mean-field theory[1, 3], the Bethe approximation[4], the correlated effective field theory[5], the renormalization group[6], series expansion methods[7], Monte Carlo methods[8] and cluster variation methods. Following Onsager's[9] solution for the square two dimensional Ising lattice, other solutions for regular

*Corresponding author. email: onofre@pq.cnpq.br, Tel.: +5535 38291954, Fax.: +5535 38291371

two-dimensional lattices have been considered, such as triangular[10, 11], honeycomb[12, 13], kagomé[14] lattices and others were explored in several works and their importance in statistical physics has waked up the search for a set of completely solvable models. Some other exact results have been obtained with restricted parameters, as investigated by Mi and Yang[16] using a non-one-to-one transformation[15].

Some half-odd-integer spin-S Ising models were already discussed in the literature[17]. Using the method proposed by Wu[18], Izmailian [19] obtained an exact solution for a spin-3/2 square lattice with only nearest-neighbor and two-body spin interactions. Izmailian and Ananikian[20] have also obtained an exact solution for a honeycomb lattice with spin-3/2. A particular case of solutions of these models were obtained by the method proposed by Joseph[21], where any spin-S could be decomposed in terms of spin-1/2. Another interesting method for mapping the spin-S lattice into a spin-1/2 lattice has been proposed by Horiguchi[22].

Mapping between models is an important tool for the study of exactly solvable models. The aim of this letter is to present the transformation of higher half-odd-integer spin-S systems into a simple spin-1/2 system.

To demonstrate this transformation we use a two dimensional mixed spin-(S,1/2) on a square Ising lattice with a quartic interaction \mathcal{L}_b as displayed in fig.1. First this mixed model can be mapped onto an exactly solvable two dimensional spin-1/2 Ising model with a quartic interaction[26] \mathcal{L}_a , such as presented in the literature[19]. Second it is also possible to transform a mixed spin lattice \mathcal{L}_b onto an effective spin-S lattice when we consider the spin-1/2 as a decorated Ising model of the lattice[23] \mathcal{L}_b . Then the classical Hamiltonian for a mixed spin-(3/2,1/2) lattice is given by

$$\mathcal{H}_{1/2,3/2} = \sum_{\langle i,j \rangle} (K_r^{(1)} S_i \sigma_j + K_r^{(3)} S_i^3 \sigma_j) + \sum_i D S_i^2, \quad (1)$$

with $\langle i, j \rangle$ meaning summation over nearest interacting neighbors on the square lattice, and the last summation is performed over all spin-3/2 sites. The coefficient $K_r^{(1)}$ is the nearest-neighbor interaction parameter of the bilinear term; $K_r^{(3)}$ corresponds to the parameter of the non-bilinear interaction, where r runs from 1 up to the coordination number (in our case up to 4); D is the single ion-anisotropy parameter acting on spin-3/2; S_i represents the spin-3/2 particle; whereas σ_i corresponds the spin-1/2 particle, with two possible values ± 1 (we use these values conveniently instead of $\pm 1/2$ for all spin σ_i).

First we would like to obtain an exactly solvable spin-(3/2,1/2) lattice \mathcal{L}_b (see fig.1); to this end, we write

the first term of the Hamiltonian (1) as $(K_r^{(1)}S_i + K_r^{(2)}S_i^3)\sigma_j$, then for spin-3/2 we have four possible values $\pm 1/2$ and $\pm 3/2$, thus the term $K_r^{(1)}S_i + K_r^{(2)}S_i^3$ can be written

$$A_r^{(1)} = \pm \frac{1}{2}K_r^{(1)} \pm \frac{1}{8}K_r^{(3)}, \quad S_i = \pm 1/2, \quad (2)$$

$$A_r^{(2)} = \pm \frac{3}{2}K_r^{(1)} \pm \frac{27}{8}K_r^{(3)}, \quad S_i = \pm 3/2, \quad (3)$$

with $r = 1, \dots, 4$.

In order to project the spin-3/2 onto spin σ with only two possible values ± 1 , we impose the condition $|A_r^{(1)}| = |A_r^{(2)}| = A_r$. Therefore we are able to find the parameters $K_r^{(1)}$ and $K_r^{(3)}$ as a function of A_r , by solving the system (2) and (3), we obtain following

$$\left\{ \begin{array}{l} K_r^{(3)} = -\frac{4}{3}A_r; \quad K_r^{(1)} = \frac{7}{3}A_r \\ \text{or} \\ K_r^{(3)} = -\frac{2}{3}A_r; \quad K_r^{(1)} = -\frac{13}{6}A_r \end{array} \right., \quad \text{in which } r = 1, \dots, 4. \quad (4)$$

The associated Boltzmann weight for a mixed spin-(3/2,1/2) lattice has a similar structure to that of the model discussed by Wu and Lin[24], the associated Boltzmann weights are given by $W(\{\sigma_r\}) = \sum_S \exp(\mathcal{H}_{1/2,3/2})$. For simplicity we consider our calculation in units of $-\beta$. Using the solution given by (4) the associated Boltzmann weights are simplified, which read as

$$\begin{aligned} w_1 &= W(+, +, +, +) = \alpha \cosh(A_1 + A_2 + A_3 + A_4), \\ w_2 &= W(+, -, +, -) = \alpha \cosh(A_1 - A_2 + A_3 - A_4), \\ w_3 &= W(+, -, -, +) = \alpha \cosh(A_1 - A_2 - A_3 + A_4), \\ w_4 &= W(+, +, -, -) = \alpha \cosh(A_1 + A_2 - A_3 - A_4), \\ w_5 &= W(+, -, +, +) = \alpha \cosh(A_1 - A_2 + A_3 + A_4), \\ w_6 &= W(+, +, +, -) = \alpha \cosh(A_1 + A_2 + A_3 - A_4), \\ w_7 &= W(+, +, -, +) = \alpha \cosh(A_1 + A_2 - A_3 + A_4), \\ w_8 &= W(-, +, +, +) = \alpha \cosh(-A_1 + A_2 + A_3 + A_4), \end{aligned} \quad (5)$$

where $\alpha = 2(e^{\frac{D}{4}} + e^{\frac{9D}{4}})$.

The lattice \mathcal{L}_b can be transformed into an effective spin-1/2 lattice \mathcal{L}_a , as shown in fig.1. Then at each site of \mathcal{L}_a there can be eight lines; we are thus led to consider an eight-vertex model \mathcal{L}_a with weights given by (5) and which satisfies the *free fermion* condition[25],

$$\omega_1\omega_2 + \omega_3\omega_4 = \omega_5\omega_6 + \omega_7\omega_8. \quad (6)$$

Consider the following Hamiltonian,

$$\mathcal{H}_{1/2} = J_0 N + \sum_{(i,j)} J_{i,j} \sigma_i \sigma_j + \sum_{\text{all tetra-}} J_{1,2,3,4} \sigma_{i_1} \sigma_{i_2} \sigma_{i_3} \sigma_{i_4} \quad (7)$$

for the effective spin-1/2 Ising model \mathcal{L}_a , which is exactly solvable using the eight-vertex model[26]. Here, (i, j) means summation over all pairs of sites over the tetrahedron (the boldface tetrahedron in Fig.1), and $J_{i,j}$ are their corresponding interacting parameters, whereas $J_{1,2,3,4}$ is the quartic interaction parameter; N represents the number of spin-3/2 sites on the lattice.

After transforming the Hamiltonian (1) into (7), we relate their parameters using the Boltzmann weights, thus to obtain

$$J_0 = \ln(2\alpha) + \frac{1}{8} \ln(w_1 w_2 w_3 w_4 w_5 w_6 w_7 w_8) \quad (8)$$

$$J_{1,2} = \frac{1}{8} \ln\left(\frac{w_1 w_4 w_5 w_8}{w_6 w_7 w_2 w_3}\right) \quad (9)$$

$$J_{2,3} = \frac{1}{8} \ln\left(\frac{w_1 w_8 w_3 w_6}{w_7 w_4 w_5 w_2}\right) \quad (10)$$

$$J_{3,4} = \frac{1}{8} \ln\left(\frac{w_1 w_6 w_7 w_4}{w_5 w_2 w_3 w_8}\right) \quad (11)$$

$$J_{4,1} = \frac{1}{8} \ln\left(\frac{w_1 w_7 w_5 w_3}{w_8 w_4 w_2 w_6}\right) \quad (12)$$

$$J_{1,3} = \frac{1}{8} \ln\left(\frac{w_1 w_7 w_2 w_8}{w_6 w_4 w_5 w_3}\right) \quad (13)$$

$$J_{2,4} = \frac{1}{8} \ln\left(\frac{w_1 w_6 w_5 w_2}{w_7 w_4 w_3 w_8}\right) \quad (14)$$

$$J_{1,2,3,4} = \frac{1}{8} \ln\left(\frac{w_1 w_4 w_2 w_3}{w_8 w_7 w_5 w_6}\right). \quad (15)$$

The partition function \mathcal{Z} of the decorated (or mixed) model in the thermodynamic limit is related to the partition function \mathcal{Z}_{8v} of the effective eight-vertex spin-1/2 model (7) by the expression $\mathcal{Z} = e^{-\beta J_0} \mathcal{Z}_{8v}$. An

analytical expression for the free energy of the *free fermion* model is well known[25] and after some manipulation, was expressed in the thermodynamic limit by

$$f = J_0 + \frac{1}{16\pi^2} \int_0^{2\pi} \int_0^{2\pi} d\theta d\phi \ln[2a + 2b \cos(\theta) + 2c \cos(\phi) + 2d \cos(\theta - \phi) + 2e \cos(\theta + \phi)], \quad (16)$$

where

$$\begin{aligned} a &= \frac{1}{2}(w_1^2 + w_2^2 + w_3^2 + w_4^2), & b &= w_1 w_3 - w_2 w_4, \\ c &= w_1 w_4 - w_2 w_3, & d &= w_3 w_4 - w_7 w_8, & e &= w_3 w_4 - w_5 w_6. \end{aligned} \quad (17)$$

The system exhibits an Ising transition at the critical points

$$w_1 + w_2 + w_3 + w_4 = 2 \max\{w_1, w_2, w_3, w_4\}. \quad (18)$$

At low temperature the system exhibits ordered states such as the ferromagnetic state (for $w_1 > w_2, w_3, w_4$), the antiferromagnetic state (for $w_2 > w_1, w_3, w_4$) and the metamagnetic state (for w_3 or $w_4 > w_1, w_2$).

Thus our goal is to transform an exactly solvable Ising model with spin-1/2 via the eight-vertex model[26] into an equivalent Ising model on the lattice with spin-3/2. For this purpose we introduce an auxiliary lattice \mathcal{L}_b , with mixed spin-1/2 and spin-3/2, the schematic transformation of which is displayed in fig.1. If we consider the spin-1/2 model as a decoration of the mixed spin model and transform it into an equivalent spin-3/2 Ising model on lattice, then we can conclude that there exist a transformation from the spin-1/2 Ising model onto the spin-3/2 Ising model, with a non-bilinear interaction and four-body interactions terms over the tetrahedron or quartic interaction.

Using the first of solution of Eq.(4), we express the Hamiltonian of the transformed spin-3/2 lattice \mathcal{L}_c in terms of four constrained parameters and one arbitrary parameter D , thus obtaining

$$\begin{aligned} \mathcal{H}_{3/2} = & J_0 N + \sum_i D S_i^2 + \sum_{(i,j)} J_{i,j} \left(\frac{49}{9} S_i S_j - \frac{28}{9} (S_i S_j^3 + S_i^3 S_j) + \frac{16}{9} S_i^3 S_j^3 \right) + \sum_{\text{all tetra-}} J_{1,2,3,4} \left(\frac{2401}{81} S_{i_1} S_{i_2} S_{i_3} S_{i_4} \right. \\ & \left. - \frac{1372}{81} S_{i_1}^3 S_{i_2} S_{i_3} S_{i_4} + \frac{784}{81} S_{i_1}^3 S_{i_2}^3 S_{i_3} S_{i_4} - \frac{448}{81} S_{i_1} S_{i_2}^3 S_{i_3}^3 S_{i_4} + \frac{256}{81} S_{i_1}^3 S_{i_2}^3 S_{i_3}^3 S_{i_4}^3 \right), \end{aligned} \quad (19)$$

where the parameters $J_{i,j}$ were already defined by Eqs.(8)-(15), and these parameters are constrained. Whereas the parameters A_1, A_2, A_3 and A_4 are free, as well as the single ion anisotropy parameter D . Summations are performed as indicated in (7).

By factorizing the terms under the summation sign, we obtain

$$\begin{aligned} \mathcal{H}_{3/2} = & J_0 N + \sum_i D S_i^2 + \sum_{\langle i,j \rangle} (J_{i,j} \frac{S_i S_j}{9} (7 - 4S_i^2)(7 - 4S_j^2)) + \\ & + \sum_{\text{all tetra-}} J_{1,2,3,4} \frac{S_{i_1} S_{i_2} S_{i_3} S_{i_4}}{81} (7 - 4S_{i_1}^2)(7 - 4S_{i_2}^2)(7 - 4S_{i_3}^2)(7 - 4S_{i_4}^2). \end{aligned} \quad (20)$$

We can reduce the Hamiltonian (20) further for both solutions in Eq.(4), using the spin transformation $\sigma^{(k)}(S)$, where this function only take two possible values ± 1 , for all values of $S = \{-3/2, -1/2, 1/2, 3/2\}$. These spin transformations read as

$$\sigma^{(k)}(S) = \begin{cases} \frac{S}{3}(7 - 4S^2); & k = 1 \\ \frac{S}{6}(13 - 4S^2); & k = 2. \end{cases} \quad (21)$$

We remark that the transformation given by (21) are the same as those obtained by Izmailian[19] using the method proposed by Wu[18] (but for another model, the spin-3/2 Ising model on a square lattice). As we can see, the previous transformation is independent of the coordination number. Therefore it may be used to yield a particular case of solution, for the exactly solvable spin-3/2 Ising model in a honeycomb lattice as was obtained by Izmailian and Ananikian[20].

We are now able to write the Hamiltonian (20) as

$$\mathcal{H}_{3/2}^{(k)} = J_0 N + \sum_i D S_i^2 + \sum_{\langle i,j \rangle} J_{i,j} \sigma_i^{(k)} \sigma_j^{(k)} + \sum_{\text{all tetra-}} J_{1,2,3,4} \sigma_{i_1}^{(k)} \sigma_{i_2}^{(k)} \sigma_{i_3}^{(k)} \sigma_{i_4}^{(k)}. \quad (22)$$

The second term of the Hamiltonian (22) may be written using a new spin variable transformation, similar to that performed by Izmailian[19]. Note that there the transformation was performed only for spin-3/2 case and for higher spin that method could become a more complex task.

We also remark that the model discussed by Izmailian[19] can be completely re-obtained, by using our simple transformation instead of Wu's[18] method.

The Hamiltonian (1) can be extended to arbitrary half-odd-integer spin-(S,1/2); the general Hamiltonian has the following form

$$\mathcal{H}_{1/2,S} = \sum_{\langle i,j \rangle} \left(K_r^{(1)} S_i + K_r^{(3)} S_i^3 + K_r^{(5)} S_i^5 \dots + K_r^{(S+\frac{1}{2})} S_i^{S+\frac{1}{2}} \right) \sigma_j + \sum_i D S_i^2, \quad (23)$$

where $K_r^{(1)}$ is the bilinear interaction parameter, and $K_r^{(u)}$ are non-bilinear interaction parameters of $S_i^u \sigma_j$, with $u = 2, \dots, S/2 + 1$ and $r = 1, \dots, 4$, thus the set of transformations for each half-odd-integer spin-S have $S + 1/2$ unknown parameters to be determined. We extend eq.(4) to the general case (23) by constructing a Vandermonde-like matrix, explicitly

$$\begin{pmatrix} \frac{1}{2} & (\frac{1}{2})^3 & (\frac{1}{2})^5 & \dots & (\frac{1}{2})^{2S} \\ \frac{3}{2} & (\frac{3}{2})^3 & (\frac{3}{2})^5 & \dots & (\frac{3}{2})^{2S} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ S & S^3 & S^5 & \dots & S^{2S} \end{pmatrix} \begin{pmatrix} K_r^{(1)} \\ K_r^{(3)} \\ \dots \\ K_r^{(S+\frac{1}{2})} \end{pmatrix} = \begin{pmatrix} A_r^{(1)} \\ A_r^{(2)} \\ \vdots \\ A_r^{(S+\frac{1}{2})} \end{pmatrix}, \quad (24)$$

where the parameters $A_r^{(u)}$ must satisfy the following identities $|A_r^{(1)}| = |A_r^{(2)}| = \dots = |A_r^{(S+\frac{1}{2})}| = A_r$, in order to project the spin-S onto spin-1/2. By inverting the Vandermonde-like matrix, we are able to obtain the solution of the algebraic system. In what follows, we show the solutions of these system of linear equations for some higher spin-S values.

For $S = 5/2$, there are three solutions:

$$\sigma^{(k)}(S) = \begin{cases} \frac{1067}{480} S - \frac{11}{12} S^3 + \frac{1}{10} S^5; & k = 1 \\ \frac{529}{240} S - \frac{5}{6} S^3 + \frac{1}{15} S^5; & k = 2 \\ \frac{1183}{480} S - \frac{23}{12} S^3 + \frac{7}{30} S^5; & k = 3. \end{cases} \quad (25)$$

For $S = 7/2$, there are four solutions:

$$\sigma^{(k)}(S) = \begin{cases} \frac{30251}{13440} S - \frac{301}{288} S^3 + \frac{61}{360} S^5 - \frac{1}{126} S^7; & k = 1 \\ \frac{60577}{26880} S - \frac{3047}{2880} S^3 + \frac{127}{720} S^5 - \frac{11}{1260} S^7; & k = 2 \\ \frac{14887}{6720} S - \frac{637}{720} S^3 + \frac{17}{180} S^5 - \frac{1}{315} S^7; & k = 3 \\ \frac{68123}{26880} S - \frac{1289}{576} S^3 + \frac{293}{720} S^5 - \frac{5}{252} S^7; & k = 4. \end{cases} \quad (26)$$

For $S = 9/2$, we obtain five solutions:

$$\sigma^{(k)}(S) = \begin{cases} \frac{5851067}{2580480} S - \frac{46573}{41472} S^3 + \frac{7501}{34560} S^5 - \frac{97}{6048} S^7 + \frac{1}{2592} S^9; & k = 1 \\ \frac{2924921}{1290240} S - \frac{813413}{725760} S^3 + \frac{3727}{17280} S^5 - \frac{239}{15120} S^7 + \frac{17}{45360} S^9; & k = 2 \\ \frac{5868067}{2580480} S - \frac{334907}{290304} S^3 + \frac{8117}{34560} S^5 - \frac{113}{6048} S^7 + \frac{43}{90720} S^9; & k = 3 \\ \frac{5725183}{2580480} S - \frac{37337}{41472} S^3 + \frac{3593}{34560} S^5 - \frac{29}{6048} S^7 + \frac{1}{12960} S^9; & k = 4 \\ \frac{6651283}{2580480} S - \frac{506017}{207360} S^3 + \frac{18341}{34560} S^5 - \frac{1237}{30240} S^7 + \frac{13}{12960} S^9; & k = 5. \end{cases} \quad (27)$$

Using the spin transformations given by eqs.(21), (25), (26) and (27), we are able to recover the transformation obtained by Joseph[21] by considering $k = 1$, whereas the remaining transformation are new.

In this report we present a transformation which maps a spin-3/2 lattice, with quartic and non-linear interactions terms, onto an effective spin-1/2 Ising model on lattice. First this transformation was carried out using an auxiliary mixed spin-(3/2,1/2) square lattice with only nearest neighbor interaction term, mapped onto an effective spin-1/2 or spin-3/2 lattice model, depending on which spin is considered to be the decoration spin. Finally, a systematic way of transformation for higher half-odd-integer spin-S is considered, inverting a Vandermonde like matrix, to obtain a families of mapping between spin-S models and spin-1/2 models, using the one-to-one transformation. Therefore we conclude that there exist a spin transformation, which can be applied to lattice models with arbitrary coordination number, including non-exactly solvable half-odd-integer spin-S models. We also recovered some results previously obtained in the literature[19, 20], as a particular case of solution.

O. Rojas. and S.M. de Souza. thanks CNPq and FAPEMIG for partial financial support.

References

- [1] M. Blume, V. J. Emery, R. B. Griffiths, Phys. Rev. A **3**, 1071 (1971).
- [2] J. Bernasconi and F. Rys, Phys. Rev. B **4**, 3045 (1982).
- [3] H. W. Capel, Physica **32**, 966 (1966); D. Mukamel and M. Blume, Phys. Rev. A **10** 61, (1974).
- [4] K. G. Chakraborty and T. Morita, Physica A **129**, 415 (1985).

- [5] T. Kaneyoshi, *Physica A* **164**, 730 (1976).
- [6] S. Krinsky and D. Furman, *Phys. Rev. B* **11**, 2602 (1975); A. N. Berker and M. Wortis, *Phys. Rev.* **14**, 4946 (1976).
- [7] D. M. Soul, M. Wortis and D. Stauffer, *Phys. Rev. B* **9**, 4964 (1974)
- [8] A. K. Jain and D. P. Landau, *Bull. Am. Phys. Soc.* **21**, 231 (1976).
- [9] Onsager, *Phys. Rev.* **65**,117 (1944).
- [10] G.F. Newell, *Phys. Rev.* **79**, 876 (1950).
- [11] K. Husimi and I. Syozi, *Prog. Theor. Phys.* **5**, 117 (1950).
- [12] T. Horiguchi, *Phys. Lett. A* **113**, 425 (1986).
- [13] K. Husimi(1) and I. Syozi,*Prog. Theor. Phys.* **5**, 341 (1950).
- [14] I. Syozi, *Prog. Theor. Phys.* **6**, 306 (1951).
- [15] M. Kolesík and L. Samaj, *Int. J. Mod. Phys. B* **6**, 1529 (1992).
- [16] X. D. Mi and Z. R. Yang, *J. Phys. A: Math. Gen.* **28**, 4883 (1995); *Phys. Rev E* **49**, 3636 (1994).
- [17] K. Tang, *J. Phys. A: Math. Gen.* **21**, L1097 (1988).
- [18] F. Y. Wu, *Phys. Lett. A* **117**, 365 (1986).
- [19] N. Sh. Izmailian, *Pis'ma Zh. Éksp. Teor. Fiz.* **63**, 270-275 (1996)
- [20] N. Sh. Izmailian and N. S. Ananikian, *Phys. Rev. B* **50**, 6829 (1994).
- [21] R. I. Joseph, *J. Phys. A:Math. Gen.* **9**, L31 (1976).
- [22] T. Horiguchi, *Physica A* **214**,452 (1995).
- [23] J. Strečka, L. Čanová and Ján Dely, *Phys. Stat. Solidi (B)* **243**, 1946 (2006).

- [24] F. Y. Wu and K. Y. Lin, J. Phys. A: Math. Gen. **20** 5737, (1987).
- [25] C. Fan and F. Y. Wu, Phys. Rev. B **2**,723 (1970).
- [26] R.J. Baxter, *Exacty Solved Models in Statistical Mechanics*, Academic Press, N.Y. 1982.




Figure 1: Schematic representation of mixed spin-($S, 1/2$) on a square lattice (\mathcal{L}_b), a square-type spin- $1/2$ Ising model (\mathcal{L}_a) and a square-type spin- S Ising model (\mathcal{L}_c)