

Beyond The Quenched (or Probe Brane) Approximation in Lattice (or Holographic) QCD

Adi Armoni

a.armoni@swan.ac.uk

*Department of Physics, Swansea University
Singleton Park, Swansea, SA2 8PP, UK.*

Abstract

We propose a method to improve the quenched approximation. The method, based on the worldline formalism, takes into account effects of quark loops. The idea is useful for both lattice simulations and for AdS/CFT (holographic) calculations. To demonstrate the method we estimate screening (string breaking) effects by a simple holographic calculation.

Introduction and conclusions. The quenched approximation is a popular method to estimate QCD quantities on the lattice. By neglecting the fermionic determinant, it is possible to carry out relatively cheap and fast calculations. In the AdS/CFT framework there exists a similar approximation called 'the probe brane approximation', where the flavor-brane backreaction is neglected (the Sakai-Sugimoto model is an example of such a setup [1]). Quenching is valid in the 't Hooft limit, where the number of colors is taken to infinity while the number of flavors is kept fixed. In the lattice framework, going beyond the quenched theory is challenging due to both conceptual and technical reasons. Similarly, in the AdS/CFT framework, it is difficult to take into account the backreaction of the flavor branes.¹

In this short letter we propose a simple method to improve the quenched approximation. The idea is to expand the fermionic determinant in powers of Wilson loops, by using the worldline formalism and to keep the leading and the sub-leading contributions. The outcome is a simple way of calculating flavor-sensitive observables in QCD.

We expect the method to be useful for both lattice simulations and holographic calculations. Below we describe our method and demonstrate its usefulness in a simple example.

The method. Consider a calculation of an observable \mathcal{O} in QCD. In the path integral formalism it can be written, after integration over the fermions, as follows

$$\langle \mathcal{O} \rangle = \frac{1}{\mathcal{Z}} \int DA_\mu \mathcal{O} \exp -S_{\text{YM}} \det (i \not{D} - m) . \quad (1)$$

The quenched approximation is obtained by omitting the fermionic determinant from the above expression (1)

$$\langle \mathcal{O} \rangle_{\text{YM}} = \frac{1}{\mathcal{Z}} \int DA_\mu \mathcal{O} \exp -S_{\text{YM}} . \quad (2)$$

Thus $\langle \mathcal{O} \rangle \approx \langle \mathcal{O} \rangle_{\text{YM}}$. For certain quantities the above approximation (2) turns out to be good. Let us propose a way of improving it.

We use the worldline formalism [3] in order to express the fermionic determinant in terms of Wilson loops. The fermionic determinant is related to

¹An interesting exception is [2] where a fully backreacted supersymmetric background was derived.

the Wilson loop as follows

$$\det(i \not{D} - m) = \exp \Gamma[A], \quad (3)$$

where

$$\begin{aligned} \Gamma[A] &= -\frac{1}{2} \int_0^\infty \frac{dT}{T} \\ &\times \int \mathcal{D}x \mathcal{D}\psi \exp \left\{ - \int_\epsilon^T d\tau \left(\frac{1}{2} \dot{x}^\mu \dot{x}^\mu + \frac{1}{2} \psi^\mu \dot{\psi}^\mu - \frac{1}{2} m^2 \right) \right\} \\ &\times \text{Tr} \mathcal{P} \exp \left\{ i \int_0^T d\tau \left(A_\mu \dot{x}^\mu - \frac{1}{2} \psi^\mu F_{\mu\nu} \psi^\nu \right) \right\}, \quad (4) \end{aligned}$$

with $x^\mu(0) = x^\mu(T)$. Thus $\Gamma[A]$ is a sum over (super)-Wilson loops. The sum is over contours of all sizes and shapes. Large contours are, however, suppressed by the quark mass, which serves as an IR cut-off. The sum can be written schematically as $\Gamma[A] = \sum_c W$. In this notation the fermionic determinant is

$$\det(i \not{D} - m) = \exp \sum_c W = \sum_n \frac{1}{n!} \left(\sum_c W \right)^n. \quad (5)$$

The quenched theory is obtained by approximating the exponent in (5) by 1. We propose to improve the approximation by keeping more terms in the above sum (5). In practise it is easy to add the first term. So, our proposal is

$$\langle \mathcal{O} \rangle \approx \langle \mathcal{O} \rangle_{\text{YM}} + \sum_c \langle \mathcal{O} W \rangle_{\text{YM}}^{\text{conn.}}. \quad (6)$$

Namely, we suggest to improve the calculation of an observable \mathcal{O} by adding to the quenched value a sum of its correlator with all possible Wilson loops. The exact weighting is according to (4). Another comment is that only connected terms contribute (terms such as $\langle \mathcal{O} \rangle \langle W \rangle$ are cancelled against the denominator of (1)).

In order to obtain a better understanding of the expansion (5) and the proposal (6) let us consider the case $\mathcal{O} = 1$ in large- N QCD, namely in $SU(N_c)$ Yang-Mills with N_f light quarks ($m \ll \Lambda_{\text{QCD}}$). The choice $\mathcal{O} = 1$,

corresponding to the vacuum energy, is not of great interest, but it will teach us about the validity of the expansion (5). The fully quenched theory contains only gluons, hence the leading large N_c contribution to the vacuum energy is $O(N_c^2)$. In order to 'improve' the calculation, let us use our prescription (6). A single Wilson loop expectation value contributes as $O(N_c N_f)$. In general, the n -th term in the expansion (5) contributes to the vacuum energy as $O(N_c^2 (N_f/N_c)^n)$. Therefore, (5) is a systematic expansion in powers of N_f/N_c . Hence the proposal (6) takes into effect the leading N_f/N_c corrections.

Example. Let us consider a more interesting example in QCD. We wish to demonstrate the screening effect of the light quarks. The observable \mathcal{O} that we choose is a circular fundamental Wilson loop of large radius R (with $R \gg \Lambda_{\text{QCD}}^{-1}$). A calculation of the Wilson loop in the quenched approximation




Figure 1: A confining Wilson loop. A holographic calculation by using an AdS with an IR cut-off background.

will yield an area law

$$\langle \mathcal{O} \rangle_{\text{YM}} = \exp -\sigma A. \quad (7)$$

The above result (7) can be obtained either by a lattice simulation, or by a lattice strong coupling expansion calculation [4], or by a holographic calculation of the Wilson loop [5] in a confining background [6]. Since the theory is quenched string breaking effects of the full theory are not visible.

Let us use our method to improve the result (7). We can demonstrate our method by using either the lattice strong coupling expansion or by a holographic calculation. We use the latter. For simplicity let us choose as a confining Yang-Mills vacuum an AdS metric with an infra-red cut-off

$$ds^2 = \frac{du^2}{u^2} + u^2 dx_i^2 ; u_{\min} < u < \infty. \quad (8)$$

A large Wilson loop in this background will exhibit an area law (7) since the string worldsheet will immediately drop from the AdS boundary to the IR cut-off $u = u_{\min}$ and will 'rest' there, see figure 1.

We now wish to calculate the second term in (6). The connected two-point function of two Wilson loops, is given by the minimal area worldsheet whose boundaries are the Wilson loops [7],

$$\langle \mathcal{O}W \rangle_{\text{YM}}^{\text{conn.}} = \exp -S_{\text{N.G.}} . \quad (9)$$

While we formally need to sum over all possible Wilson loop it is clear from the above expression (9) that the sum will be dominated by worldsheets with minimal surface. Those are given by Wilson loops which lie close to the circular Wilson loop \mathcal{O} . This is the expected effect of dynamical quarks in the fundamental representation: they create a hole in the QCD-string worldsheet. The minimal area is shown in figure 2 below. Since the metric




Figure 2: A correlation function of two Wilson loops $\langle W_1 W_2 \rangle$.

near the boundary is AdS, the minimal area is given by

$$S_{\text{N.G.}} \propto \frac{R}{L} , \quad (10)$$

where L is the difference between the radii of the two Wilson loops. Note that the minimal area is proportional to the perimeter of \mathcal{O} . Thus

$$\sum_c \langle \mathcal{O}W \rangle_{\text{YM}}^{\text{conn.}} \sim \exp -\mu R . \quad (11)$$

Since the sum is over all sizes of Wilson loops, the value of μ is expected to be $\mu \sim \Lambda_{\text{QCD}}$. Note that we work in the light quark limit. In this limit it is

'easy' to produce pairs and to screen the external charges. When the quark mass is large, the exponent $\exp -m^2 T$ in (4) suppresses large Wilson loops and it is 'difficult' to create the large Wilson loop that screens \mathcal{O} . In other words, when $m > \Lambda_{\text{QCD}}$ it is difficult to produce pairs.

Altogether the expectation value of the circular Wilson loop is

$$\langle \mathcal{O} \rangle = c_1 N_c \exp -\sigma A + c_2 N_f \exp -\mu R. \quad (12)$$

c_1 and c_2 are $O(1)$ coefficients. The prefactor N_c arises since the Wilson loop expectation value is proportional to $\dim R$ [8]. The correction is proportional to N_f because the expansion is in powers of N_f/N_c , as discussed in the previous section. The above expression (12) is exactly what we expect in QCD. In the absence of quarks, or in the limit $N_c \rightarrow \infty$, fixed N_f , the theory exhibits an area law. Incorporating quarks leads to screening. This is reflected by the second term in (12). Indeed, when $\Lambda_{\text{QCD}} R > \log \frac{N_c}{N_f}$ the second term in (12) dominates and the QCD-string breaks.

It will be interesting to carry out a lattice simulation which uses our method, in order to observe the QCD-string breaking. We expect the prescription (6) to be useful for many other calculations as well.

Acknowledgements. I wish to thank S. Hands, C. Hoyos, B. Lucini, C. Núñez, A. Patella, and A. Rago for fruitful discussions. I am supported by the PPARC advanced fellowship award.

References

- [1] T. Sakai and S. Sugimoto, "Low energy hadron physics in holographic QCD," *Prog. Theor. Phys.* **113**, 843 (2005) [arXiv:hep-th/0412141].
- [2] R. Casero, C. Nunez and A. Paredes, "Towards the string dual of $N = 1$ SQCD-like theories," *Phys. Rev. D* **73**, 086005 (2006) [arXiv:hep-th/0602027].
- [3] M. J. Strassler, "Field theory without Feynman diagrams: One loop effective actions," *Nucl. Phys. B* **385**, 145 (1992) [arXiv:hep-ph/9205205].
- [4] K. G. Wilson, "Confinement of Quarks," *Phys. Rev. D* **10**, 2445 (1974).
- [5] J. M. Maldacena, "Wilson loops in large N field theories," *Phys. Rev. Lett.* **80**, 4859 (1998) [arXiv:hep-th/9803002].

- [6] A. Brandhuber, N. Itzhaki, J. Sonnenschein and S. Yankielowicz, “Wilson loops, confinement, and phase transitions in large N gauge theories from supergravity,” JHEP **9806**, 001 (1998) [arXiv:hep-th/9803263].
- [7] D. E. Berenstein, R. Corrado, W. Fischler and J. M. Maldacena, “The operator product expansion for Wilson loops and surfaces in the large N limit,” Phys. Rev. D **59**, 105023 (1999) [arXiv:hep-th/9809188].
- [8] A. Armoni and B. Lucini, “Universality of k-string tensions from holography and the lattice,” JHEP **0606**, 036 (2006) [arXiv:hep-th/0604055].