Joint measurement of two unsharp observables of a qubit

Sixia Yu,^{1,2} Nai-le Liu,¹ Li Li,¹ and C. H. Oh²

¹Hefei National Laboratory for Physical Sciences at Microscale and Department of Modern Physics,

University of Science and Technology of China, Hefei, Anhui 230026, P.R. China

²Centre for Quantum Technologies and Physics Department,

National University of Singapore, 2 Science Drive 3, Singapore 117542

We present a single inequality as the necessary and sufficient condition for two unsharp observables of a two-level system to be jointly measurable in a single apparatus and construct explicitly the joint observables. A complementarity inequality arising from the condition of joint measurement, which generalizes Englert's duality inequality, is derived as the trade-off between the unsharpnesses of two jointly measurable observables.

PACS numbers: 03.65.Ta, 03.67.-a

Built in the standard formalism of quantum mechanics, there are mutually exclusive but equally real aspects of quantum systems, as summarized by the complementarity principle of Bohr [\[1\]](#page-3-0). Mutually exclusive aspects are often exhibited via noncommuting observables, for which the complementarity is quantitatively characterized by two kinds of uncertainty relationships, namely, the preparation uncertainty relationships (PURs) and the measurement uncertainty relationships (MURs).

The PURs stem from the semi positive definiteness of the density matrix describing the quantum state and characterize the predictability of two noncommuting observables in a given quantum state. To test PURs two different projective measurements will be performed on two identically prepared ensembles of the quantum system and these measurements cannot be performed within one experimental setup on a single ensemble.

On the other hand MURs characterize the trade-off between the precisions of unsharp measurements of two noncommuting observables in a single experimental setup. The very first effort of Heisenberg [\[2\]](#page-3-1) in deriving the uncertainty relationships was based on a simultaneous measurement of the position and momentum, with the rigorous form of MUR established recently by Werner [\[3\]](#page-3-2). In the interferometry the wave-particle duality between the path-information and the fringe visibility of interference pattern is characterized quantitatively by Englert's duality inequality [\[4\]](#page-3-3), which turns out to be originated from the joint measurability of two special unsharp observables encoding the path information and the fringe visibility [\[5](#page-3-4)]. To establish a general MUR the condition for joint measurement has to be explored, which can be turned into some kinds of MURs when equipped with proper measure of the precisions (e.g., distinguishability).

In this Letter we shall consider the joint measurability of two general unsharp observables of a qubit and derive a simple necessary and sufficient condition with joint observables explicitly constructed. We also present a MUR arising from the condition of joint measurement that generalizes Englert's duality inequality.

Joint measurability — Generally an observable is described by a positive-operator valued measure (POVM), a set of positive operators $\{O_k\}_{k=1}^K$ summed up to the identity $(O_k \geq 0)$

and $\sum_{k} O_k = I$) with K being the number of outcomes. By definition, a *joint measurement* of two observables $\{O_k\}$ and ${O'_l}$ is described by a *joint observable* ${M_{kl}}$ whose outcomes can be so grouped that

$$
O_k = \sum_l M_{kl}, \quad O'_l = \sum_k M_{kl}.\tag{1}
$$

Here we shall consider the qubits, any two-level systems such as spin-half systems or two-path interferometries. A *simple observable* $\mathcal{O}(x, \vec{m})$ refers to a most general 2-outcome POVM $\{O_{\pm}(x, \vec{m})\}$ with

$$
O_{\pm}(x,\vec{m}) = \frac{1 \pm (x + \vec{m} \cdot \vec{\sigma})}{2}.
$$
 (2)

Here $m = |\vec{m}|$ is referred to as the *sharpness* while |x| is referred to as the *biasedness*. When $|x| = 0$ the observable $\mathcal{O}(x, \vec{m})$ is called as *unbiased*, in which case the outcomes of measurement are purely random if the system is in the maximally mixed state, and when $|x| \neq 0$ the observable is referred to as *biased*, in which case priori information can be employed to make better use of the outcomes of the measurement. Positivity imposes $|x| + m \leq 1$.

Given two simple observables $\mathcal{O}(x, \vec{m})$ and $\mathcal{O}(y, \vec{n})$, it is obvious that all possible sets of four operators satisfying the marginal constraints Eq. [\(1\)](#page-0-0) are

$$
M_{\mu\nu}(Z,\vec{z}) = \frac{1 + \mu x + \nu y + \mu \nu Z + (\mu \nu \vec{z} + \vec{q}_{\mu\nu}) \cdot \vec{\sigma}}{4} \tag{3}
$$

with Z , \vec{z} being arbitrary and $\vec{q}_{\mu\nu} = \mu \vec{m} + \nu \vec{n}$ ($\mu, \nu = \pm 1$). The problem of joint measurability becomes whether there exist Z, \vec{z} such that $M_{\mu\nu}(Z, \vec{z}) \geq 0$ for all $\mu, \nu = \pm 1$. There are many partial results in special cases [\[5,](#page-3-4) [8,](#page-3-5) [9](#page-3-6)] as well as in general cases [\[6,](#page-3-7) [7](#page-3-8)]. Here we shall present a single inequality as the condition. For convenience we denote

$$
F_x = \frac{1}{2} \left(\sqrt{(1+x)^2 - m^2} + \sqrt{(1-x)^2 - m^2} \right), \text{(4a)}
$$

$$
F_y = \frac{1}{2} \left(\sqrt{(1+y)^2 - n^2} + \sqrt{(1-y)^2 - n^2} \right).
$$
 (4b)

Theorem 1 Two observables $\mathcal{O}(x, \vec{m})$ and $\mathcal{O}(y, \vec{n})$ are jointly measurable iff

$$
(1 - F_x^2 - F_y^2) \left(1 - \frac{x^2}{F_x^2} - \frac{y^2}{F_y^2} \right) \le (\vec{m} \cdot \vec{n} - xy)^2.
$$
 (5)

Due to the identities such as $x^2/F_x^2 + m^2/(1-F_x^2) = 1$ the left-hand-side of Eq. (5) can be seen to be bounded above by left-hand-sideo[f E](#page-0-1)q. (5) can be seen to be bounded above by $(mn - |xy|)^2$ so that the trivial case $s = 0$ with $s = |\vec{m} \times \vec{n}|$ is included. In the case of $x = y = 0$ Eq. [\(5\)](#page-0-1) reproduces the condition $m^2 + n^2 \leq 1 + (\vec{m} \cdot \vec{n})^2$ for unbiased observables [\[8](#page-3-5)]. When $y = 0$ the condition Eq. [\(5\)](#page-0-1) reads $F_x \sqrt{m^2 - (\vec{m} \cdot \vec{n})^2} \geq s$ which becomes simply $F_x \geq n$ for orthogonal observables where $\vec{m} \cdot \vec{n} = 0$ [\[5\]](#page-3-4).

More generally we refer a pair of observables that satisfy More generally we refer a pair of observables that satisfy $\gamma = 0$ where $\gamma = \vec{m} \cdot \vec{n} - xy$ to as *orthogonal unsharp ob-*servables. The condition of joint measurement Eq. [\(5\)](#page-0-1) becomes simply $F_x^2 + F_y^2 \ge 1$ because $x^2/F_x^2 + y^2/F_y^2 < 1$ is ensured by $mn > |xy|$. In general the condition $F_x^2 + F_y^2 \ge 1$ is sufficient for joint measurement since Eq. (5) is ensured because $(|xy| - mn)^2 \le \gamma^2$ when $mn < xy$. Specifically we refer a pair of observables that satisfy $\gamma = 0$ and $\vec{n} = \vec{n}_c$ with $n_c^2/F_x^2 + y^2/(1 - F_x^2) = 1$ to as a pair of *maximally* \overrightarrow{or} *orthogonal unsharp observables*. It is maximal because any between $\mathcal{O}(y, \vec{n})$ with $n \leq n_c$ (regardless of its direction) is jointly measurable with $\mathcal{O}(x, \vec{m})$ while all the observables $\mathcal{O}(y, \vec{n})$ with $n > n_c$ along \vec{n}_c are not jointly measurable with $\mathcal{O}(x, \vec{m}).$ \therefore generally we feler a pair of observables that satisfy $\sum_{i=1}^{n}$ inclusion with $\sum_{i=1}^{n}$ ι).

As the measure for unsharpness we take a linear combination of the sharpness and the biasedness, i.e., $D_1 = Q_1 m +$ $P_1|x|$ and $D_2 = Q_2 n + P_2|y|$ with $0 \le P_i \le Q_i$ $(i = 1, 2)$ being some constants. To measure jointly a pair of orthogonal unsharp observables there is a trade-off between the above defined unsharpnesses (since $D_1^2 + (Q_1^2 - P_1^2)F_x^2 \leq Q_1^2$)

$$
D_1^2(Q_2^2 - P_2^2) + D_2^2(Q_1^2 - P_1^2) + P_1^2 P_2^2 \le Q_1^2 Q_2^2.
$$
 (6)

Englert's duality inequality [\[4](#page-3-3)] in the case of orthogonal ob- $\frac{1}{2}$ stating inequality $\frac{1}{1}$ in the case of orthogonal or servables with one being unbiased [\[5](#page-3-4)] turns out to be a special case of the above inequality if we let $Q_1 = 1$, $P_2 = 0$ so that D_1 and D_2 become the path distinguishability and the fringe visibility respectively. Theorem 1 is derived from the following set of conditions. $f(x)$ above mequality if we fet Q_1

Theorem 2 Two observables $\mathcal{O}(x, \vec{m})$ and $\mathcal{O}(y, \vec{n})$ $(\vec{m} \times \vec{m})$ $\vec{n} \neq 0$) are jointly measurable iff either max $\{|\alpha|, |\beta|\} \geq 1$ or

$$
\sum_{\nu=\pm} |\vec{m} + \vec{n} + \nu \vec{g}| + \sum_{\nu=\pm} |\vec{m} - \vec{n} + \nu \vec{g}| \le 4, \tag{7}
$$

with $\vec{g} = \vec{m}\alpha + \vec{n}\beta$ and

$$
\alpha = \frac{1}{|\vec{m} \times \vec{n}|^2} \left[(y + \gamma x) n^2 - (x + \gamma y) \vec{m} \cdot \vec{n} \right], \quad \text{(8a)}
$$

$$
\beta = \frac{1}{|\vec{m} \times \vec{n}|^2} \left[(x + \gamma y) m^2 - (y + \gamma x) \vec{m} \cdot \vec{n} \right].
$$
 (8b)

Now let us examine the set of all observables $\mathcal{O}(y, \vec{n})$ with a given biasedness y that are jointly measurable with a fixed observable $\mathcal{O}(x, \vec{m})$. The admissible region of \vec{n} is shown in Fig. 1(a) as the union of a red- and a blue-contoured regions with boundary given by Eq. [\(5\)](#page-0-1) with equality and $|y|+n=1$. The (blue) arcs of the circle $n = 1 - |y|$ satisfying Eq. [\(5\)](#page-0-1) define a forward and a backward admissible cones around

FIG. 1: (a) The union of the blue and red contoured regions, deter-FIG. 1: (a) The union of the blue and red contoured regions, determined by $\max\{|{\alpha}|, |{\beta}|\} \geq 1$ and $R \geq 0$ respectively, represents the admissible \vec{n} in the case of $m = 0.8, x = -0.1$, and $y = 0.3$. The boundary lies between two circles $n = 1 - |y|$ and $n = n_c$. (b) The trade-off curve (solid black) between sharpnesses m and n with
The trade-off curve (solid black) between sharpnesses m and n with $x = -0.1, y = 0.2$, and $\cos \theta = 0.3$ fixed.

 \vec{m} (centered on the origin) in which all \vec{n} are admissible. If $1 - F_x^2 \le |y|$ then Eq. [\(5\)](#page-0-1) holds true and two cones overlap so that all \vec{n} are admissible as formulated as one part of conditions in [\[7](#page-3-8)]. From Lemma 3 iv) we see that Eq. [\(7\)](#page-1-0) is equivalent to $R \geq 0$ with

$$
R = 1 + x^2 + y^2 + \gamma^2 - m^2 - n^2 - |\vec{g}|^2 \tag{9}
$$

 T_{max} the sharpness m, n as T_{max} (Fig. (FF)) is Eq. which appears in an equivalent form $(Eq. (55))$ in [\[6\]](#page-3-7).

Taking the sharpnesses m, n as measures for precisions we have plotted their trade-off curve (solid black) in Fig. 1(b) with x, y , and θ fixed where θ is the angle between \vec{m} and \vec{n} . There is a critical value m_0 of the sharpness determined by Eq. [\(5\)](#page-0-1) with equality and $|y| = 1 - n$, below which there is no
constraint an $\sum_{n=1}^{\infty}$ If constraint on *n*. If

$$
(1 + \text{sgn}[xy] \cos \theta)(1 - |x|)(1 - |y|) \le 2|xy|,
$$
 (10)

m, $\begin{bmatrix} \ell \end{bmatrix}$ is defined similar simila with $\text{sgn}[f] = +1$ if $f \ge 0$ and -1 if $f < 0$, then $m_0 \ge$ $1-|x|$ (and $n_0 \geq 1-|y|$) so that there is no trade-off between m, n. Here n_0 is defined similarly as m_0 with (x, m) and (y, n) interchanged. If $m + n + |\vec{m} \pm \vec{n}| \leq 2$ every vector $\vec{g} = \vec{m}\alpha + \vec{n}\beta$ with $\max\{|\alpha|, |\beta|\} \leq 1$ satisfies Eq. [\(7\)](#page-1-0) so that there is no trade-off between x, y .

Joint unsharp observables $-$ If $s = 0$ then a joint observable of observables $\mathcal{O}(x, \vec{m})$ and $\mathcal{O}(y, \vec{n})$ is simply given by $\{O_\mu(x, \vec{m})O_\nu(y, \vec{n})\}$. If $\Delta_\tau < 0$ with $\Delta_\tau = (\vec{m} - \vec{n})O_\nu(y, \vec{n})$ $(\tau \vec{n})^2 - (x - \tau y)^2$ for some $\tau = \pm$ then $O_\eta(x, \vec{m})$ – $O_{\eta \cdot \tau}(y, \vec{n}) \geq 0$ where $\eta = \text{sgn}[x - \tau y]$. Therefore the $\mathcal{O}_{\eta \cdot \tau}(y, n) \geq 0$ where $\eta = \text{sgn}[x - iy]$. Therefore the POVM $\{0, O_{\bar{\eta}}(x, \vec{m}), O_{\eta}(x, \vec{m}) - O_{\eta \cdot \tau}(y, \vec{n}), O_{\eta \cdot \tau}(y, \vec{n})\}$ is $\sigma_{\eta}(x, m), \sigma_{\eta}(x, m) \sim \sigma_{\eta}(\tau, y, m), \sigma_{\eta}(\tau, y, m)$ a joint observable $(\bar{\eta} = -\eta)$. When $s > 0$ and $\Delta_{\pm} \ge 0$ we have:

Filave.
Theorem 3 Given observables $\mathcal{O}(x, \vec{m})$ and $\mathcal{O}(y, \vec{n})$, a) if $R \geq 0$ then ${M_{\mu\nu}(\gamma, \vec{g})}$ is a joint observable; b) if $R < 0$ and max $\{|\alpha|, |\beta|\} \ge 1$ then $\{M_{\mu\nu}(Z(\vec{z}_{\eta\tau}), \vec{z}_{\eta\tau})\}$ is a joint observable where observable where

$$
Z(\vec{z}) = \max_{\mu = \pm 1} \left\{ |\vec{z} + \mu(\vec{m} + \vec{n})| - \mu(x + y) \right\} - 1, \tag{11}
$$

$$
\vec{z}_{\eta\tau} = \vec{g} + \frac{D_{\eta\tau}(\vec{m} \times \vec{n}) \times \vec{L}_{\eta\tau}}{\vec{L}_{\eta\tau}^2 - |\vec{m} \times \vec{n}|^2},\tag{12}
$$

FIG. 2: The setup for proofs. In the plane $\mathbb P$ spanned by \vec{m} and \vec{n} there are two neighboring ellipses E_x^+ and E_y^+ and the generalized ellipse E with 4 foci $Q_{\mu\nu}$ (red curve).

with $D_{\eta\tau} = \tau A_{\eta}\alpha + \eta B_{\tau}\beta + \eta\tau\gamma - 1$, $\vec{L}_{\eta\tau} = \tau A_{\eta}\vec{n} - \eta B_{\tau}\vec{m}$, $\tau = \text{sgn}[\alpha]$ and $\eta = \text{sgn}[B_{\tau}\beta + \tau\gamma - x]$ if $|\alpha| \geq 1$, and $\eta = \text{sgn}[\beta]$ and $\tau = \text{sgn}[A_{\eta}\alpha + \eta\gamma - y]$ if $|\beta| \geq 1$, where $A_{\eta} = 1 - \eta x$ and $B_{\tau} = 1 - \tau y$.

The regions for different constructions of joint observables according to the above theorem are indicated in Fig. 1(b) whenever two observables are jointly measurable. We note that $\Delta = \min\{\Delta_{\pm}\} < 0$ infers $\max\{|\alpha|, |\beta|\} \geq 1$. Since R = $d_{\mu\nu}^2$ – $|\vec{g} - q_{\mu\nu}|^2$ for all μ, ν = \pm with $d_{\mu\nu}$ = $1 - \mu x - \nu y + \mu \nu \gamma$, we see that if $R = 0$ then $4M_{\mu\nu}(\gamma, \vec{g}) =$ $d_{\bar{\mu}\bar{\nu}} + (\mu\nu\vec{g} + \vec{q}_{\mu\nu}) \cdot \vec{\sigma}$ are proportional to some projections for all $\mu, \nu = \pm$.

Proofs — We shall at first prove 3 relevant Lemmas in which we make use of the fact that given two overlapping convex regions in a plane either their boundaries intersect or one region belongs to the other. In what follows we suppose $s > 0$. In the plane P spanned by \vec{m} and \vec{n} we denote by

$$
\mathbb{E}_x^{\mu} = \left\{ \vec{z} \in \mathbb{P} \middle| \sum_{\tau=\pm} |\vec{z} - \vec{q}_{\tau\mu}| \le 2(1 - \mu x) \right\}, (13)
$$

$$
\mathbb{E}_y^{\nu} = \left\{ \vec{z} \in \mathbb{P} \middle| \sum_{\tau=\pm} |\vec{z} - \vec{q}_{\nu\tau}| \le 2(1 - \nu y) \right\} (14)
$$

four elliptical regions with boundaries being four ellipses E^{\pm}_x and E_y^{\pm} whose semi-major and squared semi-minor axes are denoted by $A_\mu = 1 - \mu x$, $B_\nu = 1 - \nu y$ and $a_\mu = A_\mu^2 - m^2$, $b_{\nu} = B_{\nu}^2 - n^2$ respectively. Two neighboring ellipses E_x^{μ} and E_y^{ν} have one focus $Q_{\nu\mu}$ (corresponding to vector $\vec{q}_{\nu\mu}$) in common. Also we denote by

$$
\mathbb{E} = \left\{ \left| \vec{z} \in \mathbb{P} \right| \sum_{\mu,\nu=\pm} \left| \vec{z} - \vec{q}_{\mu\nu} \right| \le 4 \right\} \tag{15}
$$

an oval region with boundary being a generalized ellipse E with four foci $Q_{\mu\nu}$ with $\mu, \nu = \pm$. The condition Eq. [\(7\)](#page-1-0) becomes $\vec{g} \in \mathbb{E}$. It is easy to see that $\mathbb{J}_x := \mathbb{E}_x^+ \cap \mathbb{E}_x^- \subset$ E, $\mathbb{J}_y := \mathbb{E}_y^+ \cap \mathbb{E}_y^- \subset \mathbb{E}$ with boundaries given by $J_x =$ $(E_x^+ \cup E_x^-) \cap \mathbb{E}$ and $J_y = (E_y^+ \cup E_y^-) \cap \mathbb{E}$ respectively. Furthermore $E_x^+ \cap E_x^- \subset E \cap J_x$, $E_y^+ \cap E_y^- \subset E \cap J_y$. In Fig. 2 two neighboring ellipses with intersections and the 4-ellipse E (red curve) are shown.

Lemma 1 Two observables $\mathcal{O}(x, \vec{m})$ and $\mathcal{O}(y, \vec{n})$ are jointly measurable iff $\mathbb{J} = \mathbb{E}_x^+ \cap \mathbb{E}_x^- \cap \mathbb{E}_y^+ \cap \mathbb{E}_y^- \neq \emptyset$.

Proof If $\mathcal{O}(x, \vec{m})$ and $\mathcal{O}(y, \vec{n})$ are jointly measurable then

there exist Z and \vec{z} such that $M_{\mu\nu}(Z, \vec{z}) \geq 0$, i.e.,

$$
|\mu\nu\vec{z} + \vec{q}_{\mu\nu}| \le 1 + \mu x + \nu y + \mu\nu Z,\tag{16}
$$

for all $\mu, \nu = \pm$. As a result $\vec{z}-(\vec{z}\cdot\vec{s})\vec{s}/s^2 \in \mathbb{J}$ with $\vec{s} = \vec{m}\times\vec{n}$. If there exists $\vec{z} \in \mathbb{J}$ then Eq. [\(16\)](#page-2-0) holds true with Z given by $Z(\vec{z})$ as in Eq. [\(11\)](#page-1-1), i.e., $\{M_{\mu\nu}(Z(\vec{z}), \vec{z})\}$ is a joint observable.

Lemma 2 $\mathbb{J} \neq \emptyset$ iff either $E_x^{\mu} \cap E_y^{\nu} \cap \mathbb{E} \neq \emptyset$ for some $\mu,\nu=\pm$ or $E^{\mu}_{x}\cap E^{\nu}_{y}=\emptyset$ for all $\mu,\nu=\pm.$

Proof Sufficiency. Suppose that there exists $\vec{z} \in E_x^{\mu} \cap E_y^{\nu} \cap E_z^{\nu}$ E for some $\mu, \nu = \pm$. From $\vec{z} \in E_x^{\mu} \cap \mathbb{E}$ and $\vec{z} \in E_y^{\nu} \cap \mathbb{E}$ it follows that $\vec{z} \in \mathbb{E}_x^{\bar{\mu}}$ and $\vec{z} \in \mathbb{E}_y^{\bar{\nu}}$ respectively, which leads to $\vec{z} \in \mathbb{J}$. If $E_x^{\mu} \cap E_y^{\nu} = \emptyset$ for all μ, ν then, taking into account of of Lemma 3 ii), we have either $\mathbb{E}_x^{\mu} \subset \mathbb{E}_y^{\pm} \subset \mathbb{E}_x^{\bar{\mu}}$ or $\mathbb{E}^\nu_y\subset\mathbb{E}^\pm_x\subset\mathbb{E}^{\bar{\nu}}_y,$ i.e., either $\mathbb{J}=\mathbb{E}^\mu_x$ or $\mathbb{J}=\mathbb{E}^\nu_y$ for some $\mu,\nu,$ which is obviously not empty.

Necessity. If $\mathbb{J} \neq \emptyset$ then two convex regions \mathbb{J}_x and \mathbb{J}_y overlap. As a result we have either $J_x \cap J_y \neq \emptyset$, which means $(\exists \mu, \nu) E_x^{\mu} \cap E_y^{\nu} \cap \mathbb{E} \neq \emptyset$, or $J_x \cap J_y = \emptyset$ with either $\mathbb{J}_x \subset \mathbb{J}_y$ or $\mathbb{J}_y \subset \mathbb{J}_x$. If $\mathcal{J}_x \cap \mathcal{J}_y = \emptyset$ and $\mathbb{J}_x \subset \mathbb{J}_y$, i.e., \mathbb{J}_x lies totally within $\mathbb{J}_y \subset \mathbb{E}$, then $J_x \cap E = \emptyset$ which infers $E_x^+ \cap E_x^- =$ \emptyset , i.e., the boundaries of two overlapping regions \mathbb{E}_x^{\pm} do not intersect. As a result $(\exists \mu) \mathbb{E}_x^{\mu} \subset \mathbb{E}_x^{\bar{\mu}}$ so that $(\exists \mu) \mathbb{E}_x^{\mu} = \mathbb{J}_x \subset$ \mathbb{E}_y^{\pm} , which infers $(\exists \mu)$ $E_x^{\mu} \cap E_y^{\pm} = \emptyset$ since $J_x \cap J_y = \emptyset$. In the same manner $\mathbb{J}_y\subset \mathbb{J}_x$ with $\mathring{J}_x\cap J_y=\emptyset$ infers $(\exists \nu)\ E_y^\nu\cap$ $E_x^{\pm} = \emptyset$. In both cases, considering Lemma 3 i), we obtain $(\forall \mu, \nu) E_x^{\mu} \cap E_y^{\nu} = \emptyset.$

Lemma 3 i) $E_x^{\mu} \cap E_y^{\nu} \neq \emptyset$ iff $\Delta_{\mu \cdot \nu} \geq 0$; ii) $\mathbb{E}_x^{\mu} \subset \mathbb{E}_y^{\nu}$ infers $\mathbb{E}^{\bar{\nu}}_y\, \subset\, \mathbb{E}^{\bar{\mu}}_x;$ iii) $\left(\exists \mu,\nu\right)^{\tilde{}} E^{\mu}_x \cap E^{\nu}_y \cap \mathbb{E}\, \neq\, \emptyset$ iff either $\tilde{R}\, \geq\, 0,$ or $(\exists \mu, \nu, \tau)$ $D_{\mu\nu} \geq 0$ and $\Delta_{\tau} \geq 0$; iv) $R \geq 0$ iff $\vec{g} \in \mathbb{E}$; v) $(\forall \mu) \Delta_{\mu} < 0$ infers $(\exists \mu, \nu) D_{\mu\nu} > 0$; vi) Provided $R < 0$, $(\forall \mu, \nu) D_{\mu\nu} < 0$ iff $|\alpha| < 1$ and $|\beta| < 1$.

Proof i) Consider the straight line passing through two points $Q_{\bar{\nu}\mu}$ and $Q_{\nu\bar{\mu}}$ (dashed line in Fig. 2 for the case of $\mu = \nu = +$). If $\Delta_{\mu \cdot \nu} \geq 0$ then one intersection of E_x^{μ} (or E_y^{ν}) with the straight line will not lie in the interior of \mathbb{E}_y^{ν} (or \mathbb{E}_x^{μ} respectively) which means neither $\mathbb{E}_x^{\mu} \subset \mathbb{E}_y^{\nu}$ nor $\mathbb{E}_y^{\nu} \subset \mathbb{E}_x^{\mu}$ and hence $E_x^{\mu} \cap E_y^{\nu} \neq \emptyset$. If $\Delta_{\mu \nu} < 0$ then, e.g., $A_{\mu}^{'}-B_{\nu}$ $>$ $|\vec{m}-\mu\nu\vec{n}|$ and $\vec{z}\in E_{y}^{\nu}$ infers $|\vec{z}-\vec{q}_{\nu\mu}|+|\vec{z}-\vec{q}_{\bar{\nu}\mu}| \leq$ $2B_{\nu} + 2|\vec{m} - \mu\nu\vec{n}| < 2A_{\mu}$, i.e., $E^{\mu}_{x} \cap E^{\nu}_{y} = \emptyset$.

ii) $\mathbb{E}_x^{\mu} \subset \mathbb{E}_y^{\nu}$ is equivalent to $\Delta_{\mu \cdot \nu} \leq 0$, i.e., $\Delta_{\bar{\mu} \cdot \bar{\nu}} \leq 0$, and $A_{\mu} \leq B_{\nu}$, i.e., $B_{\bar{\nu}} \leq A_{\bar{\mu}}$.

iii) Suppose $\vec{z} \in E_x^{\mu} \cap E_y^{\nu} \cap \mathbb{E}$ for some $\mu, \nu = \pm$. Since $\vec{z} \in E^{\mu}_{x} \cap E^{\nu}_{y}$ we have $\Delta_{\mu \cdot \nu} \geq 0$, $r + |\vec{r} + 2\nu \vec{m}| = 2A_{\mu}$ and $r+|\vec{r}+2\mu\vec{n}|=2B_\nu$ where $\vec{r}=\vec{z}-\vec{q}_{\nu\mu}$ and $r=|\vec{r}|.$ It follows that $\vec{s} \times \vec{r} = \vec{K}_{\mu\nu} - r\vec{L}_{\mu\nu}$ whose square provides a quadratic equation of $r: (L_{\mu\nu}^2 - s^2)r^2 - 2r\vec{K}_{\mu\nu} \cdot \vec{L}_{\mu\nu} + K_{\mu\nu}^2 = 0$ where $\vec{K}_{\mu\nu} = \nu a_\mu \vec{n} - \mu b_\nu \vec{m}$, $L_{\mu\nu} = |\vec{L}_{\mu\nu}|$ and $K_{\mu\nu} = |\vec{K}_{\mu\nu}|$. By noticing $L^2_{\mu\nu} > s^2$ as long as $s > 0$ we obtain two solutions

$$
r_{\mu\nu}^{(\pm)} = d_{\mu\nu} + \frac{s^2 D_{\mu\nu} \pm \sqrt{s^2 a_{\mu} b_{\nu} \Delta_{\mu\nu}}}{L_{\mu\nu}^2 - s^2} \tag{17}
$$

and we denote $E_x^{\mu} \cap E_y^{\nu} = \{ \vec{z}_{\mu\nu}^{(+)}, \vec{z}_{\mu\nu}^{(-)} \}$ with $\vec{z}_{\mu\nu}^{(\pm)} = \vec{q}_{\nu\mu}$ + $\vec{r}_{\mu\nu}^{(\pm)}$ and $s^2 \vec{r}_{\mu\nu}^{(\pm)} = (\vec{K}_{\mu\nu} - r_{\mu\nu}^{(\pm)} \vec{L}_{\mu\nu}) \times \vec{s}$. The condition $(\exists \tau) \ \vec{z}_{\mu\nu}^{(\tau)} \in \mathbb{E}$, i.e., $2(A_{\mu} + B_{\nu}) - r_{\mu\nu}^{(\tau)} + |\vec{r}_{\mu\nu}^{(\tau)} + 2\vec{q}_{\nu\mu}| \leq 4$, is equivalent to $(\exists \tau)$ $r_{\mu\nu}^{(\tau)} \geq d_{\mu\nu} - \min\{d_{\bar{\mu}\bar{\nu}}, 0\}$. Due to

$$
s^{2}a_{\mu}b_{\nu}\Delta_{\mu\cdot\nu} = s^{4}D_{\mu\nu}^{2} + s^{2}R(L_{\mu\nu}^{2} - s^{2})
$$
 (18)

and Eq. [\(17\)](#page-2-1), it follows from $(\exists \tau)$ $r_{\mu\nu}^{(\tau)} \ge d_{\mu\nu}$ that either $R \ge$ 0, or $R < 0$ and $D_{\mu\nu} \geq 0$. Necessity is thus proved.

If $\Delta_{\pm} \geq 0$ then $(\forall \mu, \nu) d_{\mu\nu} \geq 0$ since $2d_{\mu\nu} = \Delta_{\mu \cdot \bar{\nu}} + a_{\mu} + a_{\mu}$ b_{ν} . Thus from $(\exists \mu, \nu)$ $D_{\mu\nu} \geq 0$ and $R < 0$ it follows that $r^{(\pm)}_{\mu\nu} \ge d_{\mu\nu}$ which infers $\vec{z}^{(\pm)}_{\mu\nu} \in \mathbb{E}$ so that $E^{\mu}_{x} \cap E^{\nu}_{y} \cap \mathbb{E} \ne \emptyset$. If $(\exists \tau) \Delta_{\tau} < 0$ and $\Delta_{\bar{\tau}} \geq 0$ then $(\forall \nu)$ $E_x^{\nu} \cap E_y^{\tau \cdot \nu} = \emptyset$ and $(\forall \nu) E_x^{\nu} \cap E_y^{\bar{\tau} \cdot \nu} \neq \emptyset$. It follows that either $\mathbb{E}_x^{\nu} \subset E_y^{\tau \cdot \nu}$ or $\mathbb{E}^{\tau\cdot\nu}_y\subset\mathbb{E}^\nu_x$, i.e., either $\mathbb{E}^{\tau\cdot\bar{\nu}}_y\subset\mathbb{E}^{\bar{\nu}}_x$ or $\mathbb{E}^{\bar{\nu}}_x\subset\mathbb{E}^{\tau\cdot\bar{\nu}}_y$. As a result either $\mathbb{J}=\mathbb{E}_x^\nu\cap\mathbb{E}_y^{\tau\cdot\bar{\nu}}$ or $\mathbb{J}=\mathbb{E}_x^{\bar{\nu}}\cap\mathbb{E}_y^{\tau\cdot\nu}$ from which it follows that $(\exists \nu) E_x^{\nu} \cap E_y^{\tilde{\tau} \cdot \tilde{\nu}} \subset \mathbb{E}$, i.e., $(\exists \tau, \nu) E_x^{\nu} \cap E_y^{\tau \cdot \tilde{\nu}} \cap \mathbb{E} \neq \emptyset$.

If $R \geq 0$ then we claim that $\Delta_\pm \geq 0,$ from which it follows immediately that $(\forall \mu, \nu)$ $E_x^{\mu} \cap E_y^{\nu} \neq \emptyset$ and $\vec{z}_{\mu\nu}^{(+)} \in \mathbb{E}$. Firstly, if $a_{\pm} = 0$ (or $b_{\pm} = 0$) then $R \ge 0$ infers $s = 0$, which is precluded. Secondly, if either ($\forall \mu, \nu$) $a_{\mu}b_{\nu} > 0$, or ($\exists \mu$) $a_{\mu} =$ 0 and $a_{\bar{\mu}} > 0$ and $b_{\pm} > 0$, or $(\exists \nu)$ $b_{\nu} = 0$ and $b_{\bar{\nu}} > 0$ and $a_{\pm} > 0$, then the claim is obviously true due to identity Eq. [\(18\)](#page-3-9). Thirdly, if $(\exists \mu, \nu)$ $a_{\mu} = b_{\nu} = 0$ and $a_{\bar{\mu}}b_{\bar{\nu}} > 0$ then $R = 0$, $D_{\mu\nu} = D_{\mu\bar{\nu}} = D_{\bar{\mu}\nu} = 0$ with $D_{\bar{\mu}\bar{\nu}} = -4$, and $\Delta_{\mu \cdot \nu} = \Delta_{\bar{\mu} \cdot \bar{\nu}} > 0$. As a result $r^{(\pm)}_{\mu \nu} = d_{\mu \nu} \geq 0$ which leads to $\Delta_{\mu\cdot\bar{\nu}}=2d_{\mu\nu}\geq 0$.

iv) If $R \ge 0$ then $(\forall \mu, \nu) d_{\mu\nu} \ge 0$ so that $(\forall \mu, \nu) |\vec{q} - \vec{q}|$ $\vec{q}_{\nu\mu} | \leq d_{\mu\nu}$, which infers $\vec{g} \in \mathbb{E}$. If $\vec{g} \in \mathbb{E}$ then $(\exists \mu) \ \vec{g} \in \mathbb{E}_x^{\mu}$. As a result $a_{\mu} - A_{\mu}d_{\mu+} = (\vec{g} - \vec{q}_{+\mu})\cdot \vec{m} \le a_{\mu} - |\vec{g} - \vec{q}_{+\mu}|A_{\mu}$ which infers either $|\vec{g} - \vec{q}_{+\mu}| \le d_{\mu+}$, i.e., $R \ge 0$, or $A_{\mu} = 0$ which leads to $R = 4y^2 \ge 0$.

v) $\Delta_{\pm} < 0$ infers $R < 0$, i.e., $(1 \pm \gamma)^2 < \Delta_{\mp} + |\vec{g}|^2$, and thus $|\vec{g}| > 1 + |\gamma|$. Let $\eta = \text{sgn}[\beta]$ and $\tau = \text{sgn}[\alpha]$ then $|\vec{g}| \leq A_{\eta}|\alpha| + B_{\tau}|\beta| \leq D_{\eta\tau} + 1 + |\gamma|$ which means $D_{\eta\tau} > 0$.

vi) If $(\forall \mu, \nu)$ $D_{\mu\nu} < 0$ then $|\alpha| < 1$ and $|\beta| < 1$ since $\max\{D_{-+}, D_{+-}\} + \max\{D_{--}, D_{++}\} < 0$. If $|\alpha| < 1$ and $|\beta|$ < 1 then $|\vec{g} - \vec{q}_{\nu\mu}| \leq A_{\mu} (1 - \nu\alpha) + B_{\nu} (1 - \mu\beta)$ which, together with $(\forall \mu, \nu) d_{\mu\nu} < |\vec{g} - \vec{q}_{\nu\mu}|$ inferred from $R < 0$, leads to $(\forall \mu, \nu)D_{\mu\nu} = d_{\mu\nu} - A_{\mu}(1-\nu\alpha) - B_{\nu}(1-\mu\beta) < 0.$

Proof of Theorem 2 From Lemmas 1 and 2 and statements i), iii), and v) of Lemma 3 it follows that two observables are jointly measurable iff either $R \ge 0$ or $(\exists \mu, \nu) D_{\mu\nu} \ge 0$ and Theorem 2 is an immediate result of statements iv) and vi) of Lemma 3.

Proof of Theorem 3 a) $R \ge 0$ is equivalent to $(\forall \mu, \nu) | \vec{q} - \vec{q}|$ $|\vec{q}_{\mu\mu}| \leq d_{\mu\nu}$, which means $(\forall \mu, \nu)$ $M_{\mu\nu}(\gamma, \vec{g}) \geq 0$.

b) From $\Delta_{\pm}\geq 0$ it follows that $(\forall \mu,\nu)\;E^{\mu}_{x}\cap E^{\nu}_{y}\neq\emptyset$ and $d_{\mu\nu} \geq 0$ and from $\max\{|\alpha|, |\beta|\} \geq 1$ and the choice of η, τ as in Theorem 3.b it follows that $D_{\eta\tau} \geq 0$. As a result $\{\vec{z}^{(\pm)}_{\eta\tau}\} =$ $E^{\eta}_x \cap E^{\tau}_y \subset \mathbb{E}$ so that $\vec{z}^{(\pm)}_{\eta \tau} \in \mathbb{J}$ (Lemma 2). Since $\mathbb J$ is convex we obtain $\vec{z}_{\eta\tau} = (\vec{z}_{\eta\tau}^{(+)} + \vec{z}_{\eta\tau}^{(-)})/2 \in \mathbb{J}$ and $M_{\mu\nu}(Z(\vec{z}_{\eta\tau}), \vec{z}_{\eta\tau})$ is a joint observable (Lemma 1).

From now on s may be 0. For simplicity we denote by Π_i $(i = 1, 2, 3, 4)$ four functions $s^2(\pm \alpha - 1)$ and $s^2(\pm \beta - 1)$ and $\Pi = \max_i {\Pi_i}$. A set of iff conditions for joint measurement reads $s^2 R \ge 0$ or $\Pi \ge 0$. We have

Lemma 4 $\Pi = 0$ infers $s^2 R \ge 0$.

Proof a) If $s > 0$ then $\Pi = 0$ infers $\max\{|\alpha|, |\beta|\} = 1$, e.g., $|\alpha| = 1$ and $|\beta| \le 1$. Thus $|\vec{g} - \vec{q}_{\nu\mu}| = (1 - \mu\beta)n \le$ $(1 - \mu \beta)B_{\nu} \le d_{\mu\nu}$ which is exactly $R \ge 0$. Here $\nu = \text{sgn}[\alpha]$ and $\mu = \text{sgn}[B_{\nu}\beta + \nu\gamma - x]$. b) If $s = 0$ then $\Pi = 0$ infers $s^2 \alpha = s^2 \beta = 0$ and thus $s^2 R = 0$.

Proof of Theorem 1 We have only to prove that Eq. [\(5\)](#page-0-1) is equivalent to either $s^2 R \ge 0$ or $\Pi \ge 0$. From the identity $s^2 R = (\gamma^2 - f_-)(f_+ - \gamma^2)$, where f_- is the l.h.s. of Eq. [\(5\)](#page-0-1) and $f_+ = f_- + \sqrt{a_+ a_- b_+ b_-}$, it follows that $s^2 R \geq 0$ is equivalent to $f_-\leq \gamma^2\leq f_+$. Thus we have only to show that $\gamma^2 \ge f_+$ infers $\Pi \ge 0$ and that $\Pi \ge 0$ infers $\gamma^2 \ge$ f –. We notice first of all that Π_i are four quadratic (or linear) functions of $c = \vec{m} \cdot \vec{n}$ by regarding x, y, m, n as parameters and Π is continuous. Case a) $F_x^2 + F_y^2 \leq 1$. In this case $mn \ge |xy|$ and $f_{\pm} \ge 0$ and $\Pi \le 0$ for $c = xy$ since $|y| \le F_y^2$ and $F_y^2(n^2 - x^2) \le m^2 n^2 - x^2 y^2$. Now that $\Pi \ge 0$ for $c =$ $\pm mn$, there exist $-mn \leq c_-\leq xy \leq c_+ \leq mn$ such that $\Pi=0$ for $c=c_\pm,$ which infers $xy\pm\sqrt{f_\mp}\leq c_\pm\leq xy\pm\sqrt{f_\pm}$ (Lemma 4). If $\gamma^2 \ge f_+$ then $c \le c_-$ or $c \ge c_+$, which ensures $\Pi \geq 0$ since all the coefficients of c^2 of Π_i $(i = 1, 2, 3, 4)$ are nonnegative. If $\Pi \ge 0$ then $c \le c_$ or $c \ge c_+$, which infers $\gamma^2 \ge f_-.$ Case b) $F_x^2 + F_y^2 \ge 1$. In this case $\gamma^2 \ge f_-$ always and we have only to show that $\gamma^2 \ge f_+$ infers $\Pi \ge 0$. If $\Pi = 0$ has no solution then $\Pi > 0$ for all *c* since $\Pi > 0$ for $c = \pm mn$. Let $c_-\leq c_+$ be its two solutions and it follows that $(c_{\pm} - xy)^2 \le f_+$. As a result if $\gamma^2 \ge f_+$ then $c \ge c_+$ or $c \leq c_$, which ensures $\Pi \geq 0$.

Remarks — We have derived a single inequality as the condition for the joint measurement of two simple qubit observables, based on which an example of MUR is established that generalizes the existing results. On finishing this work two references [\[6](#page-3-7), [7](#page-3-8)] provide two seemingly different solutions to the same problem considered here, whose equivalency can be established in an analytical or a half-numerical and halfanalytical way (see Appendix) via our results. The problems of the joint measurability of more than two observables or observables with more than 2 outcomes are left open.

Acknowledgement — This work is supported by the NNSF of China, the CAS, the Anhui Provincial Natural Science Foundation (Grant No. 070412050) and the A*STAR grant R-144-000-189-305.

- [1] N. Bohr, Nature (London) **121**, 580 (1928).
- [2] W. Heisenberg, Z. Phys. **43**, 172 (1927).
- [3] R. Werner, Quantum Inf. Comput. **4**, 546, (2004).
- [4] B.-G. Englert, Phys. Rev. Lett. **77**, 2154 (1996).
- [5] N.-L. Liu, *et al*, [arXiv:0712.3653v](http://arxiv.org/abs/0712.3653)2 [quant-ph].
- [6] P. Busch and H.-J. Schmidt, arXiv: 0802.4167 [quant-ph].
- [7] P. Stano, D. Reitzner, and T. Heinosaari, arXiv: 0802.4248 [quant-ph].
- [8] P. Busch, Phys. Rev. D **33**, 2253 (1986).
- [9] P. Busch and T. Heinosaari, arXiv: 0706.1415 [quant-ph].
- [10] S. Yu, *et al*, arXiv: 0805.1538 [quant-ph].

Comparison with known results— Here we shall formulate those results in [\[6,](#page-3-7) [7\]](#page-3-8) in our notations and and examine the boundary of admissible \vec{n} by fixing y, x, m . The same boundary means the equivalency.

SRH Theorem [\[7\]](#page-3-8) Two observables $\mathcal{O}(x, \vec{m})$ and $\mathcal{O}(y, \vec{n})$ are jointly measurable iff either

\n- (C1)
$$
\sqrt{1-|y|} \leq F_x
$$
; or
\n- (C2) $\sqrt{1-|y|} > F_x$ and $|\gamma| \geq l$; or
\n- (C3) $\sqrt{1-|y|} > F_x$ and $|\gamma| < l$ and $\sqrt{a_+ h_-} + \sqrt{a_- h_+} \geq 2s$.
\n

Here $s = |\vec{m} \times \vec{n}|$, and $\gamma = \vec{m} \cdot \vec{n} - xy$, and $a_{\pm} = (1 \mp x)^2 - m^2$, and $h_{\pm} = m^2 - (\gamma \pm y)^2$, and

$$
l = \sqrt{y^2 + m^2 - |y|(1 - x^2 + m^2)}.
$$

Remarks. The corresponding boundary is plotted in Fig. 3(a) (with the same parameters as in Fig. 1(a)). If $(C1)$ then $F_x^2 + F_y^2 \ge 1$ so that Eq. [\(5\)](#page-0-1) holds always true. If (C2) then, by noticing that the left-hand-side $f_$ of Eq. [\(5\)](#page-0-1) can be rewritten as

$$
f_{-} = \frac{(a_{+} + 2x)(b_{+} + 2y) - \sqrt{a_{-}a_{+}b_{-}b_{+}}}{2} + m^{2} + n^{2} - 1,
$$
\n(19)

we have $f_- \leq l$ so that Eq. [\(5\)](#page-0-1) holds true. If $1 - |y| > F_x^2$ and $|\gamma| < l$ then $|\gamma| < m - |y|$ so that Lemma 4.a applies and Eq. [\(20\)](#page-4-0) coincides with Eq. [\(5\)](#page-0-1). Thus we have reproduced the boundary in [\[7\]](#page-3-8) analytically.

BS Theorem [\[6\]](#page-3-7) Two observables $\mathcal{O}(x, \vec{m})$ and $\mathcal{O}(y, \vec{n})$ are jointly measurable iff either

(53) $4\Delta_{+} s^2 \leq a_+ b_+ (\vec{L}_{--}^2 - s^2)$; or (54) $4\Delta_+s^2 \leq a_-b_-(\vec{L}_{++}^2 - s^2)$; or

$$
(55) 4\Delta_+s^2 \le 2(A+B_+-c)(A-B_--c)(s^2-\vec{L}_{++}\cdot\vec{L}_{--})
$$

-(A+B_+-c)²($\vec{L}_{--}^2-s^2$)- $(A_-B_--c)^2(\vec{L}_{++}^2-s^2)$.

Here $s = |\vec{m} \times \vec{n}|$, and $\Delta_+ = (\vec{m} - \vec{n})^2 - (x - y)^2$, $a_{\pm} =$ $(1 \mp x)^2 - m^2$, and $b_{\pm} = (1 \mp y)^2 - n^2$, and,

$$
\vec{L}_{\mu\mu} = \mu(1 - \mu x)\vec{n} - \mu(1 - \mu y)\vec{m},
$$

and $A_{\pm} = 1 \mp x$ and $B_{\pm} = 1 \mp y$ and $c = \vec{m} \cdot \vec{n}$.

Remarks. Despite the facts that we have identified Eq. (55) with $R \ge 0$ (Lemma 4.b) and that the boundaries $R = 0$ and $|y| + n = 1$ and

$$
4\Delta_{+}s^2=\max_{\mu=\pm}\{a_{\mu}b_{\mu}(\vec{L}^2_{\bar{\mu}\bar{\mu}}-s^2)\}
$$

intersect at exactly where $\max\{|\alpha|, |\beta|\} = 1$ and that numerical evidences indicate that BS conditions also give rise to the same boundary, we fail to work out an analytical proof for the equivalency so far. The corresponding boundary is plotted in Fig. 3(b). The red-contoured region comes from $R \geq 0$

FIG. 3: The boundary of admissible \vec{n} arising from (a) SRH conditions; (b) BS conditions with fixed $m = 0.8$, $x = -0.1$, and $y = 0.3$

while the blue-contoured region comes from the conditions Eqs.(53,54).

Lemma 5 a) Either $R \ge 0$ or $\{|\beta| \ge 1 \text{ and } h_{\pm} \ge 0\}$ iff

$$
\sqrt{a_{+}h_{-}} + \sqrt{a_{-}h_{+}} \ge 2s.
$$
 (20)

b) Condition Eq. (55) is equivalent to $R \geq 0$.

Proof a) If $R \ge 0$ then $\vec{g} \in \mathbb{E}_x^+ \cap \mathbb{E}_x^-$ so that $h_{\pm} \ge 0$ and $|(1 \pm \beta)s| \leq \sqrt{a_{\mp}h_{\pm}}$ which infers Eq. [\(20\)](#page-4-0). If $|\beta| \geq 1$ and $h_{\pm} \ge 0$, since $4\beta s^2 = h_{+}a_{-} - h_{-}a_{+}$, then $4s^2 \le h_{+}a_{-} +$ $h_{-}a_{+}$ and Eq. [\(20\)](#page-4-0) follows. On the other hand if Eq. [\(20\)](#page-4-0) holds true then $h_{\pm} \geq 0$ and $(\exists \mu)$ $(1 - \mu \beta)|s| \leq \sqrt{a_{\mu} h_{\bar{\mu}}}$ which infers either $|\beta| \ge 1$ or $\vec{g} \in \mathbb{E}_x^{\mu}$, i.e., $R \ge 0$.

b) It follows from the identities $A_+B_+ + A_-B_- - 2c =$ $2(1 - \gamma)$ and $(A_+B_+ - c)\vec{L}_{--} + (A_-B_- - c)\vec{L}_{++} = 2(y +$ $\gamma x)\vec{n} - 2(x + \gamma y)\vec{m}$ whose length squared is equal to $4s^2|\vec{g}|^2$ and $R = (1 - \gamma)^2 - |\vec{g}|^2 - \Delta_+$.

Conclusion— We have proved the equivalency between SRH conditions [\[7](#page-3-8)] and ours analytically and the equivalency between BS conditions [\[6](#page-3-7)] and ours (so that SRH conditions) half-analytically and half-numerically.