Uncertainty relations for arbitrary measurement in terms of Rényi entropies

Alexey E. Rastegin

Department of Theoretical Physics, Irkutsk State University, Gagarin Bv. 20, Irkutsk 664003, Russia

Uncertainty relations for a pair of arbitrary measurements and for a single measurement are posed in the form of inequalities using the Rényi entropies. The formulation deals with discrete observables. Both the relations with state-dependent and state-independent bounds are presented. The entropic bounds are illustrated within the distinction between non-orthogonal states.

PACS numbers: 03.65.Ta, 03.67.-a

The Heisenberg uncertainty principle [1] is first and most known of those results that stress the primary features of the quantum world. The progress of quantum theory has lead to a few related insights such as the Bell inequality [2], the quantum Zeno effect [3], the nocloning theorem [4] and the interaction-free measurement [5]. Although these points stint our ability in manipulating quantum, they also clear novel ways. For example, the Ekert scheme of quantum cryptography is based on Bell's theorem [6]. The techniques similar to Zeno-effect behavior can be used for reducing decoherence in quantum information processing [7]. The well-known quantitative form of the uncertainty principle was given by Robertson [8]. The standard deviations of observables A and B measured in the quantum state $|\psi\rangle$ satisfy

$$\Delta \mathsf{A} \, \Delta \mathsf{B} \ge \frac{1}{2} \left| \langle \psi | [\mathsf{A}, \mathsf{B}] | \psi \rangle \right| \,. \tag{1}$$

Due to the various scenarios of measurement, more specified relations have been posed [9, 10, 11]. For example, Bohr's principle of complementarity [12] was quantified by uncertainty relations. The most wide of these were obtained in Refs. [10, 13]. In general, the key property of complementarity was studied by Kendon and Sanders [14]. Their approach follows the information-theoretic viewpoint proposed by Wootters and Zurek [15].

The recent progress in the study of quantum information processing shows that information theory gives a good base for dealing with quantum properties. The concept of entropy is much widely used in physics, both classical and quantum [16, 17]. So it is natural to state Heisenberg's principle in terms of entropies. The discussion of entropic uncertainty relations with references will be given below. Of late years, reformulations of the uncertainty principle have been posed, especially in information-theoretic terms: in terms of the skew Wigner–Yanase information [18], the Fisher information [19, 20], and the Holevo information [21]. The uncertainty relation of Landau-Pollak type [22] and the one for two unitary operators [23] should also be cited. Such relations are beyond the scope of the present work.

Now both the entropic uncertainty relations and the Rényi entropy have been widely adopted in the researches of quantum systems. For example, the entropic relations were applied to study of the entanglement [24], the locking effect [25] and the special classes of observables [26, 27]. The topic of paper by Bialynicki-Birula [28] is most close to the scope of the present work. He obtained the uncertainty relations in terms of Rényi entropies for the position-momentum and angle-angular momentum pairs. The paper [28] also contains a list of references on fruitful applications of the Rényi entropy to quantum problems. Here we mention only recent papers [29] in which the Rényi entropy of quantum state is utilized.

The aim of the present work is to formulate the uncertainty relations in terms of the Rényi entropies for arbitrary measurements. Important as the concrete observables are, they is not able to give meaning of measurement limitations in all respects. The exposition is not restricted to von Neumann measurements described by "Projector-Valued Measure" (PVM). We shall focus on generalized ones described by "Positive Operator-Valued Measure" (POVM). Recall that POVM $\{M_i\}$ is a set of positive operators M_i satisfying [30]

$$\sum_{i} \mathsf{M}_{i} = \mathbf{1} , \qquad (2)$$

where **1** is the identity operator. This improved approach to quantum measurements allows to extract more information from a quantum system than von Neumann measurements [30]. The authors of Ref. [31] showed how to perform a generalized measurement via a programmable quantum processor. There are a few ways to pose the uncertainty principle for POVM measurement [32]. In more recent paper by Massar [33], the uncertainty relations for POVM's are obtained by introducing the uncertainty operator. He also examined the entropic uncertainty relation in terms of the Shannon entropies for POVM's, whose elements are all rank one [33].

The first entropic relation was proposed by Hirschman [34]. He obtained the relation for position and momentum in terms of the Shannon entropies. Hirschman also conjectured an improvement of his result. This conjecture has been proved by Beckner [35] and by Bialynicki-Birula and Mycielski [36]. A general proper formulation of the uncertainty principle was asked by Deutsch [37]. He emphasized that the right-hand side of Eq. (1) is depended on $|\psi\rangle$. This leads to a trivial lower bound for the deviations product, namely zero for any eigenstate of either of operators A and B [38]. Deutsch obtained a lower bound on the sum of the Shannon entropies for two observables without degeneracy. This formulation has been completed by Partovi in some aspects [39]. In particular, the case of degenerate observables has been revealed. The position-momentum and angle-angular momentum pairs were examined in Ref. [39]. Improved bounds for the both pairs were then obtained in Ref. [40].

It turned out that the entropic uncertainty relation given in Ref. [37] can significantly be improved. The sharpened relation has been conjectured by Kraus [41] and then established by Maassen and Uffink [38]. In Ref. [42] Hall wrote that the proof given by Maassen and Uffink directly extends to the case of two POVM's, whose elements are all rank one. However, the formulation stated in Ref. [38] deals with two non-degenerate observables. A relevant extension for two degenerate observables has been obtained by Krishna and Parthasarathy [43]. Using Naimark's theorem, they also got a lower bound on the sum of the Shannon entropies for two generalized measurements. There is extension of the result of Ref. [38] to more than two observables [44]. This result has also been sharpened in the cases of real [45] and complex two-dimensional spaces [46, 47].

Along with the Shannon entropy, other information entropies are extensively used in the literature. One of them is so-called "Tsallis entropy." This measure introduced by Havrda and Charvát [48] became widely used in statistical mechanics after the fruitful work of Tsallis [49]. The uncertainty relations in terms of the Tsallis entropies were obtained for the position-momentum pair [50] and the spin components [51]. Another generalization of the Shannon entropy has been introduced by Rényi [52]. Larsen derived uncertainty relations in terms of the so-called purities [53]. The purity is immediately connected with the Rényi entropy of order two.

We shall now describe the notation that is used throughout the text. All logarithms are to the base two. Let $\alpha > 0$ and $\alpha \neq 1$; then the Rényi entropy of order α of probability distribution $\{p_i\}$ is defined by [52]

$$H_{\alpha} := \frac{1}{1 - \alpha} \log \left\{ \sum_{i} p_{i}^{\alpha} \right\} . \tag{3}$$

This information measure is a nonincreasing function of parameter α ; that is, if $\alpha < \beta$ then $H_{\alpha} \geq H_{\beta}$ [52]. In the limit $\alpha \to 1$ the Rényi entropy tends to the Shannon entropy $H_1 \equiv -\sum_i p_i \log p_i$. For given measurement $\{\mathsf{M}_i\}$ and state ρ , the probability of *i* th outcome is equal to $p_i = \operatorname{tr}\{\mathsf{M}_i\rho\}$ [30]. The entropy $H_{\alpha}(\mathsf{M}|\rho)$ of generated probability distribution is then defined by Eq. (3).

Let $\{M_i\}$ and $\{N_j\}$ be two POVM's. By definition, we put the function

$$f(\mathsf{M},\mathsf{N}|\psi) := \max_{ij} \frac{|\langle \psi|\mathsf{M}_i\mathsf{N}_j|\psi\rangle|}{||\mathsf{M}_i^{1/2}|\psi\rangle||\,||\mathsf{N}_j^{1/2}|\psi\rangle||} , \qquad (4)$$

where the maximum is taken over those values of labels i and j that the denominator is nonzero. In the case of mixed state ρ with spectral decomposition

$$\rho = \sum_{\lambda} \lambda |\psi_{\lambda}\rangle \langle \psi_{\lambda}| \tag{5}$$

we also define

$$f(\mathsf{M},\mathsf{N}|\rho) := \max_{\lambda} f(\mathsf{M},\mathsf{N}|\psi_{\lambda}) .$$
 (6)

For a single POVM $\{M_i\}$ we put the function

$$\phi(\mathsf{M}|\rho) := \max_{i} \operatorname{tr}\{\mathsf{M}_{i}\rho\} . \tag{7}$$

Below all the entropic bounds will be posed in terms of the defined functions. The first result of the present work is stated as follows.

Relation 1 For arbitrary two measurements $\{M_i\}$ and $\{N_j\}$, there holds

$$H_{\alpha}(\mathsf{M}|\rho) + H_{\beta}(\mathsf{N}|\rho) \ge -2\log f(\mathsf{M},\mathsf{N}|\rho) , \qquad (8)$$

where orders α and β satisfy $1/\alpha + 1/\beta = 2$.

In view of the previous result, this generalization seems to be plausible in itself. However, the proof of Relation 1 requires a considerable alteration of the techniques in many respects. Writing out these tedious details, a perception of the final results would be laboured. So the proof is given in the comprehensive paper [54], which presents a more complete account. It is valuable to get an entropic bound for a single POVM. The ordinary way is to put $\{M_i\} = \{N_j\}$ [33, 43]. But only for $\alpha \leq 1$ this way holds due to the nonincrease property [28]. Meantime, the study of the Rényi entropies of order two and three has a clear physical motivation [55]. In Ref. [54] the following relation is obtained by other method.

Relation 2 For arbitrary measurement $\{M_i\}$ and each $\alpha > 0$, there holds

$$H_{\alpha}(\mathsf{M}|\rho) \ge -\log\phi(\mathsf{M}|\rho) . \tag{9}$$

The lower bound (8) has been proved under the condition $1/\alpha + 1/\beta = 2$. Bialynicki-Birula obtained the entropic uncertainty relations for position and momentum under the same condition [28]. When orders α and β are not coupled in this way, we can write only

$$H_{\alpha}(\mathsf{M}|\rho) + H_{\beta}(\mathsf{N}|\rho) \ge -\log\left[\phi(\mathsf{M}|\rho)\phi(\mathsf{N}|\rho)\right] .$$
(10)

This inequality obtained from Eq. (9) is valid for any two POVM's and arbitrary $\alpha, \beta \in (0; +\infty)$.

The entropic bounds presented above are dependent on state before measurement. As it is already mentioned, in the Robertson relation (1) such a dependence leads to some unsuitability. In a sense, for the presented entropic relations this criticism may be refuted (for details, see [54]). Nevertheless, we can at once get the stateindependent bounds. It is easy to check [54] that

$$f(\mathsf{M},\mathsf{N}|\rho) \le \max_{ij} ||\mathsf{M}_i^{1/2}\mathsf{N}_j^{1/2}|| , \qquad (11)$$

where the norm of operator Q is defined by

$$||\mathbf{Q}|| := \max_{\langle u|u\rangle=1} ||\mathbf{Q}|u\rangle|| .$$
(12)

Hence for arbitrary state ρ we have

$$H_{\alpha}(\mathsf{M}|\rho) + H_{\beta}(\mathsf{N}|\rho) \ge -2\log\max_{ij}||\mathsf{M}_{i}^{1/2}\mathsf{N}_{j}^{1/2}|| , \quad (13)$$

where $1/\alpha + 1/\beta = 2$. Further, there holds [54]

$$H_{\alpha}(\mathsf{M}|\rho) \ge -\log\max_{i} ||\mathsf{M}_{i}|| . \tag{14}$$

If we take in Eq. (13) the two POVM's to be identical, then we just get Eq. (14). The right-hand side of Eq. (14) is an analogue of trivial zero bound for projective measurement. Of course, Relation 2 is far more sharp.

Let us discuss an example of applications of the obtained relations. Consider a game involving two parties, Alice and Bob. Alice secretly chooses a state from the set $\{|\psi_1\rangle, |\psi_2\rangle\}$ known to both parties. She then sends the chosen state to Bob. His task is to identity the state Alice has given him. Let us put $|\pm\rangle \equiv (|0\rangle \pm |1\rangle)/\sqrt{2}$. In our simple case we take $|\psi_1\rangle = |0\rangle$ and $|\psi_2\rangle = |+\rangle$. Bob can use two different strategies. The strategy developed by Helstrom [56] is not error-free. In our example, Bob's optimal measurement is described by PVM $\{N_1, N_2\}$ with elements $N_1 = |x\rangle\langle x|$ and $N_2 = |y\rangle\langle y|$, where

$$|x\rangle \equiv \cos(\pi/8) |0\rangle - \sin(\pi/8) |1\rangle , \qquad (15)$$

$$|y\rangle \equiv \sin(\pi/8) |0\rangle + \cos(\pi/8) |1\rangle . \tag{16}$$

If the outcome N_1 (the outcome N_2) is detected then Bob concludes that Alice sent the state $|\psi_1\rangle$ (the state $|\psi_2\rangle$). The above PVM minimizes the average probability of mis-identification [56] equal to $(1/2)\{\langle\psi_1|N_2|\psi_1\rangle + \langle\psi_2|N_1|\psi_2\rangle\} = (\sqrt{2}-1) 2^{-3/2}$.

Bob can also use the unambiguous discrimination proposed by Ivanovich [57], Dieks [58] and Peres [59]. If he allows the inconclusive answer then it is possible for him to perform a measurement without mis-identification. The optimal measurement minimizes the probability of inconclusive answer [59]. Note that the unambiguous discrimination has important application to the quantum cryptography. In the B92 protocol [60], Alice encode the bits 0 and 1 into two nonorthogonal pure states. In Ref. [60] Bennett described the strategy, whose efficiency is less than 50 %. The authors of Ref. [61] built the procedure based on the unambiguous discrimination. The efficiency of proposed strategy is greater than 50 %.

Let us return to the discussed game. We consider a POVM $\{M_1, M_2, M_3\}$ with elements [59]

$$\mathsf{M}_1 = \sqrt{2} \, (\sqrt{2} + 1)^{-1} \, |-\rangle \langle -| \,, \qquad (17)$$

$$\mathsf{M}_2 = \sqrt{2} \, (\sqrt{2} + 1)^{-1} \, |1\rangle \langle 1| \,, \tag{18}$$

$$\mathsf{M}_3 = \mathbf{1} - \mathsf{M}_1 - \mathsf{M}_2 \;. \tag{19}$$

If the outcome M_1 (the outcome M_2) is detected then Bob can exactly conclude that the state $|\psi_1\rangle$ (the state $|\psi_2\rangle$) has been received. Sometimes, however, the inconclusive outcome M_3 will occur, and then he will obtain no information about the received state. The probability of this answer is $|\langle\psi_1|\psi_2\rangle| = 1/\sqrt{2}$ [59].

The discussed two POVM's $\{M_i\}$ and $\{N_j\}$ have only one-rank elements. It turns out that in this case the inequality (11) is saturated regardless of state ρ [54]. So, $f(\mathsf{M},\mathsf{N}|\rho)^2 = ||\mathsf{M}_1^{1/2}\mathsf{N}_1||^2 = ||\mathsf{M}_2^{1/2}\mathsf{N}_2||^2 = 1/2$ by calculations. Relation 1 then leads to

$$H_{\alpha}(\mathsf{M}|\rho) + H_{\beta}(\mathsf{N}|\rho) \ge 1 \tag{20}$$

for any state ρ . On the contrary, the right-hand side of Eq. (10) is significantly depend on a quantum state. Let us consider this inequality for the state $|\psi_1\rangle = |0\rangle$. By calculations, we have

$$\phi(\mathsf{M}|\psi_1) = \langle 0|\mathsf{M}_3|0\rangle = 2^{-1/2} , \qquad (21)$$

$$\phi(\mathsf{N}|\psi_1) = \langle 0|\mathsf{N}_1|0\rangle = 2^{-3/2}(\sqrt{2}+1)$$
. (22)

The entropic relation (10) then gives

$$H_{\alpha}(\mathsf{M}|\psi_1) + H_{\beta}(\mathsf{N}|\psi_1) \ge 2 - \log(\sqrt{2} + 1) \approx 0.728$$
. (23)

The inequality (20) is stronger than the inequality (23). The difference 0.272 between the right-hand sides of Eq. (20) and Eq. (23) may be regarded as a manifestation of incompatibility of the two discussed measurements. We shall now compare the entropic bounds in Eqs. (9) and (14) within our example. For the state $|\psi_1\rangle$ Relation 2 gives $H_{\alpha}(\mathsf{M}|\psi_1) \geq 0.5$ according to Eq. (21). This considerably exceeds the trivial lower bound $-\log ||\mathsf{M}_3|| = \log (\sqrt{2}+1) - 1 \approx 0.272$ given by Eq. (14). We see that the state-dependent bound in Eq. (9) can be far stronger. Thus, Relation 2 ensures a nontrivial entropic bound for a single POVM.

In our example we have seen that the lower bound (8) can be more sharp than the lower bound (10). As it is shown in Ref. [54], in one's turn the lower bound (10) can be more sharp than the lower bound (8). In addition, the correctness of Eq. (10) is not limited by the condition $1/\alpha + 1/\beta = 2$. So both the bounds (8) and (10) should be considered as independent entropic relations. It was above mentioned that the bound given by Maassen and Uffink [38] has been sharpened in some cases. So there is clear scope for improvement of the lower bounds (8) and (10). These questions remain for the future.

Acknowledgments. I am grateful to an anonymous referee for helpful remarks.

- [1] W. Heisenberg, Z. Phys. **43**, 172 (1927)
- [2] J. S. Bell, Physics 1, 195 (1964)

[3] B. Misra and E. C. G. Sudarshan, J. Math. Phys. 18, 756 (1977)

- [4] W. K. Wootters and W. H. Zurek, Nature (London) 299, 802 (1982)
- [5] A. C. Elitzur and L. Vaidman, Found. Phys. 23, 987 (1993)
- [6] A. K. Ekert, Phys. Rev. Lett. 67, 661 (1991)
- [7] L. Viola and S. Lloyd, Phys. Rev. A 58, 2733 (1998)
- [8] H. P. Robertson, Phys. Rev. 34, 163 (1929)
- [9] M. Ozawa, Phys. Rev. A 67, 042105 (2003)
- [10] M. J. W. Hall, Phys. Rev. A 69, 052113 (2004)
- [11] P. Busch, T. Heinonen, and P. J. Lahti, Phys. Rep. 452, 155 (2007)
- [12] N. Bohr, Nature (London) **121**, 580 (1928)
- [13] M. Ozawa, Phys. Lett. A 320, 367 (2004)
- [14] V. Kendon and B. C. Sanders, Phys. Rev. A 71, 022307 (2005)
- [15] W. K. Wootters and W. H. Zurek, Phys. Rev. D 19, 473 (1979)
- [16] A. Wehrl, Rev. Mod. Phys. 50, 221 (1978)
- [17] V. Vedral, Rev. Mod. Phys. **74**, 197 (2002)
- S. Luo, Phys. Rev. Lett. 91, 180403 (2003); S. Luo, Phys. Rev. A 72, 042110 (2005)
- [19] S. Luo, Lett. Math. Phys. 53, 243 (2000); S. Luo and
 Z. Zhang, J. Stat. Phys. 114, 1557 (2004); P. Gibilisco,
 D. Imparato, and T. Isola, J. Stat. Phys. 130, 545 (2008)
- [20] E. Romera, P. Sánchez-Moreno, and J. S. Dehesa, J. Math. Phys. 47, 103504 (2006); P. Sánchez-Moreno, R. González-Férez, and J. S. Dehesa, New J. Phys. 8, 330 (2006)
- [21] M. Christandl and A. Winter, IEEE Trans. Inf. Theory 51, 3159 (2005)
- [22] T. Miyadera and H. Imai, Phys. Rev. A 76, 062108 (2007)
- [23] S. Massar and P. Spindel, Phys. Rev. Lett. 100, 190401 (2008)
- [24] V. Giovannetti, Phys. Rev. A 70, 012102 (2004);
 O. Gühne and M. Lewenstein, Phys. Rev. A 70, 022316 (2004)
- [25] M. A. Ballester and S. Wehner, Phys. Rev. A 75, 022319 (2007)
- [26] S. Wehner and A. Winter, arXiv: 0710.1185 [quant-ph]
- [27] J. I. de Vicente and J. Sánchez-Ruiz, Phys. Rev. A 77, 042110 (2008)
- [28] I. Bialynicki-Birula, Phys. Rev. A 74, 052101 (2006)
- [29] T. Cubitt, A. W. Harrow, D. Leung, A. Montanaro, and A. Winter, arXiv: 0712.3628 [quant-ph]; N. Schuch, M. M. Wolf, F. Verstraete, and J. I. Cirac, Phys. Rev. Lett. **100**, 030504 (2008); R. Augusiak, J. Stasińska, and P. Horodecki, Phys. Rev. A **77**, 012333 (2008); B. Dier-

ckx, M. Fannes, and C. Vandenplas, Phys. Rev. A **77**, 060302(R) (2008)

- [30] A. Peres, Quantum Theory: Concepts and Methods (Kluwer, Dordrecht, 1995)
- [31] M. Roško, V. Bužek, P. R. Chouha, and M. Hillery, Phys. Rev. A 68, 062302 (2003)
- [32] M. Ozawa, Ann. Phys. (N.Y.) 311, 350 (2004)
- [33] S. Massar, Phys. Rev. A **76**, 042114 (2007)
- [34] I. I. Hirschman, Am. J. Math. 79, 152 (1957)
- [35] W. Beckner, Ann. Math. **102**, 159 (1975)
- [36] I. Bialynicki-Birula and J. Mycielski, Commun. Math. Phys. 44, 129 (1975)
- [37] D. Deutsch, Phys. Rev. Lett. **50**, 631 (1983)
- [38] H. Maassen and J. B. M. Uffink, Phys. Rev. Lett. 60, 1103 (1988)
- [39] M. H. Partovi, Phys. Rev. Lett. **50**, 1883 (1983)
- [40] I. Bialynicki-Birula, Phys. Lett. A 103, 253 (1984)
- [41] K. Kraus, Phys. Rev. D **35**, 3070 (1987)
- [42] M. J. W. Hall, Phys. Rev. A 55, 100 (1997)
- [43] M. Krishna and K. R. Parthasarathy, Sankhya, Ser. A 64, 842 (2002)
- [44] J. Sánchez, Phys. Lett. A 173, 233 (1993)
- [45] A. J. M. Garrett and S. F. Gull, Phys. Lett. A 151, 453 (1990)
- [46] J. Sánchez-Ruiz, Phys. Lett. A **244**, 189 (1998)
- [47] G.-C. Ghirardi, L. Marinatto, and R. Romano, Phys. Lett. A 317, 32 (2003)
- [48] J. Havrda and F. Charvát, Kybernetika 3, 30 (1967)
- [49] C. Tsallis, J. Stat. Phys. **52**, 479 (1988)
- [50] A. K. Rajagopal, Phys. Lett. A 205, 32 (1995)
- [51] V. Majernik and E. Majernikova, Rep. Math. Phys. 47, 381 (2001)
- [52] A. Rényi, On Measures of Entropy and Information, Proc. Fourth Berkeley Symp. Math. Stat. Prob. (Vol. 1., University of California Press, Berkeley, 1961), p. 547
- [53] U. Larsen, J. Phys. A: Math. Gen. 23, 1041 (1990)
- [54] A. E. Rastegin, arXiv: 0807.2691 [quant-ph]
- [55] K. Życzkowski, Open Sys. Inf. Dyn. 10, 297 (2003)
- [56] C. W. Helstrom, Quantum Detection and Estimation Theory (Academic Press, New York, 1976)
- [57] I. D. Ivanovic, Phys. Lett. A **123**, 257 (1987)
- [58] D. Dieks, Phys. Lett. A **126**, 303 (1988)
- [59] A. Peres, Phys. Lett. A **128**, 19 (1988)
- [60] C. H. Bennett, Phys. Rev. Lett. 68, 3121 (1992)
- [61] A. K. Ekert, B. Huttner, G. M. Palma, and A. Peres, Phys. Rev. A 50, 1047 (1994)