

Canonical measures and dynamical systems of Bergman kernels

Hajime TSUJI*

December 19, 2010

Abstract

In this article, we construct the canonical semipositive current or the canonical measure (= the potential of the canonical semipositive current) on a smooth projective variety with nonnegative Kodaira dimension in terms of a dynamical system of Bergman kernels. This current is considered to be a generalization of a Kähler-Einstein metric and coincides the one considered independently by J. Song and G. Tian ([S-T]). The major difference between [S-T] and the present article is that they found the canonical measure in terms of Kähler-Ricci flows, while I found the canonical measure in terms of dynamical systems of Bergman kernels. Hence the present approach can be viewed as a discrete version of a Kähler-Ricci flow.

The advantage of the dynamical construction is two folds. First, it enables us to deduce the (logarithmic) plurisubharmonic variation property of the canonical measures on a projective family. Second, we can overcome the difficulty arising from the singularities of the solution of a Kähler-Ricci flow.

MSC: 53C25(32G07 53C55 58E11)

Contents

1	Introduction	2
1.1	Kähler-Einstein metrics	5
1.2	Kähler-Einstein currents	5
1.3	Iitaka fibration	6
1.4	Analytic Zariski decompositions	7
1.5	Requirement of the canonical semipositive current	7
1.6	Canonical bundle formula and the correction term for the Kähler-Einstein equations	8
1.7	Canonical Kähler currents	10
1.8	Ricci iterations	11
1.9	Dynamical construction of the canonical Kähler currents	12

*Partially supported by Grant-in-Aid for Scientific Reserch (S) 17104001

2	Construction of the canonical measure by solving Monge-Ampère equations	13
2.1	Setup	14
2.2	Smoothing of the Hodge metric $h_{L_{X/Y}}$	15
2.3	The construction of the canonical Kähler currents	18
2.4	Regularity of the canonical Kähler current	26
2.5	Generalization to general adjoint line bundles	32
3	Dynamical systems of Bergman kernels	32
3.1	The case: $D_{X/Y} = 0$	33
3.2	The case: $D_{X/Y} \neq 0$	39
3.3	Uniqueness of canonical measures	40
4	Relative version of Theorems 1.5 and 1.8	41

1 Introduction

In [T4], I have constructed a canonical Kähler-Einstein current on a smooth projective variety of general type in terms of a dynamical system of Bergman kernels originated in [T3]. This Kähler-Einstein current is the same one which has been studied in [T0, Su].

Since the same kind of dynamical systems has been defined on a smooth projective varieties of nonnegative Kodaira dimension (or even for a smooth projective variety with pseudoeffective canonical bundle) in [T3], it is natural to expect that the normalized limit of the dynamical system of Bergman kernels yields a substitute of a Kähler-Einstein metric for a smooth projective variety (of non general type) with nonnegative Kodaira dimension.

The purpose of this article is to prove that the limit satisfies the partial differential equation (see (1.25)) similar to the Kähler-Einstein equation on the base of the Iitaka fibration (not on the original variety) and give a natural generalization of the notion of Kähler-Einstein volume form. We call the (normalized) limit the canonical measure. And we call the $-\text{Ric}$ of the canonical measure (in the sense of current) the canonical semipositive current.

There are two major differences between Kähler-Einstein metrics and the canonical semipositive currents.

First of all in general the canonical semipositive current is strictly positive not on the original variety but on the base space of the Iitaka fibration. In other words, the current is the pullback of a closed generically strictly positive current on the base space of the Iitaka fibration.

Secondary although the canonical semipositive current satisfies a similar partial differential equation as a Kähler-Einstein metric on the base space of the Iitaka fibration, the equation has an additional term coming from variation of Hodge structure on the Iitaka fibration.

The objective of this generalization is to study the deformation of projective varieties with nonnegative Kodaira dimension. Actually the dynamical construction of the canonical semipositive current yields the existence of a closed semipositive current on the family which restricts the canonical semipositive current on the general fibers (Theorem 4.1). We discuss the applications of Theorem 4.1 in [T7]. And we also note that there are similar constructions of

canonical measures (canonical and supercanonical AZD's) on smooth projective varieties with pseudoeffective canonical bundles ([T5]).

After the completion of this work, I have noticed a paper of Song and Tian ([S-T]) which also constructed the canonical semipositive current from a different point of view. Actually first they have constructed the canonical semipositive current as the limit of the Kähler-Ricci flow in the case of semiample canonical bundle. Then they constructed the current which satisfies the same equation without using a Kähler-Ricci flow on a projective varieties with nonnegative Kodaira dimension whose canonical bundle is not necessarily semiample. In this sense their construction is modeled after the case of semiample canonical bundles. But in their construction, the meaning of the canonical semipositive current is not clear (although it is apparently a generalization of a Kähler-Einstein metric).

The main contribution of this article is to give a dynamical construction of the canonical semipositive currents (or the canonical measure in [S-T]) and give the authenticity to the canonical semipositive current.

The advantage of the dynamical construction is that we can overcome the difficulty arising from the singularity of the currents. For example it seems to be difficult to deduce the plurisubharmonic variation property of the canonical measures on a projective family (Theorem 4.1) by direct calculation. On the other hand Theorem 4.1 is an immediate consequence of the dynamical construction by using the logarithmic plurisubharmonic variation properties of Bergman kernels.

Also difficulty arises to study a Kähler-Ricci flow, when we consider non minimal algebraic varieties. In this case the flow of the Kähler class associated with a Kähler-Ricci flow reaches the boundary of the Kähler cone in finite time. Hence in this case it is inevitable to deal with a singular Kähler Ricci flow. But the dynamical construction (Theorem 1.8) automatically produces the canonical semipositive current (or the canonical measure) as soon as the Kodaira dimension of the variety is nonnegative. One may consider Theorem 1.8 as a discretization of a Kähler-Ricci flow¹ and it overcomes the difficulty arising from singularities.

After the completion of this paper (math.ArXiv.0805.1829), I extended the result to the case of KLT pairs ([T6]) and thought that the technique used to solve the twisted Kähler-Einstein equation (1.25) is standard. But recently it was pointed out by S. Boucksom that the existence of the solution of (1.25) does not follow from the ready made techniques because the corresponding Monge-Ampère equation has nonalgebraic singularities. Moreover I could not follow the proof of [S-T, Theorem B.2] by J. Song and G. Tian which asserts Theorem 1.5 in this paper independently. Hence I decided to publish this paper independently.

Acknowledgement I would like to express to S. Boucksom who pointed out a crucial error in the previous version. Namely he pointed out the case $D_{X/Y} \neq 0$ (cf. Section 3.2) is lacking in the previous version.

Notations

- For a real number a , $\lceil a \rceil$ denotes the minimal integer greater than or equal to a and $\lfloor a \rfloor$ denotes the maximal integer smaller than or equal to a .

¹In [T6], I have used another discretization of a Kähler-Ricci flow.

- Let X be a projective variety and let D be a Weil divisor on X . Let $D = \sum d_i D_i$ be the irreducible decomposition. We set

$$(1.1) \quad [D] := \sum [d_i] D_i, [D] := \sum [d_i] D_i.$$

- Let L be a line bundle on a compact complex manifold X . A singular hermitian metric h on L is given by

$$h = e^{-\varphi} \cdot h_0,$$

where h_0 is a C^∞ -hermitian metric on L and $\varphi \in L^1_{loc}(X)$ is an arbitrary function on X . We call φ a weight function of h .

The curvature current Θ_h of the singular hermitian line bundle (L, h) is defined by

$$\Theta_h := \Theta_{h_0} + \partial\bar{\partial}\varphi,$$

where $\partial\bar{\partial}\varphi$ is taken in the sense of current. The L^2 sheaf $\mathcal{L}^2(L, h)$ of the singular hermitian line bundle (L, h) is defined by

$$\mathcal{L}^2(L, h)(U) := \{\sigma \in \Gamma(U, \mathcal{O}_X(L)) \mid h(\sigma, \sigma) \in L^1_{loc}(U)\},$$

where U runs over the open subsets of X . In this case there exists an ideal sheaf $\mathcal{I}(h)$ such that

$$\mathcal{L}^2(L, h) = \mathcal{O}_X(L) \otimes \mathcal{I}(h)$$

holds. We call $\mathcal{I}(h)$ the **multiplier ideal sheaf** of (L, h) .

- For a closed positive $(1, 1)$ current T , T_{abc} denotes the absolutely continuous part of T .
- For a singular hermitian line bundle (F, h_F) on a compact complex manifold X of dimension n . $K(X, K_X + F, h_F)$ denotes (the diagonal part of) the Bergman kernel of $H^0(X, \mathcal{O}_X(K_X + F) \otimes \mathcal{I}(h_F))$ with respect to the L^2 -inner product:

$$(1.2) \quad (\sigma, \sigma') := (\sqrt{-1})^{n^2} \int_X h_F \cdot \sigma \wedge \bar{\sigma}',$$

i.e.,

$$(1.3) \quad K(X, K_X + F, h_F) = \sum_{i=0}^N |\sigma_i|^2,$$

where $\{\sigma_0, \dots, \sigma_N\}$ is a complete orthonormal basis of $H^0(X, \mathcal{O}_X(K_X + F) \otimes \mathcal{I}(h_F))$. It is clear that $K(X, K_X + F, h_F)$ is independent of the choice of the complete orthonormal basis.

- A line bundle L on a compact complex manifold X is said to be **pseudoeffective**, if it admits a singular hermitian metric with semipositive curvature current. A singular hermitian line bundle (L, h) is said to be pseudoeffective if the curvature current $\sqrt{-1}\Theta_h$ is semipositive. If X is a smooth projective variety, this is equivalent to the fact that $c_1(L)$ is on the closure of the effective cone.

- Let (X, D) be a pair of a normal variety and a \mathbb{Q} -divisor on X . Suppose that $K_X + D$ is \mathbb{Q} -Cartier. Let $f : Y \rightarrow X$ be a log resolution. Then we have the formula :

$$K_Y = f^*(K_X + D) + \sum a_i E_i,$$

where E_i is a prime divisor and $a_i \in \mathbb{Q}$. The pair (X, D) is said to be **subKLT**(resp. **subLC**), if $a_i > -1$ (resp. $a_i \geq -1$) holds for every i . (X, D) is said to be **KLT** (resp. **LC**), if (X, D) is subKLT(resp. subLC) and D is effective.

1.1 Kähler-Einstein metrics

Let X be a compact Kähler manifold with the Kähler form

$$(1.4) \quad \omega := \frac{\sqrt{-1}}{2} \sum_{i,j} g_{i\bar{j}} dz^i \wedge d\bar{z}^j.$$

(X, ω) is said to be Kähler-Einstein, if there exists a constant c such that

$$(1.5) \quad \text{Ric}_\omega = c\omega$$

holds, where Ric_ω denotes the Ricci form:

$$(1.6) \quad \text{Ric}_\omega := -\sqrt{-1} \partial \bar{\partial} \log \det(g_{i\bar{j}})$$

and we call ω a Kähler-Einstein form on X . If a compact complex manifold X admits a Kähler-Einstein form, then $c_1(X)$ is negative or 0 or positive. Conversely by the celebrated solution of Calabi's conjecture ([A, Y1]), for a compact Kähler manifold X , if $c_1(X)$ is negative, there exists a unique Kähler-Einstein form ω_E such that

$$(1.7) \quad -\text{Ric}_{\omega_E} = \omega_E$$

([A, Y1]) and if $c_1(X)$ is 0 in $H^2(X, \mathbb{R})$, then in every Kähler class on X , there exists a unique Kähler-Einstein form ω_E such that

$$(1.8) \quad \text{Ric}_{\omega_E} = 0$$

holds ([Y1]).

1.2 Kähler-Einstein currents

There are numerous applications of Kähler-Einstein metrics. But in general, a smooth complex projective variety does not admit a Kähler-Einstein metric, since the first Chern class is not definite in general. One way to overcome this defect is to consider Kähler-Einstein metrics allowing singularities.

In [T0], I have constructed a Kähler-Einstein current ω_E on smooth minimal algebraic variety X of general type. More precisely there exists a unique closed semipositive current ω_E such that

- (1) There exists a nonempty Zariski open subset U of X such that ω_E is a C^∞ -Kähler form on U .

- (2) $-\text{Ric}_{\omega_E} = \omega_E$ holds on U .
- (3) ω_E is absolutely continuous on X .

Later K. Sugiyama proved that there exists a Kähler-Einstein current on the canonical model of general type ([Su]). Also I have constructed a Kähler-Einstein current on an arbitrary smooth projective variety of general type (without using the finite generation of canonical rings) in [T4]. Hence for smooth projective varieties of general type, we have a substitute of a Kähler-Einstein metric.

We note that the above Kähler-Einstein current ω_E on a projective variety X of general type have the following properties : $h_E := n! (\omega_E^n)^{-1} (n = \dim X)$ is a singular hermitian metric on K_X such that the curvature current $\sqrt{-1} \Theta_{h_E}$ is a closed semipositive current and

$$(1.9) \quad H^0(X, \mathcal{O}_X(mK_X) \otimes \mathcal{I}(h_E^m)) \simeq H^0(X, \mathcal{O}_X(mK_X))$$

holds for every $m \geq 1$, i.e., h_E is an AZD of K_X (cf. Definition 1.4 below). In other words, h_E is a singular hermitian metric which extracts all the positivity of K_X .

But for smooth projective varieties of non general type, the above results do not say anything.

1.3 Iitaka fibration

The simplest way to squeeze out the positivity of canonical bundles is to use the pluricanonical systems.

Let X be a smooth projective variety. The Kodaira dimension $\text{Kod}(X)$ is defined by

$$(1.10) \quad \text{Kod}(X) := \limsup_{m \rightarrow \infty} \frac{\log \dim H^0(X, \mathcal{O}_X(mK_X))}{\log m}.$$

It is known that $\text{Kod}(X)$ is $-\infty$ or a nonnegative integer between 0 and $\dim X$.

Let X be a smooth projective variety with $\text{Kod}(X) \geq 0$. Then for a sufficiently large $m > 0$, the complete linear system $|m!K_X|$ gives a rational fibration (with connected fibers) :

$$(1.11) \quad f : X \dashrightarrow Y.$$

We call $f : X \dashrightarrow Y$ the **Iitaka fibration** of X .

The Iitaka fibration is independent of the choice of the sufficiently large m up to birational equivalence. In this sense the Iitaka fibration is unique. By taking a suitable modification, we may assume that f is a morphism and Y is smooth.

The Iitaka fibration $f : X \rightarrow Y$ satisfies the following properties:

- (1) For a general fiber F , $\text{Kod}(F) = 0$ holds.
- (2) $\dim Y = \text{Kod}(X)$.

1.4 Analytic Zariski decompositions

Let L be a pseudoeffective line bundle on a compact complex manifold X . To analyze the ring :

$$(1.12) \quad R(X, L) = \bigoplus_{m=0}^{\infty} H^0(X, \mathcal{O}_X(mL)),$$

it is useful to introduce the notion of analytic Zariski decompositions.

Definition 1.1 *Let M be a compact complex manifold and let L be a holomorphic line bundle on M . A singular hermitian metric h on L is said to be an **analytic Zariski decomposition** (AZD in short), if the followings hold.*

(1) $\sqrt{-1}\Theta_h$ is a closed positive current.

(2) for every $m \geq 0$, the natural inclusion:

$$(1.13) \quad H^0(M, \mathcal{O}_M(mL) \otimes \mathcal{I}(h^m)) \rightarrow H^0(M, \mathcal{O}_M(mL))$$

is an isomorphism. \square

Remark 1.2 *If an AZD exists on a line bundle L on a compact complex manifold M , L is pseudoeffective by the condition 1 above. \square*

It is known that for every pseudoeffective line bundle on a compact complex manifold, there exists an AZD on L (cf. [T1, T2, D-P-S]). The advantage of the AZD is that we can handle pseudoeffective line bundle L on a compact complex manifold X as a singular hermitian line bundle with semipositive curvature current as long as we consider the ring $R(X, L)$.

One may construct an AZD for a pseudoeffective line bundle on a compact complex manifold as follows. Let L be a pseudoeffective line bundle on a compact complex manifold X . Let h_0 be a C^∞ -hermitian metric on L . We set

$$(1.14) \quad h_{min} := \inf\{h \mid \text{a singular hermitian metric on } L, h \geq h_0, \sqrt{-1}\Theta_h \geq 0\}.$$

Then h_{min} is an AZD on L with minimal singularities in the following sense.

Definition 1.3 *Let L be a pseudoeffective line bundle on a compact complex manifold X . An AZD h on L is said to be an **AZD of minimal singularities**, if for any AZD h' on L , there exists a positive constant C such that*

$$(1.15) \quad h \leq C \cdot h'$$

holds. \square

In general an AZD of a pseudoeffective line bundle L on a smooth projective variety is not necessarily of minimal singularities.

1.5 Requirement of the canonical semipositive current

Let $f : X \rightarrow Y$ be the Iitaka fibration of a smooth projective variety X of nonnegative Kodaira dimension. In this article, we shall consider a canonical semipositive current, say ω_X associated with the Iitaka fibration.

It is natural to require that ω_X has the following properties :

- (1) ω_X is unique and birationally invariant, i.e., if X is birational to X' and let $\mu : X'' \rightarrow X$ and $\mu' : X'' \rightarrow X'$ be modifications from a smooth projective variety X'' . Then

$$(1.16) \quad \omega_{X''} = \mu^* \omega_X + 2\pi E = (\mu')^* \omega_{X'} + 2\pi E'$$

hold, where $E := K_{X''} - \mu^* K_X$ and $E' := K_{X''} - (\mu')^* K_{X'}$, respectively.

- (2) There exists an AZD h_K of K_X such that $\omega_X = \sqrt{-1} \Theta_{h_K}$.
- (3) There exists a closed semipositive current ω_Y on Y such that $\omega_X = f^* \omega_Y$,
- (4) There exists a nonempty Zariski open subset U of Y such that $\omega_Y|_U$ is a C^∞ -Kähler form.

1.6 Canonical bundle formula and the correction term for the Kähler-Einstein equations

To construct the AZD h_K above, we shall solve a partial differential equation. The equation is similar to the Kähler-Einstein equation : $-\text{Ric}_{\omega_E} = \omega_E$, but there are two major differences :

- (1) The equation is defined on Y not on X .
- (2) The equation has the additional term which comes from variation of Hodge structures.

For a graded ring $R := \bigoplus_{i=0}^{\infty} R_i$ and a positive integer m , we set

$$(1.17) \quad R^{(m)} := \bigoplus_{i=0}^{\infty} R_{mi}.$$

For a KLT pair (M, D) , we set

$$(1.18) \quad R(M, K_M + D) := \bigoplus_{m=0}^{\infty} \Gamma(M, \mathcal{O}_M(\lfloor m(K_M + D) \rfloor))$$

and

$$(1.19) \quad \text{Kod}(M, D) := \limsup_{m \rightarrow \infty} \frac{\log \dim \Gamma(M, \mathcal{O}_M(\lfloor m(K_M + D) \rfloor))}{\log m}.$$

Theorem 1.4 ([F-M, p.183, Theorem 5.2]) *Let (X, Δ) be a proper KLT pair with*

$$(1.20) \quad \text{Kod}(X, K_X + \Delta) = n.$$

Then there exists a n -dimensional KLT pair (Y', Δ') with $\text{Kod}(Y', \Delta') = n$, two positive integers e, e' such that

$$(1.21) \quad R(X, K_X + \Delta)^{(e)} \simeq R(Y', K_{Y'} + \Delta')^{(e')}.$$

□

Let us consider the case that $\Delta = 0$ in Theorem 1.4. Then the canonical ring $R(X, K_X)^{(e)}$ is (a subring of) the pullback of the log canonical ring of some KLT pair (Y', Δ') of log general type.

Let us explain the equation. Let $f : X \rightarrow Y$ be an Iitaka fibration such that $f_*\mathcal{O}_X(m!K_{X/Y})^{**}$ is locally free on Y for some m (hence for every sufficiently large m), where $**$ denotes the double dual. Such $f : X \rightarrow Y$ exists by [F-M, p.169, Proposition 2.2]. The divisor Δ' is related to the \mathbb{Q} -line bundle

$$(1.22) \quad L_{X/Y} := \frac{1}{m_0!} f_*\mathcal{O}_X(m_0!K_{X/Y})^{**}$$

on Y in terms of the canonical bundle formula (See [F-M] for detail. In [F-M, Section 2]), where m_0 is a sufficiently large positive integer. We note that $L_{X/Y}$ is independent of a sufficiently large m_0 (cf. [F-M, Section 2]). We call $L_{X/Y}$ the **Hodge \mathbb{Q} -line bundle** of $f : X \rightarrow Y$. $L_{X/Y}$ carries a natural singular hermitian metric $h_{L_{X/Y}}$ defined by

$$(1.23) \quad h_{L_{X/Y}}^{m_0!}(\sigma, \sigma) = \left(\int_{X_y} |\sigma|^{\frac{2}{m_0!}} \right)^{m_0!},$$

where $y \in Y, X_y := f^{-1}(y)$ and $\sigma \in L_{X/Y, y}$. It is known that $h_{L_{X/Y}}$ has semipositive curvature in the sense of current ([Ka2], [Ka3, p.174, Theorem 1.1]) by using the variation of Hodge structures. By a suitable modification of $f : X \rightarrow Y$, we may assume that there exists an decomposition:

$$(1.24) \quad L_{X/Y} = M_{X/Y} + D_{X/Y}$$

such that

- (1) $M_{X/Y}, D_{X/Y} \in \text{Div}(Y) \otimes \mathbb{Q}$,
- (2) $M_{X/Y}$ is nef, $D_{X/Y}$ is effective.
- (3) $\text{Supp } D_{X/Y}$ is a divisor with normal crossings on Y ,
- (4) $h_{L_{X/Y}}$ induces a singular hermitian metric $h_{M_{X/Y}}$ on $M_{X/Y}$ with semipositive curvature and is logarithmic growth as Lemma 2.17 below. and $D_{X/Y}$ corresponds to the singular part of the 1-st Chern current $c_1(L_{X/Y}, h_{L_{X/Y}})$.
- (5) $(Y, D_{X/Y})$ is KLT.

We call $M_{X/Y}$ the *semistable part* and $D_{X/Y}$ the *discriminant part* of $L_{X/Y}$ respectively ([F-M]).

Let Ω be a C^∞ -volume form on Y . We shall consider the following equation :

$$(1.25) \quad -\text{Ric}_{\omega_Y} + \sqrt{-1} \Theta_{h_{L_{X/Y}}} = \omega_Y,$$

where

$$(1.26) \quad \omega_Y = -\text{Ric } \Omega + \sqrt{-1} \Theta_{h_{L_{X/Y}}} + \sqrt{-1} \partial \bar{\partial} u,$$

and u is the unknown function. Here the term $\sqrt{-1} \Theta_{h_{L_{X/Y}}}$ corresponds to the boundary divisor of the KLT pair (Y', Δ') in the canonical bundle formula ([F-M,

p.183, Theorem 5.2]). The authenticity of the equation (1.25) can be verified by checking the fact that the dynamical system of Bergman kernels on X as in [T4] yields the current on Y which satisfies the equation (1.25) (cf. Theorem 1.8).

Actually I first constructed the current by using the dynamical system of Bergman kernels and then found the equation (1.25) inspired by the [F-M].

There are several difficulties to solve the equation (1.25). First of all we cannot expect that there exists a C^∞ -solution ω_Y . In fact $\Theta_{h_{L_{X/Y}}}$ is not C^∞ in general. And if $\Theta_{h_{L_{X/Y}}}$ is not C^∞ , (1.25) has no C^∞ -solution ω_Y . Moreover $h_{L_{X/Y}}$ is not of algebraic singularities (cf. Definition 2.1) in general. And even if $h_{L_{X/Y}}$ is C^∞ , u is not C^∞ in general. Secondary the solution u is not unique. But if we require that $\Omega^{-1} \cdot h_{L_{X/Y}} \cdot e^u$ is an AZD of $K_Y + L_{X/Y}$, then the solution u is actually unique and the resulting current ω_Y is nothing but the current constructed by the dynamical system of Bergman kernels (see Section 1.9 and Section 3 below).

1.7 Canonical Kähler currents

Now we shall state the existence of the canonical semipositive current on a smooth projective variety of nonnegative Kodaira dimension.

Theorem 1.5 (cf. [S-T, Theorem B.2]) *In the above notations, there exists a unique singular hermitian metric on h_K on $K_Y + L_{X/Y}$ and a nonempty Zariski open subset U in Y such that*

- (1) h_K is an AZD of $K_Y + L_{X/Y}$,
- (2) f^*h_K is an AZD of K_X , Here we have used the inclusion:

$$f^*\mathcal{O}_Y(m_0!(K_Y + L_{X/Y})) \hookrightarrow \mathcal{O}_X(m_0!K_X)$$

to identify f^*h_K with a singular hermitian metric on K_X , where m_0 is the sufficiently large positive integer used in (1.22),

- (3) h_K is C^∞ on U ,
- (4) $\omega_Y = \sqrt{-1}\Theta_{h_K}$ is a Kähler form on U ,
- (5) $-\text{Ric}_{\omega_Y} + \sqrt{-1}\Theta_{L_{X/Y}} = \omega_Y$ holds on U . \square

Although the proof of Theorem 1.5 is given in [S-T], I shall give an alternative proof in this paper for the completeness, since my original proof seems to be different from that in [S-T]. One can see the KLT version of the above theorem in [T6].

Definition 1.6 *The current ω_Y on Y is said to be the **canonical Kähler current** of the Iitaka fibration $f : X \rightarrow Y$. Also $\omega_X := f^*\omega_Y$ is said to be the **canonical semipositive current** on X . We define the measure $d\mu_{can}$ on X by*

$$(1.27) \quad d\mu_{can} := \frac{1}{n!} f^* \omega_Y^n \cdot h_{L_{X/Y}}^{-1}$$

and is said to be the **canonical measure**, where n denotes $\dim Y$. \square

The existence of the canonical Kähler current is proven in terms of solving Monge-Ampère equations. The proof given here is similar to the one of [T4, Section 5.1, Theorem 5.1]. We shall give a proof in Section 2 (see also [S-T, Section 4]).

1.8 Ricci iterations

Let $f : X \rightarrow Y$, $(L_{X/Y}, h_{L_{X/Y}})$ be as above. We assume that there exists a Zariski decomposition :

$$(1.28) \quad K_Y + L_{X/Y} = P + N \quad (P, N \in \text{Div}(X) \otimes \mathbb{Q})$$

i.e., P is semiample, N is effective and

$$(1.29) \quad H^0(Y, \mathcal{O}_X(maP)) \simeq H^0(X, \mathcal{O}(ma(K_X + L_{X/Y})))$$

holds for every $m \geq 0$, where a is the minimal positive integer such that $aL_{X/Y}, aP, aN \in \text{Div}(Y)$.

Let h_P be a C^∞ -hermitian metric on P with semipositive curvature. We set

$$(1.30) \quad U := \{x \in X \mid |\nu!P| \text{ is very ample on a neighbourhood of } x \text{ for every } \nu \gg 1\}.$$

Let a be a positive integer such that $aD \in \text{Div}(X)$. For $m \geq 0$, we shall define inductively a sequence of closed positive current $\{\omega_m\}_{m=0}^\infty$ satisfying the following conditions :

(P1) $\omega_0 = a(\sqrt{-1}\Theta_{h_P} + 2\pi[N])$, where $[N]$ denotes the current of integration over N .

(P2) The cohomology class of $[\omega_m]$ of ω_m is equal to $2\pi a \cdot c_1(K_Y + L_{X/Y})$.

(P3) ω_m is C^∞ on U .

(P4) $\{\omega_m\}_{m=0}^\infty$ satisfies the equation :

$$(1.31) \quad -\text{Ric}_{\omega_m} + \frac{a-1}{a}\omega_{m-1} + \sqrt{-1}\Theta_{h_{L_{X/Y}}} = \omega_m$$

on U for every $m \geq 1$.

(P5) We define the singular hermitian metric h_m by

$$(1.32) \quad h_m := \left(\frac{1}{n!} \omega_{m,abc}^n \right)^{-\frac{1}{a}} \cdot h_{m-1}^{\frac{a-1}{a}} \cdot h_{L_{X/Y}}^{\frac{1}{a}}$$

on $K_Y + L_{X/Y}$.

Then h_m is an AZD of $K_Y + L_{X/Y}$ for every $m \geq 1$.

Theorem 1.7 *The dynamical system $\{\omega_m\}_{m=0}^\infty$ satisfying (P1)-(P5) above exists and is unique and the limit*

$$(1.33) \quad \omega_K := \frac{1}{a} \lim_{m \rightarrow \infty} \omega_m$$

exists in C^∞ -topology on every compact subset of U (cf.(1.30)) and also in the sense of current on Y . The closed positive current ω_Y satisfies the equation :

$$-\text{Ric}_{\omega_Y} + \sqrt{-1}\Theta_{h_{L_{X/Y}}} = \omega_Y$$

on X and ω_Y is the canonical Kähler current on Y (cf. Definition 1.25). \square

The proof of Theorem 1.7 follows from the one of [T6, Theorem 3.2]. Hence we omit it.

1.9 Dynamical construction of the canonical Kähler currents

The canonical Kähler current in Theorem 1.5 can be constructed as the limit of a dynamical system as in ([T4, T6]). Actually we decompose the Ricci iteration in Theorem 1.7 as a sequence of dynamical systems of Bergman kernels. The motivation to construct such a dynamical systems is to prove the plurisubharmonic variation property of canonical measures (cf. Theorem 4.1).

Let $X, Y, f : X \rightarrow Y, (L_{X/Y}, h_{L_{X/Y}})$ as in Section 1.6, Let a be a positive integer such that $f_*\mathcal{O}_X(aK_{X/Y}) \neq 0$. Then we see that

$$(1.34) \quad H^0(X, \mathcal{O}_X(maK_X)) \simeq H^0(Y, \mathcal{O}_Y(ma(K_Y + L_{X/Y})))$$

holds for every $m \geq 0$. In particular $\text{Kod}(X) = \dim Y$ holds. Hence by (1.34), we see that $K_Y + L_{X/Y}$ is big.

Let A be a sufficiently ample line bundle on Y and let h_A be a C^∞ -hermitian metric on A with strictly positive curvature.

We shall construct the singular hermitian metrics $\{h_m\}_{m=1}^\infty$ on $K_Y + L_{X/Y}$ inductively as follows. We shall abuse the same notation as in Section 1.8, since $\{h_m\}$ is eventually the same as Theorem 1.7 (cf. Theorem 1.8 below).

Let $K_Y + L_{X/Y} = P + N$ be the Zariski decomposition as (1.28). First we set $h_1 = h_P \cdot h_N$ as in Section 1.8, where h_P is a C^∞ -hermitian metric on P with semipositive curvature and h_N be a metric on N defined as $h_N := |\sigma_N|^{-2}$, where σ_N be a multivalued holomorphic section of N with divisor N .

Suppose that we have constructed the metric h_{m-1} on $K_Y + L_{X/Y}$ for some $m \geq 2$. To construct h_m , we shall construct a sequence of singular hermitian metrics $\{h_{\ell,m}\}_{\ell=1}^\infty$ as follows.

First we set

$$(1.35) \quad K_{1,m} := K(Y, A + a(K_Y + L_{X/Y}), h_A \cdot h_{m-1}^{a-1} \cdot h_{L_{X/Y}}).$$

and

$$(1.36) \quad h_{1,m} := (K_{1,m})^{-1}.$$

We continue this process. Suppose that we have constructed $K_{\ell,m}$ and the singular hermitian metric $h_{\ell,m}$ on $A + \ell a(K_Y + L_{X/Y})$ for some $\ell \geq 1$.

$$(1.37) \quad K_{\ell+1,m} := K(Y, A + (\ell + 1)a(K_Y + L_{X/Y}), h_{\ell,m} \cdot h_{L_{X/Y}})$$

and

$$(1.38) \quad h_{\ell+1,m} := (K_{\ell+1,m})^{-1}.$$

Thus inductively we construct the sequences $\{h_{\ell,m}\}_{\ell>0}$ and $\{K_{\ell,m}\}_{m>0}$. This inductive construction is essentially the same one originated by the author in [T3]. This is the same construction as in [T6]. The following theorem asserts that the above dynamical system yields the Ricci iteration constructed in Section 1.8.

Theorem 1.8 *Let X be a smooth projective variety of nonnegative Kodaira dimension and let $f : X \rightarrow Y$ be the Iitaka fibration as above. Let $\{h_{\ell,m}\}_{\ell>0}$ be the sequence of hermitian metrics as above and let n denote $\dim Y$. Let ω_m be the Kähler current on Y defined as in Theorem 1.8. Then*

$$(1.39) \quad h_m := \liminf_{\ell \rightarrow \infty} \sqrt[\alpha \ell]{(\ell!)^n \cdot h_{\ell,m}}$$

is a singular hermitian metric on $K_Y + L_{X/Y}$ such that

$$(1.40) \quad h_m = \left(\frac{1}{n!} \omega_{m,abc}^n \right)^{-\frac{1}{\alpha}} \cdot h_{m-1}^{\frac{\alpha-1}{\alpha}} \cdot h_{L_{X/Y}}^{\frac{1}{\alpha}}$$

holds almost everywhere on Y and

$$(1.41) \quad \omega_m := \alpha \sqrt{-1} \Theta_{h_m}.$$

holds on Y . In particular h_m (and hence ω_m) is unique and is independent of the choice of A and h_A . \square

Here it may be better to replace h_m by its lower-semi-continuous envelope because of the following classical theorem.

Theorem 1.9 ([L, p.26, Theorem 5]) *Let $\{u_\alpha\}_{\alpha \in A}$ be a family of plurisubharmonic function on a domain Ω in \mathbb{C}^n . Suppose that $\{u_\alpha\}_{\alpha \in A}$ is locally uniformly bounded from above. Then the upper-semi-continuous envelope of $\sup_{\alpha \in A} u_\alpha$ is again plurisubharmonic on Ω \square*

But anyway this adjustment occurs on a set of measure 0.

2 Construction of the canonical measure by solving Monge-Ampère equations

In this section we shall prove Theorem 1.5. Although the proof of Theorem 1.5 is given in [S-T] independently, we shall give an original proof here for the completeness. Also the proof in [S-T] is quite close to the proof of [T4, Theorem 5.1]². The present proof is different from that in [S-T], in the following points :

- (1) We consider a smoothing of the Hodge metric $h_{L_{X/Y}}$ which need not be of algebraic singularities. Hence we need to consider the sequence of modified equations.
- (2) The C^0 -estimate of the solution depends on the estimate of the Hodge metric near the discriminant locus and the notion of the minimal AZD.

²The original proof of [T4, Theorem 5.1] has a gap in the proof of C^2 -regularity. I have not checked the proof in [S-T] in full detail.

The techniques used here are quite standard and have been known for more than twenty years (cf. [Su, T0]). In this sense the proof of Theorem 1.5 is not essentially new. But as in [Su], we need to require the finite generation of canonical ring ([B-C-H-M]) to prove the C^2 -regularity of the metrics on a Zariski open subset.

2.1 Setup

Let X be a smooth projective n -fold with $\text{Kod}(X) \geq 0$. And let

$$(2.1) \quad f : X \dashrightarrow Y$$

be the Iitaka fibration associated with the complete linear system $|m_0!K_X|$. By taking a suitable modifications, we shall assume the followings.

- (1) f is a morphism.
- (2) Y is smooth.
- (3) $f_*\mathcal{O}_X(m_0!K_{X/Y})^{**}$ is a line bundle on Y .
- (4) The discriminant locus D of f is a divisor with normal crossings on Y .

We define the \mathbb{Q} -line bundle $L_{X/Y}$ on Y by

$$(2.2) \quad L_{X/Y} := \frac{1}{m_0!} f_*\mathcal{O}_X(m_0!K_{X/Y})^{**}$$

and let $h_{L_{X/Y}}$ be the singular hermitian metric on $L_{X/Y}$ defined by

$$(2.3) \quad h_{L_{X/Y}}^{m_0!}(\sigma, \sigma) := \left(\int_{X_y} |\sigma|^{\frac{2}{m_0!}} \right)^{m_0!}.$$

It is clear that $h_{L_{X/Y}}$ is smooth on

$$(2.4) \quad Y^\circ := \{y \in Y \mid f \text{ is smooth over } y\} = Y \setminus D$$

and the singularity of $h_{L_{X/Y}}$ around D is described in terms of variation of Hodge structures. And we see that

$$(2.5) \quad H^0(Y, \mathcal{O}_Y(m!(K_Y + L_{X/Y})) \otimes \mathcal{I}(h_{L_{X/Y}}^{\otimes m})) \simeq H^0(Y, \mathcal{O}_Y(m!(K_Y + L_{X/Y})))$$

holds for every sufficiently large m , i.e., the L^2 -condition with respect to the singular hermitian metric $h_{L_{X/Y}}$ does not affect the global section of $m!(K_Y + L_{X/Y})$.

Let Ω be a C^∞ -volume form on Y . Let us consider the equation :

$$(2.6) \quad -\text{Ric}_{\omega_Y} + \sqrt{-1} \Theta_{h_{L_{X/Y}}} = \omega_Y$$

on Y , where

$$(2.7) \quad \omega_Y = -\text{Ric} \Omega + \sqrt{-1} \Theta_{h_{L_{X/Y}}} + \sqrt{-1} \partial \bar{\partial} u$$

for some unknown upper-semi-continuous function u bounded from above on Y . Then the above equation is equivalent to

$$(2.8) \quad \log \frac{(\omega + \sqrt{-1}\partial\bar{\partial}u)^n}{\Omega} = u,$$

where $n := \dim Y$ and

$$(2.9) \quad \omega := -\text{Ric } \Omega + \sqrt{-1}\Theta_{h_{L_{X/Y}}}.$$

We note that since $\text{Kod}(X) = \dim Y$, $K_Y + (L_{X/Y}, h_{L_{X/Y}})$ is big, i.e.,

$$(2.10) \quad \limsup_{m \rightarrow \infty} (m!)^{-n} \dim H^0(Y, \mathcal{O}_Y(m!(K_Y + L_{X/Y})) \otimes \mathcal{I}(h_{L_{X/Y}}^{m!})) > 0$$

holds, where $n = \dim Y$. The main difficulty for solving the equation (2.6) is the fact that $h_{L_{X/Y}}$ is not of algebraic singularities in the following sense.

Definition 2.1 *Let h be a singular hermitian metric on a line bundle $L_{X/Y}$. We say that h is of **algebraic singularities**, if there exists a positive integer m_0 , global holomorphic sections $\sigma_0, \dots, \sigma_N$ of $m_0 L_{X/Y}$ and a C^∞ -function ϕ such that*

$$(2.11) \quad h = e^\phi \cdot \left(\sum_{i=0}^N |\sigma_i|^2 \right)^{-\frac{1}{m_0}}$$

holds. \square

There are two ways to treat the singularity of $h_{L_{X/Y}}$ in (2.6). One way is to smooth out the singularities of $h_{L_{X/Y}}$ and the other way is to consider the metric of Poincaré growth on the complement of the discriminant locus of $f : X \rightarrow Y$. The both methods depend on the analysis of the singularities of $h_{L_{X/Y}}$ in terms of the theory of variation of Hodge structures.

2.2 Smoothing of the Hodge metric $h_{L_{X/Y}}$

The singular hermitian metric $h_{L_{X/Y}}$ is generically C^∞ , but need not be of algebraic singularities.

The singularity of $h_{L_{X/Y}}$ can be described by using variation of Hodge structures. Let a be the minimal positive integer such that $f_* \mathcal{O}_X(aK_{X/Y})^{**}$ is not 0. Then the a -th root of local holomorphic section of $f_* \mathcal{O}_X(aK_{X/Y})$ can be considered to be a family of canonical forms on the family of cyclic a -covers of the fibers. In this way the Hodge metric can be described in terms of the theory of variation of Hodge structures (cf. [Sch]). Let us assume that the discriminant locus D of $f : X \rightarrow Y$ is a divisor with normal crossings. As in [Ka1], the locally free extension of the Hodge bundle is controlled by the monodromy which is quasi-unipotent. And in this setting the local monodromy is abelian.

Definition 2.2 *Let (M, B) a pair of a complex manifold M and a divisor B with normal crossings. Let ω_P is a Kähler form on $M - B$. ω_P is said to be of Poincaré growth, if for any polydisk $\Delta^n := \{(z_1, \dots, z_n); |z_i| < 1, 1 \leq i \leq n\}$ in M such that*

$$(2.12) \quad \Delta^n \cap B = \{(z_1, \dots, z_n) \in \Delta^n | z_1 \cdots z_k = 0\},$$

there exist locally bounded positive continuous functions $a \leq b$ on Δ^n such that

$$(2.13) \quad a \left(\sum_{i=1}^k \frac{\sqrt{-1} dz_i \wedge d\bar{z}_i}{|z_i|^2 (\log |z_i|)^2} + \sum_{j=k+1}^n \sqrt{-1} dz_j \wedge d\bar{z}_j \right) \leq \omega_P \leq b \left(\sum_{i=1}^k \frac{\sqrt{-1} dz_i \wedge d\bar{z}_i}{|z_i|^2 (\log |z_i|)^2} + \sum_{j=k+1}^n \sqrt{-1} dz_j \wedge d\bar{z}_j \right)$$

hold on $\Delta^n \cap (M - B)$.

Let Ω_P be a volume form on $M - B$. Ω_P is said to be of Poincaré growth, if for any polydisk Δ^n in M such that

$$(2.14) \quad \Delta^n \cap B = \{(z_1, \dots, z_n) \in \Delta^n | z_1 \cdots z_k = 0\},$$

there exists a locally bounded positive continuous function $c(z)$ on Δ^n such that

$$(2.15) \quad \Omega_P = c(z) \cdot \left(\wedge_{i=1}^k \frac{\sqrt{-1} dz_i \wedge d\bar{z}_i}{|z_i|^2 (\log |z_i|)^2} \right) \wedge \left(\wedge_{j=k+1}^n \sqrt{-1} dz_j \wedge d\bar{z}_j \right)$$

holds on $\Delta^n \cap (M - B)$. \square

Remark 2.3 Let Ω_P be a volume form of Poincaré growth on (M, B) with M compact. If for every polydisk Δ^n as in Definiton 2.2 the function $c(z)$ above is C^2 on Δ^n , then $-\text{Ric} \Omega_P$ is of Poincaré growth in the sense that there exists a positive constant C such that

$$(2.16) \quad -C \cdot \omega_P \leq -\text{Ric} \Omega_P \leq C \cdot \omega_P,$$

where ω_P is a Kähler form of $M - B$ with Poincaré growth. This is standard and is easily verified by direct calculation. \square

Then we have the following lemma.

Lemma 2.4 There exists a positive integer m_0 such that $h_{M_{X/Y}}^{m_0!}$ is a singular hermitian metric on the line bundle $f_* \mathcal{O}_Y(m_0! K_{X/Y})^{**}$ such that with respect to a local holomorphic frame

$$(2.17) \quad h_{M_{X/Y}} = O((-\log |\sigma_D|)^q)$$

holds, where σ_D is a local defining function of D and q is a positive integer. Hence we have that the estimate

$$(2.18) \quad h_{L_{X/Y}} = O(|\sigma_D|^{-2\alpha} (-\log |\sigma_D|)^q)$$

holds for some semipositive rational number $\alpha < 1$. And the curvature $\sqrt{-1} \Theta_{h_{M_{X/Y}}}$ is dominated by a constant times a Kähler form ω_P with Poincaré growth on $Y \setminus D$. \square

Remark 2.5 Besides (2.17), we see that $h_{L_{X/Y}}$ is bounded from below by a smooth metric on $L_{X/Y}$, since $h_{L_{X/Y}}$ has semipositive curvature in the sense of current ([Ka2],[Ka3, p.174, Theorem 1.1]). This fact is essentially due to the theory of variation of Hodge structures. \square

The estimate of $h_{L_{X/Y}}$ in Lemma 2.4 follows from [Ka1] which uses the theory variation of Hodge structures due to W. Schmidt ([Sch]). And the latter estimate of $\sqrt{-1}\Theta_{h_{L_{X/Y}}}$ follows from the fact that holomorphic sectional curvature in the horizontal direction of the period domain is dominated by a negative constant ([G]) and the Yau-Royden Schwarz lemma ([Y2, R]).

In this sense $h_{L_{X/Y}}$ is very close to a smooth metric. To smooth out $h_{L_{X/Y}}$, we take a finite open covering $\mathcal{U} := \{U_\alpha\}$ of Y such that every U_α is biholomorphic to the open unit ball in \mathbb{C}^n with center O via the coordinate $z_\alpha = (z_\alpha^1, \dots, z_\alpha^n)$. Taking \mathcal{U} properly, we may and do assume z_α is a holomorphic coordinate on a larger open subset \hat{U}_α which is biholomorphic to the open ball with radius 2 in \mathbb{C}^n with center O via z_α . Let h_0 be a C^∞ -hermitian metric on the \mathbb{Q} line bundle $L_{X/Y}$. We set

$$(2.19) \quad \varphi := \log \frac{h_{L_{X/Y}}}{h_0}.$$

Let ρ be a C^∞ -function on \mathbb{C}^n such that $0 \leq \rho \leq 1$, $\text{supp } \rho$ is contained in the unit open ball in \mathbb{C}^n with center O and

$$(2.20) \quad \int_{\mathbb{C}^n} \rho(z) d\mu(z) = 1,$$

where $d\mu$ is the usual Lebesgue measure on \mathbb{C}^n . For every $0 < \delta < 1$, we set

$$(2.21) \quad \rho_\delta(z) = \delta^{-2n} \rho(z/\delta).$$

We shall take a mollification $\varphi_{\alpha,\delta}$ of $\varphi|_{U_\alpha}$ using the convolution with the mollifier ρ_δ as

$$(2.22) \quad \varphi_{\alpha,\delta} := (\varphi|_{U_\alpha}) * \rho_\delta$$

with respect to the coordinate z_α . Since z_α is a holomorphic coordinate on \hat{U}_α , $\varphi_{\alpha,\delta}$ is a well defined C^∞ -function on U_α for every $0 < \delta < 1$. Then $\varphi_{\alpha,\delta}$ converges to φ in L^1 -topology on U_α and compact uniformly in C^∞ -topology on $U_\alpha \setminus D$ as $\delta \downarrow 0$.

Let $\{\phi_\alpha\}$ be a partition of unity subordinate to \mathcal{U} . We set

$$(2.23) \quad h_{L_{X/Y},\delta} := \exp\left(\sum_{\alpha} \phi_\alpha \cdot \varphi_{\alpha,\delta}\right) \cdot h_0.$$

Then $h_{L_{X/Y},\delta}$ is a C^∞ -hermitian metric on $L_{X/Y}$ and there exists a positive constant C independent of $\delta > 0$ such that

$$(2.24) \quad \sqrt{-1}\Theta_{h_{L_{X/Y},\delta}} \leq C \cdot \omega_P$$

holds. In general $h_{L_{X/Y},\delta}$ does not have semipositive curvature. But by the construction, there exists a continuous function $e(\delta)$ on Y such that

$$(2.25) \quad \sqrt{-1}\Theta_{h_{L_{X/Y},\delta} } \geq -e(\delta) \cdot \omega_P$$

and

$$(2.26) \quad \lim_{\delta \downarrow 0} e(\delta) = 0$$

holds uniformly on Y .

2.3 The construction of the canonical Kähler currents

In this subsection, we shall prove the existence of the current ω_Y satisfying (2.6) without assuming the finite generation of canonical ring ([B-C-H-M]). This result is slightly weaker than Theorem 1.5. But the same strategy works to construct canonical Kähler-Einstein currents on LC pairs (cf. [T6]). Hence the following theorem has independent interest.

Theorem 2.6 *In the notations in Section 2.1, there exists a closed positive current ω_Y on Y such that*

- (1) ω_Y represents $2\pi c_1(K_Y + L_{X/Y})$.
- (2) $h_K := n! \left(\omega_{Y,abc}^n \right)^{-1} \cdot h_{L_{X/Y}}$ ($n = \dim Y$) is an AZD of $K_Y + L_{X/Y}$, where $\omega_{Y,abc}$ denotes the absolutely continuous part of ω_Y .
- (3) $\omega_Y = \sqrt{-1} \Theta_{h_K}$ holds on Y .
- (4) $-\text{Ric}_{\omega_Y} + \sqrt{-1} \Theta_{L_{X/Y}} = \omega_Y$ holds on Y in the sense of current, where $\text{Ric}_{\omega_Y} := \sqrt{-1} \partial \bar{\partial} \log \omega_{Y,abc}^n$. \square

The proof of Theorem 2.6 depends on the monotonicity lemma (Lemma 2.7).

Let m_0 be a sufficiently large positive integer such that for every $m \geq m_0$, $m!(K_Y + L_{X/Y})$ is a Cartier divisor on Y and $|m!(K_Y + L_{X/Y})|$ gives a birational embedding of Y . Let $\pi_m : Y_m \rightarrow Y$ be the resolution of Bs $|m!(K_Y + L_{X/Y})|$ such that for every $m > m_0$

$$(2.27) \quad \pi_m : Y_m \rightarrow Y$$

factors through $\pi_{m-1} : Y_{m-1} \rightarrow Y$. Let

$$(2.28) \quad \mu_m : Y_m \rightarrow Y_{m-1}$$

be the natural morphism. Here we may and do take $\mu_m : Y_m \rightarrow Y_{m-1}$ such that the exceptional divisor of π_m is contained in $\pi_m^{-1}(V)$. This is certainly possible by the definition of V . Hence we have an (possibly infinite) tower

$$(2.29) \quad \cdots \xrightarrow{\mu_{m+2}} Y_{m+1} \xrightarrow{\mu_{m+1}} Y_m \xrightarrow{\mu_m} Y_{m-1} \xrightarrow{\mu_{m-1}} \cdots$$

In the following proof, we shall consider this tower. But by the recent result on finite generation of canonical ring ([B-C-H-M]) we may avoid to consider an infinite tower. Namely we just need to consider one sufficiently large m_0 . This certainly simplifies the proof. The reason why we do not use the finite generation of canonical ring is that it is not essential from the analytic point of view and one may extend the theory to the case of LC pairs (cf. [T6]). Let

$$(2.30) \quad \pi_m^* |m!(K_Y + L_{X/Y})| = |P_m| + F_m$$

be the decomposition of $\pi_m^* |m!(K_Y + L_{X/Y})|$ into the free part $|P_m|$ and the fixed component F_m . Let V be the analytic subset of Y defined by:

$$V := \{y \in Y^\circ \mid y \in \cap_{m>0} \text{Bs} |m!(K_Y + L_{X/Y})| \text{ or} \\ \Phi_{|m!(K_Y + L_{X/Y})|} \text{ is not an embedding around } y \text{ for } m \gg m_0\} \\ \cup \{\text{the discriminant locus of } f\}.$$

By taking a suitable modification of Y , we may and do assume that V is a divisor with normal crossings.

There exists an effective \mathbb{Q} -divisor E_m on Y_m respectively such that the followings hold for every $m \geq m_0$.

- (1) $P_m - E_m$ is ample on Y_m .
- (2) All the coefficients of E_m are less than 1, i.e., $[E_m] = 0$.
- (3) $\text{Supp } E_m = \pi_m^{-1}(V)$.
- (4) $((m+1)!)^{-1}(P_{m+1} - E_{m+1}) - \mu_{m+1}^*(m!)^{-1}(P_m - E_m)$ is effective.

The existence of such $\{E_m\}$ follows from the definition of V and the trivial fact that for any composition of successive blowing ups

$$\varpi : \tilde{\mathbb{P}}^\nu \rightarrow \mathbb{P}^\nu$$

of a projective space \mathbb{P}^ν with smooth centers, there exists an effective \mathbb{Q} -divisor B supported on the exceptional divisors of ϖ such that $\varpi^*\mathcal{O}(1) - B$ is ample. Hence by taking a suitable successive blowing ups with smooth centres over V , if necessary, we may assume the existence of such effective \mathbb{Q} -divisors $\{E_m\}$ by considering the image of Y_m by the morphism associated with $|P_m|$ from Y_m into a projective space.

After taking such a sequence $\{E_m\}$, we replace $\{E_m\}$ by $\{2^{-m}E_m\}$. Then it has the same properties as above. And we shall denote $\{2^{-m}E_m\}$ again by $\{E_m\}$. Then instead of (4) we have

$$(2.31) \quad (4)' \quad ((m+1)!)^{-1}(P_{m+1} - E_{m+1}) - \mu_{m+1}^*(m!)^{-1}(P_m - E_m) \text{ is effective}$$

and contains $\varepsilon_m(\pi_{m+1}^{-1}V)_{red}$ for some positive number ε_m .

Let $h_{(m)}$ be a C^∞ -hermitian metric on $\pi_m^*((m!)^{-1}(P_m - E_m))$ with strictly positive curvature. We note that by (2.31) $\{h_{(m)}\}$ is getting less singular as m tends to infinity and $h_{(m+1)}$ is strictly less singular than $h_{(m)}$ along V (if we consider the metrics as singular hermitian metrics on $K_Y + L_{X/Y}$). Then

$$(2.32) \quad \Omega_{m,\delta} := h_{(m)}^{-1} \cdot (\pi_m^* h_{L_{X/Y},\delta})$$

is considered as a degenerate volume form on Y_m , where $\{h_{L_{X/Y},\delta}\}$ is the smoothing of the Hodge metric $h_{L_{X/Y}}$ as in Section 2.2. We note that $\Omega_{m,\delta}^{-1} \cdot (\pi_m^* h_{L_{X/Y},\delta})^{-1} = h_{(m)}$ is a metric with algebraic singularities on $K_Y + L_{X/Y}$.

Now we shall consider the equation :

$$(2.33) \quad (-\text{Ric } \Omega_{m,\delta} + \sqrt{-1} \pi_m^* \Theta_{h_{L_{X/Y},\delta}} + \sqrt{-1} \partial \bar{\partial} u_{m,\delta})^n = \Omega_{m,\delta} \cdot e^{u_{m,\delta}}.$$

on Y_m . Then by the definition (2.32) of $\Omega_{m,\delta}$, (2.33) is equivalent to

$$(2.34) \quad (\sqrt{-1} \Theta_{h_{(m)}} + \sqrt{-1} \partial \bar{\partial} u_{m,\delta})^n = \Omega_{m,\delta} \cdot e^{u_{m,\delta}}.$$

Then since $\sqrt{-1} \Theta_{h_{(m)}}$ is a C^∞ -Kähler form on Y_m by [Y1, p.387, Theorem 6], solving (2.33), we see that there exists a $u_{m,\delta} \in C^\infty(Y_m \setminus \pi_m^{-1}V)$ and the closed positive current:

$$(2.35) \quad \omega_{m,\delta} := -\text{Ric } \Omega_{m,\delta} + \sqrt{-1} \pi_m^* \Theta_{h_{L_{X/Y},\delta}} + \sqrt{-1} \partial \bar{\partial} u_{m,\delta}$$

on Y_m such that

- (1) $-\text{Ric}_{\omega_{m,\delta}} + \sqrt{-1} \pi_m^* \Theta_{h_{L_{X/Y},\delta}} = \omega_{m,\delta}$ holds on $Y_m - \text{Supp } E_m$,
- (2) The absolutely continuous part $\omega_{m,\delta,abc}$ of $\omega_{m,\delta}$ is closed and represents $2\pi(m!)^{-1}(P_m - E_m)$,
- (3) $(\pi_m)_* \omega_{m,\delta}$ represents the class $2\pi c_1(K_Y + L_{X/Y})$,
- (4) There exists a positive constant $C(m, \delta)$ such that

$$(2.36) \quad |u_{m,\delta}| \leq C(m, \delta)$$

holds on Y_m .

Here we note that

$$(2.37) \quad -\text{Ric } \Omega_{m,\delta} + \sqrt{-1} \pi_m^* \Theta_{h_{L_{X/Y},\delta}} = \sqrt{-1} \Theta_{h_{(m)}}$$

holds, hence it is independent of δ . If we set

$$(2.38) \quad \omega_{(m)} := \sqrt{-1} \Theta_{h_{(m)}},$$

then the equation is transcribed as :

$$(2.39) \quad \log \frac{\omega_{m,\delta}^n}{\Omega_{m,\delta}} = \log \frac{(\omega_{(m)} + \sqrt{-1} \partial \bar{\partial} u_{m,\delta})^n}{\Omega_{m,\delta}} = u_{m,\delta}.$$

Let us consider $\{\omega_{m,\delta}^n\}$ as a sequence of volume forms on $Y \setminus V^3$. And we shall identify $(\pi_m)_* \omega_{m,\delta}$ with $\omega_{m,\delta}$ on $Y \setminus V$. Hereafter we shall identify $Y \setminus V$ with a Zariski open subset of Y_m for every m and consider everything on $Y \setminus V$ (if without fear of confusion). Then by the maximum principle we have the following monotonicity lemma.

Lemma 2.7 (*Monotonicity Lemma*)

$$(2.40) \quad \omega_{m,\delta}^n \leq \omega_{m+1,\delta}^n$$

holds on $Y \setminus V$. \square

Proof of Lemma 2.7. We note that by the construction the followings hold.

- (1) The absolutely continuous parts of $\omega_{m,\delta}$ and $\omega_{m+1,\delta}$ represent $2\pi(m!)^{-1}(P_m - E_m)$ and $2\pi((m+1)!)^{-1}(P_{m+1} - E_{m+1})$ respectively.
- (2) $\mu_{m+1}^*((m!)^{-1}(F_m + E_m)) - ((m+1)!)^{-1}(F_{m+1} + E_{m+1})$ is effective and contains $\varepsilon_m(\pi_{m+1}^{-1}V)$ for some positive number ε_m .

We note that by the boundedness of $u_{m,\delta}$ and $u_{m+1,\delta}$ (cf. (2.36)) and the equation (2.39), we see that the asymptotics of $\omega_{m,\delta}^n$ and $\omega_{m+1,\delta}^n$ near V is the same as $\Omega_{m,\delta}$ and $\Omega_{m+1,\delta}$ respectively. Since by the condition (2) above, $\Omega_{m,\delta}/\Omega_{m+1,\delta}$ tends to 0 toward V , the function $\phi_{m,\delta}$ defined by

$$(2.41) \quad \phi_{m,\delta} := \log \frac{\omega_{m,\delta}^n}{\omega_{m+1,\delta}^n}$$

³Since we consider $\omega_{m,\delta}$ as a current, it seems to be more authentic to denote $\omega_{m,\delta,abc}^n$ instead of $\omega_{m,\delta}$. But we consider the equation (2.39) on $Y \setminus V$.

tends to $-\infty$ toward V . Hence there exists a point $p_m \in Y \setminus V$, where $\phi_{m,\delta}$ takes its maximum. Then

$$(2.42) \quad \sqrt{-1} \partial \bar{\partial} \phi_{m,\delta}(p_m) \leq 0$$

holds. By the equation

$$(2.43) \quad -\text{Ric}_{\omega_{k,\delta}} + \sqrt{-1} \pi_k^* \Theta_{h_{L_{X/Y},\delta}} = \omega_{k,\delta}, (k = m, m+1)$$

(2.42) implies that

$$(2.44) \quad \omega_{m,\delta}(p_m) \leq \omega_{m+1,\delta}(p_m)$$

holds. In particular $\phi_{m,\delta}(p_m) \leq 0$ holds. Hence by the definition of p_m , this implies that $\phi_{m,\delta} \leq 0$ holds on $Y \setminus V$. Hence $\omega_{m,\delta}^n \leq \omega_{m+1,\delta}^n$ holds on $Y \setminus V$. This completes the proof of Lemma 2.7. \square

Now we shall consider the uniform C^0 -estimate on every compact subset of $Y \setminus V$. Let us fix a positive integer $s \geq m_0$ and let

$$\omega_{(s)} = \sqrt{-1} \Theta_{h_{(s)}}$$

be the Kähler form defined as (2.38). We shall use $h_{(s)}$ and $\omega_{(s)}$ as standards in the following estimate. For every $m > s$, let $v_{m,\delta}$ be a C^∞ -function on $Y \setminus D$ defined by

$$(2.45) \quad v_{m,\delta} := u_{m,\delta} + \log \frac{h_{(s)}}{h_{(m)}}.$$

Then

$$(2.46) \quad \omega_{m,\delta} = \omega_{(s)} + \sqrt{-1} \partial \bar{\partial} v_{m,\delta}$$

and

$$(2.47) \quad \log \frac{(\omega_{(s)} + \sqrt{-1} \partial \bar{\partial} v_{m,\delta})^n}{\Omega_{s,\delta}} = v_{m,\delta}$$

hold. By the condition (2.31), we see that for every $m > s$, $\log(h_{(s)}/h_{(m)})$ tends to $+\infty$ toward V . And by the boundedness of $u_{m,\delta}$ (cf. (2.36)), we have the estimate:

$$(2.48) \quad -C(m,\delta) + \log \frac{h_{(s)}}{h_{(m)}} \leq v_{m,\delta} \leq C(m,\delta) + \log \frac{h_{(s)}}{h_{(m)}},$$

where $C(m,\delta)$ is the positive constant as in (2.36). Hence $v_{m,\delta}$ tends to $+\infty$ toward V . This implies that there exists a point $p_0 \in Y \setminus V$, where $v_{m,\delta}$ takes its minimum. Now we note that by (2.47),

$$(2.49) \quad \log \frac{(\omega_{(s)} + \sqrt{-1} \partial \bar{\partial} v_{m,\delta})^n}{\omega_{(s)}^n} = \int_0^1 \Delta_{(s,m,\delta,t)} v_{m,\delta} dt = v_{m,\delta} - \log \frac{\omega_{(s)}^n}{\Omega_{s,\delta}}$$

hold, where $\Delta_{(s,m,\delta,t)}$ denotes the trace of $\sqrt{-1}\partial\bar{\partial}v_{m,\delta}$ with respect to the Kähler form: $(1-t)\omega_{m,\delta} + t\omega_{(s)}$. Hence by the minimum principle,

$$(2.50) \quad v_{m,\delta}(p_0) \geq \log \frac{\omega_{(s)}^n}{\Omega_{s,\delta}}(p_0)$$

holds.

On the other hand, by (2.32) and the definition of $h_{L_{X/Y},\delta}$ (cf. (2.23)), $\Omega_{s,\delta}$ tends to 0 toward V . Hence there exists a positive constant $C_-(s)$ independent of δ such that

$$(2.51) \quad \min_{y \in Y} \log \frac{\omega_{(s)}^n}{\Omega_{s,\delta}}(y) \geq C_-(s)$$

holds. By (2.51) and (2.50), we see that

$$(2.52) \quad v_{m,\delta}(y) \geq C_-(s)$$

holds for every $y \in Y \setminus V$. By the definition of $v_{m,\delta}$ (cf. (2.45)), we have that

$$(2.53) \quad u_{m,\delta} \geq \log \frac{h_{(m)}}{h_{(s)}} + C_-(s)$$

holds. Replacing s by $t > s$, for $m > t > s$ we have

$$u_{m,\delta} \geq \log \frac{h_{(m)}}{h_{(t)}} + C_-(t)$$

and hence by (2.45), we obtain the following lemma.

Lemma 2.8 *There exists a positive constant $C_-(t)$ depending only on $t > s$ such that for every $m > t$*

$$(2.54) \quad v_{m,\delta} \geq \log \frac{h_{(s)}}{h_{(t)}} + C_-(t)$$

holds. In particular $v_{m,\delta}$ tends to infinity toward V . \square

On the other hand, we obtain the upper estimate of $u_{m,\delta}$ as follows. We may and do assume that V is a divisor with normal crossings. Let Ω_P be a volume form on $Y \setminus V$ with Poincaré growth, i.e., for every polydisk Δ^n in Y such that

$$(2.55) \quad \Delta^n \cap V = \{(z_1, \dots, z_n) \in \Delta^n \mid z_1 \cdots z_k = 0\},$$

$$(2.56) \quad \Omega_P = c \frac{|dz_1 \wedge \cdots \wedge dz_n|^2}{\prod_{i=1}^k |z_i| (\log |z_i|^2)^2},$$

where c is a positive C^∞ -function on Δ^n . Such a Ω_P can be constructed easily by using a partition of unity. We set

$$(2.57) \quad \tilde{u}_{m,\delta} := u_{m,\delta} + \log \frac{\Omega_{m,\delta}}{\Omega_P}.$$

By the condition (2.31) and the boundedness of $u_{m,\delta}$, there exists a point p'_0 on $Y \setminus V$ such that $\tilde{u}_{m,\delta}$ takes its maximum at p'_0 . Then since

$$(2.58) \quad \log \frac{\omega_{m,\delta}^n}{\Omega_P} = u_{m,\delta} + \log \frac{\Omega_{m,\delta}}{\Omega_P} = \tilde{u}_{m,\delta}$$

hold, at p'_0 we have that

$$(2.59) \quad \sqrt{-1} \partial \bar{\partial} \log \frac{\omega_{m,\delta}^n}{\Omega_P}(p'_0) \leq 0$$

holds. Hence we have the inequality:

$$(2.60) \quad -\text{Ric}_{\omega_{m,\delta}} \leq (-\text{Ric } \Omega_P)(p'_0).$$

By the equation:

$$(2.61) \quad -\text{Ric}_{\omega_{m,\delta}} + \sqrt{-1} \Theta_{h_{L_{X/Y},\delta}} = \omega_{m,\delta},$$

we see that

$$(2.62) \quad \omega_{m,\delta}(p'_0) \leq -\text{Ric } \Omega_P + \sqrt{-1} \Theta_{h_{L_{X/Y},\delta}}$$

holds. In particular,

$$(2.63) \quad \omega_{m,\delta}^n(p'_0) \leq (-\text{Ric } \Omega_P + \sqrt{-1} \Theta_{h_{L_{X/Y},\delta}})^n(p'_0)$$

holds. This implies that

$$(2.64) \quad \tilde{u}_{m,\delta} = u_{m,\delta} + \log \frac{\Omega_{m,\delta}}{\Omega_P} \leq \log \frac{(-\text{Ric } \Omega_P + \sqrt{-1} \Theta_{h_{L_{X/Y},\delta}})^n}{\Omega_P}(p'_0)$$

holds on Y . By the construction of $h_{L_{X/Y},\delta}$, Lemma 2.4 and Remark 2.3, there exists a positive constant C_+ independent of δ such that

$$(2.65) \quad \frac{(-\text{Ric } \Omega_P + \sqrt{-1} \Theta_{h_{L_{X/Y},\delta}})^n}{\Omega_P} \leq \exp(C_+)$$

holds on $Y \setminus V$. Combining (2.64) and (2.65) we have that

$$(2.66) \quad u_{m,\delta} \leq C_+ - \log \frac{\Omega_{m,\delta}}{\Omega_P}$$

and

$$(2.67) \quad e^{u_{m,\delta}} \Omega_{m,\delta} \leq \exp(C_+) \cdot \Omega_P$$

hold on Y .

On the other hand by the definition (2.32) and the definition of the smoothing $\{h_{L_{X/Y},\delta}\}$ (2.23), there exists a sequence of positive number $\{\epsilon(\delta)\}$ such that

$$(1) \quad \lim_{\delta \rightarrow 0} \epsilon(\delta) = 0 \text{ holds,}$$

(2) For every $0 < \lambda < \delta$, the inequality:

$$(2.68) \quad \Omega_{m,\delta} \leq (1 + \epsilon(\delta))\Omega_{m,\lambda}$$

holds.

Then for $0 < \lambda < \delta$ by (2.39)

$$(2.69) \quad \log \frac{(\omega_{(m)} + \sqrt{-1}\partial\bar{\partial}u_{m,\lambda})^n}{(\omega_{(m)} + \sqrt{-1}\partial\bar{\partial}u_{m,\delta})^n} = \log \frac{\Omega_{m,\lambda}}{\Omega_{m,\delta}} + (u_{m,\lambda} - u_{m,\delta})$$

holds. We note that

$$(2.70) \quad \log \frac{(\omega_{(m)} + \sqrt{-1}\partial\bar{\partial}u_{m,\lambda})^n}{(\omega_{(m)} + \sqrt{-1}\partial\bar{\partial}u_{m,\delta})^n} = \int_0^1 \tilde{\Delta}_t(u_{m,\lambda} - u_{m,\delta}) dt,$$

where $\tilde{\Delta}_t$ denotes the Laplacian with respect to $(1-t)\tilde{\omega}_{m,\lambda} + t\tilde{\omega}_{m,\delta}$. Then by (2.69),(2.70) and the maximum principle, we see that

$$(2.71) \quad u_{m,\lambda} - u_{m,\delta} \leq -\min_Y \log \frac{\Omega_{m,\lambda}}{\Omega_{m,\delta}} < \log(1 + \epsilon(\delta))$$

holds on Y . This argument is not quite right, since $u_{m,\lambda}, u_{m,\delta}$ are not C^2 on Y (although they are C^2 bounded).

To justify the argument we proceed as in the proof of [p.387, Theorem 6][Y1], i.e., we shall consider the perturbation of the equation (2.39). By the construction of $\Omega_{m,\delta}$ (cf. (2.32)) the 0-locus of $\Omega_{m,\delta}$ is the divisor $(\pi_m^* V)_{red}$ with normal crossings on Y_m . Let $(\pi_m^* V)_{red} = \sum V_j^{(m)}$ be the irreducible decomposition. Let us write $\Omega_{m,\delta}$ as

$$(2.72) \quad \Omega_{m,\delta} = \left(\prod_j \|\tau_j\|^{2a_j} \right) \cdot \tilde{\Omega}_{m,\delta},$$

where $\tilde{\Omega}_{m,\delta}$ is a nondegenerate C^∞ -volume form on Y_m and $\|\tau_j\|$ denotes the hermitian norm of a holomorphic section τ_j of the line bundle $\mathcal{O}_{Y_m}(V_j^{(m)})$ with divisor $V_j^{(m)}$ on Y_m with respect to a fixed C^∞ -hermitian metric and $\{a_j\}$ are positive rational numbers. We may and do take the factor $(\prod_j \|\tau_j\|^{2a_j})$ independent of δ . Now for $0 < \epsilon \ll 1$, we shall consider the perturbed equation:

$$(2.73) \quad \log \frac{(\omega_{(m)} + \sqrt{-1}\partial\bar{\partial}u_{m,\delta}(\epsilon))^n}{\Omega_{m,\delta}(\epsilon)} = u_{m,\delta}(\epsilon),$$

where $\Omega_{m,\delta}(\epsilon)$ is a C^∞ nondegenerate volume form on Y defined by

$$(2.74) \quad \Omega_{m,\delta}(\epsilon) = \left(\prod_j (\|\tau_j\|^2 + \epsilon)^{a_j} \right) \cdot \tilde{\Omega}_{m,\delta}.$$

Then (2.73) has a unique C^∞ -solution $u_{m,\delta}(\epsilon)$ and

$$(2.75) \quad u_{m,\delta} = \lim_{\epsilon \downarrow 0} u_{m,\delta}(\epsilon)$$

holds as in [Y1, p.387, Theorem 6] in the C^2 -norm with respect to the Kähler form $\omega_{(m)}$ on Y_m . Then replacing $u_{m,\lambda}, u_{m,\delta}$ by $u_{m,\lambda}(\varepsilon), u_{m,\delta}(\varepsilon)$ respectively and letting $\varepsilon \downarrow 0$, we may justify (2.71). Since $\{u_{m,\delta}\}$ is almost monotone decreasing as $\delta \downarrow 0$ as (2.71), we have that

$$(2.76) \quad dV_m := \frac{1}{n!} \lim_{\delta \downarrow 0} e^{u_{m,\delta}} \Omega_{m,\delta}$$

exists and by (2.39), we have

$$(2.77) \quad dV_m = \frac{1}{n!} \lim_{\delta \downarrow 0} \omega_{m,\delta,abc}^n$$

holds on Y . We note that

$$\int_Y \omega_{m,\delta,abc}^n = (m!)^{-n} (P_m - E_m)^n$$

holds. By (2.67) and the Lebesgue's bounded convergence theorem, we see that

$$(2.78) \quad \frac{1}{(2\pi)^n} \int_Y dV_m = \frac{1}{n!(m!)^n} (P_m - E_m)^n$$

holds. We set

$$(2.79) \quad \omega_m := \lim_{\delta \downarrow 0} \omega_{m,\delta}$$

in the sense of current and

$$(2.80) \quad h_m := dV_m^{-1} \cdot h_{L_{X/Y}}.$$

Then h_m is a singular hermitian metric on $K_Y + L_{X/Y}$ with semipositive curvature in the sense of current by (2.33) and (2.39). But also we may consider h_m as a singular hermitian metric \tilde{h}_m on $(m!)^{-1}(P_m - E_m)$ by the natural inclusion :

$$\mathcal{O}_{Y_m}(\ell! \pi_m^*(m!(P_m - E_m))) \hookrightarrow \mathcal{O}_{Y_m}(\ell! \pi_m^*(m!(K_Y + L_{X/Y}))),$$

where ℓ is a sufficiently large positive integer such that $\ell! \pi_m^*(m!(P_m - E_m))$ is Cartier. Now we introduce the following notion.

Definition 2.9 *Let M be a projective manifold of dimension n and let (L, h_L) be a pseudoeffective singular hermitian line bundle on X . We define the number $\mu(L, h_L)$ by*

$$\mu(L, h_L) := n! \limsup_{m \rightarrow \infty} m^{-n} h^0(M, \mathcal{O}_M(mL) \otimes \mathcal{I}(h_L^m))$$

is called the volume of (L, h_L) . \square

Remark 2.10 *This definition is easily generalized to the case of singular hermitian \mathbb{Q} -line bundles. \square*

Suppose that the Lelong number $\nu(\sqrt{-1}\Theta_{\tilde{h}_m})$ of $\sqrt{-1}\Theta_{\tilde{h}_m}$ satisfies the inequality: $\nu(\sqrt{-1}\Theta_{\tilde{h}_m}, y_m) > c$ for some $y_m \in Y_m$ and a positive number c , then by the basic property of the Lelong number, we see that

$$(2.81) \quad \mathcal{I}(\tilde{h}_m^{\ell!})_{y_m} \subseteq \mathfrak{m}_{y_m}^{\lfloor c\ell! \rfloor}$$

holds for every sufficiently large ℓ , where \mathfrak{m}_{y_m} denotes the maximal ideal at y_m . Hence the strict inequality

$$(2.82) \quad \mu((m!)^{-1}(P_m - E_m), \tilde{h}_m) < (m!)^{-n}(P_m - E_m)^n$$

holds, where $\mu((m!)^{-1}(P_m - E_m), \tilde{h}_m)$ denotes the volume of $((m!)^{-1}(P_m - E_m), \tilde{h}_m)$ (cf. Definition 3.4 below). On the other hand

$$(2.83) \quad \frac{1}{(2\pi)^n} \int_Y \omega_{m,abc}^n = \mu((m!)^{-1}(P_m - E_m), \tilde{h}_m)$$

holds by (2.78) and a theorem of Boucksom ([Bo, Proposition 3.1]). By the equality (2.78), (2.83) contradicts (2.82). Hence $\nu(\sqrt{-1}\Theta_{\tilde{h}_m})$ is identically 0 and hence \tilde{h}_m is an AZD of $(m!)^{-1}(P_m - E_m)$. By Lemma 2.7, we see that $\{dV_m\}$ is monotone increasing in m on Y , hence $\{h_m\}$ is getting less singular as m tends to infinity as metrics on $K_Y + L_{X/Y}$. Then by (2.67) and Lebesgue's bounded convergence theorem, we have the following lemma.

Lemma 2.11

$$(2.84) \quad dV_Y := \frac{1}{n!} \lim_{m \rightarrow \infty} dV_m$$

exists as a degenerate volume form on Y . And if we define the singular hermitian metric h_K on $K_Y + L_{X/Y}$ by

$$(2.85) \quad h_K := dV_Y^{-1} \cdot h_{L_{X/Y}},$$

then h_K is an AZD of $K_Y + L_{X/Y}$. \square

By the construction of ω_m ,

$$(2.86) \quad -\text{Ric}_{\omega_m} + \sqrt{-1}\pi_m^* \Theta_{h_{L_{X/Y}}} = \omega_m$$

holds for every $m \geq 1$. Then by (2.86) and Lemma 2.11, if we set

$$(2.87) \quad \omega_Y := \sqrt{-1}\Theta_{h_K},$$

then $\omega_Y := \lim_{m \rightarrow \infty} \omega_m$ holds and ω_K is a closed positive current on Y . Moreover

$$(2.88) \quad \omega_Y^n = \lim_{m \rightarrow \infty} \omega_m^n$$

holds on $Y \setminus V$ by the definition. Hence ω_Y satisfies equation:

$$(2.89) \quad -\text{Ric}_{\omega_Y} + \sqrt{-1}\Theta_{h_{L_{X/Y}}} = \omega_Y.$$

This completes the proof of Theorem 2.6. \square

2.4 Regularity of the canonical Kähler current

Here we shall prove Theorem 1.5, by using the recent result on the finite generation of canonical ring ([B-C-H-M]). By Theorem 2.6, we only need to prove the C^∞ -regularity of h_K and ω_Y on a nonempty Zariski open subset of Y .

The proof here is more or less parallel to the existence of the singular Kähler-Einstein metrics in [Su, T4] and is based on [Y1] and the idea in [T0]. But since the Hodge metric $h_{L_{X/Y}}$ is not of algebraic singularities, we need to consider the smoothing of the Hodge metric. This is the major difference. We continue to use the notations in Section 2.3. Let us start the proof of Theorem 1.5. By [B-C-H-M], we see that the canonical ring $R(X, K_X)$ (cf.(1.18)) is finitely generated. Then by the definition of $L_{X/Y}$, we see that $R(Y, K_Y + L_{X/Y})$ is finitely generated also. Hence this implies that the tower (2.29) above can be taken to be finite. Here we shall assume that (2.29) is a finite tower. Moreover taking m_0 sufficiently large, we may assume that $\mu_m : Y_m \rightarrow Y_{m-1}$ (cf.(2.28)) is identity for every $m \geq m_0$ and

$$(2.90) \quad P = \frac{1}{m!} P_m$$

is independent of m . Moreover we may and do assume that $Y_m = Y$ holds for all m . In this case, only E_m varies and we may assume that E_m is of the form

$$(2.91) \quad E_m = \frac{1}{2^m} E,$$

where E is a fixed effective \mathbb{Q} -divisor supported on V such that $P - E$ is ample.

Next we shall fix a C^∞ -hermitian metric $h_{(m)}$ with strictly positive curvature on $(m!)^{-1}(P_m - E_m)$ as in Section 2.3. Let h_P be a C^∞ -hermitian metric defined by the pull back of the Fubini-Study metric on the hyperplane bundle on \mathbb{P}^N via the morphism $\Phi_{|P_{m_0}|} : Y \rightarrow \mathbb{P}^N$ and let h_0 be a C^∞ hermitian metric on $P - E$ with strictly positive curvature on Y . And we shall take $h_{(m)}$ in the previous section as

$$(2.92) \quad h_{(m)} := h_P^{1 - \frac{1}{2^{m!}}} \cdot h_0^{\frac{1}{2^{m!}}}.$$

Let us fix $s \geq m_0$ as in the Section 2.3.

By (2.53) and (2.66), we see that there exists a positive constant C'_0 independent of m and δ such that

$$(2.93) \quad \int_Y |u_{m,\delta}| \omega_{(s)}^n < C'_0$$

holds. Then since

$$(2.94) \quad \omega_{(m)} + \sqrt{-1} \partial \bar{\partial} u_{m,\delta}$$

is a closed positive current on Y and $\omega_{(m)}$ is a C^∞ -Kähler form on Y , $u_{m,\delta}$ is an almost plurisubharmonic function on Y . By the sub-mean-value inequality for plurisubharmonic functions, we see that by (2.93) there exists a positive constant C_0 such that

$$(2.95) \quad u_{m,\delta} \leq C_0$$

holds on Y .

By Lemma 2.8, (2.95) and (2.45), we have the following lemma.

Lemma 2.12 *Let $s < t < m$ be as above. Then there exists a positive constant C_0 independent of m and δ such that for every $\delta > 0$,*

$$(2.96) \quad -C_-(t) + \log \frac{h(s)}{h(t)} \leq v_{m,\delta} \leq C_0 + \log \frac{h(s)}{h(m)},$$

hold on $Y \setminus V$, where $C_-(t)$ is the constant as in (2.54) in Lemma 2.8. \square

Now we shall estimate the C^2 -norm of $\{v_{m,\delta}\}_{\delta>0}$ on every compact subset of $Y \setminus V$.

Lemma 2.13 ([T0, p. 127, Lemma 2.2]) *We set*

$$(2.97) \quad f := \log \frac{\omega(s)^n}{\Omega}.$$

Let C be a positive number such that

$$(2.98) \quad C + \inf_{\alpha \neq \beta} R_{\alpha\bar{\alpha}\beta\bar{\beta}} > 1$$

holds on Y , where $R_{\alpha\bar{\alpha}\beta\bar{\beta}}$ denotes the bisectional curvature of $\omega(s)$.

Then

$$(2.99) \quad e^{Cv_{m,\delta}} \Delta_{m,\delta}(e^{-Cv_{m,\delta}}(n + \Delta_s v_{m,\delta})) \geq (n + \Delta_s v_{m,\delta}) \\ + \Delta_s \left(f + \log \frac{h(m)}{h(s)} \right) - (n + n^2 \inf_{\alpha \neq \beta} R_{\alpha\bar{\alpha}\beta\bar{\beta}}) \\ - C \cdot n(n + \Delta_s v_{m,\delta}) + (n + \Delta_s v_{m,\delta})^{\frac{n}{n-1}} \cdot \left(\frac{h(m)}{h(s)} \right)^{-\frac{1}{n-1}} \cdot \exp \left(-\frac{1}{n-1}(v_{m,\delta} + f) \right)$$

holds, where Δ_s denotes the Laplacian with respect to $\omega(s)$ (i.e., $\Delta_s = \text{trace}_{\omega(s)} \sqrt{-1} \partial \bar{\partial}$) and $\Delta_{m,\delta}$ denotes the Laplacian with respect to $\omega_{m,\delta}$. \square

We note that in Lemma 2.13, $C > 0$ does not depend on m and δ , but C depends on s (more precisely $\inf_{\alpha \neq \beta} R_{\alpha\bar{\alpha}\beta\bar{\beta}}$).

Lemma 2.14 *For any choice of $C > 0$ satisfying (2.98), m and δ , there exists a point $y_0 \in Y \setminus V$ where $e^{-Cv_{m,\delta}}(n + \Delta v_{m,\delta})$ takes its maximum. \square*

Proof. By Lemma 2.8 above (cf. (2.96))

$$v_{m,\delta} \geq \log \frac{h(s)}{h(t)} + C_-(t)$$

holds. We note that for every $t > s$, $h(s)/h(t)$ has pole of positive order along V by the condition (2.31). Hence by the lower estimate Lemma 2.8, $e^{-Cv_{m,\delta}}$ tends to 0 toward V . More precisely

$$(2.100) \quad e^{-Cv_{m,\delta}} \leq \exp(-C_-(t)) \left(\frac{h(t)}{h(s)} \right)^C$$

holds on $Y \setminus V$. On the other hand, we have that $\sqrt{-1}\partial\bar{\partial}v_{m,\delta}$ is bounded with respect to the Kähler form $\omega_{(m)}$ on Y as in [Y1, p.387, Theorem 6], hence also with respect to $\omega_{(s)}$, since both $\omega_{(m)}$ and $\omega_{(s)}$ are Kähler forms on Y by the assumption. Hence for any $C > 0$ satisfying (2.98) and every m and δ

$$(2.101) \quad e^{-Cv_{m,\delta}}(n + \Delta_s v_{m,\delta}) = O(1)$$

holds on Y and

$$(2.102) \quad \lim_{y \rightarrow V} (e^{-Cv_{m,\delta}}(n + \Delta_s v_{m,\delta}))(y) = 0$$

holds. Hence for any $C > 0$ satisfying (2.98) and every m and δ , there exists a point $y_0 \in Y \setminus V$ where $e^{-Cv_{m,\delta}}(n + \Delta_s v_{m,\delta})$ takes its maximum. This completes the proof of Lemma 2.14 \square

Here we have used the fact that the Monge-Ampère equation (2.33) of $u_{m,\delta}$ has algebraic singularities⁴. Then by Lemma 2.13, we have the following lemma.

Lemma 2.15 *If we take $C > 0$ satisfying (2.98), then there exists a positive constant C_2 independent of m and δ such that*

$$(2.103) \quad 0 \leq e^{-Cv_{m,\delta}(y_0)}(n + \Delta_s v_{m,\delta})(y_0) \leq C_2$$

holds. \square

Proof. By the maximal principle, we have

$$(2.104) \quad (n + \Delta_s v_{m,\delta}(y_0)) + \Delta_s \left(f + \log \frac{h_{(m)}}{h_{(s)}} \right) (y_0) - (n + n^2 \inf_{\alpha \neq \beta} R_{\alpha\bar{\alpha}\beta\bar{\beta}}) - C \cdot n(n + \Delta_s v_{m,\delta})(y_0) \\ + (n + \Delta_s v_{m,\delta}(y_0))^{\frac{n}{n-1}} \cdot \left(\frac{h_{(m)}}{h_{(s)}} \right)^{-\frac{1}{n-1}}(y_0) \cdot \exp \left(-\frac{1}{n-1} v_{m,\delta} + f \right) (y_0) \leq 0$$

holds. Then we see that there exists a positive constant C_3 independent of m and δ such that

$$(2.105) \quad n + \Delta_s v_{m,\delta}(y_0) \leq C_3 \left(1 + \left| \Delta_s \left(f + \log \frac{h_{(m)}}{h_{(s)}} \right) (y_0) \right| \right)^n$$

holds. Since f is C^∞ on Y and by the definition of $h_{(m)}$ (cf. (2.92)), there exists a positive constant C_4 independent of m such that

$$(2.106) \quad \left| \Delta_s \log \frac{h_{(m)}}{h_{(s)}} \right| < C_4$$

holds on Y . Hence by (2.105), we see that there exists a positive constant C_5 independent of m, δ and y_0 such that

$$(2.107) \quad n + \Delta_s v_{m,\delta}(y_0) \leq C_5$$

holds. Next we shall consider the factor $e^{-Cv_{m,\delta}(y_0)}$. We note that for every $t > s$, $h_{(s)}/h_{(t)}$ has pole of positive order along V by the condition (2.31).

⁴By the definition (2.32), $\Omega_{m,\delta}$ has algebraic singularities

Hence by the lower estimate Lemma 2.8, $e^{-Cv_{m,\delta}}$ tends to 0 toward V . Then by the C^0 -estimate Lemma 2.8, we see that there exists a positive constant C_6 independent of m and δ such that

$$(2.108) \quad e^{-Cv_{m,\delta}} \left(1 + \left| \Delta_s \left(f + \log \frac{h_{(m)}}{h_{(s)}} \right) \right| \right)^n \leq C_6$$

holds on $Y \setminus V$. Combining (2.105) and (2.108), by the definition of y_0 , we complete the proof of Lemma 2.15. \square

Let us take $C > 0$ satisfying (2.98) as in Lemma 2.13. By Lemma 2.15 and the definition of y_0

$$(2.109) \quad e^{-Cv_{m,\delta}} (n + \Delta_s v_{m,\delta}) \leq e^{-Cv_{m,\delta}(y_0)} (n + \Delta_s v_{m,\delta}(y_0)) \leq C_2$$

hold. Hence by (2.109) we have the inequality :

$$(2.110) \quad 0 \leq n + \Delta_s v_{m,\delta} \leq \exp(C \cdot v_{m,\delta}) \cdot C_2.$$

Estimating $\exp(C \cdot v_{m,\delta})$ from above by Lemma 2.12, (2.110) implies that there exists a positive constant C_7 independent of m and δ such that

$$(2.111) \quad n + \Delta_s v_{m,\delta} \leq C_7 \left(\frac{h_{(s)}}{h_{(m)}} \right)^C$$

holds on $Y \setminus V$. Applying the general theory of fully nonlinear elliptic equations ([Tr]), to the equation (2.47), we get a uniform higher order estimate of $\{v_{m,\delta}\}_{\delta>0}$ on every compact subset of $Y \setminus V$. Hence there exists a sequence $\{\delta_j\}$ with $\delta_j \downarrow 0$ as j tends to infinity such that

$$(2.112) \quad \omega_m := \lim_{j \rightarrow \infty} \omega_{m,\delta_j}$$

exists in C^∞ -topology on every compact subset of $Y \setminus V$. By using the diagonal argument, we may take $\{\delta_j\}$ independent of m . Then

$$(2.113) \quad \omega_m := \omega_{(s)} + \sqrt{-1} \partial \bar{\partial} v_m$$

satisfies the equation :

$$(2.114) \quad \log \frac{\omega_m^n}{\Omega_s} = v_m,$$

on $Y \setminus V$, where

$$(2.115) \quad \Omega_s := h_{(s)}^{-1} \cdot h_{L_{X/Y}}$$

and hence

$$(2.116) \quad -\text{Ric}_{\omega_m} + \sqrt{-1} \Theta_{h_{L_{X/Y}}} = \omega_m$$

holds.

Let h_{min} be an AZD of $K_Y + L_{X/Y}$ with minimal singularities as in Section 1.4 (cf. Definition 1.3). We set

$$(2.117) \quad \Omega_{min} := h_{min}^{-1} \cdot h_{L_{X/Y}}.$$

Then we have the following uniform C^0 -estimate for $\{v_m\}$.

Lemma 2.16 *There exists a positive constant C such that for every $m > s$*

$$(2.118) \quad v_m \leq C + \log \frac{\Omega_{min}}{\Omega_s}$$

holds on Y . \square

Proof of Lemma 2.16. Since

$$(2.119) \quad -\text{Ric}_{\omega_m} + \sqrt{-1}\Theta_{h_{L_{X/Y}}} = \omega_m$$

holds, we see that

$$(2.120) \quad \omega_m^n = O(\Omega_{min})$$

holds by Definition 1.3. Hence by the uniform upper bound (2.67) and Lemma 2.7, we see that there exists a positive constant C such that

$$(2.121) \quad v_m \leq C + \log \frac{\Omega_{min}}{\Omega_s}$$

holds for every $m > s$. \square

Let dV_Y be as in (2.84) in Lemma 2.11. We set

$$(2.122) \quad v := \log \frac{n! \cdot dV_Y}{\Omega_s}.$$

Then by Lemmas 2.13 and 2.16,

$$(2.123) \quad \omega_Y = \omega_{(s)} + \sqrt{-1}\partial\bar{\partial}v$$

satisfies

$$(2.124) \quad \omega_Y^n = e^v \cdot \Omega_s.$$

By Lemmas 2.13, 2.7 and 2.16, we see that taking a suitable subsequence $\{m_k\}$, if necessary, we may assume that

$$(2.125) \quad v = \lim_{k \rightarrow \infty} v_{m_k}$$

holds in C^∞ -topology on every compact subset of $Y \setminus V$. Then by the above construction (cf. (2.116))

$$(2.126) \quad -\text{Ric}_{\omega_Y} + \sqrt{-1}\Theta_{h_{L_{X/Y}}} = \omega_Y$$

holds on $Y \setminus V$.

Let us define the singular hermitian metric on $K_Y + L_{X/Y}$ by

$$(2.127) \quad h_K := (dV_Y)^{-1} \cdot h_{L_{X/Y}} = n! \cdot (e^v \cdot \Omega_s)^{-1} \cdot h_{L_{X/Y}}.$$

We shall check h_K is an AZD of $K_Y + L_{X/Y}$. First it is clear that $\sqrt{-1}\Theta_{h_K}$ is a closed semipositive current by (2.126) and the C^0 -estimate: Lemma 2.16. We note that all the coefficients of E_m is less than 1 by the construction. Then by the construction every global holomorphic section of $m! \cdot \pi_m^*(K_Y + L_{X/Y})$ is L^2 integrable on Y with respect to $h_{L_{X/Y}}^{m!} \cdot (\omega_m^n)^{-(m!-1)}$ by the Monge-Ampère equation of u_m and the almost boundedness of u_m . Then by the monotonicity of $\{\omega_{m,\delta}^n\}$ (Lemma 2.7), we see that $h_K = (dV_Y)^{-1} \cdot h_{L_{X/Y}}$ is an AZD of $K_Y + L_{X/Y}$. This completes the proof of Theorem 1.5 except the uniqueness of the canonical semipositive current. The uniqueness is the direct consequence of Theorem 1.8.

\square

2.5 Generalization to general adjoint line bundles

In the proof of Theorem 1.5, we have not used the property of the Hodge bundle $L_{X/Y}$ (cf. (1.22)) except the Poincaré growth property of the curvature of the Hodge metric $h_{X/Y}$ (cf. (1.23)). Hence without changing the proof, we have the following variant of Theorem 1.5.

Theorem 2.17 *Let Y be a smooth projective n -fold and let (L, h_L) be a \mathbb{Q} -line bundle with C^∞ -metric h_L with semipositive curvature. Suppose that $K_Y + L$ is big. Let U be the Zariski open subset of U defined by*

$$(2.128) \quad U := \{y \in Y \mid m!(K_Y + L) \text{ is very ample around } y \text{ for } m \gg 1\}.$$

Then there exist a closed positive current ω_Y on Y such that

- (1) *There exists a sequence of closed positive currents $\{\omega_m\}$ such that $\omega_m|_U$ is C^∞ and satisfies the equation*

$$(2.129) \quad -\text{Ric}_{\omega_m} + \sqrt{-1}\Theta_{h_L} = \omega_m$$

holds on U and

$$(2.130) \quad \omega_Y = \lim_{m \rightarrow \infty} \omega_m$$

in the sense of currents.

- (2)

$$(2.131) \quad h_{can} := \left(\frac{1}{n!} \omega_{Y,abc}^n \right)^{-1} \cdot h_L$$

is an AZD of $K_Y + L$.

Moreover if the log canonical ring

$$(2.132) \quad R(Y, a(K_Y + L)) = \bigoplus_{m=0}^{\infty} \Gamma(Y, \mathcal{O}_Y(ma(K_Y + L)))$$

is finitely generated, then ω_Y is C^∞ on the Zariski open subset U , where a is the minimal positive integer such that aL is a genuine line bundle. \square

3 Dynamical systems of Bergman kernels

Let $f : X \rightarrow Y$ be the Iitaka fibration and let $(L_{X/Y}, h_{L_{X/Y}})$ be the singular hermitian \mathbb{Q} -line bundle on Y as in Theorem 1.5. Let ω_Y be the canonical Kähler current on Y (cf. Definition 1.6). Then there exists a nonempty Zariski open subset U of Y such that ω_Y is a C^∞ on U and

$$(3.1) \quad -\text{Ric}_{\omega_Y} + \sqrt{-1}\Theta_{h_{L_{X/Y}}} = \omega_Y$$

constructed in Theorem 1.5.

Let $L_{X/Y} = M_{X/Y} + D_{X/Y}$ be the decomposition as (1.24). In this section we shall consider to describe the canonical Kähler current ω_Y in Theorem 1.5. First we shall consider the case : $D_{X/Y} = 0$. The general case : $D_{X/Y} \neq 0$ (Theorem 1.8) follows from entirely the same line as [T6, Section 4.2].

3.1 The case: $D_{X/Y} = 0$

First we shall consider the case that $D_{X/Y} = 0$. The reason why we consider this case is that in this case we do not need to use the Ricci iterations (cf. Section 1.8).

Let m_0 be the sufficiently large positive integer, M be a effective Cartier divisor such that $A := m_0!(K_Y + L_{X/Y}) - M$ is sufficiently ample as in Section 1.9. Let h_A be the C^∞ -hermitian metric on A . Hereafter we shall consider h_A as a singular hermitian metric on $m_0!(K_Y + L_{X/Y})$ by identifying h_A with

$$(3.2) \quad h_A/|\tau_M|^2,$$

where τ_M is a global holomorphic section of $\mathcal{O}_Y(M)$ with divisor M .

We shall construct a sequence of singular hermitian metrics $\{h_m\}_{m \geq m_0!}$ and a sequence of Bergman kernels $\{K_m\}$ as follows.

We set $h_{m_0!} := h_A$ and

$$(3.3) \quad K_{m_0!+1} := \begin{cases} K(Y, K_Y + m_0!(K_Y + L_{X/Y}), h_{m_0!}), & \text{if } a > 1 \\ K(Y, K_Y + L_{X/Y} + m_0!(K_Y + L_{X/Y}), h_{L_{X/Y}} \cdot h_{m_0!}), & \text{if } a = 1 \end{cases}$$

Let $\{h_m\}_{m \geq m_0!}$ be the corresponding dynamical system of singular hermitian metrics defined by

$$(3.4) \quad h_m := K_m^{-1}$$

as in Section 1.9⁵.

Theorem 3.1 *Let X be a smooth projective variety of nonnegative Kodaira dimension and let $f : X \rightarrow Y$ be the Iitaka fibration as above. Let $\{h_m\}_{m \geq m_0!}$ be the sequence of hermitian metrics as above and let n denote $\dim Y$. ω_Y is the canonical Kähler current on Y as in Theorem 1.5. Then*

$$(3.5) \quad h_\infty := \liminf_{m \rightarrow \infty} \sqrt[m]{(m!)^n \cdot h_m}$$

is a singular hermitian metric on $K_Y + L_{X/Y}$ such that

$$(3.6) \quad h_\infty = (2\pi)^n \cdot \left(\frac{1}{n!} \omega_{Y,abc}^n \right)^{-1} \cdot h_{L_{X/Y}}$$

holds almost everywhere on Y and

$$(3.7) \quad \omega_Y = \sqrt{-1} \Theta_{h_\infty}$$

holds on Y . In particular h_∞ (and hence ω_Y) is unique and is independent of the choice of A and h_A . \square

Now we shall prove Theorem 3.1. Let $dV_Y = \frac{1}{n!} \omega_{Y,abc}^n$ be the volume form associated with (Y, ω_Y) . This dV_Y is the same as the volume form defined as (2.84) by the proof of Theorem 1.5.

⁵Please do not confuse h_m in Section 2

Lemma 3.2

$$(3.8) \quad \limsup_{m \rightarrow \infty} h_{L_{X/Y}} \cdot \sqrt[n]{(m!)^{-n} K_m} \geq (2\pi)^{-n} dV_Y$$

holds on X . \square

Proof of Lemma 3.2. First we shall assume that $L_{X/Y}$ is a genuine line bundle on Y for simplicity. If $L_{X/Y}$ is not a genuine line bundle, we tensorize $(aL_{X/Y}, h_{L_{X/Y}}^a)$ in every a -steps instead of tensorize $(L_{X/Y}, h_{L_{X/Y}})$ every step, where a is the least positive integer such that $aL_{X/Y}$ is Cartier. But of course this is a minor technical difference. Hence we shall give a proof assuming that $L_{X/Y}$ is Cartier. The general case is left to readers to avoid inessential complication.

Let us consider the (singular) hermitian line bundle $(K_Y + L_{X/Y}, dV_Y^{-1} \cdot h_{L_{X/Y}})$ on Y . Let U be a nonempty Zariski open subset of Y such that $\omega_Y|_U$ is a C^∞ -Kähler form. Let $p \in U$ be a point. Then by the equation (1.25), there exists a holomorphic normal coordinate (U, z_1, \dots, z_n) of (Y, ω_Y) around p and a local holomorphic frame $\mathbf{e}_{L_{X/Y}}$ of $L_{X/Y}$ on U such that

$$(3.9) \quad dV_Y^{-1} \cdot h_{L_{X/Y}} = \left\{ \prod_{i=1}^n (1 - |z_i|^2) + O(\|z\|^3) \right\} \cdot 2^n \cdot |dz_1 \wedge \dots \wedge dz_n|^{-2} \cdot |\mathbf{e}_{L_{X/Y}}|^{-2}$$

and $h_{L_{X/Y}}(\mathbf{e}_{L_{X/Y}}, \mathbf{e}_{L_{X/Y}})(p) = 1$. Suppose that

$$(3.10) \quad C_{m-1} \cdot h_A^{-1} \cdot dV_Y^{m-m_0!-1} \cdot h_{L_{X/Y}}^{-(m-m_0!-1)} \leq K_{m-1}$$

holds on Y for some positive constant C_{m-1} . We note that

$$(3.11) \quad K_m(y) = \sup\{|\sigma|^2(y); \sigma \in H^0(Y, \mathcal{O}_Y(m(K_Y + L_{X/Y}))), (\sqrt{-1})^{n^2} \int_X h_{m-1} \cdot \sigma \wedge \bar{\sigma} = 1\}$$

holds for every $y \in Y$, by the extremal property of the Bergman kernel ⁶. We note that for the open unit disk $\Delta = \{t \in \mathbb{C} \mid |t| < 1\}$,

$$(3.12) \quad \sqrt{-1} \int_{\Delta} (1 - |t|^2)^m dt \wedge d\bar{t} = \frac{2\pi}{m+1}$$

holds. Then by Hörmander's L^2 -estimate of $\bar{\partial}$ -operators, we see that there exists a positive constant λ_m such that

$$(3.13) \quad (\lambda_m \cdot (2\pi)^{-n} \cdot m^n) \cdot C_{m-1} \cdot dV_Y^{m-m_0!} \leq h_{L_{X/Y}}^{m-m_0!} \cdot h_A \cdot K_m$$

with

$$(3.14) \quad \lambda_m \geq 1 - \frac{C}{\sqrt{m}},$$

where C is a positive constant independent of m .

In fact this can be verified as follows. Let $y \in Y \setminus (\text{Supp } M \cup V)$ and let

⁶This is well known. See for example, [Kr, p.46, Proposition 1.3.16].

(U, z_1, \dots, z_n) be the normal coordinate as above. We may assume that U is biholomorphic to the polydisk $\Delta^n(r)$ of radius r with center O in \mathbb{C}^n for some $0 < r < 1$ via (z_1, \dots, z_n) .

Taking $r < 1$ sufficiently small we may assume that there exists a C^∞ -function ρ on Y such that

- (1) ρ is identically 1 on $\Delta^n(r/3)$.
- (2) $0 \leq \rho \leq 1$.
- (3) $\text{Supp } \rho \subset \subset U$.
- (4) $\|\rho\| < 3/r$, where $\|\cdot\|$ denotes the pointwise norm with respect to ω_Y .

We note that by the equation (3.9), the mass of $\rho \cdot (dz_1 \wedge \dots \wedge dz_n)^m \otimes \mathbf{e}_{L_{X/Y}}^m$ concentrates around the origin as m tends to infinity. Hence by (3.12) we see that the L^2 -norm

$$(3.15) \quad \|\rho \cdot (dz_1 \wedge \dots \wedge dz_n)^m \otimes \mathbf{e}_{L_{X/Y}}^m\|$$

of $\rho \cdot (dz_1 \wedge \dots \wedge dz_n)^m \otimes \mathbf{e}_{L_{X/Y}}^m$ with respect to $(dV_Y)^{-m} \cdot h_{L_{X/Y}}^m$ and ω_Y is asymptotically

$$(3.16) \quad \|\rho \cdot (dz_1 \wedge \dots \wedge dz_n)^m \otimes \mathbf{e}_{L_{X/Y}}^m\|^2 \sim 2^{mn} \left(\frac{2\pi}{m}\right)^n$$

as m tends to infinity, where \sim means that the ratio of the both sides converges to 1 as m tends to infinity. We set

$$(3.17) \quad \phi := n\rho \log \sum_{i=1}^n |z_i|^2.$$

We may and do assume that m is sufficiently large so that

$$(3.18) \quad (m - m_0! - 1) \cdot \omega_Y + \sqrt{-1} \Theta_{h_A} + \sqrt{-1} \Theta_{h_{L_{X/Y}}} + \sqrt{-1} \partial \bar{\partial} \phi > 0$$

holds on Y . We note that $\bar{\partial}(\rho \cdot (dz_1 \wedge \dots \wedge dz_n)^m \otimes \mathbf{e}_{L_{X/Y}}^m)$ vanishes on the polydisc of radius $r/3$ with center p as above. Then by (3.16), the L^2 -norm

$$\|\bar{\partial}(\rho \cdot (dz_1 \wedge \dots \wedge dz_n)^m \otimes \mathbf{e}_{L_{X/Y}}^m)\|_\phi$$

of $\bar{\partial}(\rho \cdot (dz_1 \wedge \dots \wedge dz_n)^m \otimes \mathbf{e}_{L_{X/Y}}^m)$ with respect to

$e^{-\phi} \cdot h_A \cdot (dV_Y)^{-(m-m_0!-1)} \otimes h_{L_{X/Y}}^{(m-m_0!)}$ and ω_Y satisfies the inequality

$$(3.19) \quad \|\bar{\partial}(\rho \cdot (dz_1 \wedge \dots \wedge dz_n)^m \otimes \mathbf{e}_{L_{X/Y}}^m)\|_\phi^2 \leq C_0 \cdot \left(\frac{3}{r}\right)^{2n+2} \left(1 - \frac{r^2}{16}\right)^m 2^{mn} \left(\frac{2\pi}{m}\right)^n$$

for every m , where C_0 is a positive constant independent of m . By Hörmander's L^2 -estimate applied to the adjoint line bundle of the hermitian line bundle⁷:

$$(3.20) \quad ((m-1)(K_Y + L_{X/Y}) + L_{X/Y}, e^{-\phi} \cdot h_A \cdot dV_Y^{-(m-m_0!-1)} \cdot h_{L_{X/Y}}^{m-m_0!}),$$

⁷More precisely we apply the L^2 -estimate on a complete Kähler manifold $(U, \omega_Y + \epsilon \omega_U)$, where ω_U is a complete Kähler metric on U and $\epsilon > 0$ and then let $\epsilon \downarrow 0$.

we see that for every sufficiently large m , there exists a C^∞ - solution u of the equation ;

$$(3.21) \quad \bar{\partial}u = \bar{\partial}(\rho \cdot (dz_1 \wedge \cdots \wedge dz_n)^m \otimes \mathbf{e}_{L_{X/Y}}^m)$$

such that

$$(3.22) \quad u(p) = 0$$

and

$$(3.23) \quad \|u\|_\phi^2 \leq \frac{2}{m} \|\bar{\partial}(\rho \cdot (dz_1 \wedge \cdots \wedge dz_n)^m \otimes \mathbf{e}_{L_{X/Y}}^m)\|_\phi^2$$

hold, where $\|\cdot\|_\phi$'s denote the L^2 norms with respect to $e^{-\phi} \cdot h_A \cdot dV_Y^{-(m-m_0!-1)} \otimes h_{L_{X/Y}}^{m-m_0!}$ and ω_Y respectively. Then $\rho \cdot (dz_1 \wedge \cdots \wedge dz_n)^m \otimes \mathbf{e}_{L_{X/Y}}^m - u$ is a holomorphic section of $m(K_Y + L_{X/Y})$ such that

$$(3.24) \quad (\rho \cdot (dz_1 \wedge \cdots \wedge dz_n)^m \otimes \mathbf{e}_{L_{X/Y}}^m - u)(p) = \left((dz_1 \wedge \cdots \wedge dz_n)^m \otimes \mathbf{e}_{L_{X/Y}}^m \right)(p)$$

and

$$(3.25) \quad \begin{aligned} & \|\rho \cdot (dz_1 \wedge \cdots \wedge dz_n)^m \otimes \mathbf{e}_{L_{X/Y}}^m - u\|^2 \\ & \leq \left(1 + C_0 \cdot \left(\frac{3}{r} \right)^{2n+2} \cdot \sqrt{\frac{2}{m}} \cdot \left(1 - \frac{r^2}{16} \right)^m \right) \cdot 2^{mn} \cdot \left(\frac{2\pi}{m} \right)^n. \end{aligned}$$

Hence by the assumption of the induction (3.10) and the extremal property of Bergman kernels, this implies that there exists a positive constant C independent of m such that

$$(3.26) \quad K_m(p) \geq \left(1 - \frac{C}{\sqrt{m}} \right) \cdot m^n \cdot (2\pi)^{-n} \cdot C_{m-1} \cdot \left(h_A^{-1} \cdot h_{L_{X/Y}}^{-(m-m_0!)} \cdot dV_Y^{m-m_0!} \right)(p)$$

holds, since the point norm of $(dz_1 \wedge \cdots \wedge dz_n)^{\otimes m} \otimes \mathbf{e}_A$ at p (with respect to $h_A \cdot dV_Y^{-(m-m_0!)} \cdot h_{L_{X/Y}}^{(m-m_0!)}$) is asymptotically equal to 2^{mn} . Then by induction on m , using (3.11) and (3.13), we see that there exist a positive constant C' and a positive integer $m_1 > m_0!$ such that $C/\sqrt{m_1} < 1$ and for every $m > m_1$

$$(3.27) \quad K_m \geq C' \left(\prod_{k=m_1}^m \left(1 - \frac{C}{\sqrt{k}} \right) \right) \cdot (m!)^n \cdot (2\pi)^{-mn} \cdot h_A^{-1} \cdot h_{L_{X/Y}}^{-(m-m_0!)} \cdot dV_Y^{m-m_0!}$$

holds at p . Moving p , this implies that

$$(3.28) \quad \limsup_{m \rightarrow \infty} h_{L_{X/Y}} \cdot \sqrt[n]{(m!)^{-n} K_m} \geq (2\pi)^{-n} dV_Y$$

holds on Y . \square

On the other hand, we obtain the upper estimate by the following lemma.

Lemma 3.3

$$(3.29) \quad \int_Y h_{L_{X/Y}} \cdot K_m^{\frac{1}{m}} \leq \left(\prod_{k=m_0}^m (N_k + 1) \right)^{\frac{1}{m}} \cdot \left(\int_Y h_{L_{X/Y}} \cdot \sqrt[m]{K_{m_0}} \right)^{\frac{m_0}{m}}$$

holds, where $N_k := \dim |k(K_Y + L_{X/Y})| = \dim H^0(Y, \mathcal{O}_Y(k(K_Y + L_{X/Y}))) - 1$.

□

Proof. First we note that the trivial equality:

$$(3.30) \quad \int_Y h_{L_{X/Y}} \cdot \frac{K_m}{K_{m-1}} = N_m + 1$$

holds by the definition of K_m and the equality $h_{m-1} = 1/K_{m-1}$. Then by Hölder's inequality, we have

$$\begin{aligned} \int_Y h_{L_{X/Y}} \cdot K_m^{\frac{1}{m}} &= \int_Y h_{L_{X/Y}} \cdot \frac{K_m^{\frac{1}{m}}}{h_{L_{X/Y}} \cdot K_{m-1}^{\frac{1}{m-1}}} \cdot h_{L_{X/Y}} \cdot K_{m-1}^{\frac{1}{m-1}} \\ &\leq \left(\int_Y h_{L_{X/Y}}^m \cdot \frac{K_m}{h_{L_{X/Y}}^m \cdot K_{m-1}^{\frac{m}{m-1}}} \cdot (h_{L_{X/Y}} \cdot K_{m-1}^{\frac{1}{m-1}}) \right)^{\frac{1}{m}} \cdot \left(\int_Y h_{L_{X/Y}} \cdot K_{m-1}^{\frac{1}{m-1}} \right)^{\frac{m-1}{m}} \\ &= \left(\int_Y h_{L_{X/Y}} \cdot \frac{K_m}{K_{m-1}} \right)^{\frac{1}{m}} \cdot \left(\int_Y h_{L_{X/Y}} \cdot K_{m-1}^{\frac{1}{m-1}} \right)^{\frac{m-1}{m}} \\ &= (N_m + 1)^{\frac{1}{m}} \cdot \left(\int_Y h_{L_{X/Y}} \cdot K_{m-1}^{\frac{1}{m-1}} \right)^{\frac{m-1}{m}}. \end{aligned}$$

Hence we obtain the inequality:

$$(3.31) \quad \int_Y h_{L_{X/Y}} \cdot K_m^{\frac{1}{m}} \leq (N_m + 1)^{\frac{1}{m}} \cdot \left(\int_Y h_{L_{X/Y}} \cdot K_{m-1}^{\frac{1}{m-1}} \right)^{\frac{m-1}{m}}.$$

Continuing this process, by using

$$(3.32) \quad \int_Y h_{L_{X/Y}} \cdot K_{m-1}^{\frac{1}{m-1}} \leq (N_{m-1} + 1)^{\frac{1}{m-1}} \cdot \left(\int_Y h_{L_{X/Y}} \cdot K_{m-2}^{\frac{1}{m-2}} \right)^{\frac{m-2}{m-1}},$$

we have that

$$(3.33) \quad \int_Y h_{L_{X/Y}} \cdot (K_m)^{\frac{1}{m}} \leq \{(N_m + 1) \cdot (N_{m-1} + 1)\}^{\frac{1}{m}} \cdot \left(\int_Y h_{L_{X/Y}} \cdot (K_{m-2})^{\frac{1}{m-2}} \right)^{\frac{m-2}{m}}$$

holds. Continuing this process we obtain the lemma. □

To estimate the growth of $\{N_m\}_{m \geq m_0!}$, we introduce the following notion.

Definition 3.4 Let L be a line bundle on a compact complex manifold M of dimension n . We define the **volume** $\mu(M, L)$ of M with respect to L by

$$(3.34) \quad \mu(M, L) := n! \cdot \limsup_{m \rightarrow \infty} m^{-n} \dim H^0(M, \mathcal{O}_M(mL)).$$

□

We note that Definition 3.4 can be generalized to the case of \mathbb{Q} -line bundles in an obvious way. Using Lemma 3.3, we obtain the following lemma.

Lemma 3.5

$$(3.35) \quad \limsup_{m \rightarrow \infty} \frac{1}{(m!)^{\frac{n}{m}}} \int_Y h_{L_{X/Y}} \cdot (K_m)^{\frac{1}{m}} \leq \frac{\mu(Y, K_Y + L_{X/Y})}{n!}$$

holds. \square

Proof. By the definition of the volume $\mu(Y, K_Y + L_{X/Y})$,

$$(3.36) \quad N_m + 1 = \frac{\mu(Y, K_Y + L_{X/Y})}{n!} m^n + o(m^n)$$

holds. Then by Lemma 3.3, we see that

$$(3.37) \quad \limsup_{m \rightarrow \infty} \frac{1}{(m!)^{\frac{n}{m}}} \int_Y h_{L_{X/Y}} \cdot (K_m)^{\frac{1}{m}} \leq \frac{\mu(Y, K_Y + L_{X/Y})}{n!}$$

holds. \square

Lemma 3.6

$$(3.38) \quad \frac{1}{(2\pi)^n} \int_Y dV_Y = \frac{1}{n!} \int_Y \left(\frac{1}{2\pi} \omega_{Y,abc} \right)^n = \frac{1}{n!} \mu(Y, K_Y + L_{X/Y})$$

holds. \square

Proof of Lemma 3.6. Let $|P_m|$ be the free part of $|\pi_m^* m!(K_Y + L_{X/Y})|$, where π_m is the resolution of $\text{Bs}|m!(K_Y + L_{X/Y})|$ as in the last section (cf. (2.27)). By Fujita's theorem ([F, p.1, Theorem]), we see that

$$(3.39) \quad \lim_{m \rightarrow \infty} (m!)^{-n} P_m^n = \mu(Y, K_Y + L_{X/Y})$$

Then by (2.67), (2.83), Lemma 2.7 and (3.39), Lebesgue's bounded convergence theorem implies that

$$(3.40) \quad \mu(Y, K_Y + L_{X/Y}) = \lim_{m \rightarrow \infty} \int_Y \left(\frac{1}{2\pi} \omega_{m,abc} \right)^n = \int_Y \left(\frac{1}{2\pi} \omega_{Y,abc} \right)^n$$

hold. This implies the lemma. \square

We note that by Lemma 3.5 and the submeanvalue inequality for plurisubharmonic functions, $\{h_{L_{X/Y}} \cdot \sqrt[n]{(m!)^{-n} K_m}\}$ is a family of uniformly bounded semi-positive (n, n) forms on Y . Then by Lebesgue's bounded convergence theorem, we see that

$$(3.41) \quad \limsup_{m \rightarrow \infty} \int_Y h_{L_{X/Y}} \cdot \sqrt[n]{(m!)^{-n} K_m} = \int_Y \limsup_{m \rightarrow \infty} h_{L_{X/Y}} \cdot \sqrt[n]{(m!)^{-n} K_m}$$

holds. Combining Lemmas 3.2, 3.5 and 3.6, we have the equality:

$$(3.42) \quad \limsup_{m \rightarrow \infty} \frac{1}{(m!)^{\frac{n}{m}}} h_{L_{X/Y}} \cdot K_m^{\frac{1}{m}} = (2\pi)^{-n} dV_Y,$$

holds almost everywhere on Y . Hence by the definition of h_∞ (cf. (3.5))

$$(3.43) \quad h_\infty = \left(\limsup_{m \rightarrow \infty} \frac{1}{(m!)^{\frac{n}{m}}} K_m^{\frac{1}{m}} \right)^{-1} = (2\pi)^n \cdot dV_Y^{-1} \cdot h_{L_{X/Y}}$$

hold almost everywhere on Y . This implies the equality (3.6) in Theorem 1.8. Then by the equation (1.25) we have the equality (3.7) in Theorem 1.8:

$$\omega_Y = \sqrt{-1} \Theta_{h_\infty}.$$

This completes the proof of Theorem 1.8 assuming that $L_{X/Y}$ is Cartier. The proof of the general case can be obtained by entirely the same estimates. More precisely if $L_{X/Y}$ is not a genuine line bundle, we may have small ripple in the L^2 -estimates in the proof of Lemma 3.2, since we tensorize $(L_{X/Y}^{\otimes a}, h_{L_{X/Y}}^{\otimes a})$ every a steps. But the ripple disappears when we take the normalized limit as is easily be seen. This completes the proof of Theorem 3.1. \square

3.2 The case: $D_{X/Y} \neq 0$

In this section we shall give the proof of Theorem 1.8. The proof given below is essentially the same as the one of [T6, Theorem 4.8].

Let $K_Y + L_{X/Y} = P + N$ be the Zariski decomposition as (1.28). Let a be the positive integer such that $aP, aL_{X/Y}, aN \in \text{Div}(Y)$ as in Theorem 1.8. Then we have the following lemma.

Lemma 3.7 *Let h be an arbitrary AZD of $K_Y + L_{X/Y}$. Then for every positive integer m ,*

$$\mathcal{I}(h^{ma}) = \mathcal{I}(h^{ma-1} \cdot h_{L_{X/Y}})$$

holds. \square

Proof. We note that the followings holds:

- (1) $(Y, D_{X/Y})$ is KLT.
- (2) $h_{L_{X/Y}}$ induces a metric on $M_{X/Y}$ with logarithmic growth (Lemma 2.17).
- (3) $aP, aN, aL_{X/Y} \in \text{Div}(Y)$.
- (4) P is semiample ([B-C-H-M]).

Hence we see that

$$\mathcal{O}_Y(ma(K_Y + L_{X/Y})) \otimes \mathcal{I}(h^{ma}) \simeq \mathcal{O}_Y(maP)$$

and

$$\mathcal{O}_Y(ma(K_Y + L_{X/Y})) \otimes \mathcal{I}(h^{ma-1} \cdot h_{L_{X/Y}}) \simeq \mathcal{O}_Y(maP)$$

hold. This implies the lemma, \square

Let $\{h_{\ell, m}\}$ be the sequence of singular hermitian metrics constructed as Theorem 1.8. Then by Lemma 3.7, we have the following lemma.

Lemma 3.8 *If we take A to be sufficiently ample, then we have the followings.*

(1) $\mathcal{O}_Y(A + (\ell + 1)a(K_Y + L_{X/Y})) \otimes \mathcal{I}(h_{\ell,m} \cdot h_{L_{X/Y}})$ is globally generated for every $\ell, m \geq 1$.

(2) Let h_P is a C^∞ -hermitian metric on P and $h_N = |\sigma_N|^{-2}$ where σ_N is a multivalued holomorphic section on N with divisor N . Then

$$h_{\ell,m} = O(h_A \cdot (h_P \cdot h_N)^{\ell a})$$

holds for every $\ell, m \geq 1$.

□

Then the rest of the proof follows from the parallel argument as in Section 3.1. And it is entirely the same as the one of [T6, Theorem 4.8, Section 4.2]. □

3.3 Uniqueness of canonical measures

For the uniqueness of the canonical measure, we have the following uniqueness.

Corollary 3.9 $d\mu_{can}$ is birationally invariant. □

Proof of Corollary 3.9. Let $f : X \rightarrow Y$ be as above and let us consider the following commutative diagram :

$$\begin{array}{ccc} \tilde{X} & \xrightarrow{\pi} & X \\ \tilde{f} \downarrow & & \downarrow f \\ \tilde{Y} & \xrightarrow{\varpi} & Y \end{array}$$

where $\pi : \tilde{X} \rightarrow X, \varpi : \tilde{Y} \rightarrow Y$ are modifications. Let \tilde{A} and A be ample line bundles on \tilde{Y} and Y respectively.

Then ϖ^*A is nef and big on \tilde{Y} . Hence by Kodaira's lemma, we have that there exists a positive integers a_1, a_2 such that

$$(3.44) \quad a_1\tilde{A} - \varpi^*A, a_2\varpi^*A - \tilde{A}$$

are \mathbb{Q} -effective. Let $d\mu_{\tilde{X},can}, d\mu_{X,can}$ be canonical measures on \tilde{X} and X respectively. Then since $a_1\tilde{A} - \varpi^*A$ is \mathbb{Q} -effective, by Theorem 1.8 and its proof, we see that

$$(3.45) \quad d\mu_{\tilde{X},can} \geq \pi^*d\mu_{X,can}$$

holds. In fact this can be verified as follows. Let us fix C^∞ -hermitian metrics h_A and $h_{\tilde{A}}$ on A and \tilde{A} respectively. Let b be a sufficiently large positive integer and let

$$(3.46) \quad \tau \in H^0(\tilde{Y}, \mathcal{O}_{\tilde{Y}}(b(a_1\tilde{A} - \varpi^*A)))$$

be a nonzero section such that

$$(3.47) \quad (h_{\tilde{A}}^{a_1} \cdot \varpi^*h_A^{-1})^b(\tau, \tau) \leq 1.$$

Let $\{\tilde{K}_m\}_{m \geq 0}, \{K_m\}_{m \geq 0}$ be the dynamical systems of Bergman kernels as in Theorem 1.8, starting from $(ba_1\tilde{A}, h_{\tilde{A}}^{ba_1})$ and (bA, h_A^b) on \tilde{Y} and Y respectively. Then by the extremal property of Bergman kernels, we see that

$$(3.48) \quad \tilde{K}_{\ell,1} \geq |\tau|^2 \cdot \varpi^*K_{\ell,1}$$

holds for every $\ell \geq 1$. Hence by Theorem 1.8, we see that

$$(3.49) \quad d\mu_{\tilde{X},can} \geq \pi^* d\mu_{X,can}$$

holds.

Similarly since $a_2\varpi^*A - \tilde{A}$ is \mathbb{Q} -effective, we have the opposite inequality :

$$(3.50) \quad d\mu_{\tilde{X},can} \leq \pi^* d\mu_{X,can}$$

Hence we have that the equality

$$(3.51) \quad d\mu_{\tilde{X},can} = \pi^* d\mu_{X,can}$$

holds. This completes the poof of Corollary 3.9. \square

4 Relative version of Theorems 1.5 and 1.8

In this section we shall consider variation of the canonical measures on projective families. Our result is as follows.

Theorem 4.1 *Let $f : X \rightarrow S$ be a projective family such that X, S are smooth and f has connected fibers. Suppose that $f_*\mathcal{O}_S(mK_{X/S}) \neq 0$ for some $m > 0$. There exists a relative measure $d\mu_{can,X/S}$ such that the singular hermitian metric $h_{X/S} := d\mu_{can,X/S}^{-1}$ on $K_{X/S}$ satisfies:*

- (1) $\omega_{X/S} := \sqrt{-1}\Theta_{h_{X/S}}$ is semipositive on X .
- (2) For every smooth fiber $X_s := f^{-1}(s)$, $h_{X/S}|_{X_s}$ is well defined and is an AZD of K_{X_s} .
- (3) There exists a set T of measure 0 on S such that for every $s \in S \setminus T$, X_s is smooth and $\omega_{X/S}|_{X_s}$ is the canonical semipositive current on X_s constructed as in Theorems 1.5 and 1.8. \square

Remark 4.2 *Even for $s \in S \setminus T$, $d\mu_{can,X/S}|_{X_s}$ may not be precisely equal to the canonical measure $d\mu_{can,s}$ on X_s as a degenerate volume form on X_s . But as a measure $d\mu_{can,X/S}|_{X_s} = d\mu_{can,s}$ holds in exact sense. \square*

We call $d\mu_{can,X/S}$ **the relative canonical measure** of $f : X \rightarrow S$. Combining the logarithmic plurisubharmonicity of Bergman kernels ([B3, B-P] and [T4, Theorem 3.4]), this theorem strengthens the following famous result due to Y. Kawamata.

Theorem 4.3 *([Ka2, p.57, Theorem 1]) Let $f : X \rightarrow S$ be an algebraic fiber space. Suppose that $\dim S = 1$. Then for every positive integer m , $f_*\mathcal{O}_X(mK_{X/S})$ is a semipositive vector bundle on Y , in the sense that every quotient \mathcal{Q} of $f_*\mathcal{O}_X(mK_{X/S})$, $\deg \mathcal{Q} \geq 0$ holds. \square*

The main difference between Theorems 4.1 and 4.3 is that the semipositivity is on the total space in Theorem 4.1, while the semipositivity is on the direct image of the relative pluricanonical systems in Theorem 4.3. In [T6], we consider the relative log canonical bundle of a family of log canonical pairs. In the case of log canonical pairs, this difference becomes an essential one.

Proof of Theorems 4.1. Since the assertion is local, we may assume that S is the unit open polydisk in \mathbb{C}^n . Let m_0 be a sufficiently large positive integer and let

$$(4.1) \quad F_{m_0} := f_* \mathcal{O}_X(m_0! K_{X/S})$$

and (shrinking S , if necessary) let $\sigma_0, \dots, \sigma_{N(m)}$ be a set of global generators of F_{m_0} on S . We take the image Y of the rational map

$$(4.2) \quad \Phi_{m_0} : X - \dots \rightarrow \mathbb{P}_S^{N(m_0)}.$$

If we take m_0 sufficiently large, taking modifications of \hat{X} of X and \hat{Y} of Y respectively, we have the relative Iitaka fibration

$$(4.3) \quad \begin{array}{ccc} \hat{X} & \xrightarrow{g} & \hat{Y} \\ & \searrow f & \swarrow h \\ & & S \end{array}$$

such that \hat{X} and \hat{Y} are smooth and $g_* \mathcal{O}_{\hat{X}}(m! K_{\hat{X}/\hat{Y}})^{**}$ is a line bundle on \hat{Y} . We define the \mathbb{Q} -line bundle $L_{X/Y}$ on \hat{Y} by

$$(4.4) \quad L_{X/Y} = \frac{1}{m_0!} g_* \mathcal{O}_{\hat{X}}(m_0! K_{\hat{X}/\hat{Y}})^{**}$$

and let a be the least positive integer such that $\hat{f}_* \mathcal{O}_{\hat{X}}(a K_{\hat{X}/\hat{Y}}) \neq 0$. Hereafter we shall replace $f : X \rightarrow S$ by $\hat{f} : \hat{X} \rightarrow S$ and replace X and Y by \hat{X} and \hat{Y} respectively. This does not affect the proof of Theorem 4.1 by the birational invariance of canonical measures.

Let S° be the locus of S such that f is smooth over S° . Let A be a sufficiently ample line bundle on Y and let h_A be a C^∞ -metric on A with strictly positive curvature. Then as in Section 3, for every $s \in S^\circ$, we define the dynamical system of Bergman kernels $\{K_{\ell, m, s}\}_{\ell, m \geq 1}$ as in Section 3. We note that the Hodge metric $h_{L_{X/Y}}$ on $L_{X/Y}$ defined as in Section 3 has semipositive curvature in the sense of current on Y (not on every fiber Y_s) by ([Ka2], [Ka3, p.174, Theorem 1.1]). Then by the plurisubharmonicity of the Bergman kernel ([B3, B-P], [T4, Theorem 3.4]) of the adjoint line bundle of singular hermitian line bundle of semipositive curvature current, by induction on m , we see that

$$(4.5) \quad h_{\ell, m} := (K_{\ell, m}^*)^{-1}$$

extends to a singular hermitian metric on

$$(4.6) \quad A + a\ell(K_{Y/S} + L_{X/Y})$$

on Y and the extended metric has semipositive curvature in the sense of current, i.e. $\log K_{\ell, m}^*$ is plurisubharmonic on Y by Theorem 1.9. Then by Theorem 1.8,

$$(4.7) \quad K_m := \text{the upper semicontinuous envelope of } \limsup_{m \rightarrow \infty} \sqrt[m]{(m!)^{-n} K_m^*}$$

exists as a nontrivial $L_{X/Y}$ -valued relative volume form on Y and

$$(4.8) \quad h_m := K_m^{-1}$$

is a singular hermitian metric on $K_{Y/S} + L_{X/Y}$ with semipositive curvature current. We set

$$(4.9) \quad h_m := g^* h_m.$$

Then as before we may consider $h_{X/S}$ as a singular hermitian metric on $K_{X/S}$ with semipositive curvature current, i.e.,

$$(4.10) \quad \omega_{X/S} := \sqrt{-1} \Theta_{h_{X/S}}$$

is semipositive on X . By Theorem 1.8 and the birational invariance of the canonical semipositive current (Corollary 3.9), there exists a subset T of measure 0 on S such that $S \setminus T$ is contained in S° and for every $s \in S \setminus T$, $\omega_{X/S}|_{X_s}$ is the canonical semipositive current on X_s . Moreover for $s \in T \cap S^\circ$, we see that $h_{X/S}|_{X_s}$ is an AZD of K_{X_s} by the very definition of the upper-semi-continuous envelope. This completes the proof of Theorem 4.1. \square

References

- [A] Aubin, T.: Equation du type Monge-Ampère sur les variétés kähleriennes compactes, C.R. Acad. Paris **283** (1976), 459-464.
- [B1] Berndtsson, B.: Subharmonicity properties of the Bergman kernel and some other functions associated to pseudoconvex domains, math.CV/0505469 (2005).
- [B2] Berndtsson, B.: Curvature of vector bundles and subharmonicity of vector bundles, math.CV/050570 (2005).
- [B3] Berndtsson, B.: Curvature of vector bundles associated to holomorphic fibrations, math.CV/0511225 (2005).
- [B-P] Berndtsson, B. and Paun, M. : Bergman kernels and the pseudoeffectivity of relative canonical bundles, math.AG/0703344 (2007).
- [B-C-H-M] Birkar, C.-Cascini, P.-Hacon, C.-McKernan, J.: Existence of minimal models for varieties of log general type, arXiv:math/0610203.
- [Bo] Boucksom, S.: On the volume of a line bundle, Internat. J. Math. **13** (2002), 1043-1063.
- [D-P-S] Demailly, J.P.-Peternell, T.-Schneider, M. : Pseudo-effective line bundles on compact Kähler manifolds, International Jour. of Math. **12** (2001), 689-742.
- [F-M] Fujino, O. and Mori, S.: Canonical bundle formula, J. Diff. Geom. **56** (2000), 167-188.
- [F] Fujita, T.: Approximating Zariski decomposition of big line bundle, Kodai Math. J. **17** (1994), 1-4.
- [G] Griffiths, Ph.: Periods of integrals on algebraic manifolds III: Some global differential-geometric properties of the period mapping, Publ. Math., Inst. Hautes Etud. Sci. **38** (1970), 125-180.

- [Ka1] Kawamata, Y.: Characterization of Abelian Varieties, *Compos. Math.* **43** 253–276 (1981).
- [Ka2] Kawamata, Y.: Kodaira dimension of Algebraic fiber spaces over curves, *Invent. Math.* **66** (1982), 57-71.
- [Ka3] Kawamata, Y.: On effective non-vanishing and base-point-freeness. Kodaira’s issue. *Asian J. Math.* **4**(2000), no. 1, 173–181.
- [Kr] Krantz, S.: *Function theory of several complex variables*, John Wiley and Sons (1982).
- [L] Lelong, P.: *Fonctions Plurisousharmoniques et Formes Differentielles Positives*, Gordon and Breach (1968).
- [N] Nadel, A.M.: Multiplier ideal sheaves and existence of Kähler-Einstein metrics of positive scalar curvature, *Ann. of Math.* **132**(1990),549-596.
- [R] Royden, H.L.: The Ahlfors Schwarz lemma in several complex variables, *Comment. Math. Helv.* **55** (1980),547-558.
- [Sch] Schmid, W.: Variation of Hodge structure: the singularities of the period mapping. *Invent. math.* **22**, 211–319 (1973).
- [S-T] Song, J. and Tian, G. : Canonical measures and Kähler-Ricci flow, arXiv:0802.2570 (2008).
- [Su] Sugiyama, K.: Einstein-Kähler metrics on minimal varieties of general type and an inequality between Chern numbers. Recent topics in differential and analytic geometry, 417–433, *Adv. Stud. Pure Math.*, **18-I**, Academic Press, Boston, MA (1990).
- [Tr] Trudinger, N.S.: Fully nonlinear elliptic equation under natural structure conditions, *Trans. A.M.S.* **272** (1983), 751-769.
- [T0] Tsuji H.: Existence and degeneration of Kähler-Einstein metrics on minimal algebraic varieties of general type. *Math. Ann.* **281** (1988), no. 1, 123–133.
- [T1] Tsuji H.: Analytic Zariski decomposition, *Proc. of Japan Acad.* **61**(1992), 161-163.
- [T2] Tsuji, H.: Existence and Applications of Analytic Zariski Decompositions, *Trends in Math., Analysis and Geometry in Several Complex Variables*(Katata 1997), Birkhäuser Boston, Boston MA.(1999), 253-272.
- [T3] Tsuji, H.: Deformation invariance of plurigenera, *Nagoya Math. J.* **166** (2002), 117-134.
- [T4] Tsuji, H.: Dynamical construction of Kähler-Einstein metrics, *math.AG/0606023* (2006), *Nagoya Maht. J.* **199**, 107-122 (2010).
- [T5] Tsuji, H.: Canonical singular hermitian metrics on relative canonical bundles, *math.ArXiv0704.0566* (2007).

- [T6] Tsuji, H.: Ricci iterations and canonical Kähler currents on LC pairs, math.arXiv.0903.5445(2009).
- [T7] Tsuji, H.: Global generation of the direct images of pluri log canonical systems, arXiv.math.1012.0884 (2010).
- [Y1] Yau, S.-T.: On the Ricci curvature of a compact Kähler manifold and the complex Monge-Ampère equation, Comm. Pure Appl. Math. **31** (1978),339-441.
- [Y2] Yau, S.-T.: A general Schwarz lemma for Kähler manifolds, Amer. J. of Math. **100** (1978), 197-303.

Author's address

Hajime Tsuji

Department of Mathematics

Sophia University

7-1 Kioicho, Chiyoda-ku 102-8554

Japan

e-mail address: tsuji@mm.sophia.ac.jp or h-tsuji@h03.itscom.net