

# Erice Lectures on Black Holes and Attractors

Sergio Ferrara<sup>◇♣</sup>, Kuniko Hayakawa<sup>♡♣</sup> and Alessio Marrani<sup>♡♣</sup>

◇ *Physics Department, Theory Unit, CERN,  
CH 1211, Geneva 23, Switzerland  
sergio.ferrara@cern.ch*

♡ *Museo Storico della Fisica e  
Centro Studi e Ricerche “Enrico Fermi”  
Via Panisperna 89A, 00184 Roma, Italy*

♣ *INFN - Laboratori Nazionali di Frascati,  
Via Enrico Fermi 40, 00044 Frascati, Italy  
hayakawa, marrani@lnf.infn.it*

*Contribution to the Proceedings of the International School of Subnuclear Physics,  
45th Course: Search for the “Totally Unexpected” in the LHC era,  
Erice, Italy, 29 August – 7 September 2007, Directors: G. 't Hooft – A. Zichichi*

## Abstract

These lectures give an elementary introduction to the subject of four dimensional black holes (BHs) in supergravity and the Attractor Mechanism in the extremal case. Some thermodynamical properties are discussed and some relevant formulæ for the critical points of the BH effective potential are given. The case of Maxwell-Einstein-axion-dilaton (super)gravity is discussed in detail.

Analogies among BH entropy and multipartite entanglement of qubits in quantum information theory, as well moduli spaces of extremal BH attractors, are also discussed.

# Contents

1	Introduction	2
2	Attractor Mechanism and the Bekenstein-Hawking Entropy-Area formula	4
3	Black Holes and their Horizon Geometry	7
4	Thermodynamical Properties of Black Holes	13
5	Geodesic Action with a Constraint and Critical Points of the Black Hole Effective Potential	19
6	The Maxwell-Einstein-(Axion-)Dilaton Gravity	29
7	Black Hole Attractors in (Axion-)Dilaton (Super)gravity	33
8	Rotating Attractors	44
9	Black Hole Entropy and Quantum Entanglement	46
10	Recent Developments: Moduli Spaces of Attractors	51

# 1 Introduction

The aim of the present lecture notes is to give an elementary introduction to some aspects of black hole (BH) physics [1]–[6] in  $d = 4$  space-time dimensions, as well as an overview of some recent developments on the so-called *Attractor Mechanism* [7]– [9] for extremal BHs, as they appear in  $d = 4$  supergravity with a number  $\mathcal{N}$  of local supersymmetry [10]– [68] (for further developments, see also *e.g.* [69]–[72]).

Supergravity may be regarded as the low-energy limit (in a small curvature expansion) of some candidates for a quantum theory of gravity such as *superstring theory* [73]–[76] or *M-theory* [77, 78]. In situations where higher curvature effects may be neglected, the gravity part of the theory reduces to the Einstein-Hilbert action coupled to a certain number of matter fields, whose specific content depends on the particular low-energy theory. Typically, these fields are (*moduli*) scalars, spin 1/2 fermions, spin 1 gauge fields (Abelian in our examples) and spin 3/2 fermions, the gravitinos. The latter ones,  $\mathcal{N}$  in numbers, are the gauge fields of local supersymmetry.

In this situation, asymptotically flat charged BH solutions, within a static and spherically symmetric *Ansatz*, can be regarded as a generalization of the famous Schwarzschild BH. However, the presence of additional quantum numbers (such as charges and scalar hair) make their properties change drastically, and new phenomena appear. A novel important feature of electrically (and/or magnetically) charged BHs [79] as well as rotating ones [80] is a somewhat unconventional thermodynamical property named *extremality* (*i.e.* zero temperature, as we will see below) [6, 25, 81]. Extremal BHs are possibly stable gravitational objects with finite entropy but vanishing temperature, in which case the contribution to the gravitational energy entirely comes from the electromagnetic (charges) and rotational [26] (angular momentum/spin) attributes. *Extremality* also means that the inner (Cauchy) and outer (event) horizons do coincide, thus implying vanishing surface gravity (see Sects. 2 and 4).

The extremal situation entails a particular relation between the entropy, charges and spin, since in this case the gravitational mass is not an independent quantity. Four dimensional stationary and spherically symmetric BHs in an environment of scalar fields (typically described by a non-linear sigma model), have scalar hair (*scalar charges*), corresponding to the values of the scalars at (asymptotically flat) spatial infinity. These values may continuously vary, being an arbitrary point in the moduli space of the theory or, in a more geometrical language, a point in the target manifold of the scalar non-linear Lagrangian [7, 82]. Nevertheless, the BH entropy, as given by the Bekenstein-Hawking entropy-area formula [83], is also in this case independent on the scalar charges (“*no scalar hair*”) and it only depends on the asymptotic (generally *dyonic*) BH charges (see Sect. 3).

This apparent puzzle can be resolved thanks to the so-called *Attractor Mechanism* (see Sect. 2), a fascinating phenomenon that combines extremal BHs, dynamical systems, algebraic geometry and number theory [2]. It was firstly discovered in the context of supergravity; in a few words, in constructing extremal dyonic BHs of  $\mathcal{N} = 2$ ,  $d = 4$  supergravity coupled to vector and hypermultiplets (with no  $d = 4$  scalar potential), two phenomena occur: the hyperscalars can take arbitrary constant values, while the radial evolution of the vector multiplets’ scalars is described by a dynamical system [8, 9]. Under some mild assumptions, the scalar trajectory flows to a “fixed point”, located at the BH

event horizon, in the target (moduli) space. The “fixed point” (*i.e.* a point of vanishing *phase velocity*) represents the system in equilibrium, and it is the analogue of an *attractor* in the dynamical flow of dissipative systems. In approaching such an *attractor*, the orbits lose practically all memory of initial conditions (*i.e.* of the scalar hair), even though the dynamics is fully deterministic. The scalars at the BH horizon turn out to depend only on the dyonic (asymptotic) charges.

For  $\frac{1}{2}$ -BPS (*i.e.* supersymmetric),  $\mathcal{N} = 2$  attractors all the scalars are fixed [84], and one deals with the *attractor varieties*, which have their own interest in algebraic geometry and number theory [2]. For the so-called “large” BHs, within the Einstein approximation, the entropy can be shown to be proportional to a suitably defined *BH effective potential* (positive definite function in the moduli space) computed (for fixed BH electric and magnetic charges) at its critical point(s), reached at the horizon. All extremal static, spherically symmetric and asymptotically flat BHs in  $d = 4$  have a Bertotti-Robinson [85]  $AdS_2 \times S^2$  near-horizon geometry, with vanishing scalar curvature and conformally flat; in particular, the radius of  $AdS_2$  coincides with the radius of  $S^2$ , and it is proportional to the (square root of the) BH entropy (in turn proportional, through the Bekenstein-Hawking formula [83], to the area of the event horizon). Non-BPS (*i.e.* non-supersymmetric) (see *e.g.* [25, 40, 42, 56, 58, 59]) extremal BHs exist as well, and they also exhibit an attractor behavior. However, in this case not all vector multiplets’ scalars are stabilized in terms of BH conserved charges at the event horizon, but rather some of them remain, at least at the classical level, “flat” directions, as is the case for hyperscalars [42]. In spite of this, the entropy of non-BPS BHs enjoys the same property of their supersymmetric counterparts, namely it only depends on the dyonic BH charges [40, 42].

When the scalar manifold is an homogeneous symmetric space [25, 86] (as it is always the case for all  $\mathcal{N} > 2$ ,  $d = 4$  extended supergravities), the theory of extremal BH attractors has a beautiful connection to group theory and differential geometry. In this framework, the BPS or non-BPS nature of BH attractors can be related to the theory of orbits of the dyonic (asymptotic) charge vector [86, 87, 88]: different orbits correspond to different supersymmetry-preserving features of the fixed points of the scalar dynamics. All “non-flat” directions are *attractive*, which means that the Hessian matrix the BH effective potential is semi-positive definite [60, 89].

All issues mentioned above will be reviewed in Sects. 5-8. Thence, in Sect. 9 we report on some stunning relations recently found between pure states of multipartite entanglement of qubits in quantum information theory and extremal BH in superstring theory [3]. A particularly striking example of a tripartite entanglement of seven qubits, related to the (particular non-compact, real form of the) exceptional Lie group  $E_{7(7)}$ , the octonions and the Fano plane is considered in detail. The final Sect. 10 deals with some recent results on the classifications of extremal BH attractors in  $\mathcal{N} > 2$ -extended,  $d = 4$  supergravity, in which case “flat” directions of the effective BH potential in the target moduli-space appear for both  $\frac{1}{\mathcal{N}}$ -BPS and non-BPS attractors.


## 2 Attractor Mechanism and the Bekenstein-Hawking Entropy-Area formula

Extremal BHs with electric and magnetic charges and scalar hair are solitonic objects whose scalar degrees of freedom describe trajectories (in the radial evolution parameter) with *fixed points* (within the corresponding *basin of attraction*) [7, 8, 9]:

$$\begin{cases} \lim_{r \rightarrow r_H^+} \phi^a(r) = \phi_H^a(p, q) \equiv \phi_{fix}^a; \\ \lim_{r \rightarrow r_H^+} \frac{d\phi^a(r)}{dr} = 0. \end{cases} \quad (2.1)$$

The orbits lose all memory of the initial conditions (*i.e.* the asymptotic values  $\phi_\infty^a \equiv \lim_{r \rightarrow \infty} \phi^a(r)$ ), and the fixed (attractor) point  $\phi_H^a(p, q)$  only depends on the BH charges  $p, q$ .

Figure 1: **Realization of the Attractor Mechanism in the  $\frac{1}{2}$ -BPS dilatonic BH** [8, 81]. Independently on the set of asymptotical ( $r \rightarrow \infty$ ) scalar configurations, the near-horizon evolution of the dilatonic function  $e^{-2\phi}$  converges towards a fixed *attractor* value, which is purely dependent on the (ratio of the) quantized conserved charges of the BH. [8]



The trajectories are solutions of the field equations derived from a one-dimensional effective Lagrangian of almost-geodesic form, whose potential (computed at spatial infinity) is related to the Arnowitt-Deser-Misner (ADM) gravitational mass [90, 84] of the BH (see Eq. (5.31) below). The entropy, as given by the Bekenstein-Hawking entropy-area formula [83]

$$S_{BH} = \frac{A_H}{4} = \pi V_{BH}(\phi_H(p, q), p, q), \quad (2.2)$$

is the value of the *effective BH potential*  $V_{BH}$  [84] at the fixed attractor point  $\phi_H(p, q)$  at the BH event horizon. Note that  $S_{BH}$  is independent on  $\phi_\infty^a$  (*initial data* of the radial




Figure 2: **Minimization of the absolute value of the “central charge” function  $Z$  in  $\mathcal{M}$ .** In the picture  $z_{fix}^i(p, q)$  stands for the *attractor*, purely charge-dependent value of the scalars at the event horizon of the considered  $\frac{1}{2}$ -BPS extremal BH. The Attractor Mechanism fixes the extrema of the central charge to correspond to the discrete *fixed* points of the corresponding *attractor variety* [2]. Of course, the dependence of the central charge on scalars is shown for a given supporting BH charge configuration. [8]

scalar dynamics), which are continuous parameters. This is reasonable in view of the microscopic interpretation of  $S_{BH}$  as derived from a microstate counting [91] (see also [73, 53]):

$$S_{BH} \sim \ln N_{ms}, \quad (2.3)$$

$N_{ms}$  being the number of microscopic states realizing the considered macroscopic BH configuration. In general, *within the Ansatz of spherical symmetry*, the Bekenstein-Hawking entropy-area formula reads

$$S_{BH} = \frac{A_H}{4} \equiv \pi R_+^2, \quad (2.4)$$

where the *effective radius*  $R_+$  was introduced.

Notice that, in presence of scalars and/or for angular momentum  $J \neq 0$  (which is constant, in the stationary rotating regime),  $A_H$  is an *effective* quantity, *i.e.* (within the spherical symmetry *Ansatz*) it is not given by  $4\pi r_+^2$  (where  $r_+$  is the radius of the - *outer event* horizon), but rather by  $\pi R_+^2$ . For instance, in the static ( $J = 0$ ) *Maxwell-Einstein-(axion-)dilaton* BH ([92], treated in Sects. 6 and 7) it holds (both in *non-extremal* and *extremal* cases; see below, as well as Eq. (72) of the first of Refs. [92]) that

$$R_+^2 = R^2 \Big|_{r=\Sigma}^{r=r_+} = r_+^2 - \Sigma^2 \leq r_+^2, \quad (2.5)$$

where  $\Sigma$  denotes the *dilaton scalar charge*, and the (*squared*) *effective radial coordinate* [92]

$$R^2 \equiv r^2 - \Sigma^2 \leq r^2 \quad (2.6)$$

was introduced (see also Eqs. (5.19) and (5.20) further below).

Eq. (2.5) can be generalized to the *multi-dilaton system* (i.e. to  $\mathcal{N} = 2$ ,  $d = 4$  ungauged supergravity *minimally coupled* to a number  $n_V$  of Abelian vector multiplets [93]), where (for asymptotically flat, spherically symmetric, static BHs) the *squared effective radius* reads

$$R_+^2 = R^2 \Big|_{r=\sqrt{\frac{1}{2}G_{ab}(\phi_\infty)\Sigma^a\Sigma^b}}^{r=r_+} = r_+^2 - \frac{1}{2}G_{ab}(\phi_\infty)\Sigma^a\Sigma^b \leq r_+^2, \quad (2.7)$$

where  $G_{ab}$  and  $\Sigma^a$  respectively are the (non-singular) metric of the scalar manifold and the so-called *scalar charges* (see definition (4.7) below); in this case the (*squared*) *effective radial coordinate*  $R$  can be defined as

$$R^2 \equiv r^2 - \frac{1}{2}G_{ab}(\phi_\infty)\Sigma^a\Sigma^b \leq r^2, \quad (2.8)$$

which is nothing but the *many-moduli generalization* of Eq. (2.6) (see Eq. (5.21) below). In the *extremal* case ( $c = 0$ ; see Eq. (2.12) below) Eq. (2.7) yields the  $U$ -duality invariant  $\mathcal{I}_2$ , *quadratic* in the electric and magnetic BH charges, and *independent* on the scalar fields (see Sect. 7). For examples showing how  $J \neq 0$  affects the expression of the *effective radius* in the *extremal* case, see Sect. 8 and Refs. therein).

Furthermore, the temperature of the BH is given by the following formula [83]:

$$T_{BH} = \frac{\kappa}{2\pi}, \quad (2.9)$$

where

$$\kappa \equiv -\frac{1}{2}[\nabla_\mu \xi_\nu \nabla^\mu \xi^\nu]_{r=R_+}^{1/2} = \frac{r_0}{R_+^2}, \quad r_0 \equiv \frac{1}{2}(r_+ - r_-), \quad (2.10)$$

is the so-called *surface gravity*, and  $r_-$ ,  $\nabla_\mu$  and  $\xi_\nu$  respectively denote the radius of the inner (Cauchy) BH horizon, the Christoffel covariant derivatives and a Killing vector, which for any static metric simply reads (see the first of [92], and Refs. therein)

$$\xi_\nu = \delta_{\nu t} g_{tt}(x), \quad (2.11)$$

$x$  standing for the spatial coordinates. Eqs. (2.4) and (2.9) yield<sup>1</sup>

$$r_0 = 2S_{BH}T_{BH} \equiv c. \quad (2.12)$$

$c$  is the so-called *extremality parameter*. It vanishes for *extremal* BHs, which indeed have  $r_+ = r_- \equiv r_H$  (the Cauchy and event BH horizons do coincide). The *extremal* limit ( $c = 0$ ) of Eqs. (2.4), (2.5) and (2.7) respectively reads

$$S_{BH} = \frac{A_H}{4} \equiv \pi R_H^2; \quad (2.13)$$

$$R_H^2 = R^2 \Big|_{r=\Sigma}^{r=r_H} = r_H^2 - \Sigma^2 \leq r_H^2; \quad (2.14)$$

$$R_H^2 = R^2 \Big|_{r=\sqrt{\frac{1}{2}G_{ab}(\phi_\infty)\Sigma^a\Sigma^b}}^{r=r_H} = r_H^2 - \frac{1}{2}G_{ab}(\phi_\infty)\Sigma^a\Sigma^b \leq r_H^2, \quad (2.15)$$

---

<sup>1</sup>Eq. (2.12) fixes a typo in [84] and in the whole treatment of [22] and [89], where  $c^2 \equiv 2S_{BH}T_{BH}$ .

where the notation  $R_{+,c=0}^2 \equiv R_H^2$  has been introduced, and will be used throughout the following treatment.

In the case in which  $S_{BH} \neq 0$  (which, through Eqs. (2.2) and (2.4), expresses the non-vanishing of  $A_H$ , and thus the absence of a *naked singularity*), it necessarily implies  $T_{BH} = 0$ . Thus, *large* (*i.e.* with non-vanishing  $A_H$ ) extremal BHs necessarily have  $T_{BH} = 0$  and they are thermodynamically *stable*.

### 3 Black Holes and their Horizon Geometry

Extremal BHs can be described as supersymmetric solitons [94, 95] which interpolate between maximally symmetric geometries at spatial infinity and at the BH event horizon [96]. Concerning the near-horizon geometry of asymptotically flat, extremal  $p$ -(black) branes in  $d$  space-time dimensions, it holds that [96]

$$\lim_{r \rightarrow r_H^+} ds^2 = AdS_{p+2} \times S^{d-p-2} = \frac{SO(p+1, 2)}{SO(p+1, 1)} \times \frac{SO(d-p-1)}{SO(d-p-2)} \equiv ds_H^2. \quad (3.1)$$

The related isometries at the various relevant regimes are as follows:

- horizon isometry ( $r \rightarrow r_H^+$ ):  $SO(p+1, 2) \times SO(d-p-1)$ ;
- generic isometry ( $r_H < r < \infty$ ):  $ISO(p, 1) \times SO(d-p-1)$ ;
- asymptotic isometry ( $r \rightarrow \infty$ ):  $ISO(d-1, 1)$ ,

where  $ISO(n, 1)$  denotes the inhomogeneous Lorentz group (*i.e.* the Poincaré group) in  $n+1$  dimensions.

Some relevant examples are

- $p = 3, d = 10$ :  $ds_H^2 = AdS_5 \times S^5$  (Maldacena's *AdS/CFT* background), with isometry

$$SO(4, 2) \times SO(6) \sim SU(2, 2) \times SU(4) \sim Spin(4, 2) \times Spin(6); \quad (3.2)$$

- $p = 0, d = 4$ :  $ds_H^2 = AdS_2 \times S^2$  (Bertotti-Robinson (BR) BH metric), with isometry

$$SO(1, 2) \times SO(3) \sim SU(1, 1) \times SU(2) \sim Spin(1, 2) \times Spin(3); \quad (3.3)$$

- $p = 0, 1, d = 5$ :  $ds_{H,p=0}^2 = AdS_2 \times S^3$  (Tangherlini BH metric),  $ds_{H,p=1}^2 = AdS_3 \times S^2$  (*black string*), with isometry

$$p = 0 : SO(1, 2) \times SO(4) \sim SU(1, 1) \times SU(2) \times SU(2) \sim Spin(1, 2) \times Spin(4),$$

$$p = 1 : SO(2, 2) \times SO(3) \sim (SU(1, 1))^2 \times SU(2) \sim Spin(2, 2) \times Spin(3); \quad (3.4)$$



- $p = 1, d = 6$ :  $ds_H^2 = AdS_3 \times S^3$  (*self-dual dyonic black string*), with isometry  
 $SO(2, 2) \times SO(4) \sim (SU(1, 1))^2 \times SU(2) \times SU(2) \sim Spin(2, 2) \times Spin(4)$ . (3.5)

Notice that, introducing rotational degrees of freedom (in the stationary regime) affects the near-horizon geometry. For example  $p = 0, d = 4$  with a constant angular momentum  $J \neq 0$  yields an horizon geometry  $AdS_2 \times S^1$ , with isometry  $SO(1, 2) \times SO(2)$ .

Considering regular space-times with *physically reasonable* matter, *i.e.* with matter whose stress-energy tensor satisfies the so-called *dominant energy condition* [97]

$$T_{\mu\nu}^{matter} U^\mu V^\nu \geq 0 \quad (3.6)$$

for any pair of not space-like vector  $U^\mu$  and  $V^\nu$ , the long-standing conjecture that they must have positive ADM or Bondi [90, 98] mass  $M_{ADM} \equiv M$  has been proven in the 80's by using spinor techniques suggested by supergravity (see e.g. [97, 98, 99], as well as [100] and Refs. therein):

$$M \geq 0, \quad (3.7)$$

holding for asymptotically flat space-times, and with  $M_{ADM} = 0$  *only in the case of global flatness*.

The bound (3.7) implies that the Schwarzschild (Schw) BH, described by the metric<sup>2</sup>

$$ds_{Schw}^2 = - \left(1 - \frac{2M}{r}\right) dt^2 + \left(1 - \frac{2M}{r}\right)^{-1} dr^2 + r^2 d\Omega^2 \quad (3.8)$$

has no *naked singularity*, *i.e.* the singularity at  $r = 0$  is *covered* by the event horizon at  $r_H = 2M$ . Indeed, for Schwarzschild BH  $g_{tt} = 0$  at  $r = r_H$  and  $g_{tt} < 0$  inside the horizon ( $0 < r < r_H$ ). This means that the light-cone is inward and a light signal, emitted inside the horizon can never reach an outside observer. Thus, no communication is possible between the region  $r < r_H$  and the region  $r > r_H$ , so that the physical singularity at  $r = 0$  is *covered* by the BH event horizon. It is here worth observing that this phenomenon in General Relativity is a consequence of the necessity of extending the Newtonian potential to a relativistic theory of gravity. The BH interpretation is thus a necessary outcome of such an extension.

By defining the Nester antisymmetric tensor (two-form) [99]

$$E^{\alpha\sigma} \equiv 2 \left( \overline{\nabla_\beta \epsilon} \Gamma^{\sigma\alpha\beta} \epsilon - \bar{\epsilon} \Gamma^{\sigma\alpha\beta} \nabla_\beta \epsilon \right), \quad (3.9)$$

where  $\Gamma^{\sigma\alpha\beta} = g^{\beta\zeta} g^{\alpha\xi} \Gamma_{\zeta\xi}^\sigma$  is the Christoffel connection, and following [99, 97, 98] and [100], the application of the Gauss law yields (see *e.g.* [82]):

$$M = P_\Lambda u^\Lambda = -\frac{1}{2} \int_{S=\partial\Sigma} E^{\sigma\alpha} dS_{\sigma\alpha} = \int_\Sigma \nabla_\alpha E^{\alpha\sigma} d\Sigma_\sigma \Rightarrow M \geq 0. \quad (3.10)$$

---

<sup>2</sup>We use *Planck units* in which the Newton gravitational constant  $G$ , the speed of light in vacuum  $c$ , the Boltzmann constant  $K_B$  and the Planck constant  $\hbar$  are all put equal to 1. Moreover,  $M$  is shortcut for  $M_{ADM}$ , and  $d\Omega^2 = d\theta^2 + \sin^2\theta d\phi^2$  throughout.

As shown in [100] the generalization of the bound (3.10) to include BHs with electric and magnetic charges  $p$  and  $q$  is nothing but the so-called *BPS(-like) bound*

$$M \geq (q^2 + p^2)^{1/2}, \quad (3.11)$$

holding under the covariant generalization of the *dominant energy condition* [100]

$$T_{\mu\nu}^{matter} U^\mu V^\nu \geq [(J_\alpha^e V^\alpha)^2 + (J_\alpha^m V^\alpha)^2]^{1/2} \quad (3.12)$$

where  $J_\alpha^e$  and  $J_\alpha^m$  respectively are the electric and magnetic current vectors. The physical meaning of condition (3.12) is the requirement that the local charge density does not exceed the local matter density. The bound (3.11), generalization of the bound (3.7), is the so-called Bogomol'ny-Prasad-Sommerfeld (BPS) bound, and it is saturated iff a Dirac spinor field  $\epsilon$  exists satisfying the covariant constancy condition [100]

$$\hat{\nabla}_\mu \epsilon \equiv \nabla_\mu \epsilon - \frac{1}{4} F_{\alpha\beta} \gamma^\alpha \gamma^\beta \gamma_\mu \epsilon = 0, \quad (3.13)$$

thus making the supercovariant version of the Nester tensor [100]

$$\hat{E}^{\alpha\sigma} \equiv 2 \left( \overline{\hat{\nabla}_\beta \epsilon} \Gamma^{\sigma\alpha\beta} \epsilon - \bar{\epsilon} \Gamma_\beta^{\sigma\alpha\beta} \hat{\nabla} \epsilon \right) \quad (3.14)$$

vanish. Notice that  $\hat{\nabla}_\mu \epsilon$  is the gravitino supersymmetry variation; thus, the vanishing condition (3.13) implies the corresponding geometric background to preserve some supersymmetry.

Let us now move to consider the Reissner-Nordström (RN) electrically charged<sup>3</sup> BH, described by the metric

$$ds_{RN}^2 = - \left( 1 - \frac{2M}{r} + \frac{q^2}{r^2} \right) dt^2 + \left( 1 - \frac{2M}{r} + \frac{q^2}{r^2} \right)^{-1} dr^2 + r^2 d\Omega^2. \quad (3.15)$$

Such a metric exhibits two horizons, with radii

$$r_\pm = M \pm \sqrt{M^2 - q^2} = M \pm r_0. \quad (3.16)$$

In the *neutral limit*  $q = 0$  the RN metric (3.15) reduces to the Schwarzschild one, given by Eq. (3.8). On the other hand, in the same limit Eq. (3.16) yields  $r_- = 0$  (*physical singularity*) and  $r_+ = 2M$ , which is usually named *Schwarzschild BH event horizon*, and noted, with a slight abuse of notation, as

$$r_{+,Schw} = r_H = R_H. \quad (3.17)$$

The well definiteness of both the event horizon ( $r_+$ ) and Cauchy horizon ( $r_-$ ) radii requires the condition  $M^2 \geq q^2$ , which is nothing but the *BPS bound* (3.11) for  $p = 0$ , and it is the implementation of the *Cosmic Censorship Principle*.

---

<sup>3</sup>The generalization to include a magnetic charge  $p$  can be straightforwardly implemented by performing the shift  $q^2 \rightarrow q^2 + p^2$ .

In the extremal case [6]

$$c = 0 \Leftrightarrow r_+ = r_- \equiv r_H = M = R_H \Leftrightarrow M = |q|. \quad (3.18)$$

Thus, for RN BHs the *extremality is a necessary and sufficient condition for the saturation of the BPS bound (3.11)*. In other words, for RN BHs *extremality*  $\Leftrightarrow$  *supersymmetry*<sup>4</sup>. However, as it will be clear from the treatment below, this is no more the case in presence of scalars. It is here also worth pointing out that, as yielded by the *neutral limit*  $q = 0$  of Eqs. (3.18) and (3.16), the *extremality parameter* for Schwarzschild BH is nothing but its mass  $M$ :

$$c_{Schw} = M, \quad (3.19)$$

and thus *extremal* Schwarzschild BHs are necessarily “*small*”, *i.e.* with vanishing Bekenstein-Hawking entropy.

Anyway, (*at least*) for static, spherically symmetric, asymptotically flat, dyonic, *extremal* BHs (*i.e.* for the class we consider in the present lectures, excluding the treatment of Sect. 8) the BH event horizon radius  $r_H$  and the ADM mass  $M_{(ADM)}$  do coincide:

$$r_H = M. \quad (3.20)$$

When Eqs. (3.18) and (3.20) hold, by defining

$$\rho \equiv r - M = r - r_H \quad (3.21)$$

one obtains

$$1 - \frac{M}{r} = \left(1 + \frac{M}{\rho}\right)^{-1}, \quad (3.22)$$

and thus the extremal RN metric can be rewritten as follows:

$$ds_{RN,c=0}^2 = - \left(1 + \frac{M}{\rho}\right)^{-2} dt^2 + \left(1 + \frac{M}{\rho}\right)^2 (d\rho^2 + \rho^2 d\Omega^2). = - e^{2U} dt^2 + e^{-2U} d\vec{x}^2, \quad (3.23)$$

In such a coordinate system, the relevant regimes of the metric reads

- near-horizon:  $r \rightarrow r_H^+ \Leftrightarrow \rho \rightarrow 0^+$ ;
- physical singularity:  $r \rightarrow 0^+ \Leftrightarrow \rho \rightarrow -M^+$ ;
- Asymptotic regime:  $r \rightarrow \infty \Leftrightarrow \rho \rightarrow \infty$ .

Moreover,  $ds_{RN,c=0}^2$  acquires the general static Papapetrou-Majumdar [102] form

$$ds^2 = - e^{2U} dt^2 + e^{-2U} d\vec{x}^2, \quad (3.24)$$

---

<sup>4</sup>Recently, the name “*BPS*” has been associated to BHs admitting a first-order (thus *BPS-like*) formulation. In such a generalized sense, also extremal BHs not preserving any supersymmetry or certain non-extremal BHs (*e.g.* the ones with frozen scalars, or with no scalars at all, such as RN ones), can be called “*BPS*” ([101]; see also [36, 41]).

with  $U = U(\vec{x})$  satisfying the 3-d. D'Alembert equation

$$\Delta e^{-U(\vec{x})} = 0. \quad (3.25)$$

The general class of (static) Papapetrou, Majumdar metrics is known to admit a covariantly constant spinor [104]. Thus, extremal RN BH metric (3.23), for which

$$U(\vec{x}) = U(r) = -\ln\left(1 + \frac{M}{\rho}\right) = \ln\left(1 - \frac{M}{r}\right) = \ln\left(1 - \frac{|q|}{r}\right), \quad (3.26)$$

is a maximally supersymmetric background of  $\mathcal{N} = 2$ ,  $d = 4$  supergravity (see *e.g.* [103]), preserving 4 out of the 8 supersymmetries pertaining to the asymptotic  $d = 4$  Minkowski background. Thus, the extremal RN BH is a  $\frac{1}{2}$ -BPS solution of  $\mathcal{N} = 2$ ,  $d = 4$  supergravity.

As for the Schwarzschild case, also for the RN BH the physical singularity at  $r = 0$  is covered by an event horizon. Let us consider the radial geodesic dynamics of a point-like massless probe falling into the RN BH; in the reference frame of a distant observer, such a massless probe will travel from a radius  $r_0$  to a radius  $r$  (both bigger than  $r_+$ ) in a time given by the following formula [6]:

$$\Delta t(r) = \int_{r_0}^r \frac{dt}{dr} dr = \int_{r_0}^r \sqrt{\frac{g_{rr}}{g_{tt}}} dr \rightarrow \infty \text{ for } r \rightarrow r_+. \quad (3.27)$$

In order to determine the near-horizon geometry of an extremal RN BH, let us define a new radial coordinate as  $\tau = -\frac{1}{\rho} = \frac{1}{r_H - r}$ . Thus, the relevant regimes of the metric now reads

- near-horizon:  $r \rightarrow r_H^+ \Leftrightarrow \tau \rightarrow -\infty$ ;
- physical singularity:  $r \rightarrow 0^+ \Leftrightarrow \tau \rightarrow \frac{1}{M}^+$ ;
- asymptotic regime:  $r \rightarrow \infty \Leftrightarrow \tau \rightarrow 0^-$ .

In the near-horizon limit  $\rho \rightarrow 0^+$ , and thus  $1 + \frac{M}{\rho} \sim \frac{M}{\rho}$ ; consequently, from Eq. (3.23) one obtains

$$\lim_{\rho \rightarrow 0^+} ds_{RN, c=0}^2 = -\frac{1}{M^2 \tau^2} dt^2 + \frac{M^2}{\tau^2} (d\tau^2 + \tau^2 d\Omega^2). \quad (3.28)$$

By further rescaling  $\tau \rightarrow \frac{\tau}{M^2}$ , one finally gets

$$\lim_{\rho \rightarrow 0^+} ds_{RN, c=0}^2 = \frac{M^2}{\tau^2} (-dt^2 + d\tau^2 + \tau^2 d\Omega^2), \quad (3.29)$$

which is nothing but the  $AdS_2 \times S_2$  BR metric [85], both flat and conformally flat. Such a result is consistent with the fact that an extremal RN BH is nothing but an extremal 0-brane in  $d = 4$ .

It is also worth observing that, by introducing the physical distance coordinate  $\omega \equiv \ln \rho$ , the metric (3.28) can be rewritten as

$$\lim_{\omega \rightarrow -\infty} ds_{RN,c=0}^2 = -\frac{1}{M^2} e^{2\omega} dt^2 + M^2 d\omega^2 + M^2 d\Omega^2. \quad (3.30)$$

Thus, the physical distance from any finite  $\omega_0$  to the BH event horizon is *infinite*, because

$$\rho \rightarrow 0^+ \Leftrightarrow \omega \rightarrow -\infty. \quad (3.31)$$

As we will point out further below, this crucially discriminates the extremal case from the non-extremal one.

The extremal RN single-center metric function (3.26) can be generalized to the multi-center case as follows:

$$e^{-U(r)} = 1 + \sum_{s=1}^n \frac{M_s}{|\vec{x} - \vec{x}_s|}, \quad (3.32)$$

and the corresponding solution (3.24) can be interpreted as many non-rotating RN extremal BHs, with centers (*i.e.* physical singularities) at  $\vec{x}_s$  and masses  $M_s = (q_s^2 + p_s^2)^{1/2}$ , saturating the BPS bound (3.11). The saturation of the BPS bound allows for the gravitational attraction to equal the electro-magnetic repulsion, determining a static neutral equilibrium. Such a phenomenon is sometimes called *antigravity*. In particular, the additive nature of the solution is related to the BPS nature of such a force-free environments of (single-center) BHs.

As pointed out above, for RN BHs (in which no scalars are present) extremality is equivalent to the saturation of the BPS bound, which also implies that a Killing spinor exists [104]. As mentioned above, the near-horizon BR geometry (3.29) is conformally flat, satisfying [6]

$$\begin{aligned} \mathcal{R} &= 0 \quad (\text{vanishing scalar curvature}); \\ C_{\mu\nu\rho\sigma} &= 0 \quad (\text{vanishing Weyl tensor}); \\ D_\mu F_{\rho\sigma} &= 0 \quad (\text{covariantly constant Abelian vector field strength}), \end{aligned} \quad (3.33)$$

implying that a *doubling* of supersymmetry occurs at the event horizon of the extremal RN BH (see *e.g.* Kallosh's paper in [6], and [92]). Notice that the *doubling* also occurs at  $\rho \rightarrow \infty$ : indeed, within the asymptotic flatness *Ansatz* the asymptotic spatial limit yields the  $\mathcal{N} = 2$ ,  $d = 4$  super-Poincaré superalgebra, pertaining to the  $d = 4$  Minkowski space.

Thus, it can be stated that the *extremal* RN BH is nothing but an *extremal 0-black brane solitonic solution* of  $\mathcal{N} = 2$ ,  $d = 4$  (ungauged) supergravity, interpolating between two *maximally supersymmetric* space-time geometries [96]:

$$c = 0 : \left\{ \begin{array}{l} \text{at event horizon } (r \rightarrow r_H^+): \text{ algebra } \mathfrak{psu}(1,1|2), \text{ BR } AdS_2 \times S^2 \text{ metric;} \\ \text{at spatial infinity } (r \rightarrow \infty): \text{ algebra } \mathcal{N} = 2, d = 4 \text{ super-Poincaré, Minkowski metric.} \end{array} \right. \quad (3.34)$$

Beside  $\mathcal{N} = 2, d = 4$  super-Poincaré algebra,  $\mathfrak{psu}(1, 1|2)$  is a maximal  $\mathcal{N} = 2, d = 4$  superalgebra, which does not contain Poincaré nor other semisimple Lie algebra, but direct sum of simple Lie algebra as maximal bosonic subalgebra. Indeed, in this case the maximal bosonic subalgebra is  $\mathfrak{so}(1, 2) \oplus \mathfrak{so}(3)$  (with related maximal spin bosonic subalgebra  $\mathfrak{su}(1, 1) \oplus \mathfrak{su}(2)$ ), matching the corresponding bosonic isometry group of the BR metric (see above).

## 4 Thermodynamical Properties of Black Holes

BHs have an entropy  $S_{BH}$  and a temperature  $T_{BH}$ ; in absence of other attributes, they obey the first law of thermodynamics

$$dM = T_{BH}dS_{BH} + \dots, \quad (4.1)$$

where  $M$  is the gravitational (ADM) mass. By recalling Eqs. (2.2), (2.4), (2.9) and (2.10), the temperature and entropy are related to geometric quantities of the BH background, namely to the surface gravity  $\kappa$  and the to the horizon area  $A_H$  [6].

Thus, for a Schwarzschild BH, having  $r_H = R_H = 2M$  (see Eqs. (3.17) and (3.19)), it holds that ( $r_0 = \frac{r_H}{2} = M$ ) [80]

$$\begin{aligned} dM &= \frac{1}{8\pi}\kappa dA_H \implies M = \frac{1}{4\pi}\kappa A_H; \\ A_H &= 4\pi r_H^2 = 16\pi M^2; \\ S_{BH} &= 4\pi M^2; \\ T_{BH} &= \frac{1}{8\pi M}. \end{aligned} \quad (4.2)$$

Next, one might also consider (stationary) BHs with electric charge  $q$  (RN, mentioned above), and with (constant) angular momentum  $J$ , namely the Kerr-Newman BHs, whose *neutral limit* ( $q = 0$ ) is the Kerr BH (see also the treatment of Sect. 8; for recent advances, see *e.g.* [105, 106] and Refs. therein).

For such cases, the *first law of BH thermodynamics* gets modified as [80][107]

$$dM = T_{BH}dS_{BH} + \Omega dJ + \psi dq, \quad (4.3)$$

where  $\Omega$  is the (constant, within the assumed stationary *Ansatz*) angular velocity and  $\psi$  is the electric potential (at the BH event horizon) for the considered charged (possibly rotating) BH. The generalization to include a magnetic charge  $p$  is straightforward:

$$dM = T_{BH}dS_{BH} + \Omega dJ + \psi dq + \chi dp, \quad (4.4)$$

where  $\chi$  is the magnetic potential (at the BH event horizon).

At a finite level,  $M$ ,  $S_{BH}$ ,  $q$ ,  $p$  and  $J$  are related, in the (semi)classical Einstein approximation, by the *Smarr-integrated* form of the (generalized) first law of the thermodynamics given by Eq. (4.4), *i.e.* by [80]

$$M = 2T_{BH}S_{BH} + 2\Omega J + \psi q + \chi p, \quad (4.5)$$

which, by recalling the definition (2.12), can be rewritten as

$$M = c + 2\Omega J + \psi q + \chi p. \quad (4.6)$$

As a further generalization<sup>5</sup>, one can couple a set of (real) scalars  $\phi^a$  ( $a = 1, \dots, n$ ), parameterizing a scalar manifold  $\mathcal{M}_\phi$  with non-singular metric  $G_{ab}$ . The scalars are associated to the so-called *scalar charges*  $\Sigma^a$ , defined through an expansion at  $r \rightarrow \infty$  as [82]

$$\lim_{r \rightarrow \infty} \phi^a(r) \equiv \phi_\infty^a + \frac{1}{r} \Sigma^a + \mathcal{O}\left(\frac{1}{r^2}\right), \quad \Sigma^a = \Sigma^a(p^\Lambda, q_\Lambda, A_H, J, \phi_\infty^b). \quad (4.7)$$

Thus,  $\Sigma^a$  can be equivalently defined as follows [84] (see also [22]):

$$\Sigma^a(p^\Lambda, q_\Lambda, A_H, J, \phi_\infty^b) \equiv \lim_{\tau \rightarrow 0^-} \frac{d\phi^a(\tau)}{d\tau}. \quad (4.8)$$

The BH charges are defined as the fluxes at spatial infinity of the Abelian field strengths:

$$p^\Lambda \equiv \frac{1}{4\pi} \int_{S_\infty^2} \mathcal{F}^\Lambda, \quad q_\Lambda \equiv \frac{1}{4\pi} \int_{S_\infty^2} \mathcal{G}_\Lambda, \quad \Lambda = 1, 2, \dots, n+1, \quad (4.9)$$

where  $\mathcal{F}^\Lambda = dA^\Lambda$  and  $\mathcal{G}_\Lambda$  is the “dual” field-strength two-form [108].

Thus, Eq. (4.3) becomes [82]

$$\begin{aligned} dM &= T_{BH}dS_{BH} + \Omega dJ + \psi^\Lambda dq_\Lambda + \chi_\Lambda dp^\Lambda + \left. \frac{\partial M}{\partial \phi^a} \right|_{S_{BH}, J, q_\Lambda, p^\Lambda} d\phi^a = \\ &= T_{BH}dS_{BH} + \Omega dJ + \psi^\Lambda dq_\Lambda + \chi_\Lambda dp^\Lambda - G_{ab}(\phi_\infty) \Sigma^b d\phi^a. \end{aligned} \quad (4.10)$$

It is worth remarking that, despite the presence of the extra term  $-G_{ab}(\phi_\infty) \Sigma^b d\phi^a$  in the generalized first law of BH thermodynamics (4.10), the integrated version, *i.e.* the corresponding Smarr formula, remains (see P. Breitenlohner, Maison, and Gibbons’ paper in [6]; see also [82])

$$M = 2T_{BH}S_{BH} + 2\Omega J + \psi^\Lambda q_\Lambda + \chi_\Lambda p^\Lambda = c + 2\Omega J + \psi^\Lambda q_\Lambda + \chi_\Lambda p^\Lambda. \quad (4.11)$$

The thermodynamical quantities  $S_{BH}$  (or, equivalently by the Bekenstein-Hawking relation (2.2) or (2.4),  $A_H$ ),  $J$ ,  $q_\Lambda$ ,  $p^\Lambda$ ,  $\phi_\infty^a$  are coordinates on the state space (in the semi-classical limit of large BH charges)

$$\mathbb{R}_0^+ \times \mathbb{R}^+ \times \mathbb{R}^{2n+2} \times \mathcal{M}_{\phi_\infty}, \quad (4.12)$$

where the first factor  $\mathbb{R}_0^+$  disregards the so-called *small* BHs, which in the classical (Einstein) approximation represent *naked singularities* in the space-time, and for which the Attractor Mechanism does not hold. Furthermore,  $\mathcal{M}_{\phi_\infty}$  is the asymptotic ( $r \rightarrow \infty$ ) scalar manifold.

---

<sup>5</sup>Interesting considerations can be found in [32].

It is worth remarking that, due to the key result [82]

$$\left. \frac{\partial M}{\partial \phi^a} \right|_{S_{BH}, J, q_\Lambda, p^\Lambda} = -G_{ab}(\phi_\infty) \Sigma^b (S_{BH}, J, p^\Lambda, q_\Lambda, \phi_\infty^b), \quad (4.13)$$

for non-singular metric of the scalar manifold, the scalar charges  $\Sigma^a$  vanish iff  $\phi_\infty$ , and hence the vacuum state, is chosen to extremize the ADM mass  $M$  at fixed BH entropy  $S_{BH} = A_H/4$ , angular momentum  $J$ , and conserved electric and magnetic charges  $q_\Lambda$  and  $p^\Lambda$ .

Let us now consider the static case ( $J = 0$ ), described by the bosonic Lagrangian density [84]

$$-\frac{\mathcal{L}}{\sqrt{-g}} = \frac{1}{2} \mathcal{R} - \frac{1}{2} G_{ab} \partial_\mu \phi^a \partial_\nu \phi^b g^{\mu\nu} - (\text{Im} \mathcal{N}_{\Lambda\Sigma}) F^\Lambda F^\Sigma - (\text{Re} \mathcal{N}_{\Lambda\Sigma}) F^\Lambda \tilde{F}^\Sigma. \quad (4.14)$$

Under the convexity condition<sup>6</sup>

$$\nabla_a \partial_b V_{BH} \geq 0, \quad (4.15)$$

requiring the (semi-)positive definiteness of the covariant Hessian of the so-called *BH effective potential* [84]

$$\begin{aligned} V_{BH}(\phi, p, q) &\equiv -\frac{1}{2} \Gamma^T M \Gamma; \\ \Gamma^T &\equiv (p^\Lambda, q_\Lambda); \\ M &\equiv \begin{pmatrix} \text{Im} \mathcal{N} + \text{Re} \mathcal{N} (\text{Im} \mathcal{N})^{-1} \text{Re} \mathcal{N} & -\text{Re} \mathcal{N} (\text{Im} \mathcal{N})^{-1} \\ -(\text{Im} \mathcal{N})^{-1} \text{Re} \mathcal{N} & (\text{Im} \mathcal{N})^{-1} \end{pmatrix}, \end{aligned} \quad (4.16)$$

in [82] it was proved that

$$\Sigma^a = 0 \Leftrightarrow \left. \frac{\partial M}{\partial \phi^a} \right|_{S_{BH}, J, q_\Lambda, p^\Lambda} = 0 \Leftrightarrow \begin{cases} \frac{d\phi^a(\tau)}{d\tau} = 0 \quad \forall \tau \in \mathbb{R}^- \Leftrightarrow \phi^a \neq \phi^a(\tau); \\ \phi^a = \phi_\infty^a = \phi_H(p, q) : \frac{\partial V_{BH}(\phi, p, q)}{\partial \phi^a} = 0. \end{cases} \quad (4.17)$$

In words, the scalar charges vanish iff the extremal BH is actually double-extremal, *i.e.* if the scalars are *constant* (*independent* on  $\tau$ ); moreover, their constant values are not arbitrary, but they are chosen to extremize  $V_{BH}(\phi, p, q)$  (at fixed, supporting BH charge configuration). Thus (at fixed, supporting BH charge configuration  $p^\Lambda, q_\Lambda$ , corresponding to a fixed value of the BH entropy  $S_{BH}(p, q)$ ), under the condition (4.15) it holds that

$$\Sigma^a = 0 \Leftrightarrow \phi_\infty^a = \phi_H(p, q), \quad \forall a = 1, \dots, n, \quad \forall \tau \in \mathbb{R}^-, \quad (4.18)$$

with  $\phi_H(p, q)$  fulfilling the *criticality conditions* ( $\partial_a \equiv \frac{\partial}{\partial \phi^a}$ )

$$\partial_a V_{BH}(\phi, p, q)|_{\phi=\phi_H(p, q)} = 0. \quad (4.19)$$

---

<sup>6</sup> $\nabla_a$  is the Christoffel covariant derivative, which acquires also a Kähler connection component when the scalars parameterize a (special) Kähler manifold, such as in  $\mathcal{N} = 2, d = 4$  supergravity coupled to Abelian vector supermultiplets (see *e.g.* [108] and Refs. therein).



When evaluated at the critical point of  $V_{BH}$  given by purely charge-dependent scalar configuration  $\phi_H(p, q)$ , condition (4.15) is nothing but the requirement that the  $\phi_H(p, q)$  actually corresponds to a *stable* (up to some *massless Hessian modes*) critical point of  $V_{BH}$ , and thus that it gives rise to an *attractor* in a strict sense.

## First Order Formalism

However, one can go a step further beyond the proof of [82] (which however, beside implying some loss of generality, holds also in the non extremal case), and, in the extremal case, relate directly the scalar charges to the critical points of a suitably generalized real, positive superpotential  $\mathcal{W}$  ([41]; see below, Eqs. (5.42) and (5)).

Indeed, in [36, 41] a first order formalism for static, spherically symmetric and asymptotically flat extremal BHs was introduced, based on the *generalized (fake) superpotential*  $\mathcal{W}$ . For  $c = 0$  the second order ordinary differential equation satisfied by the scalars  $\phi^a$ , *i.e.* [84]

$$\frac{\nabla^2 \phi^a(\tau)}{\nabla \tau^2} = \frac{d^2 \phi^a(\tau)}{d\tau^2} + \Gamma_{bc}{}^a(\phi(\tau)) \frac{d\phi^b(\tau)}{d\tau} \frac{d\phi^c(\tau)}{d\tau} = G^{ab}(\phi(\tau)) \frac{\partial V_{BH}(\phi, p, q)}{\partial \phi^b} e^{2U(\tau)} \quad (4.20)$$

have been shown to reduce to the following first order one<sup>7</sup> [41]:

$$\frac{d\phi^a(\tau)}{d\tau} = 2e^{U(\tau)} G^{ab}(\phi(\tau)) \frac{\partial \mathcal{W}(\phi, p, q)}{\partial \phi^b}. \quad (4.21)$$

The relation between  $V_{BH}$  and  $\mathcal{W}$  is given by [36, 41]

$$V_{BH} = \mathcal{W}^2 + 2G^{ab}(\partial_a \mathcal{W}) \partial_b \mathcal{W}, \quad (4.22)$$

yielding that

$$\partial_a V_{BH} = 2 [\delta_a^b \mathcal{W} + 2G^{bc}(\nabla_a \partial_c \mathcal{W})] \partial_b \mathcal{W} = 2 [\delta_a^b \mathcal{W} + 2G^{bc}(\partial_a \partial_c \mathcal{W} - \Gamma_{ac}{}^f \partial_f \mathcal{W})] \partial_b \mathcal{W}. \quad (4.23)$$

Eq. (4.21) is *BPS-like*; for a given  $\mathcal{W}$ , it relates the evolution of the scalar fields to the partial derivative of  $\mathcal{W}$  itself with respect to the scalar fields. For a fixed, supporting BH charge configuration, the extrema of  $\mathcal{W}$  in  $\mathcal{M}_\phi$  are fixed points in the radial evolution of  $\phi^a$ . Furthermore, Eq. (4.23) implies that for a fixed supporting BH charge configuration, the extrema of  $\mathcal{W}$  in  $\mathcal{M}_\phi$  are also extrema of  $V_{BH}$  in  $\mathcal{M}_\phi$ .

Now, by recalling the definition (4.8) and performing the asymptotic limit of the first order differential Eq. (4.21), one obtains the general expression of the *contravariant* scalar charges in the extremal environment under consideration<sup>8</sup>:

$$\Sigma^a(\phi_\infty, p, q) = 2 \lim_{\tau \rightarrow 0^-} e^{U(\tau)} G^{ab}(\phi(\tau)) \partial_b \mathcal{W}(\phi(\tau), p, q) = 2G^{ab}(\phi_\infty) \frac{\partial \mathcal{W}(\phi_\infty, p, q)}{\partial \phi_\infty^b}, \quad (4.24)$$

---

<sup>7</sup>Some subtleties (however immaterial for  $c = 0$ ) are given in footnote 3 of [41].

<sup>8</sup>We assume

$$(\partial_a \mathcal{W}(\phi, p, q))_\infty \equiv \lim_{\tau \rightarrow 0^-} \frac{\partial \mathcal{W}(\phi(\tau), p, q)}{\partial \phi^a(\tau)} = \frac{\partial \mathcal{W}(\phi_\infty, p, q)}{\partial \phi_\infty^a}.$$

where minimal regularity conditions in order to split the asymptotical limit have been understood, and in the last step the asymptotic boundary condition for  $U(\tau)$  have been used. Similarly, *covariant* scalar charges read

$$\Sigma_a(\phi_\infty, p, q) = 2 \lim_{\tau \rightarrow 0^-} e^{U(\tau)} \partial_a \mathcal{W}(\phi(\tau), p, q) = 2 \frac{\partial \mathcal{W}(\phi_\infty, p, q)}{\partial \phi_\infty^a}, \quad (4.25)$$

Eqs. (4.24) and (4.25), holding true for extremal, static, spherically symmetric and asymptotically flat BHs (without any assumption on the Hessian matrix of  $V_{BH}$ ), relates the (vanishing of the) scalar charges  $\Sigma^a$  to (the critical points of) the generalized superpotential  $\mathcal{W}$ .

Now, since the explicit forms of  $\mathcal{W}$  and of its critical points in  $\mathcal{M}_{\phi(\infty)}$  have been explicitly determined for some  $\mathcal{N} > 2$ -extended,  $d = 4$  supergravities [41] and for some examples of  $\mathcal{N} = 2$ ,  $d = 4$  supergravity<sup>9</sup>, the explicit forms of  $\Sigma^a$  and of the geometrical *loci* of their zeros in  $\mathcal{M}_{\phi(\infty)}$  (for a given, supporting BH charge configuration) for all such cases are currently known. It is worth noticing that  $\mathcal{W}$  varies depending on the class of extremal BH attractors under consideration (and it is not unique inside the same class, too), thus the expression of  $\Sigma^a$  will be dependent on the considered class of attractor flows.

It is then clear from the reasoning above that the scalar charges  $\Sigma^a$  can vanish also in *non-double-extremal* BHs (*i.e.* in BHs with non-trivial scalar dynamics), when a fine-tuning of the asymptotical values  $\phi_\infty^a$  of the scalars is performed such that  $\phi_\infty^a = \phi_H^a(p, q)$ , with  $\phi_H^a(p, q)$  satisfying the *criticality conditions* of  $\mathcal{W}$  (and thus, through Eq. (4.23), of  $V_{BH}$ ) in  $\mathcal{M}_{\phi(\infty)}$  (for a given, supporting BH charge configuration). In other words, in order to make the  $\Sigma^a$ 's vanish, the *initial data* of the radial scalar dynamics must be fine-tuned to coincide with the near-horizon, *attracted* scalar configurations  $\phi_H^a(p, q)$  (at least within the same *basin of attraction*).

As anticipated above, an interesting feature of such an environment with scalars and/or electric and magnetic charges concerns extremality:  $M$  and  $S_{BH}$  can be finite and non-vanishing, with  $T_{BH} = 0$ . In other words, extremal *large* BHs with non-vanishing ADM mass and entropy can exist!

The simplest example is provided by the RN BH treated above, in which  $n = 0$ . Let us now focus on its thermodynamical properties; by recalling Eqs. (2.4), (2.9) and (2.10), and putting  $\Sigma = 0$ , one obtains that

$$T_{BH} = \frac{\kappa}{2\pi} = \frac{1}{2\pi} \frac{r_0}{R_+^2} = \frac{c}{2S_{BH}}, \quad (4.26)$$

---

<sup>9</sup>Namely, the  $\mathcal{N} = 2$ ,  $d = 4$  models treated so far are:

- 1) the so-called  $t^3$  model, in the *electric* BH charge configuration  $(p^0, p^1 = 0, q_0 = 0, q_1)$  [36];
- 2) the so-called *stu* model, in the *electric* BH charge configuration  $(p^0, p^i = 0, q_0 = 0, q_i)$  ( $i = 1, 2, 3$ ) [36];
- 3) the models based on the sequence  $\frac{SU(1, n_V)}{U(1) \times SU(n_V)}$  of irreducible homogeneous symmetric special Kähler manifolds with quadratic prepotential [41] (for  $n_V = 1$ , see also [36]);
- 4) the model related to the degree three Jordan algebra on the quaternions  $J_3^{\mathbb{H}}$  (dual to the  $\mathcal{N} = 6$ ,  $d = 4$  case) [41].

where (recalling Eqs. (2.12), (2.4) and (3.16) and perform the shift  $q^2 \rightarrow q^2 + p^2$ )

$$\begin{aligned}
c &\equiv r_0 = \sqrt{M^2 - q^2 - p^2}; \\
S_{BH} &= \pi(2M^2 - q^2 - p^2 + 2M\sqrt{M^2 - p^2 - q^2}); \\
&\quad \Downarrow \\
r_+^2 = R_+^2 &= 2M^2 - q^2 - p^2 + 2M\sqrt{M^2 - p^2 - q^2}.
\end{aligned} \tag{4.27}$$

Thus, as stated above, for RN BHs the *extremality* (*i.e.*  $c = 0$ ) is a *necessary and sufficient* condition for the *saturation* of the BPS bound (3.11), and thus for the (*maximally*, in the present case) *supersymmetric nature* of the considered background. Indeed, Eqs. (4.27) yield

$$c = 0 \Leftrightarrow R_H = r_H = M = (q^2 + p^2)^{1/2} \implies S_{BH}|_{c=0} = \pi(q^2 + p^2). \tag{4.28}$$

Consequently, *large* extremal RN BHs are ( $\frac{1}{2}$ -)BPS, and they have  $T_{BH} = 0$ . As given by Eq. (4.28), the *small BH limit* is reached in the degenerate limit  $p \rightarrow 0$  and  $q \rightarrow 0$ .

By further introducing (constant) angular momentum  $J$ , one may consider the stationary, rotating uncharged (Kerr) or charged (Kerr-Newman) BHs. At a finite level,  $M$ ,  $S_{BH}$ ,  $q$ ,  $p$  and  $J$  are related, in the (semi)classical Einstein approximation, by the *Smarr-integrated* form of the (generalized) first law of the thermodynamics given by Eq. (4.4), *i.e.* by Eqs. (4.5) and (4.6). Eqs. (3.16) and (2.4) are generalized as

$$\begin{aligned}
r_{\pm} &= M \pm \sqrt{M^2 - q^2 - p^2 - \frac{J^2}{M^2}}; \\
R_+^2 &= r_+^2 + \frac{J^2}{M^2} \geq r_+^2; \\
S_{BH} &= \frac{A_H}{4} = \pi R_+^2 = \pi \left( r_+^2 + \frac{J^2}{M^2} \right),
\end{aligned} \tag{4.29}$$

and thus

$$\begin{aligned}
S_{BH} &= \pi \left[ 2M^2 - q^2 - p^2 + 2M \left( M^2 - q^2 - p^2 - \frac{J^2}{M^2} \right)^{1/2} \right]; \\
&\quad \Downarrow \\
R_+^2 &= 2M^2 - q^2 - p^2 + 2M \left( M^2 - q^2 - p^2 - \frac{J^2}{M^2} \right)^{1/2},
\end{aligned} \tag{4.30}$$

whose inversion reads

$$\begin{aligned}
M^2 &= \frac{\pi J^2}{S_{BH}} + \frac{\pi (q^2 + p^2)^2}{4S_{BH}} + \frac{S_{BH}}{4\pi} + \frac{(q^2 + p^2)}{2} = \\
&= \left( \frac{J}{R_+} \right)^2 + \left( \frac{q^2 + p^2}{2R_+} \right)^2 + \left( \frac{R_+}{2} \right)^2 + \frac{(q^2 + p^2)}{2}.
\end{aligned} \tag{4.31}$$

Notice that for (constant, in the considered stationary rotating regime) angular momentum  $J \neq 0$ ,  $A_H$  is an *effective* quantity, *i.e.* (within the spherical symmetry *Ansatz*)

it is not given by  $\pi r_+^2$ , but rather by  $\pi R_+^2$ . In particular, the second of Eqs. (4.29) and Eq. (4.30) yields that  $R_+ \geq r_+$ . Thus, the (stationary) rotating regime actually *increases* the BH radius relevant for the computation of the entropy through the Bekenstein-Hawking (semi)classical formula (2.4).

Other useful relations among the various geometric and thermodynamical quantities read [80]

$$T_{BH} = \frac{1}{32M} \left[ \frac{1}{\pi} - \frac{4\pi J^2}{S_{BH}^2} - \frac{\pi(q^2 + p^2)^2}{S_{BH}^2} \right] = \frac{1}{32\pi M} \left[ 1 - \left( \frac{2J}{R_+} \right)^2 - \left( \frac{q^2 + p^2}{R_+^2} \right)^2 \right]; \quad (4.32)$$

$$\Omega = \frac{\pi J}{MS_{BH}} = \frac{J}{MR_+^2}; \quad (4.33)$$

$$\psi = \frac{1}{2M} \left( q + \frac{\pi q^3}{S_{BH}} \right) = \frac{q}{2M} \left[ 1 + \left( \frac{q}{R_+} \right)^2 \right]. \quad (4.34)$$

The expression of the *extremality parameter* given by the first of Eqs. (4.27) is suitably generalized as

$$c \equiv r_0 = \sqrt{M^2 - q^2 - p^2 - \frac{J^2}{M^2}}, \quad (4.35)$$

and thus extremality implies (see [26] and Refs. therein)

$$c = 0 \Leftrightarrow \begin{cases} M^2 = r_H^2 = \frac{(q^2 + p^2)}{2} + \sqrt{\frac{(q^2 + p^2)^2}{4} + J^2}; \\ J = M \sqrt{M^2 - q^2 - p^2}; \end{cases} \quad (4.36)$$

↓

$$S_{BH,c=0} = \pi \left( r_H^2 + \frac{J^2}{M^2} \right) = \pi \sqrt{(q^2 + p^2)^2 + 4J^2},$$

and thus the (*squared*) *effective radius* for extremal Kerr-Newman BH reads

$$R_{+,c=0}^2 \equiv R_H^2 = r_H^2 + \frac{J^2}{M^2} = \sqrt{(q^2 + p^2)^2 + 4J^2}. \quad (4.37)$$

## 5 Geodesic Action with a Constraint and Critical Points of the Black Hole Effective Potential

Let us now reconsider the system described by the (bosonic) Lagrangian density (4.14). In such a framework, the expression (4.35) of the extremality parameter gets modified as follows (see below) [84]:

$$M^2 + \frac{1}{2}G_{ab}(\phi_\infty)\Sigma^a\Sigma^b - V_{BH}(\phi_\infty^b, p^\Lambda, q_\Lambda) = c^2, \quad (5.1)$$

which in the extremal case ( $c = 0$ ) becomes

$$M^2 = V_{BH}(\phi_\infty^b, p^\Lambda, q_\Lambda) - \frac{1}{2}G_{ab}(\phi_\infty)\Sigma^a\Sigma^b. \quad (5.2)$$

Let us also recall that  $c$  is defined by (2.12) to be  $c = 2S_{BH}T_{BH}$ . Thus, in presence of scalars it follows that in general it depends on the asymptotical values  $\phi_\infty^a$  of the scalars (through the dependence of both  $S_{BH}$  and  $T_{BH}$  on  $\phi_\infty^a$ 's; consistently with the holding of Eq. (5.1), we consider the case  $J = 0$ ):

$$c = 2S_{BH}(\phi_\infty, p, q)T_{BH}(\phi_\infty, p, q), \quad (5.3)$$

and Eq. (5.1) can be rewritten as follows:

$$M^2 + \frac{1}{2}G_{ab}(\phi_\infty)\Sigma^a\Sigma^b - V_{BH}(\phi_\infty^b, p^\Lambda, q_\Lambda) = 4S_{BH}^2(\phi_\infty, p, q)T_{BH}^2(\phi_\infty, p, q). \quad (5.4)$$

By differentiating with respect to  $\phi_\infty^c$  and recalling the *Metric Postulate* in the asymptotic scalar manifold  $\mathcal{M}_{\phi_\infty}$ , one gets<sup>10</sup>

$$\begin{aligned} M \frac{\partial M}{\partial \phi_\infty^c} + \frac{1}{2}G_{ab}(\phi_\infty)\Sigma^a\nabla_{c,\infty}\Sigma^b - \frac{1}{2}\frac{\partial V_{BH}(\phi_\infty, p, q)}{\partial \phi_\infty^c} &= \\ &= 2c \left[ T_{BH}(\phi_\infty, p, q) \frac{\partial S_{BH}(\phi_\infty, p, q)}{\partial \phi_\infty^c} + S_{BH}(\phi_\infty, p, q) \frac{\partial T_{BH}(\phi_\infty, p, q)}{\partial \phi_\infty^c} \right], \end{aligned} \quad (5.5)$$

where  $\nabla_{c,\infty}$  is the (Christoffel) covariant derivative in  $\mathcal{M}_{\phi_\infty}$ :

$$\nabla_{c,\infty}\Sigma^b = \frac{\partial \Sigma^b}{\partial \phi_\infty^c} + \Gamma_{ca}{}^b(\phi_\infty)\Sigma^a. \quad (5.6)$$

When  $c$  is constant *globally* in  $\mathcal{M}_{\phi_\infty}$ , it trivially holds that

$$\frac{\partial c}{\partial \phi_\infty^a} = 0 \Leftrightarrow T_{BH}(\phi_\infty, p, q) \frac{\partial S_{BH}(\phi_\infty, p, q)}{\partial \phi_\infty^c} + S_{BH}(\phi_\infty, p, q) \frac{\partial T_{BH}(\phi_\infty, p, q)}{\partial \phi_\infty^c} = 0, \quad (5.7)$$

and Eq. (5.5) becomes

$$\begin{aligned} \frac{\partial M}{\partial \phi_\infty^c} &= \frac{1}{2M} \left[ \frac{\partial V_{BH}(\phi_\infty, p, q)}{\partial \phi_\infty^c} - G_{ab}(\phi_\infty)\Sigma^a\nabla_{c,\infty}\Sigma^b \right] = \\ &= \frac{\left[ \frac{\partial V_{BH}(\phi_\infty, p, q)}{\partial \phi_\infty^c} - G_{ab}(\phi_\infty)\Sigma^a\nabla_{c,\infty}\Sigma^b \right]}{2\sqrt{V_{BH}(\phi_\infty, p, q) - \frac{1}{2}G_{ab}(\phi_\infty)\Sigma^a\Sigma^b + c^2}}, \end{aligned} \quad (5.8)$$

---

<sup>10</sup>This corrects Eq. (23) of [82].

where in the last step of Eq. (5.8) we used Eq. (5.1). In the extremal case ( $c = 0$ ) Eq. (5.8) simplifies to

$$c = 0 : \frac{\partial M}{\partial \phi_\infty^c} = \frac{\left[ \frac{\partial V_{BH}(\phi_\infty, p, q)}{\partial \phi_\infty^c} - G_{ab}(\phi_\infty) \Sigma^a \nabla_{c, \infty} \Sigma^b \right]}{2 \sqrt{V_{BH}(\phi_\infty, p, q) - \frac{1}{2} G_{ab}(\phi_\infty) \Sigma^a \Sigma^b}}. \quad (5.9)$$

By recalling the expression (4.24) of the scalar charges and the relation (4.22) between  $V_{BH}$  and the generalized superpotential  $\mathcal{W}$  (both holding in the considered framework for  $c = 0$ ), one consistently obtains the second of Eqs. (5), giving the general expression of the (squared) ADM mass  $M^2$  in the extremal case ( $G^{ab}(\phi_\infty) \equiv G_\infty^{ab}$ ,  $G_{ab}(\phi_\infty) \equiv G_{ab, \infty}$ ,  $\mathcal{W}^2(\phi_\infty, p, q) \equiv \mathcal{W}_\infty^2$ ,  $V_{BH}(\phi_\infty, p, q) \equiv V_{BH, \infty}$ ; see also footnote 8):

$$c = 0 : \begin{cases} \frac{1}{2} G_{ab, \infty} \Sigma^a \Sigma^b = 2 G_\infty^{ab} (\partial_a \mathcal{W})_\infty (\partial_b \mathcal{W})_\infty ; \\ V_{BH, \infty} = \mathcal{W}_\infty^2 + 2 G_\infty^{ab} (\partial_a \mathcal{W})_\infty (\partial_b \mathcal{W})_\infty ; \end{cases} \implies M^2 = V_{BH, \infty} - \frac{1}{2} G_{ab, \infty} \Sigma^a \Sigma^b = \mathcal{W}_\infty^2. \quad (5.10)$$

By differentiating with respect to  $\phi_\infty^a$ , from such a result it trivially follows:

$$\frac{\partial M}{\partial \phi_\infty^a} = \frac{\partial \mathcal{W}_\infty}{\partial \phi_\infty^a} \quad (5.11)$$

which by recalling Eq. (5.9), consistently with Eqs. (4.23) and (4.24), yields an expression relating the critical points of  $V_{BH}(\phi_\infty, p, q)$ ,  $\mathcal{W}^2(\phi_\infty, p, q) \equiv \mathcal{W}_\infty^2$  and  $\Sigma^b(\phi_\infty, p, q)$  in  $\mathcal{M}_{\phi_\infty}$ :

$$c = 0 : \frac{\partial V_{BH}(\phi_\infty, p, q)}{\partial \phi_\infty^c} = \frac{\partial \mathcal{W}_\infty^2}{\partial \phi_\infty^c} + G_{ab}(\phi_\infty) \Sigma^a \nabla_{c, \infty} \Sigma^b. \quad (5.12)$$

As mentioned above, since the explicit form of  $\mathcal{W}$  has been explicitly determined for some  $\mathcal{N} > 2$ -extended,  $d = 4$  supergravities [41] and for some examples of  $\mathcal{N} = 2$ ,  $d = 4$  supergravity [36, 41], the explicit form of the ADM mass  $M$  (for a given, supporting BH charge configuration) for all such cases is currently known.

It is worth noticing that in *double-extremal* BHs (having  $c = 0$  and for which  $\Sigma^a = 0 \forall a = 1, \dots, n$ ) the relation (5.1) simply becomes

$$M^2 = V_{BH}(\phi_H(p, q), p, q), \quad (5.13)$$

expressing the fact that the ADM mass only depends on charges through the horizon value of  $V_{BH}$ , a trivial fact by recalling that the scalars are constant (independent on  $r$ )!

The most general class for which the above expression holds is the one of spherically symmetric, static, asymptotically flat BHs specified by the metric *Ansatz* ([84][92]; see also [89] and [41]):

$$ds^2 = -e^{2U} dt^2 + e^{-2U} \gamma_{mn} dx^m dx^n \equiv -e^{2U} dt^2 + e^{-2U} \left( \frac{c^4}{\sinh^4(c\tau)} d\tau^2 + \frac{c^2}{\sinh^2(c\tau)} d\Omega^2 \right), \quad (5.14)$$

The evolution coordinate  $\tau$  is related to the radial coordinate  $r$  by the relation<sup>11</sup>

$$\left(\frac{dr}{d\tau}\right)^2 = \frac{c^4}{\sinh^4(c\tau)} = (r - r_-)^2 (r - r_+)^2 = (r - M + c)^2 (r - M - c)^2 = [(r - M)^2 - c^2]^2, \quad (5.15)$$

whose integration yields the relation

$$r = -c \coth(c\tau) + M. \quad (5.16)$$

In the *extremal* case ( $c = 0$ ), since  $r_+ = r_- = r_H = M$ , the relation (5.16) reduces to the definition of  $\tau \equiv \frac{1}{r_H - r} = \frac{1}{M - r}$  previously introduced, and the metric (5.14) becomes

$$\begin{aligned} ds_{c=0}^2 &= -e^{2U} dt^2 + e^{-2U} [dr^2 + (r - r_H)^2 d\Omega^2] = -e^{2U} dt^2 + e^{-2U} [dr^2 + (r - M)^2 d\Omega^2] = \\ &= -e^{2U} dt^2 + \frac{e^{-2U}}{\tau^2} \left[ \frac{d\tau^2}{\tau^2} + d\Omega^2 \right] = -e^{2U} dt^2 + e^{-2U} [d\rho^2 + \rho^2 d\Omega^2], \end{aligned} \quad (5.17)$$

where in the last step we recalled the definition (3.21). Notice that Eq. (5.17) is nothing but the static Papapetrou-Majumdar class of metrics (3.24)-(3.25).

The relation (5.15) allows one to rewrite the metric (5.14) as follows:

$$\begin{aligned} ds^2 &= -e^{2U} dt^2 + e^{-2U} [dr^2 + (r - r_-)(r - r_+) d\Omega^2] = \\ &= -e^{2U} dt^2 + e^{-2U} \{dr^2 + [(r - M)^2 - c^2] d\Omega^2\}. \end{aligned} \quad (5.18)$$

By comparing such an expression with the expression of  $ds^2$  given below Eq. (22) of the first of Refs. [92], one obtains that the (*squared*) *effective radial coordinate*  $R^2$  introduced in the first of Refs. [92] can also be written as follows:

$$R^2 = e^{-2U} (r - r_-)(r - r_+) = e^{-2U} [(r - M)^2 - c^2], \quad (5.19)$$

which in the particular case of the *Maxwell-Einstein-(axion-)dilaton* BH treated in [92] reads (both in *non-extremal* and *extremal* cases; see Eq. (25) of the first of Refs. [92], and Eq. (2.6))

$$R^2 = r^2 - \Sigma^2. \quad (5.20)$$

Remarkably, in the *multi-dilaton system* such an expression enjoys the following generalization (see the treatment of Sect. 7, as well as Eq. (2.8)):

$$R^2 = r^2 - \frac{1}{2} G_{ab}(\phi_\infty) \Sigma^a \Sigma^b. \quad (5.21)$$

In the *non-extremal* case ( $c \neq 0$ ), the requirement of *non-vanishing, finite*  $A_H$  (“*large*” BH) implies the near-horizon limit of the scale function  $U$  to be [18] [89]

$$\begin{aligned} \lim_{\tau \rightarrow -\infty} e^{-2U} &= \left(\frac{A_H}{4\pi}\right) \frac{\sinh^2(c\tau)}{c^2} \sim R_+^2 \frac{e^{-2c\tau}}{c^2} = \\ &= \left(\frac{A_H}{4\pi}\right) \frac{1}{(r - r_-)(r - r_+)} = \frac{R_+^2}{(r - r_-)(r - r_+)}, \end{aligned} \quad (5.22)$$

<sup>11</sup>This corrects a mistake in Eq. (4.1.8) of [22], and a typo in Eq. (4) of [41].

where the relation (5.15) was used, and  $R_+$  is the *effective BH radius* defined by Eq. (2.4) (which, as pointed out above, in presence of scalars and/or for  $J \neq 0$  does *not* coincide with the radius  $r_+$  of the outer -event- horizon).

Thus, the near-horizon limit of the particular, spherically symmetric, asymptotically flat, static Papapetrou-Majumdar metric (5.14) (or (5.18)) reads

$$\begin{aligned}
\lim_{r \rightarrow r_H^+} ds^2|_{c \neq 0} &= - \left( \frac{4\pi}{A_H} \right) (r - r_-) (r - r_+) dt^2 + \left( \frac{A_H}{4\pi} \right) \frac{dr^2}{(r - r_-) (r - r_+)} + \left( \frac{A_H}{4\pi} \right) d\Omega^2 = \\
&= - \frac{(r - r_-) (r - r_+)}{R_+^2} dt^2 + \frac{R_+^2}{(r - r_-) (r - r_+)} dr^2 + R_+^2 d\Omega^2 = \\
&= - \frac{\pi (r - r_-) (r - r_+)}{S_{BH}} dt^2 + \frac{S_{BH}}{\pi (r - r_-) (r - r_+)} dr^2 + R_+^2 d\Omega^2 \sim \\
&\sim - \frac{c^2}{R_+^2} e^{2c\tau} dt^2 + R_+^2 \frac{e^{-2c\tau}}{c^2} dr^2 + R_+^2 d\Omega^2. \tag{5.23}
\end{aligned}$$

By defining  $\hat{\rho} \equiv e^{c\tau}$ , one obtains that

$$\lim_{\tau \rightarrow -\infty} ds^2|_{c \neq 0} = - \left( \frac{\hat{\rho} c}{R_+} \right)^2 dt^2 + R_+^2 (d\hat{\rho}^2 + d\Omega^2). \tag{5.24}$$

Thus, in the non-extremal case, the near-horizon geometry is not flat nor conformally flat; moreover, the distance in physical coordinate  $\hat{\rho}$  of any point from the horizon  $\hat{\rho}_H = 0$  is *finite*.

On the other hand, in the extremal case ( $c = 0$ , and thus  $r_+ = r_- \equiv r_H$ ), in order to have a non-vanishing, finite  $A_H$ ,  $U(\tau)$  should behave near the event horizon as follows:

$$\lim_{\tau \rightarrow -\infty} e^{-2U} = \frac{R_H^2}{(r - r_H)^2} = R_H^2 \tau^2 = \frac{R_H^2}{\rho^2}, \tag{5.25}$$

where we recall once again the notation  $R_{+,c=0}^2 \equiv R_H^2$ , and thus the near-horizon limit of the extremal metric (5.17) reads

$$\begin{aligned}
\lim_{\tau \rightarrow -\infty} ds_{c=0}^2 &= - \frac{(r - r_H)^2}{R_H^2} dt^2 + \frac{R_H^2}{(r - r_H)^2} [dr^2 + (r - r_H)^2 d\Omega^2] = \\
&= - R_H^2 \tau^2 dt^2 + R_H^2 \left[ \frac{d\tau^2}{\tau^2} + d\Omega^2 \right] = - \frac{\rho^2}{R_H^2} dt^2 + R_H^2 \left[ \frac{d\rho^2}{\rho^2} + d\Omega^2 \right]. \tag{5.26}
\end{aligned}$$

As noticed in [18], the physical difference between the extremal and non-extremal class of metrics (5.14) gives an hint why in the former case the *Attractor Mechanism* occurs, contrarily to the second case. In the extremal case, a scalar field  $\phi(\omega)$  (where  $\omega$  is the physical distance coordinate for  $c = 0$ , defined above as  $\omega \equiv \ln \rho = -\ln(-\tau)$ ) loses memory of the initial conditions of its radial dynamical evolution when

$$\frac{d\phi(\omega)}{d\omega} \rightarrow 0, \tag{5.27}$$






Figure 3: **On the left:** Euclidean section of the near-horizon “cigar” geometry of a *large* non-extremal BH ( $T_{BH} \neq 0$ ). **On the right:** Euclidean section of the *infinite throat* near-horizon geometry of a *large* extremal BH ( $T_{BH} = 0$ ). [18]

and this exactly happens in the near-horizon limit  $r \rightarrow r_H^+ \Leftrightarrow \omega_H = -\infty$ , meaning that the throat of an extremal (static, spherically symmetric, asymptotically flat) BH is *infinite*.

This actually does *not* occur in the non-extremal case, in which the non-extremal physical coordinate of the event (outer) horizon is  $\widehat{\rho}_H = 0$ , and

$$\lim_{\widehat{\rho} \rightarrow 0^+} \frac{d\phi(\widehat{\rho})}{d\widehat{\rho}} \rightarrow 0. \quad (5.28)$$

In other words, the Attractor Mechanism does *not* hold, because the scalar fields  $\phi^a(\widehat{\rho})$  do *not have enough (radial) time to loose memory of the asymptotical data of their evolution* [18].

As shown in [84] (and related Refs. therein), in general all the evolution equations for the gravitational degree of freedom  $U(\tau)$ , scalar fields  $\phi^a(\tau)$ , and electric and magnetic field strengths (whose unique non-vanishing components respectively are, within the considered *Ansätze* on space-time symmetries,  $F_{t\tau}^\Lambda = \partial_\tau \psi^\Lambda$  and  $G_{t\tau\Lambda} = \partial_\tau \chi_\Lambda$ ) can be derived in a one-dimensional Lagrangian formulation, obtained by performing a dimensional reduction based on the space-time symmetries (staticity, spherical symmetry, asymptotical flatness) of the considered background (5.14), yielding the following (almost geodesic) effective Lagrangian:

$$\mathcal{L}(U(\tau), \phi^a(\tau), p^\Lambda, q_\Lambda) = \left( \frac{dU(\tau)}{d\tau} \right)^2 + \frac{1}{2} G_{ab}(\phi(\tau)) \frac{d\phi^a}{d\tau} \frac{d\phi^b}{d\tau} + e^{2U(\tau)} V_{BH}(\phi(\tau), p^\Lambda, q_\Lambda), \quad (5.29)$$

along with the constraint

$$\left( \frac{dU(\tau)}{d\tau} \right)^2 + \frac{1}{2} G_{ab}(\phi(\tau)) \frac{d\phi^a}{d\tau} \frac{d\phi^b}{d\tau} - e^{2U(\tau)} V_{BH}(\phi(\tau), p^\Lambda, q_\Lambda) = c^2. \quad (5.30)$$

Taking into account that the asymptotical limit of the gravitational scale function reads

$$\lim_{\tau \rightarrow 0^-} U(\tau) \rightarrow M\tau, \quad (5.31)$$

the asymptotical limit of the constraint (5.30) yields, by recalling the definition (4.7) of the scalar charges, the relation (5.1). On the other hand, by noticing that (under minimal requirements of regularity) Eq. (5.22) yields [82]

$$\lim_{\tau \rightarrow -\infty} e^{2U} \sim c^2 \frac{e^{2c\tau}}{R_+^2} \Leftrightarrow \lim_{\tau \rightarrow -\infty} U(\tau) \sim c\tau \Leftrightarrow \lim_{\tau \rightarrow -\infty} \frac{dU(\tau)}{d\tau} \sim c, \quad (5.32)$$

and considering that the near-horizon limit of  $\frac{d\phi^a}{d\tau}$  is given by

$$\lim_{\tau \rightarrow -\infty} \frac{d\phi^a(\tau)}{d\tau} \sim e^{c\tau}, \quad (5.33)$$

it is immediate to check that the near-horizon limit of the constraint (5.30) yields nothing but the trivial identity  $c^2 = c^2$ .

It is worth remarking that the asymptotic relation (5.1) does *not* rely on supersymmetry at all, and holds under the general assumptions made on the field content, couplings, boundary conditions and space-time symmetries of the physical system being considered.

On the other hand, the derivation of the so-called *BPS(-like) bound* (3.11) requires both supersymmetry and duality invariance [100]. A conceivable generalization of such a bound in the considered framework embedded in  $\mathcal{N} = 2$ ,  $d = 4$  (ungauged) supergravity is

$$M_{ADM}^2 \geq |Z|^2(\phi_\infty, p^\Lambda, q_\Lambda), \quad (5.34)$$

where  $Z = Z(\phi, p, q)$  is the  $\mathcal{N} = 2$ ,  $d = 4$  *central charge function* (in the real parametrization of the scalar manifold). In particular, under the assumptions made  $|Z|^2(\phi_\infty, p, q)$  is the squared absolute value of the central charge of the  $\mathcal{N} = 2$ ,  $d = 4$  Poincaré superalgebra.

It is worth pointing out that for  $\mathcal{N} > 2$ -extended,  $d = 4$  (ungauged) supergravities  $|Z|$  can be generally replaced by the absolute value of the largest eigenvalue of the central charge matrix; indeed, generally the absolute values of the eigenvalues of such a matrix can be ordered *uniquely, i.e.* as functions of the scalars, with the ordering eventually depending only on the considered supporting BH charge configuration. In  $\mathcal{N} = 2$ ,  $d = 4$  (ungauged) supergravity coupled to  $n_V$  vector multiplets, the bound (5.34) can eventually be extended by a chain of (not necessarily strict) inequalities, involving the squared absolute values of the (covariant) derivatives of the *central charge function*  $Z$ , *i.e.* the so-called *matter charges*. By disregarding  $M_{ADM}^2$ , such a chain of inequalities generally hold for any value of the radial coordinate  $r$ , the ordering eventually depending only on the considered supporting BH charge configuration.

Now, by using the (supersymmetry-independent) asymptotical relation (5.1), Eq. (5.34) would imply that in  $\mathcal{N} = 2$ ,  $d = 4$  supergravity (under the assumptions made) it holds that

$$V_{BH}(\phi_\infty, p^\Lambda, q_\Lambda) - \frac{1}{2}G_{ab}(\phi_\infty)\Sigma^a\Sigma^b + c^2 \geq |Z|^2(\phi_\infty, p^\Lambda, q_\Lambda). \quad (5.35)$$

By also recalling that the BH effective potential in  $\mathcal{N} = 2$ ,  $d = 4$  (ungauged) supergravity reads<sup>12</sup> (in the real parametrization of the scalars)

$$V_{BH}(\phi, p^\Lambda, q_\Lambda) = |Z|^2(\phi, p^\Lambda, q_\Lambda) + G^{ab}(\phi)(D_a Z)(\phi, p^\Lambda, q_\Lambda) D_b Z(\phi, p^\Lambda, q_\Lambda), \quad (5.36)$$

one thus would achieve the following result<sup>13</sup>:

$$\begin{aligned} & G^{ab}(\phi_\infty)(D_a Z)(\phi_\infty, p^\Lambda, q_\Lambda) D_b Z(\phi_\infty, p^\Lambda, q_\Lambda) - \frac{1}{2} G_{ab}(\phi_\infty) \Sigma^a \Sigma^b + c^2 = \\ & = G^{ab}(\phi_\infty) [D_a Z(\phi_\infty, p, q) D_b Z(\phi_\infty, p, q) - \frac{1}{2} \Sigma_a \Sigma_b] + c^2 \geq 0, \end{aligned} \quad (5.37)$$

where  $\Sigma_a \equiv G_{ab}(\phi_\infty) \Sigma^b$ .

In the treatment given further below, the *BPS(-like) bound* (5.34), or equivalently the bound (5.37), will be shown to hold for the  $((U(1))^6 \rightarrow (U(1))^2$  truncation of the Maxwell-Einstein-(axion-)dilaton supergravity, corresponding to a particular 1-modulus model of  $\mathcal{N} = 2$ ,  $d = 4$  supergravity (see Sects. 6 and 7). Furthermore, it is worth pointing out that in [18] the bounds (5.34) and (5.37) are claimed to hold in general for  $c = 0$ .

Extremal BPS BHs have  $c = 0$  and they saturate the bounds (5.34) and (5.37), because  $G^{ab}(\phi_\infty) [D_a Z(\phi_\infty, p, q) D_b Z(\phi_\infty, p, q) - \frac{1}{2} \Sigma_a \Sigma_b] = 0$  due to the *first order BPS equations* (see *e.g.* [84]). However, by assuming the bounds (5.34) and (5.37) to hold (also) for  $c \neq 0$ , it is conceivable that they are saturated in some particular cases. Consequently, (beside non-extremal non-BPS) also non-extremal BPS BHs might exist.

In the case of *frozen* scalars (trivial radial dynamics:  $\frac{d\phi^a(r)}{dr} = 0 \forall a = 1, \dots, n$  and  $\forall r \in [r_H, \infty)$ )  $\Sigma_a = 0$ , and thus the BPS bound (5.37) simplifies to

$$G^{ab}(\phi_\infty)(D_a Z)(\phi_\infty, p^\Lambda, q_\Lambda) D_b Z(\phi_\infty, p^\Lambda, q_\Lambda) + c^2 \geq 0. \quad (5.38)$$

For  $c = 0$  one can have the well known *double-extremal BPS* BHs for which (see *e.g.* [84])

$$D_a Z(\phi(r), p^\Lambda, q_\Lambda) = 0 \forall a = 1, \dots, n, \quad \forall r \in [r_H, \infty), \quad (5.39)$$

but also *double-extremal non-BPS* BHs, having

$$G^{ab}(\phi_\infty)(D_a Z)(\phi_\infty, p^\Lambda, q_\Lambda) D_b Z(\phi_\infty, p^\Lambda, q_\Lambda) > 0, \quad (5.40)$$

whose first example was explicitly constructed in [27]). On the other hand, for  $c \neq 0$  one can have *double-non-extremal non-BPS* BHs for which the strict inequality holds in (5.38). Notice that the (strict) positive definiteness of the metric of the scalar manifold prevents the bound (5.38) to be saturated for  $c \neq 0$ ; thus, *double-non-extremal BHs or non-extremal BHs with no scalars at all* (such as non-extremal RN BHs) *should never be BPS*<sup>14</sup>.

<sup>12</sup>We use covariant derivatives, because the real parametrization of the scalars inherits the *Kähler* structure of the complex parametrization. Indeed, vector multiplets' scalars of  $\mathcal{N} = 2$ ,  $d = 4$  span (*special*) *Kähler* manifolds.

<sup>13</sup>The extremal limit ( $c = 0$ ) of inequality (5.37) corrects and clarifies Eq. (2.11) of [18].

<sup>14</sup>As pointed out above, the clash with some results in recent literature ([101]; see also [36, 41]) is just apparent, since a different, slightly generalized meaning, is given to "BPS" therein, as "admitting a *first-order* (and thus *BPS-like*) formulation".

In the considered framework, the effective (squared) radius relevant for the computation of the BH entropy is given by the following generalization of the first of Eqs. (4.29):

$$r_{\pm} = M \pm \sqrt{M^2 - V_{BH}(\phi_{\infty}, p, q) + \frac{1}{2}G_{ab}(\phi_{\infty})\Sigma^a\Sigma^b}. \quad (5.41)$$

Furthermore, as mentioned above, in [36, 41] a first order formalism for extremal (and some cases of non-extremal [41]) static, spherically symmetric BHs was developed, based on a positive definite *generalized superpotential*  $\mathcal{W}$ . In the extremal case  $\mathcal{W}$  is a function only of the conserved BH electric and magnetic charges and of the scalars, and it does *not* depend explicitly on the radial evolution coordinate  $\tau$ :

$$\mathcal{W} = \mathcal{W}(\phi(\tau), p, q). \quad (5.42)$$

By exploiting the relation between  $\mathcal{W}$  and  $V_{BH}$ , and using the first order differential Eqs. satisfied by  $\mathcal{W}$ , in [41] it was proved that in the *extremal* case

$$\lim_{\tau \rightarrow -\infty} \mathcal{W}(\phi(\tau), p, q) = R_H(p, q) = \sqrt{V_{BH}(\phi_H(p, q), p, q)} \quad (\text{near-horizon limit});$$

$$\lim_{\tau \rightarrow 0^-} \mathcal{W}(\phi(\tau), p, q) = M_{ADM}(\phi_{\infty}, p, q) = r_H(\phi_{\infty}, p, q) \quad (\text{asymptotical limit});$$

$$\frac{d\mathcal{W}(\phi(\tau), p, q)}{d\tau} \geq 0 \quad (\tau\text{-monotonicity of } \mathcal{W})$$

↓

$$r_H(\phi_{\infty}, p, q) = M_{ADM}(\phi_{\infty}, p, q) = \lim_{\tau \rightarrow 0^-} \mathcal{W}(\phi(\tau), p, q) \geq \sqrt{V_{BH}(\phi_H(p, q), p, q)} = R_H(p, q). \quad (5.43)$$

The first two equations give the general expressions of the *effective radius*  $R_H(p, q)$  (defined by Eq. (2.4)) and of the ADM mass  $M_{ADM}(\phi_{\infty}, p, q) = r_H(\phi_{\infty}, p, q)$  (see Eq. (3.20)) for static, spherically symmetric, asymptotically flat, dyonic *extremal* BHs.

Thus,  $\mathcal{W}$  given by Eq. (5.42) is a positive *monotonic* function, *decreasing* from the *moduli-dependent* value  $M_{ADM}(\phi_{\infty}, p, q) = r_H(\phi_{\infty}, p, q)$  at  $r \rightarrow \infty$ , towards the *moduli-independent* value  $R_H(p, q) = \sqrt{V_{BH}(\phi_H(p, q), p, q)}$  at the event BH horizon radius  $r = r_H$ . Consequently, in the extremal case  $\mathcal{W}^2$  appears to be the suitable candidate for the *C-function*<sup>15</sup>, whose existence was shown in [16].

In Sect. 7 we will prove that the inequality given in the fourth line of (5) can actually be further specialized in the *multi-dilaton system* (*i.e.* in  $\mathcal{N} = 2$  quadratic  $d = 4$  supergravity), in which it holds that

$$\begin{aligned} \frac{S_{BH, c=0}(p, q)}{\pi} &\equiv R_H^2(p, q) = r_H^2(\phi_{\infty}, p, q) - \frac{1}{2}G_{ab}(\phi_{\infty})\Sigma^a(\phi_{\infty}, p, q)\Sigma^b(\phi_{\infty}, p, q) = \\ &= \mathcal{W}^2(\phi_{\infty}, p, q) - 2G^{ab}(\phi_{\infty}) \lim_{\tau \rightarrow 0^-} (\partial_a \mathcal{W}(\phi(\tau), p, q)) \partial_b \mathcal{W}(\phi(\tau), p, q). \end{aligned} \quad (5.44)$$

---

<sup>15</sup>Notice that, when going beyond the Einstein approximation, the uniqueness of the *C-function* does not generally hold any more. For example, in *Lovelock gravity* [109] (at least) two independent *C-functions* can be determined.

This is nothing but the *extremal limit*  $c = 0$ , expressed in the framework of *first order formalism*, of Eq. (2.7), or equivalently the *many-moduli generalization* of the *extremal case* of formula holding for the so-called (*axion-*)*dilaton extremal BH* [92], given by Eq. (2.5) or Eq. (6.10) below; furthermore, in the last step Eq. (5.10) was used (for the assumptions on the limit, see footnotes 8 and 16).

It is here worth pointing out that in the *non-extremal case* (*i.e.*  $c \neq 0$ ) the generalization from the *one-modulus* formula (6.10) to the *many-moduli* formula (see Eq. (2.7))

$$\frac{S_{BH,c \neq 0}(\phi_\infty, p, q)}{\pi} \equiv R_+^2(\phi_\infty, p, q) = r_+^2(\phi_\infty, p, q) - \frac{1}{2}G_{ab}(\phi_\infty)\Sigma^a(\phi_\infty, p, q)\Sigma^b(\phi_\infty, p, q) \quad (5.45)$$

is only *guessed*, but at present cannot be rigorously proved. Indeed, for static, spherically symmetric, asymptotically flat dyonic *non-extremal* BHs a *first order formalism* is currently unavailable, so there is no way to compute the scalar charges (beside the direct integration of the Eqs. of motion of the scalars, as far as we know feasible only for the (*axion-*)*dilaton* BH [92]).

The crucial feature expressed by Eq. (5.44) is the *disappearance* of the dependence on the (*asymptotical*) *moduli*  $\phi_\infty$  in the combination  $r_H^2 - \frac{1}{2}G_{ab}\Sigma^a\Sigma^b$  of quantities  $r_H^2$  and  $G_{ab}\Sigma^a\Sigma^b$ , which separately are *moduli-dependent*.

Let us now recall the (guessed generalization of the) BPS bound (5.34), which in the considered context is equivalent, through the relation (5.1), to the bound (5.37). The first order formalism for extremal BHs introduced in [36, 41] might provide, through the second limit of Eq. (5) and the knowledge of an explicit expression for  $\mathcal{W}$ , a proof of the  $c = 0$  limit of inequalities (5.34) and (5.37) in some of the  $\mathcal{N} \geq 2$ ,  $d = 4$  (ungauged) supergravities. However, let us recall once again that currently an expression of  $\mathcal{W}$  for a generic  $\mathcal{N} = 2$ ,  $d = 4$  supergravity is unavailable.

Summarizing, one ends up with four sets of relations, which can be used to study the *BH thermodynamics* (at *equilibrium*, for  $c = 0$ ) and its interconnection with *attractors* of the scalar dynamics (and *supersymmetry*, when the bosonic system under consideration is embedded in *supergravity*). Let us recall once again that the most general environment considered in the present work is given by stationary (rotating with constant angular momentum), spherically symmetric and asymptotically flat BHs.

The mentioned four sets of relations, whose *interplay* is (partially) shown by the treatment given above, are:

1) the spatial asymptotic ( $r \rightarrow \infty \Leftrightarrow \tau \rightarrow 0^-$ ) limit (5.1) of the *constraint* (5.30) of the effective one-dimensional Lagrangian density (5.29). It holds true under the assumed asymptotic boundary conditions for  $U(\tau)$  and  $\phi^a(\tau)$ , for  $c \in \mathbb{R}^+$  and  $J = 0$ ;

2) the general expression of  $r_\pm^2$  (and of the *effective squared radius*  $R_\pm^2$ ) in terms of the other thermodynamical quantities (thence, directly yielding the value of the Bekenstein-Hawking BH entropy);

3) the relation given by the *Smarr-integrated*, finite version of the (*generalized*) *first law of BH thermodynamics* (in presence of scalars);

4) the asymptotical inequality (5.34), which is the proposed generalization of the *BPS bound* (3.11) [100] to the presence of scalars in  $\mathcal{N} = 2$ ,  $d = 4$  ungauged supergravity. In turn, let us recall that the *BPS bound* (3.11) is the generalization of the so-called *Theorem*

of *Positivity of Energy* (3.7)-(3.10) in the presence of electric charge  $q$  and magnetic charge  $p$  pertaining to the metric background (absence of scalars). It is worth remarking that the saturation of the *BPS bound* always implies the preservation of some (amount of the) supersymmetries, out of the all the ones pertaining to the asymptotically flat space-time background. The *BPS bounds* (3.11) and (5.34) hold for  $J = 0$ , and, differently from the relations of points 1, 2 and 3, their derivation relies on supersymmetry and duality invariance (see [100] and Refs. therein). As treated above, by combining the bound (5.34) with the asymptotic relation (5.1), one gets the  $c$ -parametrized bound (5.37), whose extremal limit ( $c = 0$ ) in [18] is stated to hold true in general in the considered framework. In Sect. 7 we will prove the extremal limit ( $c = 0$ ) of the bounds (5.34) and (5.37) to hold in the  $((U(1))^6 \rightarrow (U(1))^2$  truncation of the) Maxwell-Einstein-(axion-)dilaton supergravity, and we will further elaborate on the non-extremal case, as well.

## 6 The Maxwell-Einstein-(Axion-)Dilaton Gravity

The bosonic part of the  $d = 4$  Lagrangian density is [110]

$$-\frac{\mathcal{L}}{\sqrt{-g}} = (\partial_\mu \phi)^2 + e^{4\phi}(\partial_\mu a)^2 + e^{-2\phi} \sum_{\Lambda=1}^6 F_{\mu\nu}^\Lambda F_{\Lambda}^{\mu\nu} + \frac{1}{2}a \sum_{\Lambda=1}^6 F_{\mu\nu}^\Lambda \tilde{F}_{\Lambda}^{\mu\nu} + \frac{1}{2}\mathcal{R}. \quad (6.1)$$

It contains 6 Maxwell fields (for a total  $(U(1))^6$  gauge-invariance) and a dilaton-axial scalar

$$s \equiv a + ie^{-2\phi}. \quad (6.2)$$

The Lagrangian density (6.1) is the bosonic part of  $\mathcal{N} = 4$ ,  $d = 4$  *pure* supergravity (a sector of heterotic superstring compactified on a six-torus  $T^6$ ; see *e.g.* [111, 112]). By truncating  $(U(1))^6 \rightarrow (U(1))^2$ , the bosonic part of  $\mathcal{N} = 2$ ,  $d = 4$  supergravity (coupled to one vector multiplet,  $n_V = 1$ ) with *minimal coupling* (the so-called  $t^2$  model) [93] is recovered.

The coupled set of field equations for such a system in a static, spherically symmetric, asymptotically flat dyonic BH background described by the metric (5.14) were studied and solved in (the first Ref. of) [92]. In such a paper, the considered effective BH potential was

$$V_{BH}(\phi, q, p) = e^{2\phi}q^2 + e^{-2\phi}p^2 = Q^2 + P^2, \quad (6.3)$$

where  $V_{BH}(\phi, q, p) \equiv V_{BH}(\phi, a = 0, q, p)$  (see Eq. (7.1) below), and the *dressed charges*

$$Q \equiv qe^\phi, \quad P \equiv pe^{-\phi} \quad (6.4)$$

were introduced. The charges  $q$  and  $p$  respectively are an electric and a magnetic charge pertaining to two different  $U(1)$ 's inside  $(U(1))^6$ ; all the other charges are chosen to vanish, and this allows to put the axion field  $a$  to zero (in the extremal case, such a procedure will be clear from the treatment given below, based on the BH effective potential).

The scalar charge of the dilaton  $\phi$  is found to be (see the first Ref. of [92], and [114])

$$\Sigma^\phi \equiv \Sigma = \frac{e^{-2\phi_\infty}p^2 - e^{2\phi_\infty}q^2}{2M} = \frac{P_\infty^2 - Q_\infty^2}{2M}, \quad (6.5)$$

where  $\phi_\infty$  is the value of the dilaton at spatial infinity ( $r \rightarrow \infty$ ), and  $M$  is the ADM mass of the BH. By recalling Eq. (5.1), one obtains ( $V_{BH,\infty} \equiv V_{BH}(\phi_\infty, q, p)$ ):

$$M^2 + \Sigma^2 - V_{BH,\infty} = c^2 \Leftrightarrow M^2 + \frac{(P_\infty^2 - Q_\infty^2)}{4M^2} - Q_\infty^2 - P_\infty^2 = c^2;$$

$$\Updownarrow$$

$$M = \sqrt{V_{BH,\infty} - \Sigma^2 + c^2}. \quad (6.6)$$

By noticing that Eq. (4.17) [82] or Eqs. (4.24) and (4.23) [41] imply  $\Sigma$  to vanish at the critical point(s) of  $V_{BH(\infty)}$ , Eqs. (6.5) and (6.6) allow one to compute the *dilaton charge* as follows<sup>16</sup> ( $\partial \equiv \frac{\partial}{\partial \phi}$ ; also recall Eq. (5.5)):

$$\Sigma^2(\phi_\infty, p, q) = \frac{1}{2}(V_{BH,\infty} + c^2) \left[ 1 - \sqrt{1 - \frac{1}{4} \left( \frac{(\partial V_{BH})_\infty}{V_{BH,\infty} + c^2} \right)^2} \right]. \quad (6.7)$$

In the extremal case ( $c = 0$ ) Eq. (6.7) simplifies to

$$\Sigma^2(\phi_\infty, p, q)|_{c=0} = \frac{V_{BH,\infty}}{2} \left[ 1 - \sqrt{1 - \left( \frac{1}{2} (\partial \ln V_{BH})_\infty \right)^2} \right]. \quad (6.8)$$

By recalling Eq. (7.7), the inner (Cauchy) and outer (horizon) event radii are given for the considered (charge configuration of the) Maxwell-Einstein-(axion-)dilaton BH read [92]

$$r_\pm(\phi_\infty, p, q) = M \pm c = M \pm \sqrt{M^2 - V_{BH,\infty}(\phi_\infty, p, q) + \Sigma^2(\phi_\infty, p, q)} =$$

$$= M \pm \sqrt{M^2 - P_\infty^2 - Q_\infty^2 + \frac{(P_\infty^2 - Q_\infty^2)^2}{4M^2}}. \quad (6.9)$$

The *squared effective radius* reads [92]

$$R_+^2(\phi_\infty, p, q) = r_+^2(\phi_\infty, p, q) - \Sigma^2(\phi_\infty, p, q) = M^2 + c^2 + 2cM - \Sigma^2(\phi_\infty, p, q) =$$

$$= 2M^2 - P_\infty^2 - Q_\infty^2 + \sqrt{[2M^2 - (P_\infty + Q_\infty)^2][2M^2 - (P_\infty - Q_\infty)^2]}. \quad (6.10)$$

---

<sup>16</sup>We assume

$$(\partial V_{BH})_\infty \equiv \lim_{\tau \rightarrow 0^-} \frac{\partial V_{BH}(\phi(\tau), p, q)}{\partial \phi(\tau)} = \frac{\partial V_{BH}(\phi_\infty, p, q)}{\partial \phi_\infty}.$$

Also notice that Eq. (6.7) is constrained by the *reality condition*

$$1 - \frac{1}{4} \left( \frac{(\partial V_{BH})_\infty}{V_{BH,\infty} + c^2} \right)^2 \geq 0.$$

Thus, Eq. (2.4) yields the BH entropy to be

$$\begin{aligned}
S_{BH}(\phi_\infty, p, q) &= \frac{A_H(\phi_\infty, p, q)}{4} = \pi R_+^2(\phi_\infty, p, q) = \\
&= \pi \left[ 2M^2 - P_\infty^2 - Q_\infty^2 + \sqrt{[2M^2 - (P_\infty + Q_\infty)^2][2M^2 - (P_\infty - Q_\infty)^2]} \right].
\end{aligned} \tag{6.11}$$

By recalling the definition (2.12) of the extremality parameter, one can compute the BH temperature<sup>17</sup>:

$$\begin{aligned}
T_{BH}(\phi_\infty, p, q) &= \frac{c}{2S_{BH}(\phi_\infty, p, q)} = \\
&= \frac{\sqrt{M^2 - P_\infty^2 - Q_\infty^2 + \frac{(P_\infty^2 - Q_\infty^2)^2}{4M^2}}}{2\pi \left[ 2M^2 - P_\infty^2 - Q_\infty^2 + \sqrt{[2M^2 - (P_\infty + Q_\infty)^2][2M^2 - (P_\infty - Q_\infty)^2]} \right]}.
\end{aligned} \tag{6.12}$$

Let us now consider the *extremal* case ( $c = 0$ ). Thus, one obtains the following expression of the ADM mass:

$$c = 0; \tag{6.13}$$

$\Downarrow$

$$M^4 - (P_\infty^2 + Q_\infty^2)M^2 + \frac{1}{4}(P_\infty^2 - Q_\infty^2)^2 = 0; \tag{6.14}$$

$\Downarrow$

$$M_\pm^2 = \frac{(|P_\infty| \pm |Q_\infty|)^2}{2} \Leftrightarrow M_\pm = \frac{||P_\infty| \pm |Q_\infty||}{\sqrt{2}}, \tag{6.15}$$

and, by using Eq. (6.5), of the *dilaton scalar charge*:

$$\Sigma_{\pm, c=0} = \frac{P_\infty^2 - Q_\infty^2}{2M_\pm} = \frac{(|P_\infty| + |Q_\infty|)(|P_\infty| - |Q_\infty|)}{\sqrt{2}||P_\infty| \pm |Q_\infty||}; \tag{6.16}$$

$\Downarrow$

$$|\Sigma_\pm|_{c=0} = \frac{|P_\infty^2 - Q_\infty^2|}{2M_\pm} = \frac{||P_\infty| + |Q_\infty|| ||P_\infty| - |Q_\infty||}{\sqrt{2}||P_\infty| \pm |Q_\infty||} = \frac{||P_\infty| \mp |Q_\infty||}{\sqrt{2}}. \tag{6.17}$$

Consequently, the BH entropy in the extremal case reads ( $R_{+, c=0}^2 \equiv R_H^2$ )

$$\begin{aligned}
S_{BH, c=0} &= \frac{A_{H, c=0}}{4} = \pi R_H^2 = \pi [M_\pm^2 - \Sigma_\pm^2(\phi_\infty, p, q)]_{c=0} = \\
&= 2\pi [\pm |P_\infty Q_\infty|] = 2\pi |pq|,
\end{aligned} \tag{6.18}$$

---

<sup>17</sup>Notice we divide by  $S_{BH}(\phi_\infty, p, q)$ , because we always assume it to be strictly positive (by the Bekenstein-Hawking entropy-area formula (2.2) (or (2.4)), this is equivalent to consider *large* BHs, *i.e.* BHs with non-vanishing  $A_H$ ).



from which follows that only the branches  $M_+$  and  $\Sigma_+$  of Eqs. (6.15) and (6.17) are admissible (the branch “ $-$ ” yields negative entropy). Furthermore, let us remark that the dependence on the asymptotical value  $\phi_\infty$  of the dilaton drops out from the expression of the BH entropy  $S_{BH}|_{c=0}$  in the extremal case, as it has to be:

$$\frac{\partial S_{BH,c=0}}{\partial \phi_\infty} = 0. \quad (6.19)$$

It is worth mentioning that the formula (6.18) has an  $(SU(1,1) \times SO(n))$ -invariant generalization given by

$$S_{BH,c=0} = 2\pi \sqrt{p^2 q^2 - (p \cdot q)^2}, \quad (6.20)$$

where  $p^2 \equiv p \cdot p$ ,  $q^2 \equiv q \cdot q$  and  $p \cdot q \equiv p^\Lambda q_\Lambda = p^\Lambda q^\Sigma \delta_{\Lambda\Sigma}$ , here  $\Lambda$  ranging  $1, \dots, n$ , with the scalar product  $\cdot$  defined by  $\delta_{\Lambda\Sigma}$ , the  $n$ -dim. Euclidean metric. In turn, formula (6.20) is a particular case ( $m = 0$ , or  $n = 0$ ) of the more general,  $(SU(1,1) \times SO(n, m))$ -invariant expression

$$S_{BH,c=0} = 2\pi \sqrt{p^2 q^2 - (p \cdot q)^2}, \quad (6.21)$$

with  $\Lambda$  ranging  $1, \dots, n, n+1, \dots, n+m$ , and the scalar product  $\cdot$  defined by  $\eta_{\Lambda\Sigma}$ , the Lorentzian metric with signature  $(n, m)$ .

By putting  $m = 2$  (or  $n = 2$ ) in Eq. (6.21), one obtains the  $(SU(1,1) \times SO(n, 2))$ -invariant formula of the BH entropy for the  $\mathcal{N} = 2$ ,  $d = 4$  supergravity coupled to  $n_V$  vector multiplets whose scalar manifold is the sequence of reducible homogeneous symmetric special Kähler manifolds (with *cubic* prepotential)  $\frac{SU(1,1)}{U(1)} \times \frac{SO(2,n)}{SO(2) \times SO(n)}$  ( $n_V = n+1$ ,  $n \in \mathbb{N}$ ) (see *e.g.* [25] and Refs. therein).

On the other hand, by choosing  $m = 6$  (or  $n = 6$ ) in Eq. (6.21), one obtains the  $(SU(1,1) \times SO(6, n))$ -invariant formula of the BH entropy for  $\mathcal{N} = 4$ ,  $d = 4$  supergravity coupled to  $n$  matter multiplets, whose scalar manifold is the sequence of reducible homogeneous symmetric manifolds  $\frac{SU(1,1)}{U(1)} \times \frac{SO(6,n)}{SO(6) \times SO(n)}$  (see *e.g.* [89] and Refs. therein).

When considering *pure*  $\mathcal{N} = 4$ ,  $d = 4$  supergravity (*i.e.*  $n = 0$ : no matter multiplets), one thus obtains  $\Lambda = 1, \dots, 6$  and  $SU(1,1)$  as overall symmetry; the bosonic sector of the theory is described by the Lagrangian density (6.1). By further truncating the gauge group  $(U(1))^6 \rightarrow (U(1))^2$  (and thus restricting to  $\Lambda = 1, 2$ ), one can interpret  $SU(1,1)$  as the  $U$ -duality group [113] of the  $\mathcal{N} = 2$ ,  $d = 4$  supergravity *minimally coupled* [93] to  $n_V = 1$  vector multiplet (the so-called  $t^2$  model), whose scalar manifold is<sup>18</sup>  $\frac{SU(1,1)}{U(1)}$ , which is the  $n = 1$  element of the sequence of irreducible homogeneous symmetric special Kähler manifolds (with quadratic prepotential)  $\frac{SU(1,n)}{U(1) \times SU(n)}$  ( $n_V = n$ ,  $n \in \mathbb{N}$ ) (see *e.g.* [25] and Refs. therein). Consistently, this is the theory described by the Lagrangian (6.1) when

---

<sup>18</sup>Notice that there is another  $\mathcal{N} = 2$ ,  $d = 4$  supergravity coupled to  $n_V = 1$  vector multiplet, whose complex scalar spans the rank-1 homogeneous symmetric special Kähler manifold  $\frac{SU(1,1)}{U(1)}$ , endowed with cubic holomorphic prepotential (the so-called  $t^3$  model). However, such a manifold is *not* an element of the sequence  $\frac{SU(1,1)}{U(1)} \times \frac{SO(2,n)}{SO(2) \times SO(n)}$ , but rather it is an isolated case (see *e.g.* [115]).

In this sense, the quadratic sequence  $\frac{SU(1,n)}{U(1) \times SU(n)}$  is the only one (among all the homogeneous - symmetric - special Kähler manifolds) to admit a consistent *pure theory* (*i.e.* *no matter multiplets*) limit.

On the other hand, the first ( $n = 1$ ) element of the cubic sequence  $\frac{SU(1,1)}{U(1)} \times \frac{SO(2,n)}{SO(2) \times SO(n)}$  is the 2-moduli model based on  $\left(\frac{SU(1,1)}{U(1)}\right)^2$  (the so-called  $st^2$  model).

$\Lambda = 1, 2$  (in particular, in the symplectic basis in which the holomorphic prepotential reads  $F(X) = -iX^0X^1$ , see *e.g.* [36]). Indeed, when all BH charges  $(p^1, p^2, q_1, q_2)$  are switched on, the full fledged form of the BH entropy (6.18) reads

$$\begin{aligned} S_{BH,c=0} &= 2\pi\sqrt{((p^1)^2 + (p^2)^2)(q_1^2 + q_2^2) - (p^1q_1 + p^2q_2)^2} = \\ &= 2\pi\sqrt{(p^1)^2q_2^2 + (p^2)^2q_1^2 - 2p^1q_1p^2q_2} = 2\pi|p^1q_2 - p^2q_1|, \end{aligned} \quad (6.22)$$

Notice that Eq. (6.22) reduces to Eq. (6.18) for  $(p \equiv p^1, q \equiv q_2, p^2 = 0, q_1 = 0)$  or  $(p \equiv p^2, q \equiv q_1, p^1 = 0, q_2 = 0)$ . Moreover, it corresponds to the  $(SU(1,1) \times SO(n))$ -invariant formula (6.20) for  $n = 2$ . Indeed, this value of  $n$  is the only one for which, as shown by Eq. (6.22), the quantity under the square root is a perfect square for a generic BH charge configuration, thus reproducing the quadratic invariant  $\mathcal{I}_2$  of the (1-modulus,  $n = 1$  element of the) quadratic  $\mathcal{N} = 2, d = 4$  sequence (see *e.g.* [25] and Refs. therein).

## 7 Black Hole Attractors in (Axion-)Dilaton (Super)gravity

Let us analyze now the model described by the Lagrangian density (6.1) in the extremal case ( $c = 0$ ) within the formalism based on the BH effective potential  $V_{BH}$ . In the general case in which all electric and magnetic BH charges are switched on,  $V_{BH}$  reads<sup>19</sup> [89] ( $\Lambda = 1, \dots, 6$ )

$$\begin{aligned} V_{BH}(\phi, a, p^\Lambda, q_\Lambda) &= e^{2\phi}(sp_\Lambda - q_\Lambda)(\bar{s}p^\Lambda - q^\Lambda) = \\ &= e^{2\phi}[(a + ie^{-2\phi})p_\Lambda - q_\Lambda][(a - ie^{-2\phi})p^\Lambda - q^\Lambda] = \\ &= (e^{2\phi}a^2 + e^{-2\phi})p^2 + e^{2\phi}q^2 - 2ae^{2\phi}p \cdot q, \end{aligned} \quad (7.1)$$

where the definition (6.2) was used. By computing the criticality conditions of  $V_{BH}$  given by Eq. (7.1), one obtains the following stabilization equations for the axion  $a$  and the dilaton  $\phi$  [89]:

The critical points are :

$$\left. \frac{\partial V_{BH}(\phi, a, p, q)}{\partial a} \right|_{(\phi, a) = (\phi_H(p, q), a_H(p, q))} = 0 \iff a_H(p, q) = \frac{p \cdot q}{p^2}; \quad (7.2)$$

$$\begin{aligned} &\left. \frac{\partial V_{BH}(\phi, a, p, q)}{\partial \phi} \right|_{(\phi, a) = (\phi_H(p, q), a_H(p, q))} = \\ &= -e^{-4\phi}p^2 + q^2 - a_H(p, q)p \cdot q = -e^{-4\phi}p^2 + q^2 - \frac{(p \cdot q)^2}{p^2} = 0; \\ &\quad \Updownarrow \\ &e^{-2\phi_H(p, q)} = \frac{\sqrt{p^2q^2 - (p \cdot q)^2}}{p^2}. \end{aligned} \quad (7.3)$$

<sup>19</sup>Eq. (7.1) fixes a typo in Eq. (225) of [89].

Thus, the Bekenstein-Hawking BH entropy can be computed to be

$$S_{BH}(p, q) = \frac{A_H(p, q)}{4} = \pi V_{BH}(\phi_H(p, q), a_H(p, q), p, q) = 2\pi \sqrt{p^2 q^2 - (p \cdot q)^2}, \quad (7.4)$$

which corresponds to formula (6.20) for  $n = 6$ , and whose truncation to  $n = 2$  gives formula (6.22). By further putting  $p^1 = 0, q_2 = 0$  (or  $p^2 = 0, q_1 = 0$ ) and defining  $p^2 \equiv p, q_1 \equiv q$  (or  $p^1 \equiv p, q_2 \equiv q$ ), the expression (6.18) for the BH entropy is obtained.

In other words, the Maxwell-Einstein-dilaton  $d = 4$  gravity is given by the  $(U(1))^6 \rightarrow (U(1))^2$  truncation of the (bosonic sector of)  $\mathcal{N} = 4, d = 4$  supergravity, in a charge configuration in which

$$p \cdot q \equiv p^\Lambda q_\Lambda = p^1 q_1 + p^2 q_2 = 0. \quad (7.5)$$

Such a constraint implies, by the stabilization Eqs. (7.2)-(7.3), the axion to be *frozen out* ( $a_H(p, q) = 0$ ), and the dilaton to be stabilized as follows:

$$e^{-2\phi_H(p, q)} \Big|_{p \cdot q = 0} = \left| \frac{q}{p} \right|. \quad (7.6)$$

Notice that for the BH charge configuration (7.5) the *vanishing axion* solution  $a(r) = 0 \forall r \in [r_H, \infty)$  is a particular solution of the axionic Eq. of motion (actually also for  $c \neq 0$ , in which case  $r_H$  is usually understood as the radius of the outer - event - horizon). In general, the condition (7.5) is necessary but not sufficient for the choice of the *vanishing axion* solution.

Furthermore, as anticipated in Sect. 2 (see Eq. (2.7)), Eq. (6.10) can be generalized (*at least*) for the *multi-dilaton system* whose scalar manifold is  $\frac{SU(1, n)}{U(1) \times SU(n)}$  ( $n_V = n, n \in \mathbb{N}$ ) (which is nothing but the  $\mathcal{N} = 2, d = 4$  *quadratic sequence* introduced above) as follows (here and below  $\phi$  denotes the whole set of scalars; see also Eq. (2.7)):

$$\begin{aligned} R_+^2 &= R^2 \Big|_{r = \sqrt{\frac{1}{2} G_{ab}(\phi_\infty) \Sigma^a \Sigma^b}}^{r=r_+} = r_+^2 - \frac{1}{2} G_{ab}(\phi_\infty) \Sigma^a \Sigma^b = \\ &= 2M^2 - V_{BH}(\phi_\infty, p, q) + 2M \sqrt{M^2 - V_{BH}(\phi_\infty, p, q) + \frac{1}{2} G_{ab}(\phi_\infty) \Sigma^a \Sigma^b} = \\ &= R_+^2(\phi_\infty, p, q), \end{aligned} \quad (7.7)$$

where the *effective radial coordinate*  $R$  is given by Eqs. (5.21) (or (2.8)). Thus, the BH entropy is

$$\begin{aligned} S_{BH}(\phi_\infty, p, q) &= \frac{A_H(\phi_\infty, p, q)}{4} = \pi R_+^2(\phi_\infty, p, q) = \pi \left[ r_+^2(\phi_\infty, p, q) - \frac{1}{2} G_{ab}(\phi_\infty) \Sigma^a \Sigma^b \right] = \\ &= \pi \left[ 2M^2 - V_{BH}(\phi_\infty, p, q) + 2M \sqrt{M^2 - V_{BH}(\phi_\infty, p, q) + \frac{1}{2} G_{ab}(\phi_\infty) \Sigma^a \Sigma^b} \right]. \end{aligned} \quad (7.8)$$

Due to the positive definiteness of  $G_{ab}$ , Eq. (7.7) yields  $R_+ \leq r_+$  (see also Eq. (2.7)). Thus, the presence of scalars actually *decreases* the BH radius relevant for the computation of the entropy through the Bekenstein-Hawking (semi)classical formula (2.4). In particular,

$$R_+ (= R_H) = r_+ (= r_- = r_H) \quad (7.9)$$

only for *double-extremal* (static, spherically symmetric, asymptotically flat) dyonic BHs. On the other hand, in the *double-non-extremal* case (see comment below Eq. (5.40)) within the same class of BHs the expressions in brackets in Eq. (7.9) do *not* hold.

In the extremal case:

$$c = 0 \Leftrightarrow M^2 = V_{BH}(\phi_\infty, p, q) - \frac{1}{2}G_{ab}(\phi_\infty)\Sigma^a\Sigma^b; \quad (7.10)$$

consequently

$$r_{+,c=0}^2 = r_{-,c=0}^2 \equiv r_H^2 = V_{BH}(\phi_\infty, p, q) - \frac{1}{2}G_{ab}(\phi_\infty)\Sigma^a\Sigma^b = M^2; \quad (7.11)$$

$$\begin{aligned} R_{+,c=0}^2 &\equiv R_H^2 = \left[ r_+^2 - \frac{1}{2}G_{ab}(\phi_\infty)\Sigma^a\Sigma^b \right]_{c=0} = \left[ M^2 - \frac{1}{2}G_{ab}(\phi_\infty)\Sigma^a\Sigma^b \right]_{c=0} = \\ &= \left[ 2M^2 - V_{BH}(\phi_\infty, p, q) \right]_{c=0} = \left[ V_{BH}(\phi_\infty, p, q) - G_{ab}(\phi_\infty)\Sigma^a\Sigma^b \right]_{c=0}. \end{aligned} \quad (7.12)$$

By recalling Eqs. (4.22) and (4.24), one obtains

$$\begin{aligned} R_{+,c=0}^2 &\equiv R_H^2 = \mathcal{W}^2(\phi_\infty, p, q) - 2G^{ab}(\phi_\infty)(\partial_a\mathcal{W})(\phi_\infty, p, q)(\partial_b\mathcal{W})(\phi_\infty, p, q) = \\ &= \mathcal{W}^2(z_\infty, \bar{z}_\infty, p, q) - 4G^{i\bar{j}}(z_\infty, \bar{z}_\infty)(\partial_i\mathcal{W})(z_\infty, \bar{z}_\infty, p, q)(\bar{\partial}_{\bar{j}}\mathcal{W})(z_\infty, \bar{z}_\infty, p, q) = \\ &= R_H^2(p, q) \end{aligned} \quad (7.13)$$

where  $\mathcal{W}$  is the superpotential of the first order formulation for extremal BHs [36, 41] (see also the treatment of Sects. 4 and 5). Notice that in the second line a complex parametrization of the scalar manifold has been introduced, which is convenient for the *multi-dilaton system* under consideration, whose scalar manifold is the homogeneous symmetric special Kähler space  $\frac{SU(1,n)}{U(1)\times SU(n)}$ .

By using the explicit expressions of  $\mathcal{W}_{\frac{1}{2}\text{-BPS}}$  and  $\mathcal{W}_{\text{non-BPS}(Z=0)}$  obtained in [41] (for the case  $n = 1$  see also [36]), recalling that  $C_{ijk} = 0$  for the case at hand, and using the differential relations of special Kähler geometry (see *e.g.* [108], and Refs. therein), one obtains the following results:

1) for the  $\frac{1}{2}$ -BPS attractor flow:

$$r_{H, \frac{1}{2}-BPS}^2(z_\infty, \bar{z}_\infty, p, q) = M_{ADM, \frac{1}{2}-BPS}^2(z_\infty, \bar{z}_\infty, p, q) = \mathcal{W}_{\frac{1}{2}-BPS}^2(z_\infty, \bar{z}_\infty, p, q) = \lim_{\tau \rightarrow 0^-} |Z|^2; \quad (7.14)$$

$$\begin{aligned} \Sigma_{i, \frac{1}{2}-BPS}(z_\infty, \bar{z}_\infty, p, q) &= 2 \lim_{\tau \rightarrow 0^-} \left[ \partial_i \mathcal{W}_{\frac{1}{2}-BPS}(z(\tau), \bar{z}(\tau), p, q) \right] = \\ &= 2 \lim_{\tau \rightarrow 0^-} \left[ \nabla_i \mathcal{W}_{\frac{1}{2}-BPS}(z(\tau), \bar{z}(\tau), p, q) \right] = \\ &= \lim_{\tau \rightarrow 0^-} \left[ \sqrt{\frac{\bar{Z}}{Z}} D_i Z \right] = \lim_{\tau \rightarrow 0^-} \left[ \sqrt{\frac{\bar{Z}}{Z}} \left( \partial_i Z + \frac{1}{2} (\partial_i K) Z \right) \right]; \end{aligned} \quad (7.15)$$

$$R_{+, c=0, \frac{1}{2}-BPS}^2 \equiv$$

$$\begin{aligned} R_{H, \frac{1}{2}-BPS}^2 &= \lim_{\tau \rightarrow 0^-} \left[ \begin{array}{l} \mathcal{W}_{\frac{1}{2}-BPS}^2(z(\tau), \bar{z}(\tau), p, q) + \\ -4G^{i\bar{j}}(z(\tau), \bar{z}(\tau)) \left( \partial_i \mathcal{W}_{\frac{1}{2}-BPS} \right)(z(\tau), \bar{z}(\tau), p, q) \cdot \\ \cdot \left( \bar{\partial}_{\bar{j}} \mathcal{W}_{\frac{1}{2}-BPS} \right)(z(\tau), \bar{z}(\tau), p, q) \end{array} \right] = \\ &= \mathcal{I}_2(p, q) = V_{BH}|_{(z_H, \bar{z}_H)_{\frac{1}{2}-BPS}} = \\ &= R_{H, \frac{1}{2}-BPS}^2(p, q) = \frac{S_{BH, \frac{1}{2}-BPS}(p, q)}{\pi}; \end{aligned} \quad (7.16)$$

2) for the non-BPS ( $Z = 0$ ) attractor flow

$$\begin{aligned}
r_{H,non-BPS}^2(z_\infty, \bar{z}_\infty, p, q) &= M_{ADM,non-BPS}^2(z_\infty, \bar{z}_\infty, p, q) = \\
&= \mathcal{W}_{non-BPS}^2(z_\infty, \bar{z}_\infty, p, q) = \lim_{\tau \rightarrow 0^-} \left[ G^{i\bar{j}}(D_i Z) \bar{D}_{\bar{j}} \bar{Z} \right];
\end{aligned} \tag{7.17}$$

$$\begin{aligned}
\Sigma_{i,non-BPS}(z_\infty, \bar{z}_\infty, p, q) &= 2 \lim_{\tau \rightarrow 0^-} [\partial_i \mathcal{W}_{non-BPS}(z(\tau), \bar{z}(\tau), p, q)] = \\
&= 2 \lim_{\tau \rightarrow 0^-} [\nabla_i \mathcal{W}_{non-BPS}(z(\tau), \bar{z}(\tau), p, q)] = \\
&= \lim_{\tau \rightarrow 0^-} \left[ \frac{\bar{Z} D_i Z}{\sqrt{G^{i\bar{j}}(D_i Z) \bar{D}_{\bar{j}} \bar{Z}}} \right] = \\
&= \lim_{\tau \rightarrow 0^-} \left[ \frac{\bar{Z} (\partial_i Z + \frac{1}{2} (\partial_i K) Z)}{\sqrt{G^{i\bar{j}} (\partial_i Z + \frac{1}{2} (\partial_i K) Z) (\bar{\partial}_{\bar{j}} \bar{Z} + \frac{1}{2} (\bar{\partial}_{\bar{j}} K) \bar{Z})}} \right];
\end{aligned} \tag{7.18}$$

$$\begin{aligned}
R_{+,c=0,non-BPS}^2 &\equiv R_{H,non-BPS}^2 = \\
&= \lim_{\tau \rightarrow 0^-} \left[ \begin{aligned} &\mathcal{W}_{non-BPS}^2(z(\tau), \bar{z}(\tau), p, q) + \\ &-4G^{i\bar{j}}(z(\tau), \bar{z}(\tau)) (\partial_i \mathcal{W}_{non-BPS})(z(\tau), \bar{z}(\tau), p, q) \cdot \\ &\cdot (\bar{\partial}_{\bar{j}} \mathcal{W}_{non-BPS})(z(\tau), \bar{z}(\tau), p, q) \end{aligned} \right] = \\
&= -\mathcal{I}_2(p, q) = V_{BH}|_{(z_H, \bar{z}_H)_{non-BPS}} = \\
&= R_{H,non-BPS}^2(p, q) = \frac{S_{BH,non-BPS}(p, q)}{\pi}.
\end{aligned} \tag{7.19}$$

Here  $D_i$  denotes the Kähler- $U(1)$  and  $H = U(1) \times SU(n)$ -covariant derivative, whereas  $\mathcal{I}_2(p, q)$  is the invariant of the  $U$ -duality group  $SU(1, n)$ , *quadratic* in charges (see *e.g.* [89], and Refs. therein). Notice that  $|Z|^2$  and  $G^{i\bar{j}}(D_i Z) \bar{D}_{\bar{j}} \bar{Z}$  are the only independent ( $H = U(1) \times SU(n)$ )-invariants.

Eqs. (7.16) and (7.19) are consistent, because the  $\frac{1}{2}$ -BPS- and non-BPS ( $Z = 0$ )- supporting BH charge configurations in the (extremal) *multi-dilaton system* are respectively defined by the quadratic constraints  $\mathcal{I}_2(p, q) > 0$  and  $\mathcal{I}_2(p, q) < 0$ .

Notice that in the extremality regime ( $c = 0$ ) the *effective radius*  $R_H$ , and thus  $A_H$  and the Bekenstein-Hawking entropy  $S_{BH}$  are *independent* on the particular vacuum or ground state of the considered theory, *i.e.* on  $\phi_\infty^a$  (or equivalently on  $(z_\infty, \bar{z}_\infty)$ ), but rather they depend *only* on the electric and magnetic charges  $q_\Lambda$  and  $p^\Lambda$ , which are *conserved*

due to the overall  $(U(1))^{n+1}$  gauge-invariance. The independence on  $\phi_\infty^a$  is of crucial importance for the consistency of the *microscopic state counting interpretation* of  $S_{BH}$ , as well as for the overall consistency of the macroscopic thermodynamic picture of the BH. However, it is worth recalling that the ADM mass  $M_{(ADM)}$  generally does depend on  $\phi_\infty^a$  *also in the extremal case*.

## Black Hole Attractors and Supersymmetry

Let us now analyze the supersymmetry-preserving features of (extremal) BH attractors of the Maxwell-Einstein-(axion-)dilaton system, when it is embedded in supergravity.

We start by defining the two following quantities [92][112]

$$Z_1(\phi, p, q) \equiv \frac{(Q + P)}{\sqrt{2}} = \frac{qe^\phi + pe^{-\phi}}{\sqrt{2}}, \quad (7.20)$$

$$Z_2(\phi, p, q) \equiv \frac{(Q - P)}{\sqrt{2}} = \frac{qe^\phi - pe^{-\phi}}{\sqrt{2}}. \quad (7.21)$$

In  $\mathcal{N} = 4, d = 4$  supergravity [110],  $Z_1$  and  $Z_2$  are nothing but the eigenvalues of the skew-diagonal(ized) central charge matrix  $Z^{AB}$ , arising in the  $\mathcal{N} = 4, d = 4$  superalgebra ( $\alpha, \beta = 1, 2, A, B = 1, \dots, 4; \epsilon$  is the 2-dim. symplectic metric):

$$\{Q_\alpha^A, Q_\beta^B\} = \epsilon_{\alpha\beta} Z^{AB}; \quad (7.22)$$

$$Z_{SD}^{AB} = \begin{pmatrix} Z_1 & 0 \\ 0 & Z_2 \end{pmatrix} \otimes \epsilon, \quad \epsilon \equiv \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}. \quad (7.23)$$

When truncating down to the *minimally coupled* 1-modulus  $\mathcal{N} = 2, d = 4$  supergravity (and thus after dropping two gravitinos out from the supergravity multiplet)  $Z_1$  and  $Z_2$  can respectively be interpreted as follows:

$$Z_1 = Z, \quad Z_2 = \partial_\phi Z, \quad (7.24)$$

where  $Z = Z(\phi, p, q)$  is the *central charge function* of  $\mathcal{N} = 2, d = 4$  supergravity. Its asymptotical values for  $r \rightarrow r_H^+$  and  $r \rightarrow \infty$  respectively define the central charges of the  $\mathfrak{psu}(1, 1|2)$  (only for  $c = 0$ ) and  $\mathcal{N} = 2, d = 4$  Poincaré superalgebras (here  $A, B = 1, 2$ ; recall Eq. (3.34), which actually holds for any static, spherically symmetric, asymptotically flat, extremal BH)

$$c = 0 : \{Q_\alpha^A, Q_\beta^B\}_{\mathfrak{psu}(1,1|2)} = \varepsilon_{\alpha\beta} \varepsilon^{AB} Z(\phi_H(p, q), p, q); \quad (7.25)$$

$$c \in \mathbb{R}^+ : \{Q_\alpha^A, Q_\beta^B\}_{\mathcal{N}=2, d=4 \text{ Poincaré}} = \varepsilon_{\alpha\beta} \varepsilon^{AB} Z(\phi_\infty, p, q). \quad (7.26)$$

Notice that the central charges  $Z_1$  and  $Z_2$  are real, because we have dropped the axion field  $a$ , consistently with the charge constraint  $p \cdot q = 0$ .





preserving features will be discussed further below) the BH entropy reads ( $R_{+,c=0}^2 \equiv R_H^2$ )

$$\begin{aligned}
S_{BH,c=0} &= \frac{A_{H,c=0}}{4} = \pi R_H^2 = \pi [M^2 - \Sigma^2(\phi_\infty, p, q)]_{c=0} = \\
&= \pi [Z_1(\phi_\infty, q, p) + Z_2(\phi_\infty, q, p)] [Z_1(\phi_\infty, q, p) - Z_2(\phi_\infty, q, p)] = \\
&= 2\pi |pq|, \tag{7.36}
\end{aligned}$$

consistently with the result (6.18). Let us remark once again that *the dependence on the asymptotical value  $\phi_\infty$  of the dilaton drops out from the expression of the BH entropy  $S_{BH}|_{c=0}$ !*

For the chosen BH charge configuration, the considered extremal BH is a *non-degenerate*  $\frac{1}{4}$ -BPS state in *pure*  $\mathcal{N} = 4$ ,  $d = 4$  supergravity [88] (which actually is the unique *non-degenerate* state in such a theory), for which it holds

$$S_{BH,c=0}(p, q) = \pi Z_1^2(\phi_H(p, q), p, q); \tag{7.37}$$

$$Z_2(\phi_H(p, q), p, q) = 0. \tag{7.38}$$

It is worth pointing out that such a result can be generalized for the *non-degenerate*  $\frac{1}{\mathcal{N}}$ -BPS states in all  $\mathcal{N} > 2$ -extended,  $d = 4$  pure supergravities as follows [8]:

$$S_{BH,c=0}(p, q) = \pi |Z_1|^2(\phi_H(p, q), p, q); \tag{7.39}$$

$$Z_2(\phi_H(p, q), p, q) = \dots = Z_{[\mathcal{N}/2]}(\phi_H(p, q), p, q) = 0, \tag{7.40}$$

where  $Z_1, Z_2, \dots, Z_{[\mathcal{N}/2]}$  are the eigenvalues of the central charge matrix  $Z_{AB}$  ( $A, B = 1, \dots, \mathcal{N}$ ), and the square brackets denote the integer part.

In the  $\mathcal{N} = 2$ ,  $d = 4$ ,  $n_V = 1$  *minimally coupled* supergravity interpretation, one obtains that the  $\frac{1}{4}$ -BPS  $\mathcal{N} = 4$ ,  $d = 4$  extremal BH treated above corresponds to both  $\frac{1}{2}$ -BPS and non-BPS ( $Z = 0$ )  $\mathcal{N} = 2$ ,  $d = 4$  extremal BHs. Indeed, the criticality condition for  $V_{BH}$  reads

$$\begin{aligned}
\partial_\phi V_{BH} &= 2Z\partial_\phi Z + 2(\partial_\phi Z)\partial_\phi^2 Z = 4Z\partial_\phi Z = \\
&= 2(P + Q)(P - Q) = 2(pe^{-\phi} + qe^\phi)(pe^{-\phi} - qe^\phi) = 0, \tag{7.41}
\end{aligned}$$

where the relation

$$\partial_\phi^2 Z = Z, \tag{7.42}$$

yielded by the definitions (7.20) and (7.21) and by the identifications (7.24), was used.

Thus, the extremal BH attractors can be classified as follows:

$$\mathcal{N} = 2, d = 4, \frac{1}{2}\text{-BPS} :$$

$$\left\{ \begin{array}{l} Z \left( \phi_{H, \frac{1}{2}\text{-BPS}}(p, q), p, q \right) \equiv Z_{\frac{1}{2}\text{-BPS}}(p, q) \neq 0; \\ (\partial_\phi Z) \left( \phi_{H, \frac{1}{2}\text{-BPS}}(p, q), p, q \right) = 0 \Leftrightarrow \begin{cases} \phi_{H, \frac{1}{2}\text{-BPS}}(p, q) = \frac{1}{2} \ln \left( \frac{p}{q} \right); \\ pq > 0; \end{cases} \end{array} \right. \quad (7.43)$$

$$\Downarrow \\ Z_{\frac{1}{2}\text{-BPS}}(p, q) = \sqrt{2} p e^{-\phi_{H, \frac{1}{2}\text{-BPS}}(p, q)} = \text{sgn}(p) \sqrt{2pq} = \text{sgn}(p) \sqrt{2|pq|}; \quad (7.44)$$

$$\Downarrow \\ S_{BH, \frac{1}{2}\text{-BPS}}(p, q) = \pi V_{BH} \left( \phi_{H, \frac{1}{2}\text{-BPS}}(p, q), p, q \right) = \pi Z_{\frac{1}{2}\text{-BPS}}^2(p, q) = 2\pi pq = 2\pi |pq|; \quad (7.45)$$

$$\mathcal{N} = 2, d = 4, \text{non-BPS} (Z = 0) :$$

$$\left\{ \begin{array}{l} (\partial_\phi Z) \left( \phi_{H, \text{non-BPS}}(p, q), p, q \right) \equiv (\partial_\phi Z)_{\text{non-BPS}}(p, q) \neq 0; \\ Z \left( \phi_{H, \text{non-BPS}}(p, q), p, q \right) = 0 \Leftrightarrow \begin{cases} \phi_{H, \text{non-BPS}}(p, q) = \frac{1}{2} \ln \left( -\frac{p}{q} \right); \\ pq < 0; \end{cases} \end{array} \right. \quad (7.46)$$

$$\Downarrow \\ (\partial_\phi Z)_{\text{non-BPS}}(p, q) = \text{sgn}(p) \sqrt{-2pq} = \text{sgn}(p) \sqrt{2|pq|}; \quad (7.47)$$

$$\Downarrow \\ S_{BH, \text{non-BPS}}(p, q) = \pi V_{BH} \left( \phi_{H, \text{non-BPS}}(p, q), p, q \right) = \pi (\partial_\phi Z)_{\text{non-BPS}}^2(p, q) = -2\pi pq = 2\pi |pq|. \quad (7.48)$$

Thus, for the considered  $\mathcal{N} = 2$  BH charge configuration  $(p^1, p^2, q_1, q_2) : p \cdot q = 0$ , one obtains that:

$$\exp \left[ -2\phi_{H, \frac{1}{2}\text{-BPS}}(p, q) \right] = \exp \left[ -2\phi_{H, \text{non-BPS}}(p, q) \right] = \left| \frac{q}{p} \right|; \quad (7.49)$$

$$Z_{\frac{1}{2}\text{-BPS}}(p, q) = (\partial_\phi Z)_{\text{non-BPS}}(p, q) = \text{sgn}(p) \sqrt{2|pq|}; \quad (7.50)$$

$$S_{BH, \frac{1}{2}\text{-BPS}}(p, q) = S_{BH, \text{non-BPS}}(p, q) = 2\pi |pq|. \quad (7.51)$$

The unique discrimination between the two classes of attractors is due to the range of the supporting BH charges:  $pq > 0$  supports  $\frac{1}{2}$ -BPS attractors, whereas  $pq < 0$  supports the non-BPS ( $Z = 0$ ) ones. The case  $pq = 0$  must be disregarded, because it yields  $A_{H, \frac{1}{2}\text{-BPS}} = A_{H, \text{non-BPS}} = 0$  (“small” extremal BH), and it does not allow to consistently stabilize the dilaton at the BH horizon, as well.

Now, the first branch of Eq. (7.34) trivially yields the saturation of the  $c = 0$  limit of the (guessed generalization of the) BPS bound (5.34) which, through the asymptotical relation (5.2), yields the  $c = 0$  limit of the bound (5.37). It is thus clear that the first branch of Eqs. (7.34) and (7.35) corresponds to the  $\mathcal{N} = 2$ ,  $d = 4$   $\frac{1}{2}$ -BPS attractor solution given by Eqs. (7.43) and (7.45).

It remains to show that for the second branch, which thus<sup>20</sup> necessarily belongs to the  $\mathcal{N} = 2$ ,  $d = 4$  non-BPS ( $Z = 0$ ) attractor solution given by Eqs. (7.46) and (7.48), the strict inequality of the  $c = 0$  limit of the bound (5.34) (and thus of the bound (5.37)) holds. We can do this even without solving analytically the equation of motion for the dilaton  $\phi(r)$  in the extremal case (see *e.g.* - the first Ref. of - [92]). Indeed, by using the definition (6.4) and Eq. (6.15), and recalling that only its “+” branch is admissible (see below Eq. (6.18)), one gets the following expression of the extremal ADM squared mass:

$$M^2 = \frac{(|P_\infty| + |Q_\infty|)^2}{2} = \frac{(|p|e^{-\phi_\infty} + |q|e^{\phi_\infty})^2}{2}. \quad (7.52)$$

On the other hand, Eqs. (7.20) and (7.24) yield the following expression for the asymptotical limit of the squared absolute value of the *central charge function* (*i.e.* for the central charge of the  $\mathcal{N} = 2$ ,  $d = 4$  Poincaré superalgebra):

$$|Z|^2(\phi_\infty, p, q) = Z^2(\phi_\infty, p, q) = \frac{(P_\infty + Q_\infty)^2}{2} = \frac{(pe^{-\phi_\infty} + qe^{\phi_\infty})^2}{2}. \quad (7.53)$$

Thus, it is immediate to conclude that, depending on the (reciprocal) signs of the BH supporting charges  $p$  and  $q$ , it holds that

$$\begin{aligned} pq > 0 : M^2 &= |Z|^2(\phi_\infty, p, q); \\ pq < 0 : M^2 &> |Z|^2(\phi_\infty, p, q). \end{aligned} \quad (7.54)$$

Such a result is consistent with the fact that, as given by Eqs. (7.43)-(7.45) and (7.46)-(7.48),  $pq > 0$  supports the  $\frac{1}{2}$ -BPS attractors, whereas  $pq < 0$  supports the non-BPS ( $Z = 0$ ) ones. Summarizing, one can finally state that the (first) branch  $M = |Z|(\phi_\infty, p, q)$  of Eq. (7.34) is the  $\frac{1}{2}$ -BPS one, the (second) branch  $M = |\partial_\phi Z|(\phi_\infty, p, q)$  of Eq. (7.34) is the non-BPS ( $Z = 0$ ) one.

In general, it seems conceivable that (as stated above and in [18]) the BPS bounds (5.34) and (5.37) generally hold in extremal case ( $c = 0$ ), *regardless* of the asymptotical values  $\phi_\infty^a$  of the scalar fields.

---

<sup>20</sup>We assume the *absence* of more than one *basin of attraction* in the radial dynamics of the dilaton in the considered extremal dilaton BH supported by electric and magnetic charges constrained by  $p \cdot q = 0$  [81, 92].

As a further illustrative example, let us consider the *double-extremal* dilaton BH (which indeed is a possible solution of the equation of motion [81, 92]), *i.e.* for the case in which

$$\frac{d\phi}{dr} = 0 \quad \forall r \in [r_H, \infty) \quad (7.55)$$

$$\begin{cases} \frac{1}{2}\text{-BPS} : \phi(r) = \phi_\infty = \phi_{H,BPS}(p, q) = \frac{1}{2} \ln\left(\frac{p}{q}\right), \quad pq > 0; \\ \text{non-BPS} (Z = 0) : \phi(r) = \phi_\infty = \phi_{H,non-BPS}(p, q) = \frac{1}{2} \ln\left(-\frac{p}{q}\right), \quad pq < 0. \end{cases} \quad (7.56)$$

Let us start by considering the  $\frac{1}{2}$ -BPS case; since  $|\partial_\phi Z|(\phi_\infty, p, q) = |\partial_\phi Z|(\phi_{H,BPS}, p, q) = 0$ , it is clear that the second branch of Eq. (7.34) *cannot* be the BPS one, because it would yield  $M = 0$ , which, within the assumptions formulated (see [100] and Refs. therein), holds iff the space-time is (globally) flat, which is *not* the case. Thus, it holds that

$$M_{\frac{1}{2}\text{-BPS}}(\phi_\infty, p, q) = M_{\frac{1}{2}\text{-BPS}}(\phi_{H,BPS}(p, q), p, q) = |Z|(\phi_{H,BPS}(p, q), p, q) = \sqrt{2pq}. \quad (7.57)$$

The reasoning gets mirrored for the non-BPS ( $Z = 0$ ) case, for which  $|Z|(\phi_\infty, p, q) = |Z|(\phi_{H,non-BPS}, p, q) = 0$ . Consequently, since the considered metric background is not (globally) flat, the second branch of Eq. (7.34) *cannot* be the non-BPS ( $Z = 0$ ) one. Also, the check of the second relation of (7.54) in such a case is trivial, since

$$\begin{aligned} M_{non-BPS}(\phi_\infty, p, q) &= M_{non-BPS}(\phi_{H,non-BPS}(p, q), p, q) = \\ &= |\partial_\phi Z|(\phi_{H,non-BPS}(p, q), p, q) = \\ &= \sqrt{-2pq} > |Z|(\phi_{H,non-BPS}(p, q), p, q) = 0. \end{aligned} \quad (7.58)$$

On the other hand, Eqs. (7.21) and (7.24) yield the following expression for the asymptotical limit of the squared absolute value of the derivative of the *central charge function* (*i.e.* for the so-called asymptotical *matter charge*):

$$\begin{aligned} |D_\phi Z|^2(\phi_\infty, p, q) &= |\partial_\phi Z|^2(\phi_\infty, p, q) = (\partial_\phi Z)^2(\phi_\infty, p, q) = \\ &= \frac{(-P_\infty + Q_\infty)^2}{2} = \frac{(-pe^{-\phi_\infty} + qe^{\phi_\infty})^2}{2}. \end{aligned} \quad (7.59)$$

By comparing such a result with Eq. (7.53), one thus obtains

$$\begin{aligned} pq > 0 : |Z|^2(\phi_\infty, p, q) &> |D_\phi Z|^2(\phi_\infty, p, q); \\ pq < 0 : |D_\phi Z|^2(\phi_\infty, p, q) &> |Z|^2(\phi_\infty, p, q). \end{aligned} \quad (7.60)$$

Finally, let us consider the generic, *non-extremal* case ( $c \neq 0$ ). By recalling Eqs. (6.6), (7.27) and (7.28), one gets ( $|Z|_\infty^2 \equiv |Z|^2(\phi_\infty, q, p)$ ,  $|D_\phi Z|_\infty^2 \equiv |D_\phi Z|^2(\phi_\infty, q, p)$ ); recall

that  $\partial_\phi Z = D_\phi Z$ , due to the considered BH charge configuration such that  $p \cdot q = 0$ )

$$M^2 = V_{BH}(\phi_\infty, q, p) - \Sigma^2 + c^2 = |Z|_\infty^2 + |D_\phi Z|_\infty^2 - \frac{|Z|_\infty^2 |D_\phi Z|_\infty^2}{M} + c^2;$$

$$\Updownarrow$$

$$M_\pm^2 = \frac{|Z|_\infty^2}{2} \left\{ \left[ 1 + \frac{(|D_\phi Z|_\infty^2 + c^2)}{|Z|_\infty^2} \right] \left[ 1 \pm \sqrt{1 - \left[ \frac{2|Z|_\infty |D_\phi Z|_\infty}{|Z|_\infty^2 + |D_\phi Z|_\infty^2 + c^2} \right]^2} \right] \right\}, \quad (7.61)$$

where  $|Z|_\infty^2 > 0$  (otherwise, *i.e.* for  $|Z|_\infty^2 = 0$ , the strict inequality of BPS bounds (5.34) and (5.37) holds) and the necessary condition

$$[|Z|_\infty^2 + |D_\phi Z|_\infty^2 + c^2]^2 - 4|Z|_\infty^2 |D_\phi Z|_\infty^2 \geq 0, \quad (7.62)$$

were assumed to hold. Eqs. (7.61) implies that in the generic *non-extremal* case ( $c \neq 0$ , within  $|Z|_\infty^2 > 0$  and the consistency condition (7.62)) the BPS bounds (5.34) and (5.37) do hold true depending on the asymptotical dilaton value  $\phi_\infty$  and on the value of the extremality parameter  $c$ . Notice that, in the extremal limit  $c \rightarrow 0$  one achieves the results obtained above.

In general, it seems conceivable that in the generic *non-extremal* case ( $c \neq 0$ , within suitable consistency conditions) the BPS bounds (5.34) and (5.37) hold true depending, *in a model-dependent way*, on the asymptotical values  $\phi_\infty^a$  of the scalar fields and on the very value of the extremality parameter  $c$ .

## 8 Rotating Attractors

Let us now briefly report some results about the *extremal* rotating case.

In the stationary rotating regime ( $J \neq 0$  constant), an asymptotically flat *extremal* BH has a near-horizon geometry  $AdS_2 \times S^1$  (see *e.g.* [26] and Refs. therein). The most general *Ansatz* for the near-horizon metric consistent with the  $SO(2, 1) \times SO(2)$  symmetry reads

$$ds^2 = v_1(\theta) \left( -r^2 dt^2 + \frac{dr^2}{r^2} \right) + d\theta^2 + v_2(\theta) (d\phi - \alpha r dt)^2, \quad (8.1)$$

where  $\alpha \in \mathbb{R}$ , and  $v_1(\theta)$  and  $v_2(\theta)$  are functions of the angle  $\theta \in [0, \pi]$ . By a reparametrization of such a coordinate, the metric can be recast in the following form [26]:

$$ds^2 = \Xi^2(\theta) e^{2\psi(\theta)} \left( -r^2 dt^2 + \frac{dr^2}{r^2} + d\theta^2 \right) + e^{-2\psi(\theta)} (d\phi - \alpha r dt)^2, \quad (8.2)$$

where  $\Xi(\theta)$  and  $\psi(\theta)$  are suitably defined functions of  $\theta$ . The resulting BH entropy reads

$$S_{BH} = 16\pi^2 a(p, q, J), \quad (8.3)$$

where  $a(p, q, J)$  is a certain function of the BH charges  $p^\Lambda$ ,  $q_\Lambda$  and of  $J$ . By the very definition (2.4) of *effective radius* (and recalling  $R_{+,c=0}^2 \equiv R_H^2$ ), it thus holds that

$$R_H^2(p, q, J) = 16\pi a(p, q, J). \quad (8.4)$$

The near-horizon *extremal* Kerr BH metric is given by

$$\alpha = 1; \quad (8.5)$$

$$\Xi(\theta) = \frac{J}{8\pi} \sin \theta; \quad (8.6)$$

$$e^{-2\psi(\theta)} = \frac{J}{4\pi} \frac{\sin^2 \theta}{1 + \cos^2 \theta}; \quad (8.7)$$

$$a = \frac{J}{8\pi}, \quad (8.8)$$

yielding [116]

$$S_{BH,Kerr} = 2\pi J \Leftrightarrow R_{H,Kerr}^2(J) = 2J. \quad (8.9)$$

On the other hand, the near-horizon *extremal* Kerr-Newman BH metric (with electric and magnetic charge  $q$  and  $p$ ) is given by

$$\alpha = \frac{J}{\sqrt{J^2 + \left(\frac{q^2+p^2}{8\pi}\right)^2}}; \quad (8.10)$$

$$\Xi(\theta) = a \sin \theta; \quad (8.11)$$

$$e^{-2\psi(\theta)} = \frac{2a \sin^2 \theta}{1 + \cos^2 \theta + \frac{(q^2+p^2)\sin^2 \theta}{64\pi^2 a}}; \quad (8.12)$$

$$a = \frac{\sqrt{J^2 + \left(\frac{q^2+p^2}{8\pi}\right)^2}}{8\pi}, \quad (8.13)$$

yielding [116]

$$S_{BH,Kerr-Newman} = 2\pi \sqrt{J^2 + \left(\frac{q^2+p^2}{8\pi}\right)^2}; \quad (8.14)$$

$\Downarrow$

$$R_{H,Kerr-Newman}^2(p, q, J) = 2\sqrt{J^2 + \left(\frac{q^2+p^2}{8\pi}\right)^2}. \quad (8.15)$$

Up to a rescaling of BH charges by  $\frac{1}{2\sqrt{\pi}}$  and an overall rescaling of  $ds^2$  by  $\frac{1}{16\pi}$ , in the static limit ( $J \rightarrow 0$ ), the near-horizon Kerr-Newman metric (8.10)-(8.13) gives back the BR metric (3.29) which, as shown above, is the near-horizon of every static, spherically symmetric, asymptotically flat, *extremal* BH (and in this particular case, of the RN extremal BH with electric and magnetic charge  $q$  and  $p$ ):

$$\alpha = 0, \quad \Xi(\theta) = a \sin \theta, \quad e^{-2\psi(\theta)} = a \sin^2 \theta, \quad a = \frac{(q^2+p^2)}{64\pi^2}; \quad (8.16)$$

$\Downarrow$

$$ds^2 = \frac{(q^2+p^2)}{64\pi^2} \left( -r^2 dt^2 + \frac{dr^2}{r^2} + d\Omega^2 \right) = \frac{1}{16\pi} ds_{BR}^2 \Big|_{M^2 = \frac{q^2+p^2}{4\pi}}. \quad (8.17)$$

In [26], beside the *double-extremal* case, various explicit examples of stationary, asymptotically flat rotating BHs with non-trivial scalar dynamics were treated. Among them, let us here mention the generalization of the *extremal* Kerr-Newman BH to supergravity given by heterotic compactification on a six-torus  $T^6$ . This in general determines *matter-coupled*  $\mathcal{N} = 4$ ,  $d = 4$  supergravity [111, 112], which can be consistently truncated to *pure*  $\mathcal{N} = 4$ ,  $d = 4$  supergravity (see discussion at the start of Sect. 6) or also to  $\mathcal{N} = 2$ ,  $d = 4$  reducible symmetric *cubic* sequence  $\frac{SU(1,1)}{U(1)} \times \frac{SO(2,n)}{SO(2) \times SO(n)}$ , in the symplectic basis with maximum non-compact manifest symmetry  $SO(2, n)$  [117, 118], in which the pre-potential does *not* exist at all (see *e.g.* [108] and Refs. therein). In the so-called *ergo-free* branch, the corresponding near-horizon metric reads [26]

$$ds^2 = \frac{1}{8\pi} \sqrt{\mathcal{I}_4 - J^2 \cos^2 \theta} \left( -r^2 dt^2 + \frac{dr^2}{r^2} + d\theta^2 \right) + \frac{1}{8\pi} \frac{(\mathcal{I}_4 - J^2)}{\sqrt{\mathcal{I}_4 - J^2 \cos^2 \theta}} \sin^2 \theta \left( d\phi + \frac{J}{\sqrt{\mathcal{I}_4 - J^2}} r dt \right)^2, \quad (8.18)$$

with entropy

$$S_{BH} = 2\pi \sqrt{\mathcal{I}_4 - J^2} \Leftrightarrow R_H^2 = 2\sqrt{\mathcal{I}_4 - J^2}, \quad (8.19)$$

where  $\mathcal{I}_4$  is the (unique) *quartic* invariant of the relevant  $U$ -duality group ( $SU(1, 1) \times SO(6, n)$  for  $\mathcal{N} = 4$ ,  $d = 4$  theory; see also the discussion in Sect. 6):

$$\mathcal{I}_4 \equiv p^2 q^2 - (p \cdot q)^2. \quad (8.20)$$

## 9 Black Hole Entropy and Quantum Entanglement

Sometimes, it happens that two very different areas of theoretical physics share the same mathematics. This may eventually lead to the realization that they are, in fact, dual descriptions of the same physical phenomena, or it may not. Anyway, this fact often leads to new insights in both areas. Recent papers [3] have established an intriguing analogy between the entropy of certain  $d = 4$  supersymmetric BHs in string theory and the entanglement measure in quantum information theory.

### $\mathcal{N} = 2$ , $d = 4$ *stu* Black Holes and the Tripartite Entanglement of Three Qubits

In quantum information theory, the three qubit system (Alice, Bob, Charlie) is described by the quantum state

$$|\Psi\rangle = a_{ABC} |ABC\rangle, \quad (9.1)$$

where  $A = 0, 1$ , so the corresponding Hilbert space has dimension  $2^3 = 8$ . The numbers  $a_{ABC} \in \mathbb{C}$  transforms as the fundamental representation  $(\mathbf{2}, \mathbf{2}, \mathbf{2})$  of  $SL(2, \mathbb{C})_A \times$

$SL(2, \mathbb{C})_B \times SL(2, \mathbb{C})_C$ . The so-called *tripartite entanglement* of the quantum state (9.1) is measured by the *3-tangle* [119, 120]

$$\tau_3(ABC) \equiv 4|Det(a_{ABC})|, \quad (9.2)$$

where  $Det(a_{ABC})$  is *Cayley's hyperdeterminant* [121]

$$Det(a_{ABC}) \equiv -\frac{1}{2}\epsilon^{A_1 A_2} \epsilon^{B_1 B_2} \epsilon^{A_3 A_4} \epsilon^{B_3 B_4} \epsilon^{C_1 C_4} \epsilon^{C_2 C_3} a_{A_1 B_1 C_1} a_{A_2 B_2 C_2} a_{A_3 B_3 C_3} a_{A_4 B_4 C_4}, \quad (9.3)$$

which is invariant under  $SL(2, \mathbb{C})_A \times SL(2, \mathbb{C})_B \times SL(2, \mathbb{C})_C$  and under a *triality* interchanging  $A$ ,  $B$  and  $C$  (-indices).

In the context of stringy BHs, the 8 components of  $a_{ABC}$  are the 4 electric and 4 magnetic charges of an extremal BH in the so-called (3-moduli) *stu* model of  $\mathcal{N} = 2$ ,  $d = 4$  supergravity [112, 122], and thus they take real (or integer, at the quantized level) values. The  $U$ -duality group of the *stu* model is  $(SU(1, 1))^3 \sim (SL(2, \mathbb{R}))^3$  (or  $(SL(2, \mathbb{Z}))^3$ , at the quantized level).

The Bekenstein-Hawking entropy  $S_{BH}$  of such a BH was firstly calculated in [122]. The connection to quantum information theory arises by noting (see the first Ref. of [3]) that  $S_{BH}$  can also be expressed in terms of *Cayley's hyperdeterminant*:

$$S_{BH} = \pi \sqrt{|Det(a_{ABC})|}. \quad (9.4)$$

Thus, one can establish a *dictionary* between the classification of various *entangled states* (the so-called *separable A-B-C*, *bipartite entangled of type A-BC*, *B-CA*, *C-AB*, *tripartite entangled W*, *tripartite entangled GHZ* ones) and the classification of various *small* and *large* BPS and non-BPS *stu* BHs [3]. For example, the *canonical GHZ* state [123]

$$|\Psi\rangle \sim \frac{1}{\sqrt{2}} (|111\rangle + |000\rangle), \quad (9.5)$$

having  $Det(a_{ABC}) > 0$  corresponds to a *large* non-BPS *stu* BH. The *tripartite entangled W* state

$$|\Psi\rangle \sim \frac{1}{\sqrt{3}} (|100\rangle + |010\rangle + |001\rangle) \quad (9.6)$$

having  $Det(a_{ABC}) = 0$  corresponds to a *small* BPS *stu* BH. The *GHZ* state

$$|\Psi\rangle = \frac{1}{2} (-|000\rangle + |011\rangle + |101\rangle + |110\rangle) \quad (9.7)$$

having  $Det(a_{ABC}) < 0$  corresponds to a *large* BPS *stu* BH.

## $\mathcal{N} = 2$ , $d = 4$ Axion-Dilaton Black Holes and the Bipartite Entanglement of Two Qubits

An even structurally simpler framework (see the second Ref. of [3]) in which the analogy between stringy BHs and quantum information theory works is provided by the two qubit system (Alice and Bob), described by the quantum state

$$|\Psi\rangle = a_{AB} |AB\rangle, \quad (9.8)$$



where  $A = 0, 1$ , so the corresponding Hilbert space has dimension  $2^2 = 4$ . The numbers  $a_{AB} \in \mathbb{C}$  transform as the fundamental representation  $(\mathbf{2}, \mathbf{2})$  of  $SL(2, \mathbb{C})_A \times SL(2, \mathbb{C})_B$ . The so-called *bipartite entanglement* of the quantum state (9.8) is measured by the *2-tangle*

$$\tau_2(AB) \equiv C^2(AB), \quad (9.9)$$

where

$$C(AB) \equiv 2|\det(a_{AB})| \quad (9.10)$$

is the so-called *concurrence*.  $\det(a_{AB})$  is invariant under  $SL(2, \mathbb{C})_A \times SL(2, \mathbb{C})_B$  and under a *duality* interchanging  $A$  and  $B$ (-indices).

In the context of stringy BHs, the 4 components of  $a_{AB}$  are the 2 electric and 2 magnetic charges of an extremal BH in the so-called (1-modulus)  $t^2$  model of  $\mathcal{N} = 2$ ,  $d = 4$  supergravity. Thus, in such an interpretation the  $a_{AB}$ 's take real (or integer, at the quantized level) values. As pointed out in Sects. 6 and 7, the  $t^2$  model also corresponds to the  $(U(1))^6 \rightarrow (U(1))^2$  truncation of the so-called Maxwell-Einstein-axion-dilaton BH, with entropy

$$S_{BH} = \pi|\det(a_{AB})|. \quad (9.11)$$

For example, the so-called *Bell* state

$$|\Psi\rangle = \frac{1}{\sqrt{2}}(|00\rangle + |11\rangle) \quad (9.12)$$

with  $\det(a_{AB}) > 0$  corresponds to a *large* non-BPS axion-dilaton BH.

## $\mathcal{N} = 8$ , $d = 4$ Black Holes and the Tripartite Entanglement of Seven Qubits

In  $\mathcal{N} = 8$ ,  $d = 4$  supergravity the (BH) charge vector contains 28 electric and 28 magnetic charges, and it sits in the fundamental representation  $\mathbf{56}$  of the  $U$ -duality group  $E_{7(7)}$ . The BH hole entropy reads [124]

$$S_{BH} = \pi\sqrt{|\mathcal{J}_4|}, \quad (9.13)$$

where  $\mathcal{J}_4$  is Cartan's quartic  $E_7$ -invariant [125], which can be written as follows:

$$\mathcal{J}_4 = P^{ij}Q_{jk}P^{kl}Q_{li} - \frac{1}{4}P^{ij}Q_{ij}P^{kl}Q_{kl} + \frac{1}{96}(\epsilon^{ijklmnop}Q_{ij}Q_{kl}Q_{mn}Q_{op} + \epsilon_{ijklmnop}P^{ij}P^{kl}P^{mn}P^{op}), \quad (9.14)$$

$P^{ij}$  and  $Q_{jk}$  being  $8 \times 8$  antisymmetric charge matrices.

The qubit interpretation of  $\mathcal{J}_4$  arises from realizing the existence of the following embedding:

$$E_7(\mathbb{C}) \supseteq [SL(2, \mathbb{C})]^7, \quad (9.15)$$

under which the  $\mathbf{56}$  of  $E_7(\mathbb{C})$  branches as

$$\begin{aligned} \mathbf{56} \rightarrow & (\mathbf{2}, \mathbf{2}, \mathbf{1}, \mathbf{2}, \mathbf{1}, \mathbf{1}, \mathbf{1}) + (\mathbf{1}, \mathbf{2}, \mathbf{2}, \mathbf{1}, \mathbf{2}, \mathbf{1}, \mathbf{1}) + (\mathbf{1}, \mathbf{1}, \mathbf{2}, \mathbf{2}, \mathbf{1}, \mathbf{2}, \mathbf{1}) + (\mathbf{1}, \mathbf{1}, \mathbf{1}, \mathbf{2}, \mathbf{2}, \mathbf{1}, \mathbf{2}) + \\ & + (\mathbf{2}, \mathbf{1}, \mathbf{1}, \mathbf{1}, \mathbf{2}, \mathbf{2}, \mathbf{1}) + (\mathbf{1}, \mathbf{2}, \mathbf{1}, \mathbf{1}, \mathbf{1}, \mathbf{2}, \mathbf{2}) + (\mathbf{2}, \mathbf{1}, \mathbf{2}, \mathbf{1}, \mathbf{1}, \mathbf{1}, \mathbf{2}). \end{aligned} \quad (9.16)$$




Figure 4: **The  $E_7$  entanglement diagram.** Each of the seven vertices A,B,C,D,E,F,G represents a qubit and each of the seven triangles ABD, BCE, CDF, DEG, EFA, FGB, GAC describes a *tripartite entanglement*. [3]

Such a branching decomposition suggests the introduction of a system of seven qubits (Alice, Bob, Charlie, Daisy, Emma, Fred and George), described by the quantum state

$$|\Psi\rangle = a_{ABD} |ABD\rangle + b_{BCE} |BCE\rangle + c_{CDF} |CDF\rangle + d_{DEG} |DEG\rangle + e_{EFA} |EFA\rangle + f_{FGB} |FGB\rangle + g_{GAC} |GAC\rangle, \quad (9.17)$$

where  $A = 0, 1$ , yielding that the corresponding Hilbert space has dimension  $7 \times 2^3 = 56$ . The  $a, b, c, d, e, f, g$  transform as a **56** of  $E(\mathbb{C})$ .

The entanglement of the system of seven qubits can be represented by a heptagon where the vertices A,B,C,D,E,F,G represent the seven qubits and the seven triangles ABD, BCE, CDF, DEG, EFA, FGB, GAC represent the *tripartite entanglement* (see Fig. 4). Alternatively, one can use the *Fano plane*, (see Fig. 5) corresponding to the multiplication table of the octonions  $\mathbb{O}$ .

The so-called *tripartite entanglement of the seven qubits system* described by the state (9.17) is measured by the *3-tangle*

$$\tau_3(ABCDEFG) \equiv 4|\mathcal{J}_4|, \quad (9.18)$$




Figure 5: The *Fano plane* has seven points, representing the seven qubits, and seven lines (the circle counts as a line) with three points on every line, representing the *tripartite entanglement*, and three lines through every point. [3]

with

$$\mathcal{J}_4 \sim \begin{cases} a^4 + b^4 + c^4 + d^4 + e^4 + f^4 + g^4 + \\ +2[a^2b^2 + b^2c^2 + c^2d^2 + d^2e^2 + e^2f^2 + f^2g^2 + g^2a^2 + a^2c^2 + b^2d^2 + c^2e^2 + \\ +d^2f^2 + e^2g^2 + f^2a^2 + g^2b^2 + a^2d^2 + b^2e^2 + c^2f^2 + d^2g^2 + e^2a^2 + f^2b^2 + g^2c^2] + \\ +8[bcdf + cdeg + defa + efgb + fgac + gabd + abce], \end{cases} \quad (9.19)$$

where products like

$$\begin{aligned} a^4 &\equiv (ABD)(ABD)(ABD)(ABD) = \\ &= \epsilon^{A_1A_2} \epsilon^{B_1B_2} \epsilon^{D_1D_4} \epsilon^{A_3A_4} \epsilon^{B_3B_4} \epsilon^{D_2D_3} a_{A_1B_1D_1} a_{A_2B_2D_2} a_{A_3B_3D_3} a_{A_4B_4D_4} \end{aligned} \quad (9.20)$$

exclude four individuals (here Charlie, Emma, Fred and George), products like

$$\begin{aligned} a^2b^2 &\equiv \frac{1}{2}(ABD)(ABD)(BCE)(BCE) = \\ &= \frac{1}{2}\epsilon^{A_1A_2}\epsilon^{B_1B_3}\epsilon^{D_1D_2}\epsilon^{B_2B_4}\epsilon^{C_3C_4}\epsilon^{E_3E_4}a_{A_1B_1D_1}a_{A_2B_2D_2}b_{B_3C_3E_3}b_{B_4C_4E_4} \end{aligned} \quad (9.21)$$

exclude two individuals (here Charlie and Emma), and products like

$$\begin{aligned} abce &\equiv (ABD)(BCE)(CDF)(EFA) = \\ &= \epsilon^{A_1A_4}\epsilon^{B_1B_2}\epsilon^{C_2C_3}\epsilon^{D_1D_3}\epsilon^{E_2E_4}\epsilon^{F_3F_4}a_{A_1B_1D_1}b_{B_2C_2E_2}c_{C_3D_3F_3}e_{E_4F_4A_4} \end{aligned} \quad (9.22)$$

exclude one individual (here George).

Analogously to the cases treated above, *large* non-BPS, *small* BPS and *large* ( $\frac{1}{8}$ -)BPS  $\mathcal{N} = 8$ ,  $d = 4$  BHs correspond to states with  $\mathcal{J}_4 < 0$ ,  $\mathcal{J}_4 = 0$  and  $\mathcal{J}_4 > 0$ , respectively.

## 10 Recent Developments: Moduli Spaces of Attractors

In  $\mathcal{N} = 2$  homogeneous (not necessarily symmetric) and  $\mathcal{N} > 2$ -extended,  $d = 4$  supergravities the Hessian matrix of  $V_{BH}$  at its critical points is in general semi-positive definite, eventually with some vanishing eigenvalues (*massless Hessian modes*), which actually are *flat* directions of  $V_{BH}$  itself [39, 42]. Thus, it can be stated that for all supergravities based on homogeneous scalar manifolds the critical points of  $V_{BH}$  which are *non-degenerate* (*i.e.* for which it holds  $V_{BH} \neq 0$ ) all are *stable*, up to some eventual *flat* directions.

The Attractor Equations of  $\mathcal{N} = 2$ ,  $d = 4$  supergravity coupled to  $n_V$  Abelian vector multiplets may have *flat* directions in the non-BPS cases [39, 42], but *not* in the  $\frac{1}{2}$ -BPS one [84]. Indeed, in the  $\frac{1}{2}$ -BPS case (satisfying  $Z \neq 0$ ,  $D_i Z = 0 \forall i = 1, \dots, n_V$ ) the covariant  $2n_V \times 2n_V$  Hessian matrix of  $V_{BH}$  reads [84]

$$\left( D_{\hat{i}} D_{\hat{j}} V_{BH} \right)_{\mathcal{N}=2, \frac{1}{2}\text{-BPS}} = \frac{1}{2} |Z|_{\frac{1}{2}\text{-BPS}} \begin{pmatrix} 0 & G_{i\bar{j}} \\ G_{j\bar{i}} & 0 \end{pmatrix}_{\frac{1}{2}\text{-BPS}}, \quad (10.1)$$

where hatted indices can be either holomorphic or anti-holomorphic; thus, as far as the metric  $G_{i\bar{j}}$  of the scalar manifold is strictly positive definite, Eq. (10.1) yields that no *massless*  $\frac{1}{2}$ -BPS *Hessian modes* arise out.

Tables 1 and 2 respectively list the moduli spaces of non-BPS  $Z \neq 0$  and non-BPS  $Z = 0$  attractors for homogeneous symmetric  $\mathcal{N} = 2$ ,  $d = 4$  special geometries, for which a complete classification is available [42]. Notice that the non-BPS  $Z \neq 0$  moduli spaces are nothing but the symmetric real special scalar manifolds of the corresponding  $\mathcal{N} = 2$ ,  $d = 5$  supergravity.

Within the symmetric  $\mathcal{N} = 2$ ,  $d = 4$  supergravities, there are some remarkable models in which no non-BPS *flat* directions exist at all.

The unique  $n_V = 1$  models are the so-called  $t^2$  and  $t^3$  models; they are based on the rank-1 scalar manifold  $\frac{SU(1,1)}{U(1)}$ , but with different holomorphic prepotential functions.

	$\frac{\widehat{H}}{h}$	r	$dim_{\mathbb{R}}$
$\mathbb{R} \oplus \Gamma_n, n \in \mathbb{N}$	$SO(1, 1) \otimes \frac{SO(1, n-1)}{SO(n-1)}$	$1(n = 1)$ $2(n \geq 2)$	$n$
$J_3^{\mathbb{O}}$	$\frac{E_{6(-26)}}{F_{4(-52)}}$	2	6
$J_3^{\mathbb{H}}$	$\frac{SU^*(6)}{USp(6)}$	2	14
$J_3^{\mathbb{C}}$	$\frac{SL(3, \mathbb{C})}{SU(3)}$	2	8
$J_3^{\mathbb{R}}$	$\frac{SL(3, \mathbb{R})}{SO(3)}$	2	5

Table 1: **Moduli spaces of non-BPS  $Z \neq 0$  critical points of  $V_{BH, \mathcal{N}=2}$  in  $\mathcal{N} = 2, d = 4$  homogeneous symmetric supergravities. They are the  $\mathcal{N} = 2, d = 5$  homogeneous symmetric real special manifolds [42]**

As mentioned at the end of Sect. 6, the  $t^2$  model is the first element ( $n = 1$ ) of the sequence of irreducible symmetric special Kähler manifolds  $\frac{SU(1, n)}{U(1) \times SU(n)}$  ( $n_V = n, n \in \mathbb{N}$ ) (see *e.g.* [25] and Refs. therein), endowed with *quadratic* prepotential. Let us recall once again that the bosonic sector of the  $t^2$  model is given by the  $(U(1))^6 \rightarrow (U(1))^2$  truncation of Maxwell-Einstein-axion-dilaton (super)gravity, treated in Sects. 6 and 7. On the other hand, the  $t^3$  model has *cubic* prepotential; it is an *isolated case* in the classification of symmetric special Kähler manifolds (see *e.g.* [115]), but can be thought also as the  $s = t = u$  *degeneration* of the  $stu$  model. It is worth pointing out that the  $t^2$  and  $t^3$  models are based on the same rank-1 special Kähler manifold, with different constant scalar curvature, which respectively can be computed to be (see *e.g.* [35] and Refs. therein)

$$\begin{aligned} \frac{SU(1,1)}{U(1)}, t^2 \text{ model} : \mathcal{R} = -2; \\ \frac{SU(1,1)}{U(1)}, t^3 \text{ model} : \mathcal{R} = -\frac{2}{3}. \end{aligned} \tag{10.2}$$

Beside the BPS attractors, the  $t^2$  model admits only non-BPS  $Z = 0$  critical points of  $V_{BH}$  with no *flat* directions. Analogously, the  $t^3$  model admits only non-BPS  $Z \neq 0$  critical points of  $V_{BH}$  with no *flat* directions.

For  $n_V > 1$ , the non-BPS  $Z \neq 0$  critical points of  $V_{BH}$ , if any, all have *flat* directions, and thus a related moduli space (see Table 1). However, models with no non-BPS  $Z = 0$  *flat* directions at all and  $n_V > 1$  exist, namely they are the first and second element ( $n = 1$ ,

	$\frac{\tilde{H}}{h} = \frac{\tilde{H}}{h' \otimes U(1)}$	r	$dim_{\mathbb{C}}$
<i>quadratic sequence</i> $n \in \mathbb{N}$	$\frac{SU(1,n-1)}{U(1) \otimes SU(n-1)}$	1	$n - 1$
$\mathbb{R} \oplus \Gamma_n, n \in \mathbb{N}$	$\frac{SO(2,n-2)}{SO(2) \otimes SO(n-2)}, n \geq 3$	$1(n = 3)$ $2(n \geq 4)$	$n - 2$
$J_3^{\mathbb{O}}$	$\frac{E_{6(-14)}}{SO(10) \otimes U(1)}$	2	16
$J_3^{\mathbb{H}}$	$\frac{SU(4,2)}{SU(4) \otimes SU(2) \otimes U(1)}$	2	8
$J_3^{\mathbb{C}}$	$\frac{SU(2,1)}{SU(2) \otimes U(1)} \otimes \frac{SU(1,2)}{SU(2) \otimes U(1)}$	2	4
$J_3^{\mathbb{R}}$	$\frac{SU(2,1)}{SU(2) \otimes U(1)}$	1	2

Table 2: **Moduli spaces of non-BPS  $Z = 0$  critical points of  $V_{BH, \mathcal{N}=2}$  in  $\mathcal{N} = 2, d = 4$  homogeneous symmetric supergravities. They are (non-special) homogeneous symmetric Kähler manifolds [42]**

2) of the sequence of reducible symmetric special Kähler manifolds  $\frac{SU(1,1)}{U(1)} \times \frac{SO(2,n)}{SO(2) \times SO(n)}$  ( $n_V = n+1, n \in \mathbb{N}$ ) (see *e.g.* [25] and Refs. therein), *i.e.* the so-called  $st^2$  and  $stu$  models, respectively. The  $stu$  model, which is relevant for the analogy between stringy extremal BHs and quantum information theory treated in Sect. 9, has two non-BPS  $Z \neq 0$  *flat* directions, spanning the moduli space  $SO(1,1) \times SO(1,1)$  (*i.e.* the scalar manifold of the  $stu$  model in  $d = 5$ ), but *no* non-BPS  $Z = 0$  *massless Hessian modes* at all. On the other hand, the  $st^2$  model (which can be thought as the  $t = u$  *degeneration* of the  $stu$  model) has one non-BPS  $Z \neq 0$  *flat* direction, spanning the moduli space  $SO(1,1)$  (*i.e.* the scalar manifold of the  $st^2$  model in  $d = 5$ ), but *no* non-BPS  $Z = 0$  *flat* direction at all. The  $st^2$  is the "*smallest*" symmetric model exhibiting a non-BPS  $Z \neq 0$  *flat* direction.

Concerning the "*smallest*" symmetric models exhibiting a non-BPS  $Z = 0$  *flat* direction they are the second ( $n = 2$ ) element of the sequence  $\frac{SU(1,n)}{U(1) \times SU(n)}$  and the third ( $n = 3$ ) element of the sequence  $\frac{SU(1,1)}{U(1)} \times \frac{SO(2,n)}{SO(2) \times SO(n)}$ . In both cases, the unique non-BPS  $Z = 0$  *flat* direction spans the non-BPS  $Z = 0$  moduli space  $\frac{SU(1,1)}{U(1)} \sim \frac{SO(2,1)}{SO(2)}$  (see Table 2), whose local geometrical properties however differ in the two cases (for the same reasons

	$\frac{1}{\mathcal{N}}$ -BPS orbits $\frac{G}{\mathcal{H}}$	non-BPS, $Z_{AB} \neq 0$ orbits $\frac{G}{\mathcal{H}}$	non-BPS, $Z_{AB} = 0$ orbits $\frac{G}{\mathcal{H}}$
$\mathcal{N} = 3$	$\frac{SU(3,n)}{SU(2,n)}$	—	$\frac{SU(3,n)}{SU(3,n-1)}$
$\mathcal{N} = 4$	$\frac{SU(1,1)}{U(1)} \otimes \frac{SO(6,n)}{SO(4,n)}$	$\frac{SU(1,1)}{SO(1,1)} \otimes \frac{SO(6,n)}{SO(5,n-1)}$	$\frac{SU(1,1)}{U(1)} \otimes \frac{SO(6,n)}{SO(6,n-2)}$
$\mathcal{N} = 5$	$\frac{SU(1,5)}{SU(3) \otimes SU(2,1)}$	—	—
$\mathcal{N} = 6$	$\frac{SO^*(12)}{SU(4,2)}$	$\frac{SO^*(12)}{SU^*(6)}$	$\frac{SO^*(12)}{SU(6)}$
$\mathcal{N} = 8$	$\frac{E_{7(7)}}{E_{6(2)}}$	$\frac{E_{7(7)}}{E_{6(6)}}$	—

Table 3: **Charge orbits of the real, symplectic  $R$  representation of the  $U$ -duality group  $G$  supporting BH attractors with non-vanishing entropy in  $3 \leq \mathcal{N} \leq 8$ ,  $d = 4$  supergravities [60]**

holding for the  $t^2$  and  $t^3$  models treated above).

In  $\mathcal{N} > 2$ -extended,  $d = 4$  supergravities there are *flat* directions of  $V_{BH}$  at both its *non-degenerate* BPS and non-BPS critical points. Group-theoretically, this is due to the fact that the corresponding supporting BH *charge orbits* always have a *non-compact* stabilizer [42, 60]. The BPS *flat* directions can be interpreted in terms of left-over hypermultiplets' scalar degrees of freedom in the truncation down to the  $\mathcal{N} = 2$ ,  $d = 4$  theories [126, 39]. In Tables 3 and 4 all *charge orbits* and the corresponding moduli spaces of attractor solution in  $\mathcal{N} > 2$ -extended,  $d = 4$  supergravities are reported [60].

We conclude by pointing out that all the reported results hold at the classical, Einstein supergravity level. It is conceivable that the *flat* directions of classical *non-degenerate* extremal BH attractors will be removed (*i.e.* lifted) by quantum (perturbative and non-perturbative) corrections (such as the ones coming from higher-order derivative contributions to the gravity and/or gauge sector) to the *classical* effective BH potential  $V_{BH}$ . Consequently, *at the quantum* (perturbative and non-perturbative) *level*, *no moduli spaces for attractor solutions might exist at all* (and therefore also *the actual attractive nature of the critical points of  $V_{BH}$  might be destroyed*). *However, this might not be the case for  $\mathcal{N} = 8$ .*

In presence of *quantum* lifts of *classically flat* directions of the Hessian matrix of  $V_{BH}$  at its critical points, in order to answer to the key question: '*Do extremal BH attractors (in a strict sense) survive the quantum level?*', it is thus crucial to determine whether such lifts originate Hessian modes with *positive* squared mass (corresponding to *attractive*

	$\frac{1}{\mathcal{N}}$ -BPS moduli space $\frac{\mathcal{H}}{\mathfrak{h}}$	non-BPS, $Z_{AB} \neq 0$ moduli space $\frac{\widehat{\mathcal{H}}}{\widehat{\mathfrak{h}}}$	non-BPS, $Z_{AB} = 0$ moduli space $\frac{\widetilde{\mathcal{H}}}{\widetilde{\mathfrak{h}}}$
$\mathcal{N} = 3$	$\frac{SU(2,n)}{SU(2) \otimes SU(n) \otimes U(1)}$	—	$\frac{SU(3,n-1)}{SU(3) \otimes SU(n-1) \otimes U(1)}$
$\mathcal{N} = 4$	$\frac{SO(4,n)}{SO(4) \otimes SO(n)}$	$SO(1,1) \otimes \frac{SO(5,n-1)}{SO(5) \otimes SO(n-1)}$	$\frac{SO(6,n-2)}{SO(6) \otimes SO(n-2)}$
$\mathcal{N} = 5$	$\frac{SU(2,1)}{SU(2) \otimes U(1)}$	—	—
$\mathcal{N} = 6$	$\frac{SU(4,2)}{SU(4) \otimes SU(2) \otimes U(1)}$	$\frac{SU^*(6)}{USp(6)}$	—
$\mathcal{N} = 8$	$\frac{E_{6(2)}}{SU(6) \otimes SU(2)}$	$\frac{E_{6(6)}}{USp(8)}$	—

Table 4: **Moduli spaces of BH attractors with non-vanishing entropy in  $3 \leq \mathcal{N} \leq 8$ ,  $d = 4$  supergravities** ( $\mathfrak{h}$ ,  $\widehat{\mathfrak{h}}$  and  $\widetilde{\mathfrak{h}}$  are maximal compact subgroups of  $\mathcal{H}$ ,  $\widehat{\mathcal{H}}$  and  $\widetilde{\mathcal{H}}$ , respectively) [60]

directions) or with *negative* squared mass (*i.e. tachyonic, repeller* directions).

The fate of the unique non-BPS  $Z \neq 0$  flat direction of the  $st^2$  model in presence of the most general class of quantum perturbative corrections consistent with the axionic-shift symmetry has been studied in [127], showing that, as intuitively expected, the *classical solutions get lifted at the quantum level*. Interestingly, in [127] it is found the *quantum lift occurs more often towards repeller directions* (thus destabilizing the whole critical solution, and *destroying the attractor in strict sense*), rather than towards *attractive* directions. The same behavior may be expected for the unique non-BPS  $Z = 0$  flat direction of the  $n = 2$  element of the quadratic irreducible sequence and the  $n = 3$  element of the cubic reducible sequence (see above).

Generalizing to the presence of more than one *flat* direction, this would mean that *only a (very) few classical attractors do remain attractors in strict sense at the quantum level*; consequently, *at the quantum* (perturbative and non-perturbative) *level the “landscape” of extremal BH attractors should be strongly constrained and reduced*.

## Acknowledgments

The original parts of the contents of these lectures result from collaborations with L. Andrianopoli, S. Bellucci, R. D’Auria, E. Gimon, M. Trigiante, and especially M. J. Duff, G. Gibbons, M. Günaydin, R. Kallosh and A. Strominger, which are gratefully



acknowledged.

A. M. would like to thank the Department of Physics, Theory Unit Group at CERN, where part of this work was done, for kind hospitality and stimulating environment.

The work of S.F. has been supported in part by the European Community Human Potential Programme under contract MRTN-CT-2004-503369 “*Constituents, Fundamental Forces and Symmetries of the Universe*” in association with LNF-INFN and in part by DOE grant DE-FG03-91ER40662, Task C.

The work of K.H. and A.M. has been supported by Museo Storico della Fisica e Centro Studi e Ricerche “*Enrico Fermi*”, Rome, Italy, in association with INFN-LNF.

## References

- [1] B. De Witt, C. De Witt eds.: *Black Holes* (Gordon and Breach, New York, 1973); S. W. Hawking, W. Israel: *General Relativity* (Cambridge University Press, Cambridge, 1979); R. M. Wald: *General Relativity* (University of Chicago Press, Chicago, 1984).
- [2] G. W. Moore: *Les Houches lectures on strings and arithmetic*, arXiv:hep-th/0401049; M. R. Douglas, R. Reinbacher, S. T. Yau: *Branes, bundles and attractors: Bogomolov and beyond*, arXiv:math.ag/0604597.
- [3] M. J. Duff, *String Triality, Black Hole Entropy and Cayley’s Hyperdeterminant*, Phys. Rev. **D76**, 025017 (2007), hep-th/0601134; R. Kallosh, A. Linde: *Strings, black holes, and quantum information*, Phys. Rev. **D73**, 104033 (2006); P. Levey: *Stringy black holes and the geometry of entanglement*, Phys. Rev. **D74**, 024030 (2006); M. J. Duff, S. Ferrara: *E7 and the tripartite entanglement of seven qubits*, Phys. Rev. **D76**, 025018 (2007), arXiv:quant-ph/0609227; P. Levey: *Strings, black holes, the tripartite entanglement of seven qubits and the Fano plane*, Phys. Rev. **D75**, 024024 (2007), arXiv:hep-th/0610314; M. J. Duff and S. Ferrara, *Black hole entropy and quantum information*, arXiv:hep-th/0612036; L. Borsten, D. Dahanayake, M. J. Duff, W. Rubens and H. Ebrahim: *Wrapped branes as qubits*, arXiv:0802.0840.
- [4] S. W. Hawking, R. Penrose: *The singularities of gravitational collapse and cosmology*, Proc. Roy. Soc. Lond. **A314**, 529 (1970).
- [5] R. Penrose: *Gravitational collapse: the role of general relativity*, Riv. Nuovo Cim. **1**, 252 (1969); Gen. Rel. Grav. **34**, 1141 (2002); R. Penrose: in *General Relativity, an Einstein Centenary Survey*, ed. by S. W. Hawking, W. Israel (Cambridge University Press, Cambridge, 1979).
- [6] G. Gibbons: in *Unified theories of Elementary Particles. Critical Assessment and Prospects*, Proceedings of the Heisenberg Symposium, Munchen, Germany, 1981, ed. by P. Breitenlohner, H. P. Durr, Lecture Notes in Physics **160** (Springer-Verlag, Berlin, 1982); G. W. Gibbons: in *Supersymmetry, Supergravity and Related Topics*, Proceedings of the XVth GIFT International Physics (Girona, Spain 1984), ed. by F. del Aguila, J. de Azcárraga, L. Ibáñez, (World Scientific, Singapore, 1985), p.

- 147; P. Breitenlohner, D. Maison, G. W. Gibbons: *4-dimensional black holes from Kaluza-Klein theories*, Commun. Math. Phys. **120**, 295 (1988); R. Kallosh: *Supersymmetric black holes*, Phys. Lett. **B282**, 80 (1992); R. R. Khuri, T. Ortín: *Supersymmetric black holes in  $\mathcal{N}=8$  supergravity*, Nucl. Phys. **B467**, 355 (1996); A. Sen: *Black-Hole Solutions in Heterotic String Theory on a Torus*, Nucl. Phys. **B440**, 421 (1995); A. Sen: *Quantization of dyon charge and electric-magnetic duality in string theory*, Phys. Lett. **B303**, 22 (1993); A. Sen: *Extremal Black-Holes and Elementary String States*, Mod. Phys. Lett. **A10**, 2081 (1995); M. Cvetič, C. M. Hull: *Black holes and U-duality*, Nucl. Phys. **B480**, 296 (1996); M. Cvetič, I. Gaida: *Duality-invariant non-extreme black holes in toroidally compactified string theory*, Nucl. Phys. **B505**, 291 (1997); M. Cvetič, D. Youm: *General Static Spherically Symmetric Black Holes of Heterotic String on a Six Torus*, arXiv:hep-th/9512127; M. Cvetič and A. A. Tseytlin: *Solitonic strings and BPS saturated dyonic black holes*, Phys. Rev. **D53**, 5619 (1996) [Erratum-ibid. **D55**, 3907 (1997)], hep-th/9512031.
- [7] S. Ferrara, R. Kallosh, A. Strominger:  *$\mathcal{N}=2$  extremal black holes*, Phys. Rev. **D52**, 5412 (1995).
- [8] S. Ferrara, R. Kallosh: *Supersymmetry and attractors*, Phys. Rev. **D54**, 1514 (1996); S. Ferrara, R. Kallosh: *Universality of supersymmetric attractors*, Phys. Rev. **D54**, 1525 (1996).
- [9] A. Strominger: *Macroscopic entropy of  $\mathcal{N}=2$  extremal black holes*, Phys. Lett. **B383**, 39 (1996).
- [10] A. Sen, *Black Hole Entropy Function and the Attractor Mechanism in Higher Derivative Gravity*, JHEP **09**, 038 (2005), hep-th/0506177.
- [11] K. Goldstein, N. Iizuka, R. P. Jena and S. P. Trivedi, *Non-Supersymmetric Attractors*, Phys. Rev. **D72**, 124021 (2005), hep-th/0507096.
- [12] A. Sen, *Entropy Function for Heterotic Black Holes*, JHEP **03**, 008 (2006), hep-th/0508042.
- [13] R. Kallosh, *New Attractors*, JHEP **0512**, 022 (2005), hep-th/0510024.
- [14] P. K. Tripathy and S. P. Trivedi, *Non-Supersymmetric Attractors in String Theory*, JHEP **0603**, 022 (2006), hep-th/0511117.
- [15] A. Giriyavets, *New Attractors and Area Codes*, JHEP **0603**, 020 (2006), hep-th/0511215.
- [16] K. Goldstein, R. P. Jena, G. Mandal and S. P. Trivedi, *A C-Function for Non-Supersymmetric Attractors*, JHEP **0602**, 053 (2006), hep-th/0512138.
- [17] M. Alishahiha and H. Ebrahim, *Non-supersymmetric attractors and entropy function*, JHEP **0603**, 003 (2006), hep-th/0601016.
- [18] R. Kallosh, N. Sivanandam and M. Soroush, *The Non-BPS Black Hole Attractor Equation*, JHEP **0603**, 060 (2006), hep-th/0602005.

- [19] B. Chandrasekhar, S. Parvizi, A. Tavanfar and H. Yavartanoo, *Non-supersymmetric attractors in  $\mathcal{R}^2$  gravities*, JHEP **0608**, 004 (2006), [hep-th/0602022](#).
- [20] J. P. Hsu, A. Maloney and A. Tomasiello, *Black Hole Attractors and Pure Spinors*, JHEP **0609**, 048 (2006), [hep-th/0602142](#).
- [21] S. Bellucci, S. Ferrara and A. Marrani, *On some properties of the Attractor Equations*, Phys. Lett. **B635**, 172 (2006), [hep-th/0602161](#).
- [22] S. Bellucci, S. Ferrara and A. Marrani, *Supersymmetric Mechanics. Vol.2: The Attractor Mechanism and Space-Time Singularities* (LNP **701**, Springer-Verlag, Heidelberg, 2006).
- [23] S. Ferrara and R. Kallosh, *On  $\mathcal{N}=8$  attractors*, Phys. Rev. D **73**, 125005 (2006), [hep-th/0603247](#).
- [24] M. Alishahiha and H. Ebrahim, *New attractor, Entropy Function and Black Hole Partition Function*, JHEP **0611**, 017 (2006), [hep-th/0605279](#).
- [25] S. Bellucci, S. Ferrara, M. Günaydin and A. Marrani, *Charge Orbits of Symmetric Special Geometries and Attractors*, Int. J. Mod. Phys. **A21**, 5043 (2006), [hep-th/0606209](#).
- [26] D. Astefanesei, K. Goldstein, R. P. Jena, A. Sen and S. P. Trivedi, *Rotating Attractors*, JHEP **0610**, 058 (2006), [hep-th/0606244](#).
- [27] R. Kallosh, N. Sivanandam and M. Soroush, *Exact Attractive non-BPS STU Black Holes*, Phys. Rev. **D74**, 065008 (2006), [hep-th/0606263](#).
- [28] P. Kaura and A. Misra, *On the Existence of Non-Supersymmetric Black Hole Attractors for Two-Parameter Calabi-Yau's and Attractor Equations*, Fortsch. Phys. **54**, 1109 (2006), [hep-th/0607132](#).
- [29] G. L. Cardoso, V. Grass, D. Lüst and J. Perz, *Extremal non-BPS Black Holes and Entropy Extremization*, JHEP **0609**, 078 (2006), [hep-th/0607202](#).
- [30] J. F. Morales and H. Samtleben, *Entropy function and attractors for AdS black holes*, JHEP **0610**, 074 (2006), [hep-th/0608044](#).
- [31] S. Bellucci, S. Ferrara, A. Marrani and A. Yeranyan, *Mirror Fermat Calabi-Yau Threefolds and Landau-Ginzburg Black Hole Attractors*, Riv. Nuovo Cim. **029**, 1 (2006), [hep-th/0608091](#).
- [32] D. Astefanesei, K. Goldstein and S. Mahapatra, *Moduli and (un)attractor black hole thermodynamics*, [hep-th/0611140](#).
- [33] G.L. Cardoso, B. de Wit and S. Mahapatra, *Black hole entropy functions and attractor equations*, JHEP **0703**, 085 (2007) [hep-th/0612225](#).
- [34] R. D'Auria, S. Ferrara and M. Trigiante, *Critical points of the Black-Hole potential for homogeneous special geometries*, JHEP **0703**, 097 (2007), [hep-th/0701090](#).

- [35] S. Bellucci, S. Ferrara and A. Marrani, *Attractor Horizon Geometries of Extremal Black Holes*, contribution to the Proceedings of the XVII SIGRAV Conference, 4–7 September 2006, Turin, Italy, [hep-th/0702019](#).
- [36] A. Ceresole and G. Dall’Agata, *Flow Equations for Non-BPS Extremal Black Holes*, *JHEP* **0703**, 110 (2007), [hep-th/0702088](#).
- [37] L. Andrianopoli, R. D’Auria, S. Ferrara and M. Trigiante, *Black Hole Attractors in  $\mathcal{N}=1$  Supergravity*, *JHEP* **0707**, 019 (2007), [hep-th/0703178](#).
- [38] K. Saraikin and C. Vafa, *Non-supersymmetric Black Holes and Topological Strings*, *Class. Quant. Grav.* **25**, 095007 (2008), [hep-th/0703214](#).
- [39] S. Ferrara and A. Marrani,  *$\mathcal{N}=8$  non-BPS Attractors, Fixed Scalars and Magic Supergravities*, *Nucl. Phys.* **B788**, 63 (2008), [arXiv:0705.3866](#).
- [40] S. Nampuri, P. K. Tripathy and S. P. Trivedi, *On The Stability of Non-Supersymmetric Attractors in String Theory*, *JHEP* **0708**, 054 (2007), [arXiv:0705.4554](#).
- [41] L. Andrianopoli, R. D’Auria, E. Orazi, M. Trigiante, *First Order Description of Black Holes in Moduli Space*, *JHEP* **0711**, 032 (2007), [arXiv:0706.0712](#).
- [42] S. Ferrara and A. Marrani, *On the Moduli Space of non-BPS Attractors for  $\mathcal{N}=2$  Symmetric Manifolds*, *Phys. Lett.* **B652**, 111 (2007), [arXiv:0706.1667](#).
- [43] D. Astefanesei and H. Yavartanoo, *Stationary black holes and attractor mechanism*, *Nucl. Phys.* **B794**, 13 (2008), [arXiv:0706.1847](#).
- [44] G. L. Cardoso, A. Ceresole, G. Dall’Agata, J. M. Oberreuter, J. Perz, *First-order flow equations for extremal black holes in very special geometry*, *JHEP* **0710**, 063 (2007), [arXiv:0706.3373](#).
- [45] A. Misra and P. Shukla, *‘Area codes’, large volume (non-)perturbative alpha-prime and instanton: Corrected non-supersymmetric (A)dS minimum, the ‘inverse problem’ and ‘fake superpotentials’ for multiple-singular-loci-two-parameter Calabi-Yau’s*, [arXiv:0707.0105](#).
- [46] A. Ceresole, S. Ferrara and A. Marrani, *4d/5d Correspondence for the Black Hole Potential and its Critical Points*, *Class. Quant. Grav.* **24**, 5651 (2007), [arXiv:0707.0964](#).
- [47] M. M. Anber and D. Kastor, *The Attractor mechanism in Gauss-Bonnet gravity*, *JHEP* **0710**, 084 (2007), [arXiv:0707.1464](#).
- [48] Y. S. Myung, Y.-W. Kim and Y.-J. Park, *New attractor mechanism for spherically symmetric extremal black holes*, *Phys. Rev.* **D76**, 104045 (2007), [arXiv:0707.1933](#).
- [49] S. Bellucci, A. Marrani, E. Orazi and A. Shcherbakov, *Attractors with Vanishing Central Charge*, *Phys. Lett.* **B655**, 185 (2007), [arXiv:0707.2730](#).

- [50] K. Hotta and T. Kubota, *Exact Solutions and the Attractor Mechanism in Non-BPS Black Holes*, Prog. Theor. Phys. **118N5**, 969 (2007), [arXiv:0707.4554](#).
- [51] X. Gao, *Non-supersymmetric Attractors in Born-Infeld Black Holes with a Cosmological Constant*, JHEP **0711**, 006 (2007), [arXiv:0708.1226](#).
- [52] S. Ferrara and A. Marrani, *Black Hole Attractors in Extended Supergravity*, contribution to the Proceedings of *13th International Symposium on Particles, Strings and Cosmology (PASCOS 07)*, London, England, 2-7 Jul 2007, AIP Conf. Proc. **957**, 58 (2007), [arXiv:0708.1268](#).
- [53] A. Sen, *Black Hole Entropy Function, Attractors and Precision Counting of Microstates*, [arXiv:0708.1270](#).
- [54] A. Belhaj, L.B. Drissi, E.H. Saidi and A. Segui,  *$\mathcal{N}=2$  Supersymmetric Black Attractors in Six and Seven Dimensions*, Nucl. Phys. **B796**, 521 (2008), [arXiv:0709.0398](#).
- [55] L. Andrianopoli, S. Ferrara, A. Marrani and M. Trigiante, *Non-BPS Attractors in 5d and 6d Extended Supergravity*, Nucl. Phys. **B795**, 428 (2008), [arXiv:0709.3488](#).
- [56] D. Gaiotto, W. Li and M. Padi, *Non-Supersymmetric Attractor Flow in Symmetric Spaces*, JHEP **0712**, 093 (2007), [arXiv:0710.1638](#).
- [57] S. Bellucci, S. Ferrara, A. Marrani and A. Shcherbakov, *Splitting of Attractors in 1-modulus Quantum Corrected Special Geometry*, JHEP **0802**, 088 (2008), [arXiv:0710.3559](#).
- [58] E. G. Gimon, F. Larsen and J. Simon, *Black Holes in Supergravity: the non-BPS Branch*, JHEP **0801**, 040 (2008), [arXiv:0710.4967](#).
- [59] D. Astefanesei, H. Nastase, H. Yavartanoo and S. Yun, *Moduli flow and non-supersymmetric AdS attractors*, JHEP **0804**, 074 (2008), [arXiv:0711.0036](#).
- [60] S. Bellucci, S. Ferrara, R. Kallosh and A. Marrani, *Extremal Black Hole and Flux Vacua Attractors*, contribution to the Proceedings of the Winter School on Attractor Mechanism 2006 (SAM2006), 20-24 March 2006, INFN-LNF, Frascati, Italy, [arXiv:0711.4547](#).
- [61] R.-G. Cai and D.-W. Pang, *A Note on exact solutions and attractor mechanism for non-BPS black holes*, JHEP **0801**, 046 (2008), [arXiv:0712.0217](#).
- [62] M. Huebscher, P. Meessen, T. Ortín and S. Vaulà, *Supersymmetric  $\mathcal{N}=2$  Einstein-Yang-Mills monopoles and covariant attractors*, [arXiv:0712.1530](#).
- [63] W. Li: *Non-Supersymmetric Attractors in Symmetric Coset Spaces*, contribution to the Proceedings of 3rd School on Attractor Mechanism (SAM 2007), Frascati, Italy, 18-22 Jun 2007, [arXiv:0801.2536](#).

- [64] S. Bellucci, S. Ferrara, A. Marrani and A. Yeranyan: *d = 4 Black Hole Attractors in  $\mathcal{N} = 2$  Supergravity with Fayet-Iliopoulos Terms*, arXiv:0802.0141.
- [65] E. H. Saidi, *BPS and non BPS 7D Black Attractors in M-Theory on K3*, arXiv:0802.0583.
- [66] E. H. Saidi, *On Black Hole Effective Potential in 6D/7D  $\mathcal{N} = 2$  Supergravity*, arXiv:0803.0827.
- [67] E. H. Saidi and A. Segui, *Entropy of Pairs of Dual Attractors in six and seven Dimensions*, arXiv:0803.2945.
- [68] S. Bellucci, S. Ferrara and A. Marrani, *Attractors in Black*, contribution to the Proceedings of the 3rd RTN Workshop “*Constituents, Fundamental Forces and Symmetries of the Universe*”, 1–5 October 2007, Valencia, Spain, arXiv:0805.1310.
- [69] H. Ooguri, A. Strominger and C. Vafa: *Black Hole Attractors and the Topological String*, Phys. Rev. **D70**, 106007 (2004), hep-th/0405146.
- [70] H. Ooguri, C. Vafa and E. Verlinde: *Hartle-Hawking wave-function for flux compactifications: the Entropic Principle*, Lett. Math. Phys. **74**, 311 (2005), hep-th/0502211.
- [71] M. Aganagic, A. Neitzke and C. Vafa: *BPS microstates and the open topological string wave function*, hep-th/0504054.
- [72] S. Gukov, K. Saraikin and C. Vafa: *The Entropic Principle and Asymptotic Freedom*, Phys. Rev. **D73**, 066010 (2006), hep-th/0509109.
- [73] For reviews on black holes in superstring theory see e.g.: J. M. Maldacena: *Black-Holes in String Theory*, hep-th/9607235; A. W. Peet: *TASI lectures on black holes in string theory*, arXiv:hep-th/0008241; B. Pioline: *Lectures on black holes, topological strings and quantum attractors*, Class. Quant. Grav. **23**, S981 (2006); A. Dabholkar: *Black hole entropy and attractors*, Class. Quant. Grav. **23**, S957 (2006).
- [74] For recent reviews see: J. H. Schwarz: *Lectures on superstring and M-theory dualities*, Nucl. Phys. Proc. Suppl. **B55**, 1 (1997); M. J. Duff: *M-theory (the theory formerly known as strings)*, Int. J. Mod. Phys. **A11**, 5623 (1996); A. Sen: *Unification of string dualities*, Nucl. Phys. Proc. Suppl. **58**, 5 (1997).
- [75] J. H. Schwarz, A. Sen: *Duality symmetries of 4D heterotic strings*, Phys. Lett. **B312**, 105 (1993); J. H. Schwarz, A. Sen: *Duality Symmetrical Actions*, Nucl. Phys. **B411**, 35 (1994).
- [76] M. Gasperini, J. Maharana, G. Veneziano: *From trivial to non-trivial conformal string backgrounds via  $O(d, d)$  transformations*, Phys. Lett. **B272**, 277 (1991); J. Maharana, J. H. Schwarz: *Noncompact Symmetries in String Theory*, Nucl. Phys. **B390**, 3 (1993).
- [77] E. Witten: *String Theory Dynamics in Various Dimensions*, Nucl. Phys. **B443**, 85 (1995).

- [78] J. H. Schwarz: *M-theory extensions of T duality*, arXiv:hep-th/9601077; C. Vafa: *Evidence for F-theory*, Nucl. Phys. **B469**, 403 (1996).
- [79] G. Nordström: *On the energy of gravitational field in Einstein's theory*, Proc. Kon. Ned. Akad. Wet. **20**, 1238 (1918); H. Reissner: *Über die Eigengravitation des elektrischen Feldes nach der Einsteinschen Theorie*, Ann. Physik **50**, 106 (1916).
- [80] L. Smarr: *Mass Formula for Kerr Black Hole*, Phys. Rev. Lett. **30**, 71 (1973).
- [81] R. Kallosh, T. Ortín: *Charge quantization of axion-dilaton black holes*, Phys. Rev. **D48**, 742 (1993); E. Bergshoeff, R. Kallosh, T. Ortín: *Stationary axion/dilaton solutions and supersymmetry*, Nucl. Phys. **B478**, 156 (1996).
- [82] G. Gibbons, R. Kallosh and B. Kol: *Moduli, Scalar Charges, and the First Law of Black Hole Thermodynamics*, Phys. Rev. Lett. **77**, 4992 (1996).
- [83] S. W. Hawking: *Gravitational Radiation from Colliding Black Holes*, Phys. Rev. Lett. **26**, 1344 (1971); J. D. Bekenstein: *Black Holes and Entropy*, Phys. Rev. **D7**, 2333 (1973).
- [84] S. Ferrara, G. W. Gibbons and R. Kallosh: *Black holes and critical points in moduli space*, Nucl. Phys. **B500**, 75 (1997).
- [85] B. Bertotti: *Uniform Electromagnetic Field in the Theory of General Relativity*, Phys. Rev. **116**, 1331 (1959); I. Robinson: *A solution of the Maxwell-Einstein equations*, Bull. Acad. Pol. Sci. Ser. Sci. Math. Astron. Phys. **7**, 351 (1959).
- [86] S. Ferrara and M. Günaydin: *Orbits of Exceptional Groups, Duality and BPS States in String Theory*, Int. J. Mod. Phys. **A13**, 2075 (1998), hep-th/9708025.
- [87] S. Ferrara and M. Günaydin: *Orbits and attractors for  $\mathcal{N}=2$  Maxwell-Einstein supergravity theories in five dimensions*, Nucl. Phys. **B759**, 1 (2006), hep-th/0606108.
- [88] S. Ferrara and J. M. Maldacena: *Branes, central charges and U-duality invariant BPS conditions*, Class. Quant. Grav. **15**, 749 (1998), hep-th/9706097.
- [89] L. Andrianopoli, R. D'Auria, S. Ferrara and M. Trigiante: *Extremal Black Holes in Supergravity*, in : *String Theory and Fundamental Interactions*, M. Gasperini and J. Maharana eds. (LNP, Springer, Berlin-Heidelberg, 2007), hep-th/0611345.
- [90] R. Arnowitt, S. Deser and C. W. Misner: *Canonical Variables for General Relativity*, Phys. Rev. **117**, 1595 (1960).
- [91] A. Strominger, C. Vafa: *Microscopic origin of the Bekenstein-Hawking entropy*, Phys. Lett. **B379**, 99 (1996); C. G. Callan, J. M. Maldacena: *Brane dynamics from the Born-Infeld action*, Nucl. Phys. **B472**, 591 (1996); G. T. Horowitz, A. Strominger: *Counting States of Near-Extremal Black Holes*, Phys. Rev. Lett. **77**, 2368 (1996); R. Dijkgraaf, E. P. Verlinde, H. L. Verlinde: *BPS spectrum of the five-brane and black hole entropy*, Nucl. Phys. **B486**, 77 (1997); D. M. Kaplan,

- D. A. Lowe, J. M. Maldacena, A. Strominger: *Microscopic entropy of  $\mathcal{N}=2$  extremal black holes*, Phys. Rev. **D55**, 4898 (1997); J. M. Maldacena:  *$\mathcal{N}=2$  extremal black holes and intersecting branes*, Phys. Lett. **B403**, 20 (1997); J. M. Maldacena, A. Strominger, E. Witten: *Black hole entropy in M-Theory*, JHEP **9712**, 002 (1997); M. Bertolini, M. Trigiante: *Microscopic entropy of the most general four-dimensional BPS black hole*, JHEP **0010**, 002 (2000).
- [92] R. Kallosh, A. D. Linde, T. Ortín, A. W. Peet, A. Van Proeyen: *Supersymmetry as a cosmic censor*, Phys. Rev. **D46**, 5278 (1992); R. Kallosh, T. Ortín, A. W. Peet: *Entropy and action of dilaton black holes*, Phys. Rev. **D47**, 5400 (1993); R. Kallosh, A. W. Peet: *Dilaton black holes near the horizon*, Phys. Rev. **D46**, 5223 (1992).
- [93] J. F. Luciani: *Coupling of  $O(2)$  Supergravity with Several Vector Multiplets*, Nucl. Phys. **B132**, 325 (1978).
- [94] The literature on this topic is quite extended. As a general review, see *e.g.* the lecture notes: K. Stelle: *Lectures on Supergravity p-Branes*, presented at 1996 ICTP Summer School, Trieste, [arXiv:hep-th/9701088](https://arxiv.org/abs/hep-th/9701088).
- [95] M. J. Duff, R. R. Khuri, J. X. Lu: *String solitons*, Phys. Rep. **259**, 213 (1995).
- [96] G. W. Gibbons, P. K. Townsend: *Vacuum interpolation in supergravity via super p-branes.*, Phys.Rev.Lett.71, 3754, (1993).
- [97] S. W. Hawking: *The conservation of matter in general relativity*, Comm. Math. Phys. **18**, 301 (1970).
- [98] S. Deser and C. Teitelboim: *Supergravity Has Positive Energy*, Phys. Rev. Lett. **39**, 249 (1977).
- [99] J. M. Nester: *A new gravitational energy expression with a simple positivity proof*, Phys. Lett. **A83**, 241 (1981).
- [100] G.W.Gibbons and C.H.Hull: *A Bogomol'ny bound for general relativity and solitons in  $\mathcal{N}=2$  supergravity*, Phys. Lett. **B109**, 190 (1982).
- [101] C. M. Miller, K. Schalm and E. J. Weinberg: *Nonextremal black holes are BPS*, Phys. Rev. **D76**, 044001 (2007), [hep-th/0612308](https://arxiv.org/abs/hep-th/0612308).
- [102] A. Papapetrou: Proc. R. Irish Acad. **A51**, 191 (1947); S. D. Majumdar: Phys. Rev. **72**, 930 (1947).
- [103] R. D'Auria, S. Ferrara and P. Fré, *Special and Quaternionic Isometries: General Couplings in  $\mathcal{N}=2$  Supergravity and the Scalar Potential*, Nucl. Phys. **B359**, 705 (1991); L. Andrianopoli, M. Bertolini, A. Ceresole, R. D'Auria, S. Ferrara, P. Frè and T. Magri,  *$\mathcal{N}=2$  Supergravity and  $\mathcal{N}=2$  Super Yang-Mills Theory on General Scalar Manifolds : Symplectic Covariance, Gaugings and the Momentum Map*, J. Geom. Phys. **23**, 111 (1997), [hep-th/9605032](https://arxiv.org/abs/hep-th/9605032); L. Andrianopoli, M. Bertolini, A. Ceresole, R. D'Auria, S. Ferrara and P. Fré, *General Matter Coupled  $\mathcal{N}=2$  Supergravity*, Nucl. Phys. **B476**, 397 (1996), [hep-th/9603004](https://arxiv.org/abs/hep-th/9603004).



- [104] K. P. Tod: *All metrics admitting super-covariantly constant spinors*, Phys. Lett. **B121**, 241 (1983).
- [105] J. L. Alvarez, H. Quevedo and A. Sanchez, *Unified geometric description of black hole thermodynamics*, Phys. Rev. **D77**, 084004 (2008), [arXiv:0801.2279](#).
- [106] G. Ruppeiner, *Thermodynamic curvature and phase transitions in Kerr-Newman black holes*, [arXiv:0802.1326](#).
- [107] G. Ruppeiner: *Stability and fluctuations in black hole thermodynamics*, Phys. Rev. **D75**, 024037 (2007).
- [108] A. Ceresole, R. D'Auria and S. Ferrara: *The Symplectic Structure of  $\mathcal{N}=2$  Supergravity and Its Central Extension*, Talk given at ICTP Trieste Conference on Physical and Mathematical Implications of Mirror Symmetry in String Theory, Trieste, Italy, 5-9 June 1995, Nucl. Phys. Proc. Suppl. **46** (1996), [hep-th/9509160](#).
- [109] M. M. Anber and D. Kastor: *C-Functions in Lovelock Gravity*, [arXiv:0802.1290](#).
- [110] E. Cremmer, J. Scherk and S. Ferrara,  *$SU(4)$  Invariant Supergravity Theory*, Phys. Lett. **B74**, 61 (1978).
- [111] J. Maharana and J. H. Schwarz, *Noncompact Symmetries in String Theory*, Nucl. Phys. **B390**, 3 (1993), [hep-th/9207016](#).
- [112] M. J. Duff, J. T. Liu and J. Rahmfeld: *Four-dimensional string/string/string triality*, Nucl. Phys. **B459**, 125 (1996), [hep-th/9508094](#).
- [113] C. M. Hull, P. K. Townsend: *Unity of Superstring Dualities*, Nucl. Phys. **B438**, 109 (1995).
- [114] D. Garfinkle, G. T. Horowitz and A. Strominger: *Charged black holes in string theory*, Phys. Rev. **D43**, 3140 (1991) [Erratum-ibid. **D45**, 3888 (1992)].
- [115] S. Cecotti, S. Ferrara and L. Girardello: *Geometry of Type II Superstrings and the Moduli of Superconformal Field Theories*, Int. J. Mod. Phys. **A4**, 2475 (1989).
- [116] J. M. Bardeen and G. T. Horowitz: *The extreme Kerr throat geometry: a vacuum analog of  $AdS_2 \times S^2$* , Phys. Rev. **D60**, 104030 (1999), [hep-th/9905099](#).
- [117] E. Calabi and E. Visentini, Ann. Math. **71**, 472 (1960).
- [118] A. Ceresole, R. D'Auria, S. Ferrara and A. Van Proeyen, *Duality Transformations in Supersymmetric Yang-Mills Theories coupled to Supergravity*, Nucl. Phys. **B444**, 92 (1995), [hep-th/9502072](#).
- [119] V. Coffman, J. Kundu and W. Wooters: *Distributed entanglement*, Phys. Rev. **A61** (2000) 52306, [arXiv:quant-ph/9907047](#).
- [120] A. Miyake and M. Wadati: *Multipartite entanglement and hyperdeterminants*, Quant. Info. Comp. **2** (Special), 540 (2002), [arXiv:quant-ph/0212146](#).

- [121] A. Cayley: *On the theory of linear transformations*, Camb. Math. J. **4**, 193 (1845).
- [122] K. Behrndt, R. Kallosh, J. Rahmfeld, M. Shmakova and W. K. Wong: *STU black holes and string triality*, Phys. Rev. **D54**, 6293 (1996), [arXiv:hep-th/9608059](#).
- [123] D. M. Greenberger, M. Horne and A. Zeilinger, in: *Bell's Theorem, Quantum Theory and Conceptions of the Universe*, ed. M. Kafatos (Kluwer, Dordrecht, 1989).
- [124] R. Kallosh and B. Kol:  *$E(7)$  symmetric area of the black hole horizon*, Phys. Rev. **D53**, 5344 (1996).
- [125] E. Cartan: *Œuvres complètes*, (Editions du Centre National de la Recherche Scientifique, Paris, 1984); E. Cremmer and B. Julia: *The  $SO(8)$  Supergravity*, Nucl. Phys. **B159**, 141 (1979).
- [126] L. Andrianopoli, R. D'Auria and S. Ferrara:  *$U$  invariants, black hole entropy and fixed scalars*, Phys. Lett. **B403**, 12 (1997), [hep-th/9703156](#).
- [127] S. Bellucci, S. Ferrara, A. Marrani and A. Shcherbakov, *Quantum Lift of Non-BPS Flat Directions*, to appear.