

RAMANUJAN-TYPE SUPERCONGRUENCES

WADIM ZUDILIN

ABSTRACT. We present several supercongruences that may be viewed as p -adic analogues of Ramanujan-type series for $1/\pi$ and $1/\pi^2$, and prove three of these examples.

1. INTRODUCTION

Throughout the paper, the letter p is used for primes. It was observed by L. Van Hamme [12] that several Ramanujan's and Ramanujan-like formulas [11] for $1/\pi$,

$$\begin{aligned} \sum_{n=0}^{\infty} \frac{(\frac{1}{2})_n^3}{n!^3} (4n+1)(-1)^n &= \frac{2}{\pi}, \\ \sum_{n=0}^{\infty} \frac{(\frac{1}{2})_n^3}{n!^3} (6n+1) \frac{1}{4^n} &= \frac{4}{\pi}, \\ \sum_{n=0}^{\infty} \frac{(\frac{1}{2})_n^3}{n!^3} (6n+1) \frac{(-1)^n}{8^n} &= \frac{2\sqrt{2}}{\pi}, \\ \sum_{n=0}^{\infty} \frac{(\frac{1}{2})_n^3}{n!^3} (42n+5) \frac{1}{64^n} &= \frac{16}{\pi}, \end{aligned} \tag{1}$$

admit very nice conjectural p -analogues:

$$\sum_{n=0}^{p-1} \frac{(\frac{1}{2})_n^3}{n!^3} (4n+1)(-1)^n \stackrel{?}{\equiv} \left(\frac{-1}{p}\right) p \pmod{p^3} \quad \text{for } p > 2, \tag{2}$$

$$\sum_{n=0}^{p-1} \frac{(\frac{1}{2})_n^3}{n!^3} (6n+1) \frac{1}{4^n} \stackrel{?}{\equiv} \left(\frac{-1}{p}\right) p \pmod{p^3} \quad \text{for } p > 2, \tag{3}$$

$$\sum_{n=0}^{p-1} \frac{(\frac{1}{2})_n^3}{n!^3} (6n+1) \frac{(-1)^n}{8^n} \stackrel{?}{\equiv} \left(\frac{-2}{p}\right) p \pmod{p^3} \quad \text{for } p > 2, \tag{4}$$

Date: October 14, 2008.

2000 *Mathematics Subject Classification.* Primary 11Y55, 33C20, 33F10; Secondary 11B65, 11D88, 11F33, 11F85, 11S80, 12H25, 40G99, 65-05, 65B10.

The work was supported by a fellowship of the Max Planck Institute for Mathematics (Bonn) and by project no. R-146-000-103-112 of the National University of Singapore Academic Research Fund.

$$\sum_{n=0}^{p-1} \frac{\left(\frac{1}{2}\right)_n^3}{n!^3} (42n+5) \frac{1}{64^n} \stackrel{?}{\equiv} 5 \left(\frac{-1}{p}\right) p \pmod{p^3} \quad \text{for } p > 2, \quad (5)$$

where $\left(\frac{\cdot}{p}\right)$ and $(a)_n = \Gamma(a+n)/\Gamma(n)$ denote the Legendre (quadratic residue) symbol and the Pochhammer symbol, respectively. Note that since the p -adic order of $\left(\frac{1}{2}\right)_n/n!$ is 1 for $n = \frac{p+1}{2}, \dots, p-1$, one may compute the sums in (2)–(5) for n up to $\frac{p-1}{2}$ (and this is how the four supercongruences¹ (2)–(5) were conjectured in [12]); for example, an equivalent form of (2) is [12, (B.2)]

$$\sum_{n=0}^{(p-1)/2} \frac{\left(\frac{1}{2}\right)_n^3}{n!^3} (4n+1) (-1)^n \stackrel{?}{\equiv} (-1)^{(p-1)/2} p \pmod{p^3} \quad \text{for } p > 2. \quad (6)$$

Van Hamme himself was able to prove (2) modulo p (not p^3); recently E. Mortenson [10] gave a proof for (2). As far as we know, no progress has been done towards proving the other congruences (3), (4), and (5).

Remark 1. In fact, one can state all these p -supercongruences as infinite sums by replacing $n!$ in the denominators of the summands by $(-1)^n \Gamma_p(n+1)$, where Γ_p is Morita's p -adic gamma function [8]. For instance, the congruence (2) takes the form

$$\sum_{n=0}^{\infty} \frac{\left(\frac{1}{2}\right)_n^3}{\Gamma_p(n+1)^3} (4n+1) \equiv (-1)^{(p-1)/2} p \pmod{p^3} \quad \text{for } p > 2. \quad (7)$$

It seems that *all* known Ramanujan-type formulas for $1/\pi$ admit similar p -analogues. One example, which is not given in the list [12], is an analogue of the formula

$$\sum_{n=0}^{\infty} \frac{\left(\frac{1}{2}\right)_n \left(\frac{1}{4}\right)_n \left(\frac{3}{4}\right)_n}{n!^3} (20n+3) \frac{(-1)^n}{2^{2n}} = \frac{8}{\pi}; \quad (8)$$

it reads as

$$\sum_{n=0}^{p-1} \frac{\left(\frac{1}{2}\right)_n \left(\frac{1}{4}\right)_n \left(\frac{3}{4}\right)_n}{n!^3} (20n+3) \frac{(-1)^n}{2^{2n}} \stackrel{?}{\equiv} 3 \left(\frac{-1}{p}\right) p \pmod{p^3} \quad \text{for } p > 2. \quad (9)$$

Even more, some recently discovered, by J. Guillera, formulas for $1/\pi^2$ (see [3]–[5] and [14]), like

$$\sum_{n=0}^{\infty} \frac{\left(\frac{1}{2}\right)_n^3 \left(\frac{1}{4}\right)_n \left(\frac{3}{4}\right)_n}{n!^5} (120n^2 + 34n + 3) \frac{1}{2^{4n}} = \frac{32}{\pi^2}, \quad (10)$$

possess such analogues as well:

$$\sum_{n=0}^{p-1} \frac{\left(\frac{1}{2}\right)_n^3 \left(\frac{1}{4}\right)_n \left(\frac{3}{4}\right)_n}{n!^5} (120n^2 + 34n + 3) \frac{1}{2^{4n}} \stackrel{?}{\equiv} 3p^2 \pmod{p^5} \quad \text{for } p > 2. \quad (11)$$

In Section 3 we present a (far from exhaustive) list of many other supercongruences which are analogues of the examples indicated in our recent mini-survey [14]; there one can find the analogue (31) of the formula for $1/\pi^3$ experimentally discovered by

¹A congruence modulo a prime p is said to be a supercongruence if it happens to hold modulo some higher power of p .

B. Gourevich (see [14]). Unfortunately, all these congruences remain conjectures, except the already mentioned example (2) and, as we show below, the congruences (9) and (11). We do not dispose of a suitable general method to prove our experimentally discovered supercongruences. But at least in the three cases we can use the WZ-method [2], [3], [5] which results in very simple proofs given in Section 2 of the following results.

Theorem 1. *The congruence in (6) is valid.*

Theorem 1 is the main result in [10], but our proof is essentially different from the one in [10] and shorter.

Theorem 2. *The following congruence is true:*

$$\sum_{n=0}^{p-1} \frac{\left(\frac{1}{2}\right)_n^3 \left(\frac{1}{2}\right)_{2n}}{n!^5} (120n^2 + 34n + 3) \frac{1}{2^{6n}} \equiv 3p^2 \pmod{p^5} \quad \text{for } p > 2; \quad (12)$$

it is equivalent to (11).

Theorem 3. *We have*

$$\sum_{n=0}^{p-1} \frac{\left(\frac{1}{2}\right)_n \left(\frac{1}{2}\right)_{2n}}{n!^3} (20n + 3) \frac{1}{2^{4n}} \equiv 3(-1)^{(p-1)/2} p \pmod{p^3} \quad \text{for } p > 2, \quad (13)$$

which is equivalent to (9).

2. PROOFS BY THE WZ-MACHINERY

The original argument of D. Zeilberger and S. Ekhad [2], applied there to proving the identity (1) (which, in fact, starts its history from Bauer's work [1] in 1859, much earlier than Ramunujan's birth), was later developed in Guillera's works [3], [5]. It is based on introducing suitable WZ-pairs and creative telescoping. This method has rather strong limitations (and it always requires a preliminary human guess) but it is the only one known so far for proving the hypergeometric formulas for $1/\pi^2$, like (10). In our proof of Theorems 1, 2, and 3 we do not need an intelligent guess, since we can simply borrow the Zeilberger–Ekhad–Guillera WZ-pairs for (1), (10), and (8), respectively.

Proof of Theorem 1. Introduce the rational functions

$$F(n, k) = (-1)^{n+k} (4n + 1) \frac{\left(\frac{1}{2}\right)_n^2 \left(\frac{1}{2}\right)_{n+k}}{\left(1\right)_n^2 \left(1\right)_{n-k} \left(\frac{1}{2}\right)_k^2},$$

$$G(n, k) = -\frac{(2n-1)^2}{2(n-k)(4n-3)} F(n-1, k) = (-1)^{n+k} \cdot 2 \cdot \frac{\left(\frac{1}{2}\right)_n^2 \left(\frac{1}{2}\right)_{n+k-1}}{\left(1\right)_{n-1}^2 \left(1\right)_{n-k} \left(\frac{1}{2}\right)_k^2}$$

in two parameters n and k . They form a WZ-pair, namely, they satisfy

$$F(n, k-1) - F(n, k) = G(n+1, k) - G(n, k). \quad (14)$$

Indeed,

$$\frac{F(n, k-1)}{F(n, k)} = -\frac{(k - \frac{1}{2})^2}{(n + k - \frac{1}{2})(n - k + 1)}, \quad \frac{F(n, k)}{G(n, k)} = \frac{(4n + 1)(n + k - \frac{1}{2})}{2n^2},$$

$$\frac{G(n, k+1)}{G(n, k)} = -\frac{(n + \frac{1}{2})^2(n + k - \frac{1}{2})}{n^2(n - k + 1)},$$

and it is routine to verify the identity

$$-\frac{(k - \frac{1}{2})^2(4n + 1)}{2(n - k + 1)n^2} - \frac{(4n + 1)(n + k - \frac{1}{2})}{2n^2} = -\frac{(n + \frac{1}{2})^2(n + k - \frac{1}{2})}{n^2(n - k + 1)} - 1$$

which is the result of division of both sides of (14) by $G(n, k)$.

Summing (14) over $n = 0, 1, \dots, \frac{p-1}{2}$, we obtain

$$\sum_{n=0}^{(p-1)/2} F(n, k-1) - \sum_{n=0}^{(p-1)/2} F(n, k) = G(\frac{p+1}{2}, k) - G(0, k) = G(\frac{p+1}{2}, k). \quad (15)$$

Furthermore, for $k = 1, 2, \dots, \frac{p-1}{2}$ we have

$$G(\frac{p+1}{2}, k) = (-1)^{(p+1)/2+k} \cdot 2 \cdot \left(\frac{(\frac{1}{2})_{(p-1)/2}}{(1)_{(p-1)/2}} \right)^2 \frac{(\frac{p-1}{2} + \frac{1}{2})^2 (\frac{1}{2})_{(p+1)/2+k-1}}{(1)_{(p+1)/2-k} (\frac{1}{2})_k^2}$$

$$= (-1)^{(p+1)/2+k} \cdot 2^{-p} \binom{p-1}{\frac{p-1}{2}}^2 p^2 \cdot \frac{(\frac{1}{2})_{(p+1)/2+k-1}}{(1)_{(p+1)/2-k} (\frac{1}{2})_k^2} \equiv 0 \pmod{p^3},$$

since $(\frac{1}{2})_{(p+1)/2+k-1}$ is divisible by $(\frac{1}{2})_{(p+1)/2}$, hence by p , while the denominator is coprime to p . Comparing this result with (15) we see that

$$\sum_{n=0}^{(p-1)/2} F(n, 0) \equiv \sum_{n=0}^{(p-1)/2} F(n, 1) \equiv \sum_{n=0}^{(p-1)/2} F(n, 2) \equiv \dots \equiv \sum_{n=0}^{(p-1)/2} F(n, \frac{p-1}{2}) \pmod{p^3}. \quad (16)$$

On the other hand,

$$\sum_{n=0}^{(p-1)/2} F(n, \frac{p-1}{2}) = F(\frac{p-1}{2}, \frac{p-1}{2}) = (4 \cdot \frac{p-1}{2} + 1) \frac{(\frac{1}{2})_{p-1}}{(1)_{(p-1)/2}^2}$$

$$= 2 \frac{(\frac{1}{2})_p}{(1)_{(p-1)/2}^2} = p \cdot 2^{-2(p-1)} \binom{2p-1}{p-1} \binom{p-1}{\frac{p-1}{2}}. \quad (17)$$

It remains to use the well-known congruences

$$\binom{2p-1}{p-1} \equiv 1 \pmod{p^3}$$

due to J. Wolstenholme (see [13]) and

$$\binom{p-1}{\frac{p-1}{2}} \equiv (-1)^{(p-1)/2} 2^{2(p-1)} \pmod{p^3} \quad (18)$$

due to F. Morley [9], although they are true modulo p^2 only if $p = 3$, to conclude that the expression in (17) is congruent to

$$\sum_{n=0}^{(p-1)/2} F(n, \frac{p-1}{2}) \equiv (-1)^{(p-1)/2} p \pmod{p^4}$$

(for our purposes we need the latter modulo p^3), and the desired result follows from (16). \square

Proof of Theorem 2. The validity of (12) for $p = 3$ is checked by hand; therefore, we may assume that $p \geq 5$.

The rational functions

$$F(n, k) = (120n^2 - 84nk + 34n - 10k + 3) \frac{(\frac{1}{2})_n^3 (\frac{1}{2})_{2n+k}}{2^{6n} (1)_n^3 (1)_{n-k}^2 (\frac{1}{2})_k^3},$$

$$G(n, k) = \frac{(\frac{1}{2})_n^3 (\frac{1}{2})_{2n+k-1}}{2^{6n-8} (1)_{n-1}^3 (1)_{n-k}^2 (\frac{1}{2})_k^3}$$

satisfy (14), since after division of both sides by $G(n, k)$ one only needs to verify that

$$\begin{aligned} & \frac{(120n^2 - 84nk + 118n - 10k + 13)(k - \frac{1}{2})^3}{256n^3(n - k + 1)^2} \\ & - \frac{(120n^2 - 84nk + 34n - 10k + 3)(2n + k - \frac{1}{2})}{256n^3} \\ & = \frac{(2n + k - \frac{1}{2})(2n + k + \frac{1}{2})(n + \frac{1}{2})^3}{64n^3(n - k + 1)^2} - 1. \end{aligned}$$

Summing (14), this time over $n = 0, 1, \dots, p-1$, we obtain

$$\sum_{n=0}^{p-1} F(n, k-1) - \sum_{n=0}^{p-1} F(n, k) = G(p, k) - G(0, k) = G(p, k).$$

Using

$$G(p, k) = \frac{(\frac{1}{2})_p^3 (\frac{1}{2})_{2p+k-1}}{2^{6p-8} (1)_{p-1}^3 (1)_{p-k}^2 (\frac{1}{2})_k^3} \equiv 0 \pmod{p^5}$$

for $k = 1, 2, \dots, \frac{p-1}{2}$, we deduce as above that

$$\sum_{n=0}^{p-1} F(n, 0) \equiv \sum_{n=0}^{p-1} F(n, \frac{p-1}{2}) \pmod{p^5}.$$

In addition, for $n = \frac{p+1}{2}, \frac{p+1}{2} + 1, \dots, p-1$ we have

$$\begin{aligned} \text{ord}_p \frac{(\frac{1}{2})_n^3 (\frac{1}{2})_{2n+(p-1)/2}}{2^{6n} (1)_n^3 (1)_{n-(p-1)/2}^2 (\frac{1}{2})_{(p-1)/2}^3} &= \text{ord}_p (\frac{1}{2})_n^3 (\frac{1}{2})_{2n+(p-1)/2} \\ &\geq \text{ord}_p (\frac{1}{2})_{(p+1)/2}^3 (\frac{1}{2})_{p+1+(p-1)/2} = 5; \end{aligned}$$

thus

$$F(n, \frac{p-1}{2}) \equiv 0 \pmod{p^5}$$

for these values of n , and we obtain

$$\sum_{n=0}^{p-1} F(n, \frac{p-1}{2}) = \sum_{n=(p-1)/2}^{p-1} F(n, \frac{p-1}{2}) \equiv F(\frac{p-1}{2}, \frac{p-1}{2}) \pmod{p^5}.$$

Finally, note that

$$F(\frac{p-1}{2}, \frac{p-1}{2}) = 3p(3p-2) \frac{(\frac{1}{2})_{3(p-1)/2}}{2^{3(p-1)} (1)_{(p-1)/2}^3} = 3p \cdot 2^{-6(p-1)} \frac{(3p-2)!}{(\frac{3(p-1)}{2})! (\frac{p-1}{2})!^3}$$

and

$$2^{-6(p-1)} \frac{(3p-2)!}{(\frac{3(p-1)}{2})! (\frac{p-1}{2})!^3} \equiv p \pmod{p^4}, \quad (19)$$

the congruence following from the lemma below, to arrive at the desired claim (12). \square

Lemma 1. *For a prime $p \geq 5$ take $n = (p-1)/2$. Then*

$$\frac{(6n+1)!}{(3n)! n!^3} \equiv p \cdot 2^{12n} \pmod{p^4}.$$

Proof. For $p > 3$, we have

$$\begin{aligned} \frac{(6n+1)!}{(3n)!} &= \frac{(6n+2)!}{2 \cdot (3n+1)!} = (2n+1) \prod_{j=1}^n (2p-j)(2p+j)(3p-j) \\ &= pn!^3 \prod_{j=1}^n \left(1 - \frac{2p}{j}\right) \left(1 + \frac{2p}{j}\right) \left(1 - \frac{3p}{j}\right) \\ &= pn!^3 \left(1 - 3pH_n + \frac{9}{2}p^2H_n^2 - \frac{17}{2}p^2H'_n + O(p^3)\right), \end{aligned}$$

where

$$H_n = \sum_{l=1}^n \frac{1}{l} \quad \text{and} \quad H'_n = \sum_{l=1}^n \frac{1}{l^2} \equiv 0 \pmod{p}.$$

Thus,

$$\frac{(6n+1)!}{(3n)! n!^3} \equiv p \left(1 - 3pH_n + \frac{9}{2}p^2H_n^2\right) \pmod{p^4}. \quad (20)$$

It remains to use a result of E. Lehmer [6, Eq. (45)],

$$H_n \equiv -2Q_p(2) + pQ_p(2)^2 \pmod{p^2}, \quad \text{where} \quad Q_p(2) = \frac{2^{p-1} - 1}{p},$$

which may be written in the form

$$Q_p(2) \equiv -\frac{1}{2}H_n + \frac{1}{8}pH_n^2 \pmod{p^2},$$

equivalently,

$$2^{p-1} \equiv 1 - \frac{1}{2}pH_n + \frac{1}{8}p^2H_n^2 \pmod{p^3}.$$

Raising both sides of the congruence to the sixth power we obtain

$$2^{12n} = 2^{6(p-1)} \equiv 1 - 3pH_n + \frac{9}{2}p^2H_n^2 \pmod{p^3}.$$

The comparison of this with (20) leads to the desired claim. \square

Proof of Theorem 3. This time the choice of a WZ-pair satisfying (14) is as follows:

$$F(n, k) = (20n - 2k + 3) \frac{(-1)^{n+k} \left(\frac{1}{2}\right)_n \left(\frac{1}{2}\right)_{2n+k}}{2^{4n} (1)_n^2 (1)_{n-k} \left(\frac{1}{2}\right)_k^2},$$

$$G(n, k) = \frac{(-1)^{n+k} \left(\frac{1}{2}\right)_n \left(\frac{1}{2}\right)_{2n+k-1}}{2^{4n-6} (1)_{n-1}^2 (1)_{n-k} \left(\frac{1}{2}\right)_k^2}.$$

Since $G(p, k) \equiv 0 \pmod{p^3}$ for $k = 1, 2, \dots, \frac{p-1}{2}$, by applying (14) we conclude that

$$\sum_{n=0}^{p-1} F(n, 0) \equiv \sum_{n=0}^{p-1} F\left(n, \frac{p-1}{2}\right) \pmod{p^3}.$$

Furthermore, $F(n, \frac{p-1}{2}) \equiv 0 \pmod{p^3}$ for $n = \frac{p+1}{2}, \dots, p-1$, hence the latter sum reduces to the single term $F(\frac{p-1}{2}, \frac{p-1}{2}) \pmod{p^3}$. Finally,

$$\begin{aligned} F\left(\frac{p-1}{2}, \frac{p-1}{2}\right) &= 3(3p-2) \frac{\left(\frac{1}{2}\right)_{3(p-1)/2}}{2^{2(p-1)} (1)_{(p-1)/2} \left(\frac{1}{2}\right)_{(p-1)/2}} \\ &= 3 \cdot 2^{-6(p-1)} \frac{(3p-2)!}{\left(\frac{3(p-1)}{2}\right)! \left(\frac{p-1}{2}\right)!^3} \cdot 2^{2(p-1)} \binom{p-1}{\frac{p-1}{2}}^{-1} \end{aligned}$$

and application of (18) and (19) finishes the proof of (13). \square

3. EXPERIMENTAL DATABASE AND CONCLUSION

To convince the reader that there is a general pattern for p -supercongruences analogous to Ramanujan-type series for $1/\pi$ and their generalizations, we provide a list of several examples (the corresponding original formulas, some of them conjectural as well, are all represented in [14]):

$$\sum_{n=0}^{p-1} \frac{\left(\frac{1}{2}\right)_n \left(\frac{1}{6}\right)_n \left(\frac{5}{6}\right)_n}{n!^3} (5418n + 263) \frac{(-1)^n}{80^{3n}} \stackrel{?}{\equiv} 263 \left(\frac{-15}{p}\right) p \pmod{p^3} \quad \text{for } p > 5, \quad (21)$$

$$\sum_{n=0}^{p-1} \frac{\left(\frac{1}{2}\right)_n \left(\frac{1}{4}\right)_n \left(\frac{3}{4}\right)_n}{n!^3} (21460n + 1123) \frac{(-1)^n}{882^{2n}} \stackrel{?}{\equiv} 1123 \left(\frac{-1}{p}\right) p \pmod{p^3} \quad \text{for } p > 7, \quad (22)$$

$$\sum_{n=0}^{p-1} \frac{\left(\frac{1}{2}\right)_n \left(\frac{1}{4}\right)_n \left(\frac{3}{4}\right)_n}{n!^3} (26390n + 1103) \frac{1}{99^{4n}} \stackrel{?}{\equiv} 1103 \left(\frac{-2}{p}\right) p \pmod{p^3} \quad \text{for } p > 11, \quad (23)$$

$$\sum_{n=0}^{p-1} \frac{(\frac{1}{2})_n (\frac{1}{3})_n (\frac{2}{3})_n}{n!^3} (14151n + 827) \frac{(-1)^n}{500^{2n}} \stackrel{?}{\equiv} 827 \left(\frac{-3}{p}\right) p \pmod{p^3} \quad \text{for } p > 5, \quad (24)$$

$$\sum_{n=0}^{p-1} \frac{(\frac{1}{2})_n^5}{n!^5} (20n^2 + 8n + 1) \frac{(-1)^n}{2^{2n}} \stackrel{?}{\equiv} p^2 \pmod{p^5} \quad \text{for } p > 3, \quad (25)$$

$$\sum_{n=0}^{p-1} \frac{(\frac{1}{2})_n^5}{n!^5} (820n^2 + 180n + 13) \frac{(-1)^n}{2^{10n}} \stackrel{?}{\equiv} 13p^2 \pmod{p^5} \quad \text{for } p > 3, \quad (26)$$

$$\sum_{n=0}^{p-1} \frac{(\frac{1}{2})_n (\frac{1}{4})_n (\frac{3}{4})_n (\frac{1}{6})_n (\frac{5}{6})_n}{n!^5} (1640n^2 + 278n + 15) \frac{(-1)^n}{2^{10n}} \stackrel{?}{\equiv} 15 \left(\frac{3}{p}\right) p^2 \pmod{p^5} \\ \text{for } p > 3, \quad (27)$$

$$\sum_{n=0}^{p-1} \frac{(\frac{1}{2})_n (\frac{1}{4})_n (\frac{3}{4})_n (\frac{1}{3})_n (\frac{2}{3})_n}{n!^5} (252n^2 + 63n + 5) \frac{(-1)^n}{48^n} \stackrel{?}{\equiv} 5p^2 \pmod{p^5} \quad (28) \\ \text{for } p > 3,$$

$$\sum_{n=0}^{p-1} \frac{(\frac{1}{2})_n (\frac{1}{3})_n (\frac{2}{3})_n (\frac{1}{6})_n (\frac{5}{6})_n}{n!^5} (5418n^2 + 693n + 29) \frac{(-1)^n}{80^{3n}} \stackrel{?}{\equiv} 29 \left(\frac{5}{p}\right) p^2 \pmod{p^5} \\ \text{for } p > 5, \quad (29)$$

$$\sum_{n=0}^{p-1} \frac{(\frac{1}{2})_n (\frac{1}{8})_n (\frac{3}{8})_n (\frac{5}{8})_n (\frac{7}{8})_n}{n!^5} (1920n^2 + 304n + 15) \frac{1}{7^{4n}} \stackrel{?}{\equiv} 15 \left(\frac{7}{p}\right) p^2 \pmod{p^5} \\ \text{for } p > 7, \quad (30)$$

$$\sum_{n=0}^{p-1} \frac{(\frac{1}{2})_n^7}{n!^7} (168n^3 + 76n^2 + 14n + 1) \frac{1}{2^{6n}} \stackrel{?}{\equiv} \left(\frac{-1}{p}\right) p^3 \pmod{p^7} \quad (31) \\ \text{for } p > 2,$$

$$\sum_{n=0}^{p-1} \sum_{k=0}^n \binom{n}{k}^4 \cdot (4n+1) \frac{1}{6^{2n}} \stackrel{?}{\equiv} \left(\frac{-15}{p}\right) p \pmod{p^2} \quad (32) \\ \text{for } p > 3,$$

$$\sum_{n=0}^{p-1} \sum_{k=0}^{\lfloor n/3 \rfloor} (-1)^{n-k} 3^{n-3k} \frac{(3k)!}{k!^3} \binom{n}{3k} \binom{n+k}{k} \cdot (4n+1) \frac{1}{81^n} \stackrel{?}{\equiv} \left(\frac{-3}{p}\right) p \pmod{p^3} \quad (33) \\ \text{for } p > 3,$$

$$\sum_{n=0}^{p-1} \sum_{k=0}^n \binom{n}{k}^2 \binom{2k}{k} \binom{2n-2k}{n-k} \cdot (5n+1) \frac{1}{64^n} \stackrel{?}{\equiv} \left(\frac{-3}{p}\right) p \pmod{p^3} \quad (34) \\ \text{for } p > 2,$$

$$\sum_{n=0}^{p-1} \binom{2n}{n}^2 \sum_{k=0}^n \binom{2k}{k}^2 \binom{2n-2k}{n-k}^2 \cdot (36n^2 + 12n + 1) \frac{1}{2^{10n}} \stackrel{?}{\equiv} p^2 \pmod{p^3} \quad (35)$$

for $p > 2$.

Remark 2. Replacing the finite sums on n in (32)–(35) by infinite sums following the ‘recipe’ in Remark 1 (that is, replacing some factorials by the corresponding values of the p -adic gamma function as in (7)) looks like a tricky problem. It seems that this is the place where we lose some ‘natural’ powers of p in (32) and (35).

We find it quite unfortunate that, in the other cases when we have proofs of the original Ramanujan-type series for $1/\pi$ and $1/\pi^2$ by creative telescoping, we cannot recover our argument in Section 2. These are the supercongruences (3)–(5), (25) and (26); the choices of the first member in the corresponding WZ-pair (again borrowed from [3], [5]) are

$$F(n, k) = (6n - 2k + 1) \frac{\left(\frac{1}{2}\right)_n \left(\frac{1}{2}\right)_{n+k} \left(\frac{1}{2}\right)_{n-k}}{2^{2n} (1)_n^2 (1)_{n-k} \left(\frac{1}{2}\right)_k},$$

$$F(n, k) = (6n - 2k + 1) \frac{(-1)^{n+k} \left(\frac{1}{2}\right)_{n+k} \left(\frac{1}{2}\right)_{n-k}^2}{2^{3n-k} (1)_n^2 (1)_{n-k}},$$

$$F(n, k) = (84n^2 - 56nk + 4k^2 + 52n - 12k + 5) \frac{(-1)^k \left(\frac{1}{2}\right)_n \left(\frac{1}{2}\right)_{n+k} \left(\frac{1}{2}\right)_{n-k}^2}{2^{4n} (1)_n^2 (1)_{2n-k+1}},$$

$$F(n, k) = (20n^2 - 12nk + 8n - 2k + 1) \frac{(-1)^{n+k} \left(\frac{1}{2}\right)_n^3 \left(\frac{1}{2}\right)_{n+k} \left(\frac{1}{2}\right)_{n-k}}{2^{2n} (1)_n^3 (1)_{n-k}^2 \left(\frac{1}{2}\right)_k^2},$$

$$F(n, k) = (3280n^4 - 4592n^3k + 2160n^2k^2 - 336nk^3 + 4000n^3 - 3816n^2k$$

$$+ 1008nk^2 - 40k^3 + 1592n^2 - 884nk + 92k^2 + 232n - 62k + 13)$$

$$\times \frac{(-1)^{n+k} \left(\frac{1}{2}\right)_n^3 \left(\frac{1}{2}\right)_{n+k} \left(\frac{1}{2}\right)_{n-k}^3}{2^{6n} (1)_n^3 (1)_{2n-k+1}^2},$$

respectively. Nevertheless, following the lines of the above proofs of Theorems 1 and 2, we can show the validity of (3) modulo p^2 , of (4) and (5) modulo p , of (25) modulo p^4 , and of (26) modulo p^2 .

We wonder whether there exists a deep general theory behind all these Ramanujan-type supercongruences, like the theory of overconvergent p -adic modular forms, which could replace the theory of modular forms used in the proofs of Ramanujan’s formulas for $1/\pi$.

We also wonder whether the methods of the classical theory of hypergeometric transformations can be applied to prove the supercongruences mentioned above. We have at least two successful examples of their application: D. McCarthy and R. Osburn [7] use them to prove a different supercongruence conjectured in [12], and E. Mortenson [10] gives a similar proof of the supercongruence (2).

Acknowledgements. This work was written during the author’s visits in the Fakultät für Mathematik, Universität Wien (March 2008), and the Department of Mathematics, National University of Singapore (May 2008). The author would like

to thank Heng Huat Chan and Christian Krattenthaler for useful conversations on the subject. The author is also thankful to Frits Beukers, Jonathan Sondow, Jan Stienstra and Don Zagier for helpful comments, and to Eric Mortenson for pointing out the references [10] and [12] which became one of the starting points of this project.

REFERENCES

- [1] G. BAUER, Von den Coefficienten der Reihen von Kugelfunctionen einer Variablen, *J. Reine Angew. Math.* **56** (1859), 101–121.
- [2] S. B. EKHAD and D. ZEILBERGER, A WZ proof of Ramanujan’s formula for π , *Geometry, Analysis, and Mechanics*, J. M. Rassias (ed.), World Scientific, Singapore (1994), 107–108.
- [3] J. GUILLERA, [Some binomial series obtained by the WZ-method](#), *Adv. in Appl. Math.* **29**:4 (2002), 599–603.
- [4] J. GUILLERA, [About a new kind of Ramanujan-type series](#), *Experiment. Math.* **12**:4 (2003), 507–510.
- [5] J. GUILLERA, [Generators of some Ramanujan formulas](#), *Ramanujan J.* **11**:1 (2006), 41–48.
- [6] E. LEHMER, [On congruences involving Bernoulli numbers and the quotients of Fermat and Wilson](#), *Ann. Math. (2)* **39**:2 (1938), 350–360.
- [7] D. MCCARTHY and R. OSBURN, A p -adic analogue of a formula of Ramanujan, *Arch. Math. (Basel)* (to appear); [arXiv: 0708.3307 \[math.NT\]](#) (2007).
- [8] Y. MORITA, A p -adic analogue of the Γ -function, *J. Fac. Sci. Univ. Tokyo Sect. IA* **22** (1975), 255–266.
- [9] F. MORLEY, [Note on the congruence \$2^{4n} \equiv \(-1\)^n\(2n\)!/\(n!\)^2\$, where \$2n + 1\$ is a prime](#), *Ann. Math.* **9** (1895), 168–170.
- [10] E. MORTENSON, [A \$p\$ -adic supercongruence conjecture of Van Hamme](#), *Proc. Amer. Math. Soc.* **136**:12 (2008), 4321–4328.
- [11] S. RAMANUJAN, Modular equations and approximations to π , *Quart. J. Math. Oxford Ser. (2)* **45** (1914), 350–372; Reprinted in *Collected papers of Srinivasa Ramanujan*, G.H. Hardy, P.V. Sechu Aiyar, and B.M. Wilson (eds.), Cambridge University Press, Cambridge (1927) & Chelsea Publ., New York (1962), 23–39.
- [12] L. VAN HAMME, [Some conjectures concerning partial sums of generalized hypergeometric series](#), *p -adic functional analysis* (Nijmegen, 1996), Lecture Notes in Pure and Appl. Math. **192**, Dekker, New York (1997), 223–236.
- [13] J. WOLSTENHOLME, On certain properties of prime numbers, *Quart. J. Math. (Oxford)* **5** (1862), 35–39.
- [14] W. ZUDILIN, Ramanujan-type formulae for $1/\pi$: A second wind?, *Modular forms and string duality* (Banff, June 2006), N. Yui, H. Verrill, and C.F. Doran (eds.), Fields Inst. Commun. Ser. **54**, Amer. Math. Soc. & Fields Inst., Providence, RI (2008), 179–188; [arXiv: 0712.1332 \[math.NT\]](#) (2007).

MAX-PLANCK-INSTITUT FÜR MATHEMATIK, VIVATSGASSE 7, D-53111 BONN, GERMANY
 WWW-address: <http://wain.mi.ras.ru/>