

COAMOEBAS OF COMPLEX ALGEBRAIC PLANE CURVES AND THE LOGARITHMIC GAUSS MAP

MOUNIR NISSE

ABSTRACT. The coamoeba of any complex algebraic plane curve V is its image in the real torus under the argument map. The area counted with multiplicity of the coamoeba of any algebraic curve in $(\mathbb{C}^*)^2$ is bounded in terms of the degree of the curve. We show in this Note that up to multiplication by a constant in $(\mathbb{C}^*)^2$, the complex algebraic plane curves whose coamoebas are of maximal area (counted with multiplicity) are defined over \mathbb{R} , and their real loci are Harnack curves possibly with ordinary real isolated double points (c.f. [MR-00]). In addition, we characterize the complex algebraic plane curves such that their coamoebas contain no extra-piece.

CONTENTS

| | |
|--|---|
| 1. Introduction | 1 |
| 2. Preliminaries | 3 |
| 3. Area Counted with Multiplicity of Coamoebas and the Alga map | 6 |
| 4. Characterization of Coamoebas of Plane Curves Without Extra-Piece | 8 |
| References | 9 |

1. INTRODUCTION

There is a very strong relationship between the amoeba and coamoeba of a complex algebraic hypersurface. Given an amoeba of a hypersurface, it is well known by Mikael Passare and Hans Rullgård [PR-04] that the spine of the amoeba is equipped with a structure of tropical hypersurface which is dual to some convex subdivision of the Newton polytope. We also know that amoeba reaches infinity by several tentacles [V2-02]. The coamoeba covers the hyperplanes orthogonal to the external edges of some subdivision of the Newton polytope by several *extra-pieces*. An extra-piece goes away from some hyperplane orthogonal to some inner edge of the subdivision of the Newton polytope. These hyperplanes are equipped with a frames spanned by the normal vectors to the sides of the element of the subdivision containing those edges, and these frames determine the position of the coamoeba relatively to these hyperplanes.

The purpose of this Note is to describe the relations and the similarities which exist between amoebas and coamoebas of complex algebraic plane curves. For a polynomial

in two variables $f \in \mathbb{C}[z, w]$, let V be the curve of its zero set in $(\mathbb{C}^*)^2$ and Δ its Newton polygon, which is the convex hull in \mathbb{R}^2 of the index of the non zero coefficients of f . The amoeba $\mathcal{A} \subset \mathbb{R}^2$ of the curve V is the image of V by the logarithmic map Log ; where $\text{Log} : (\mathbb{C}^*)^2 \rightarrow \mathbb{R}^2$ is the map defined by $\text{Log}(z, w) = (\log \|z\|, \log \|w\|)$ (c.f. [GKZ-94]).

The amoeba's complement has a finite number of convex connected components, corresponding to domains of convergence of the Laurent series expansions of the rational function $\frac{1}{f}$. We know that the spine Γ of the amoeba \mathcal{A} has a structure of a tropical curve in \mathbb{R}^2 (proved by M. Passare and H. Rullgård in 2000 [PR-04], and independently by G. Mikhalkin in 2000). In addition the spine of the amoeba is dual to some coherent (i.e. convex) subdivision τ of the integer convex polygon Δ . It is shown by M. Forsberg, M. Passare and A. Tsikh that the set of vertices of τ is in bijection with the set of complement components of \mathcal{A} in \mathbb{R}^2 [FPT-00]. It was shown by Mikael Passare and Hans Rullgård in [PR-04] that the amoebas of complex algebraic plane curves have finite area, and the upper bound on the area is in terms of the the degree of the curve. In addition, it was shown by Grigory Mikhalkin and Hans Rullgård, that up to multiplication by a constant in $(\mathbb{C}^*)^2$ the algebraic curves in $(\mathbb{C}^*)^2$ whose amoebas are of maximal area are defined over \mathbb{R} and, furthermore their real locus are a Harnack curves possibly with ordinary real isolated double points (c.f. [MR-00]). The coamoeba $co\mathcal{A} \subset (S^1)^2$ of the curve V is the image of V under the argument mapping $\text{Arg} : (z, w) \mapsto (e^{i\arg(z)}, e^{i\arg(w)})$. This terminology has been introduced by M. Passare and A. Tsikh in 2000. It is shown in [N2-07] that the complement components of the closure in the flat torus of the coamoeba of a complex algebraic hypersurface of complex dimension $n - 1$ and defined by a polynomial f with Newton polytope Δ are convex and their number don't exceed $n! \text{Vol}(\Delta)$.

In this Note, we will prove a similar properties of the coamebas of the complex algebraic plane curves. More precisely we characterize the curves such that the area of their coamoeb counted with multiplicity is maximal. Using the fact that the set of critical point Crit_{Arg} of the argument map is equal to that of the logarithmic map which is the inverse image by the logarithmic Gauss map of the real projective line, and some properties of extra-pieces introduced in [N1-06] and [N3-07], we obtain the following results.

Theorem 1.1. *Let V be a complex algebraic plane curve defined by a polynomial with Newton polygon Δ . Then the following statements are equivalent:*

- (i) *The inverse image by the map $pr : co\mathcal{A} \setminus \text{Arg}(\text{Crit}_{\text{Arg}}) \rightarrow T \setminus pr \circ \text{Arg}(\text{Crit}_{\text{Arg}})$ of any point in the image is equal to $2 \text{Area}(\Delta)$ (i.e., the map pr is $2 \text{Area}(\Delta) - to - 1$),*
- (ii) *the coamoeba of V has no extra-piece.*

Theorem 1.2. *Let V be a complex algebraic plane curve defined by a polynomial with Newton polygon Δ . Then the area counted with multiplicity of the coamoeba of V cannot exceed $2\pi^2 \text{Area}(\Delta)$, and we have the following equivalent statements:*

- (i) $\text{Area}_{\text{mult}}(\text{co}\mathcal{A}) = 2\pi^2 \text{Area}(\Delta)$,
- (ii) *The curve V is real up to multiplication by a constant in \mathbb{C}^* , and its real part $\mathbb{R}V$ is a Harnack curve possibly with ordinary real isolated double points.*

The extra-pieces are subsets of the coamoeba with no-vanishing area and such that their boundary has a no discrete component contained in the set of critical values of the argument map. We can remark that if we assume that the closure in the torus of the coamoeba is not all the torus, then the no-linear components of the boundary of the coamoeba are contained in the boundary of some extra-pieces (for more details see [N1-06] or [N3-07]). They play the role of the tentacles of the amoeba which correspond to the external edges of the subdivision τ dual to the spine of the amoeba. Except that extra-pieces correspond to some inner edges of the subdivision τ .

2. PRELIMINARIES

Let V be an algebraic hypersurface in $(\mathbb{C}^*)^n$ defined by the polynomial

$$f(z) = \sum_{\alpha \in \text{supp}(f)} a_\alpha z^\alpha, \quad z^\alpha = z_1^{\alpha_1} z_2^{\alpha_2} \dots z_n^{\alpha_n}, \quad (1)$$

where a_α are non-zero complex numbers and $\text{supp}(f) = \{\alpha \in \mathbb{Z}^n \mid a_\alpha \neq 0\}$ is the support of f , and we denote by Δ the Newton polytope of f (i.e., the convex hull in \mathbb{R}^n of $\text{supp}(f)$). We assume in this Note that the components of α are natural.

2.1. Gauss map and the critical points of the complex logarithmic map.

We use here the definition of the Gauss map of an hypersurface in the algebraic torus given by Kapranov in [K1-91].

Definition 2.2. Let $(\mathbb{C}^*)^n$ be the algebraic torus and $V \subset (\mathbb{C}^*)^n$ be an algebraic hypersurface. For each $z \in (\mathbb{C}^*)^n$ let $l_z : h \mapsto z.h$ be the translation by z . The *Gauss map* of the hypersurface V is the rational map $\gamma : V \rightarrow \mathbb{C}\mathbb{P}^{n-1}$ taking a smooth point $z \in V$ to the hyperplane $d(l_{z^{-1}})(T_z V) \subset T_1((\mathbb{C}^*)^n)$, where $\mathbb{C}\mathbb{P}^{n-1} = \mathbb{P}(T_1((\mathbb{C}^*)^n))$

$$\gamma : V \rightarrow \mathbb{C}\mathbb{P}^{n-1}$$

$$z \mapsto T_z V \subset T_z((\mathbb{C}^*)^n) \xrightarrow{t_{z^{-1}}^*} t_{z^{-1}}^*(T_z V) \subset T_1((\mathbb{C}^*)^n)$$

$$z \mapsto [t_{z^{-1}}^*(T_z V)]$$

where $[t_{z^{-1}}^*(T_z V)]$ is the class of $t_{z^{-1}}^*(T_z V)$ in $\mathbb{C}\mathbb{P}^{n-1}$.

We assume from now that $V \subset (\mathbb{C}^*)^2$ is an algebraic curve defined by a polynomial f and nowhere singular. Let $z \in V$, $U \subset (\mathbb{C}^*)^2$ be a small open neighborhood of z

and $\text{Log}_U : U \longrightarrow \mathbb{C}^2$ be a choice of a branch of the holomorphic logarithm function and then we take the image of $U \cap V$ by Log_U . Hence the vector $\gamma(z) \in \mathbb{C}^2$ (looking at it as $T_{\text{Log}_U(z)}\mathbb{C}^2$) tangent to $\text{Log}_U(U \cap V)$ at the point $\text{Log}_U(z)$ is a complex vector, which is independent of the choice of the holomorphic branch. So it defines a map from $U \cap V$ (a rational map from all V) to \mathbb{CP}^1 , called the *logarithmic Gauss map* of the curve V . Explicitly one have:

$$\begin{aligned} \gamma : V &\longrightarrow \mathbb{CP}^1 \\ z &\longmapsto [\gamma(z)] \end{aligned}$$

where $[\gamma(z)]$ denotes the class of the vector $\gamma(z)$ in \mathbb{CP}^1 . We decompose the identity as follow:

$$U \cap V \xrightarrow{\text{Log}_U} \mathbb{C}^2 \xrightarrow{\exp} (\mathbb{C}^*)^2 \xrightarrow{f} \mathbb{C}$$

Let us look now to the amoeba \mathcal{A} of the curve V , and more precisely to $\text{Log}(U \cap V)$ (where here $\text{Log}(z_1, z_2) = (\log \|z_1\|, \log \|z_2\|)$). It's clear that $\text{Log}(U \cap V) = \text{Re} \circ \text{Log}_U(U \cap V)$ where $\text{Re}(z_1, z_2) = (\text{Re}(z_1), \text{Re}(z_2))$ and Re denotes the real part. So if $t = \text{Log}_U(u)$ where $u = (u_1, u_2) \in U$, then $dt = \frac{du}{u}$, and we obtains

$$\begin{aligned} \gamma(z) &= \frac{df(\exp(t))}{dt} \Big|_{t=\text{Log}_U(z)} \\ &= (u_1 \frac{\partial f}{\partial u_1}(u), u_2 \frac{\partial f}{\partial u_2}(u)) \Big|_{u=z} \\ &= (z_1 \frac{\partial f}{\partial z_1}(z), z_2 \frac{\partial f}{\partial z_2}(z)) \end{aligned}$$

Hence a point $z \in V$ is a critical point of the function $\text{Log} = \text{Re} \circ \text{Log}_U$ is equivalent to saying that the tangent space of $\text{Log}_U(U \cap V)$ at the point $\text{Log}_U(z)$ contains a non zero imaginary vector v , which means that $T_{\text{Log}_U(z)}(\text{Log}_U(U \cap V))$ is the complexification of a real line and then it is invariant under complex conjugation so $\gamma(z) \in \mathbb{RP}^1 \subset \mathbb{CP}^1$. We have the following Proposition and its proof given by Mikael Passare at IMJ, l'institut de Mathématiques de Jussieu in *Ecole d'été en Géométrie Tropicale* in 2006 [P-06]:

Proposition 2.3. *Let $V \subset (\mathbb{C}^*)^n$ be an algebraic hypersurface and γ be the logarithmic Gauss map. If $\text{Crit}(\text{Log}|_V)$ (resp. $\text{Crit}(\text{Arg}|_V)$) denotes the set of critical points of the logarithmic (resp. argument) mapping, then we have the following equalities: $\text{Crit}(\text{Log}|_V) = \text{Crit}(\text{Arg}|_V) = \gamma^{-1}(\mathbb{RP}^{n-1})$.*

Proof Let f be the polynomial defining the hypersurface V , $\text{reg}(V)$ be the set of regular points of V , and γ the logarithmic Gauss map

$$\begin{aligned} \gamma : \text{reg}(V) &\longrightarrow \mathbb{CP}^{n-1} \\ (z_1, \dots, z_n) &\longmapsto (z_1 \frac{\partial f}{\partial z_1} : \dots : z_n \frac{\partial f}{\partial z_n}) \end{aligned}$$

In local coordinates $(w_1, \dots, w_n) = (\text{Log } z_1, \dots, \text{Log } z_n)$ one has $z_j \frac{\partial}{\partial z_j} = \frac{\partial}{\partial w_j}$, so $\gamma(e^w)$ is the normal direction to the set $\{W \mid f(e^W) = 0\}$. Put $w_j = x_j + i\theta_j$, and let $C = A + iB = (A_1 + iB_1, \dots, A_n + iB_n)$ be a complex normal at W . We know that the complex tangent plane is given by the set of vectors C such that $\langle C, W \rangle = 0$, which is equivalently to say that we have the following equations

$$\begin{cases} \langle A, X \rangle = \langle B, \Theta \rangle \\ \langle B, X \rangle = -\langle A, \Theta \rangle \end{cases}$$

where $X = (x_1, \dots, x_n)$ and $\Theta = (\theta_1, \dots, \theta_n)$. The maps $W \mapsto X$ and $W \mapsto \Theta$ are submersion if A, B are linearly independent over \mathbb{R} . On the other hand, if $B = kA$ then we can see that we have the following

$$\langle A, X \rangle = k \langle A, \Theta \rangle = -k \langle B, X \rangle = -k^2 \langle A, X \rangle$$

which is equivalently to $\langle A, X \rangle = \langle B, \Theta \rangle = 0$. ■

2.4. Complex tropical curves.

Let h be a strictly positive real number and H_h be the self diffeomorphism of $(\mathbb{C}^*)^2$ defined by :

$$H_h : \begin{array}{ccc} (\mathbb{C}^*)^2 & \longrightarrow & (\mathbb{C}^*)^2 \\ (z_1, \dots, z_n) & \longmapsto & (|z|^h \frac{z}{|z|}, |w|^h \frac{w}{|w|}). \end{array}$$

which defines a new complex structure on $(\mathbb{C}^*)^2$ denoted by $J_h = (dH_h) \circ J \circ (dH_h)^{-1}$ where J is the standard complex structure.

A J_h -holomorphic curve V_h is a curve holomorphic with respect to the J_h complex structure on $(\mathbb{C}^*)^2$. It is equivalent to say that $V_h = H_h(V)$ where $V \subset (\mathbb{C}^*)^2$ is an holomorphic curve for the standard complex structure J on $(\mathbb{C}^*)^2$.

Recall that the Hausdorff distance between two closed subsets A, B of a metric space (E, d) is defined by:

$$d_{\mathcal{H}}(A, B) = \max\{\sup_{a \in A} d(a, B), \sup_{b \in B} d(A, b)\}.$$

In this Note we take $E = \mathbb{R}^2 \times (S^1)^2$, with the distance defined as the product of the Euclidean metric on \mathbb{R}^2 and the flat metric on $(S^1)^2$. We have the following definition given by G. Mikhalkin in [M2-04].

Definition 2.5. A complex tropical curve $V_\infty \subset (\mathbb{C}^*)^2$ is the limit (with respect to the Hausdorff metric on compact sets in $(\mathbb{C}^*)^2$) of a sequence of a J_h -holomorphic curves $V_h \subset (\mathbb{C}^*)^2$ when h tends to zero.

2.6. External hyperplanes and extra-pieces.

In this subsection we consider a complex algebraic hypersurface V defined by a polynomial f with Newton polytope Δ . We denote by \mathcal{A} the amoeba of V and by Γ its spine. Let us denote by τ the subdivision of Δ dual to Γ , and assume that f is as (1) (c.f. [RST-05] and [PR-04] for more details).

Definition 2.7. Let \mathbb{R}^n be the universal covering of the real torus $(S^1)^n$. Let α and β in the support of f . A hypersurface $H_{\alpha\beta} \subset \mathbb{R}^n$ is called *codual* (or corresponding) to an edge $E_{\alpha\beta}$ if it is given by the following equation:

$$\arg(a_\alpha) - \arg(a_\beta) + \langle \alpha - \beta, x \rangle = \pi.$$

In addition if $E_{\alpha\beta}$ is an external edge of τ (i.e., $E_{\alpha\beta}$ is a proper edge of the Newton polytope Δ), then $H_{\alpha\beta}$ is called an *external hyperplane*.

We denote by $\text{Critv}(\text{Arg})$ the set of critical values of the argument map.

Definition 2.8. An *extra-piece* is a connected component \mathcal{C} of $co\mathcal{A} \setminus \text{Critv}(\text{Arg})$ such that the boundary of its closure $\partial\overline{\mathcal{C}}$ is not contained in the union of hyperplanes.

This means that its boundary contains at least one component (smooth) in the set of critical values of the argument map.

3. AREA COUNTED WITH MULTIPLICITY OF COAMOEBAS AND THE ALGA MAP

For a polynomial in two variables $f \in \mathbb{C}[z, w]$, let V be the curve of its zero set in $(\mathbb{C}^*)^2$. We use the Alga map introduced by G. Mikhalkin and A. Okounkov in [MO-07] defined as follow. Let $(\mathbb{Z}_2)^2$ be the subgroup of real points of the torus $(S^1)^2$, and take now the quotient group $T = (S^1)^2 / (\mathbb{Z}_2)^2$. Let us denote by Alg the composition of the argument map with the projection $pr : (S^1)^2 \rightarrow T$. Let $\mathcal{R} \subset co\mathcal{A}$ be a subset of the regular values of the argument map. We define the *multiplicity* of \mathcal{R} as the cardinality of the inverse image by the argument map of any point in \mathcal{R} .

Lemma 3.1. *Let V be a complex algebraic plane curve defined by a polynomial f with Newton polygon Δ . Then the area of the coamoeba $co\mathcal{A}$ of V counted with multiplicity and denoted by $\text{Area}_{mult}(co\mathcal{A})$ cannot exceed $2\pi^2 \text{Area}(\Delta)$.*

Proof Let $f(z, w) = \sum_{(i,j) \in \Delta \cap \mathbb{Z}^2} a_{ij} z^i w^j$ be the polynomial defining the curve V . If the

subdivision τ dual to the spine of the amoeba of the curve is a triangulation, then by the deformation of the curve given by $f_t(z, w) = \sum_{(i,j) \in \Delta \cap \mathbb{Z}^2} a_{ij} (et)^{\nu(i,j)} z^i w^j$ with $t \in]0; \frac{1}{e}]$

with $\nu(i, j) = -c_{(i,j)}$ the constant defined by Passare and Rullgård in [PR-04] we have the result. Assume on the contrary it is not the case. It is well known that there exists a polynomial g with the same support as f such that the arguments of its coefficients are the same as those of f (see [R-00]), and the subdivision of Δ dual to the spine of the amoeba of the curve defined by g is a triangulation. The result is immediate

after deforming the coefficients of g on those of f (the deformation can be given by $(1-u)a_{\alpha,g} + ua_{\alpha}$ with $u \in [0; 1]$ and $a_{\alpha,g}$ the coefficient of g with index α). ■

Lemma 3.2. *Let $V \subset (\mathbb{C}^*)^2$ be a no singular algebraic curve defined by a polynomial f such that $\mathbb{R}V$ is a Harnack curve. Then the coamoeba $\text{co}\mathcal{A}$ of V is equal to the coamoeba of the complex tropical curve $V_{\infty,f}$.*

Proof Let f be the polynomial defining the curve V which we assume with real coefficients. Consider $\{f_t\}_{t \in]0; \frac{1}{e}]}$ the family of polynomials $\{f_t\}$ defined as above. We claim that the coamoebas of the curves V_{f_t} for sufficiently small t and those with t sufficiently close to $\frac{1}{e}$ are the same. Indeed, let $V_{\infty,f}$ be the complex tropical curve which is the limit when t tends to zero of the J_t -holomorphic curves $H_t(V_{f_t})$ (abusive notation, because here we have precisely H_h with $h = -\frac{1}{\log t}$ if we take the notation of 2.4), and Γ_{∞} be the tropical curve which is the limit of the amoebas $\mathcal{A}_{H_t(V_{f_t})}$. We know that the set of critical values of the argument map in our case is contained in the subgroup $(\mathbb{Z}_2)^2$ of the real points of the torus (recall that for a Harnack curves we have $\text{Crit}_{\text{Arg}} = \gamma^{-1}(\mathbb{R}\mathbb{P}^1) = \mathbb{R}V$ see [M1-00]). We can remark that the arguments of the coefficients of f_t are equal to those of f , and then the coamoeba of the truncation of f_t to any element of the subdivision τ is the same as that of the truncation of f to the same element. Using Lemma 1.3 and 1.4 [N4-08], one can deduce that there was only one possibility. ■

Remark 3.3.

- (a) Let $V \subset (\mathbb{C}^*)^2$ be a no singular algebraic curve defined by a polynomial f such that $\mathbb{R}V$ is a Harnack curve. Let $V_{\infty,f}$ and Γ_{∞} as above and v be a vertex of Γ_{∞} , then by the tropical localization we can see that $\text{Arg}(\text{Log}^{-1}(v) \cap V_{\infty,f})$ is just two triangles of area $\pi^2/2$, and whose vertices are real and pairwise identified (they are real because those points depends on the coefficients of f , see section 3.6 in [N1-06], which are real, and there area is equal to $\pi^2/2$ because the linear part of the affine linear surjection sending the standard simplex to the simplex Δ_v dual to the vertex v is in $SL_2(\mathbb{Z})$). Let v_1 and v_2 be two adjacent vertices of Γ_{∞} with $E_{v_1v_2}$ the edge between them. Let U be a small open neighborhood of the edge $E_{v_1v_2}$, and $H_{v_1v_2}$ be the line in the torus corresponding to the edge $E_{v_1v_2}^*$ in the triangulation of Δ dual to $E_{v_1v_2}$. By tropical localization (Viro's principle see [V-90]) and Kapranov's Theorem (see [N1-06] and [K2-00]), we claim that the set $\text{Arg}(V_{\infty,f} \cap \text{Log}^{-1}(U))$ contains the line $H_{v_1v_2}$ in its interior. Indeed, on the contrary, this means that $H_{v_1v_2}$ is contained in the critical values of the argument map, which contradict the fact that the only critical values of the map Arg are in the subset of real points of the torus $(S^1)^2$.




FIGURE 1. The first picture in the left is the Harnack case and the two others are no-Harnack; extra-pieces and imaginary critical values of the argument map appear in the no-Harnack case.

- (b) Let $\overset{\circ}{D}_{std}$ be the interior of one triangle of the coamoeba of the standard line defined by $z + w + 1 = 0$. If Δ' is a primitive simplex i.e., of area $\frac{1}{2}$, and L is the linear part of the affine linear surjection of \mathbb{R}^2 sending the standard simplex to Δ' , then for any $v \in (\mathbb{Z}_2)^2 \subset (S^1)^2$ and x, y two different points in $\overset{\circ}{D}_{std}$ we have $\tau_v \circ {}^tL^{-1}(x) \neq {}^tL^{-1}(y)$ where τ_v is the translation in the torus by the vector v . Indeed, assume that there exist two different points x and y in $\overset{\circ}{D}_{std}$ and $v \in (\mathbb{Z}_2)^2$ such that $\tau_v \circ {}^tL^{-1}(x) = {}^tL^{-1}(y)$, then $x - y = {}^tL(v)$. So $x - y$ is a vector of the form $-\begin{pmatrix} k\pi \\ k'\pi \end{pmatrix}$ which is impossible. Hence for any simplex Δ' of minimal area in \mathbb{R}^2 , we have $\overline{pr \circ {}^tL_{\Delta'}^{-1}(co\mathcal{A}_{std})} = T$.

4. CHARACTERIZATION OF COAMOEBAS OF PLANE CURVES WITHOUT EXTRA-PIECE

Proof of Theorem 1.1. The implication (i) \Rightarrow (ii) is obvious and the inverse is an immediate issue of Remarks 3.3 (a) and (b). \blacksquare

We say that a map is at most p -to-1 if the inverse image of any point in the target has at most p points.

Lemma 4.1. *Let V be a complex algebraic plane curve defined by a polynomial f with Newton polygon Δ , and let $co\mathcal{A}$ be its coamoeba. Then if we have the equality $\text{Area}_{mult}(co\mathcal{A}) = 2\pi^2 \text{Area}(\Delta)$, then the map $\text{Log}_V : V \rightarrow \mathbb{R}^2$ is at most 2-to-1.*

Proof Let Γ be the spine of the amoeba of V (viewed as tropical curve), and τ the subdivision of Δ dual to Γ . If we take the same notation as in the proof of Lemma 3.2, let $V_{\infty, f}$ be the complex tropical curve equal to the limit of $H_t(V_{f_t})$ when t tends to zero (with respect to the Hausdorff metric on compacts of $(\mathbb{C}^*)^2 \approx \mathbb{R}^2 \times (S^1)^2$ equipped with the product of the standard Euclidean metric and the flat metric on the torus). Assume that the map $\text{Log}|_V : V \rightarrow \mathbb{R}^2$ is not at most 2-to-1. Let us denote by F (resp. F_c) the set of critical values of the logarithmic (resp. argument) map, and we set $\text{co}\mathcal{A} \setminus F_c = \cup_{i=1}^l \mathcal{R}_i$. By Proposition 2.3, if $\text{Arg}^{-1}(\mathcal{R}_i) = \cup_{j=1}^{n_i} \mathcal{R}_{ij}$ (n_i is the multiplicity of the region \mathcal{R}_i) then for any i and j the set $\text{Log}(\mathcal{R}_{ij})$ is a connected component of $\mathcal{A} \setminus F$. We have one of the two following cases: (i) the tropical curve Γ has at least one vertex v of multiplicity strictly greater than one (the multiplicity of a vertex v is the area of a parallelogram spanned by two primitive vectors in the direction of any two edges adjacent to v , see [M2-04]); (ii) there exist at least two adjacent vertices v_i and v_j of Γ such that if U is a small open neighborhood of the edge $E_{v_1 v_2}$ joining v_1 and v_2 , then $\overline{\text{Arg}(\text{Log}^{-1}(U) \cap V_{\infty, f})}$ don't contain the line $H_{v_1 v_2}$ corresponding to the edge $E_{v_1 v_2}$. In the two cases (i) and (ii) we have $\text{Area}_{\text{mult}}(\text{co}\mathcal{A}) < 2\pi^2 \text{Area}(\Delta)$. Indeed, the first case means that the area of the coamoeba of the truncation of f to the element Δ_v of τ dual to v is such that $\text{Area}_{\text{mult}}(\text{co}\mathcal{A}_{f^{\Delta_v}}) < 2\pi^2 \text{Area}(\Delta_v)$, because in this case the truncation f^{Δ_v} is maximally sparse or it has some coefficients which has no contribution on the spine of the amoeba and then the coamoeba of the truncation has some extra-pieces. The second case means that we have extra-pieces and then $\text{Area}_{\text{mult}}(\text{co}\mathcal{A}) < 2\pi^2 \text{Area}(\Delta)$ (see Remark 3.3 (b)). ■

Proof of Theorem 1.2. The implication (ii) \Rightarrow (i) is obvious after Lemma 3.2 and Remark 3.3. (a), and the implication (i) \Rightarrow (ii) is an immediate issue of Lemma 4.3 and Theorem 1 in [MR-00]. ■

REFERENCES

- [FPT-00] M. FORSBERG, M. PASSARE AND A. TSIKH, *Laurent determinants and arrangements of hyperplane amoebas*, Advances in Math. **151**, (2000), 45-70.
- [GKZ-94] I. M. GELFAND, M. M. KAPRANOV AND A. V. ZELEVINSKI, *Discriminants, resultants and multidimensional determinants*, Birkhäuser Boston 1994.
- [K1-91] M. M. KAPRANOV, *A characterization of A-discriminantal hypersurfaces in terms of the logarithmic Gauss map*, Math. Ann. **290** (1991), 277-285.
- [K2-00] M. M. KAPRANOV, *Amoebas over non-Archimedean fields*, Preprint 2000.
- [M1-00] G. MIKHALKIN, *Real algebraic curves, moment map and amoebas*, Ann. of Math. **151** (2000), 309-326.
- [M2-04] G. MIKHALKIN, *Enumerative Tropical Algebraic Geometry In \mathbb{R}^2* , J. Amer. Math. Soc. **18**, (2005), 313-377.
- [MO-07] G. MIKHALKIN AND A. OKOUNKOV, *Geometry of Planar Log-Fronts*, Moscow Math. J. **7**, (2007), 507-531.
- [MR-00] G. MIKHALKIN AND RULLGÄRD, *Amoebas of maximal area*, Internat. Math. Res. Notices **9**, (2001), 441-451.

- [N1-06] M. NISSE, *Maximally sparse polynomials have solid amoebas*, Preprint 2006, <http://fr.arxiv.org/pdf/0704.2216>
- [N2-07] M. NISSE, *Coamoebas of Complex Algebraic Hypersurfaces*, Preprint (2007).
- [N3-07] M. NISSE, *Amoebas and coamoebas, relations and similarities*, Preprint (2008).
- [N4-08] M. NISSE, *Complex Tropical Localization*, In Preparation.
- [P-06] M. PASSARE, *Basic Amoeba Theory*, Ecole d'été en Géométrie Tropicale, Institut de Mathématiques de Jussieu, (2006).
- [PR-04] M. PASSARE AND H. RULLGÅRD, *Amoebas, Monge-Ampère measures, and triangulations of the Newton polytope*, Duke Math. J. **121**, (2004), 481-507.
- [RST-05] Richter-Gebert, Sturmfels, Theobald J. RICHTER-GEBERT, B. STURMFELS ET T. THEOBALD, *First steps in tropical geometry*, Idempotent mathematics and mathematical physics, Contemp. Math., **377**, (2005), 289-317, Amer. Math. Soc., Providence, RI, 2005.
- [R-00] H. RULLGÅRD, *Stratification des espaces de polynômes de Laurent et structure de leurs amibes*, C. R. Acad. Sci. (2000), 355-358.
- [V-90] O. VIRO, *Patchworking real algebraic varieties*, preprint: <http://www.math.uu.se/oleg>; Arxiv: AG/0611382
- [V2-02] O. VIRO, *Wath is ... An Amoeba?*, Notices of the AMS, **49**, (2002), 916-917.

INSTITUT DE MATHÉMATIQUES DE JUSSIEU (UMR 7586), UNIVERSITÉ PIERRE ET MARIE CURIE, ANALYSE ALGÈBRE, 175, RUE DU CHEVALERET,, 75013 PARIS
E-mail address: nisse@math.jussieu.fr