© Indian Academy of Sciences

PRAMANA — journal of physics

# Quantum logic gates using Coherent Population Trapping states

ASHOK VUDAYAGIRI, \*

<sup>a</sup>School of Physics, University of Hyderabad Hyderabad 500046

**Abstract.** A scheme is proposed for achieving a Controlled Phase gate using interaction between atomic spin dipoles. Further, the spin states are prepared in a Coherent Population Trap states, which are robust against perturbations, laser fluctuations etc. And we show that single qubit and two qubit operations can easily be obtained in this scheme. The scheme is also robust against decoherences due to spontaneous emissions as the the CPT states used are dressed states formed out of Zeeman sublevels of ground states of the bare atom. However, certain practical issues are of concern in actually obtaining the scheme, which are also discussed at the end of this paper.

Keywords. Coherent Population Trap, Quantum Computation, Controlled phase gate

PACS Nos. 03.67.Lx,32.80.Qk,32.90+a,42.50.Ct

Conventional computers handle information in the form of bits - which take up values 0 or 1. 'Quantum Computers' on the other hand, use Quantum bits (qubits), which can be prepared in states 0, 1 or any superposition of the two. Algorithms of quantum computation exploit this unique feature of quantum mechanical system so as to solve certain class of computational problems with lesser number of steps. [1]. Hence a race to produce a reliable, robust and scalable Quantum Mechanical system which can be used as gates for Quantum Logic. There have been several attempts in the past to prepare such a system, using NMR of large molecules, quantum dot structures, ions in linear traps or neutral atoms in Optical lattices [2, 13], each system withs its own benefits and drawbacks. One of the major requirement for design of any QC system is that they should be robust and reliable while interactions between any two of them should be on-demand. One such system is proposed here which involve neutral atoms prepared in Coherent Population Trap (CPT) states. It is shown in this paper that such systems can be easily prepared and manipulated and it is possible to build one-qubit and two-qubit gates using them. Since CPT states are 'dark states' of the atom-light interaction, the atoms prepared in such states will not interact with the light any more [3, 4]. Nor will they evolve in time, since they are already eigenstates of the full Hamiltonian that consists of atomic as well as interaction terms.

In this communication, a configuration involving Zeeman sublevels of <sup>87</sup>Rb atom is considered, which exhibits two different CPT states which can be mapped to two qubits 0 and 1. It is shown that robust states can be prepared and single qubit and two - qubit operations can be performed using magnetic dipole interactions.

<sup>\*</sup>avsp@uohyd.ernet.in



**Figure 1.** (a) Schematic of laser arrangement. HWP is halfwave plate, PBS is polarizing beam splitter and M are mirrors and Mg are magnets to provide the weak field. (b) Energy level configuration of system used in the setup. Details of the notations are in detail in the text.

#### 1. the configuration

We consider the transition between  $|5S_{1/2}, F = 1\rangle$  and  $|5P_{1/2}, F = 1\rangle$  of <sup>87</sup>Rb. <sup>1</sup>, coupled by a two lasers, which are of same frequency but polarized orthogonal to each other one in plane containing quantization axis 'z' and other in the 'xy' plane. Following the selection rules [5] they both couple transitions between different Zeeman sublevels.

As shown in figure 1b. the beam  $E_z = \mathcal{E}_z \exp[i(\omega t - k.\{x,y\})]$  couples  $\Delta m_F = 0$  transitions between levels labeled  $|g_+\rangle \leftrightarrow |e_+\rangle$  and  $|g_-\rangle \leftrightarrow |e_-\rangle$ . The other beam, with is plane of polarization in xy plane can be considered as a combination of  $\sigma_+$  and  $\sigma_-$  beams coupling  $\Delta m_F = \pm 1$  transitions  $|g_{\pm}\rangle \leftrightarrow |e_0\rangle$  and  $|g_0\rangle \leftrightarrow |e_{\pm}\rangle$ .  $|g_0\rangle \leftrightarrow |e_0\rangle$  is not coupled by the  $E_z$  laser due to the vanishing Clebsch-Gordon coefficients. Both  $E_z$  and  $E_p$  beams can be derived from a same laser source by use of a halfwave plate and a polarizing beam splitter as shown in figure 1a. The ratio of values of  $E_{p,z}$  can be controlled by rotating the halfwave plate HWP.

When only the  $E_p$  beam is present, the configuration is the well known  $\Lambda$  system made up of  $|g_{-}\rangle \leftrightarrow |e_0\rangle \leftrightarrow |g_{+}\rangle$ . The steady state solution of this situation is the Coherent Population Trapping (CPT) state  $|\Psi_{-}\rangle = (1/\sqrt{2})[|g_{-}\rangle - |g_{+}\rangle]$  [3]. It is interesting to note that  $|\Psi_{-}\rangle$  is the CPT state even when there exists another CPT configuration - the V form of  $|e_{-}\rangle \leftrightarrow |g_0\rangle \leftrightarrow |e_{+}\rangle$ , and competes with the  $\Lambda$ . Our numerical results confirm this fact and it will be shown in a forthcoming communication. However, in the light of the argument presented in reference [4], one can undertand this as a result of atoms trickling from one dressed state to other, eventually reaching the the state  $|\Psi_{-}\rangle$ . On the other hand, when only  $E_z$  beam is present, then all the atoms in  $|g_{\pm}\rangle$  will be optically pumped out and eventually reach  $|g_0\rangle$ . This is a trap state for the  $E_z$  beam. The two trap states  $|\Psi_0\rangle = |g_0\rangle$ and  $|\Psi_{-}\rangle$  can now be mapped to the qubit states  $|\Psi_0\rangle = |0\rangle$  and  $|\Psi_{-}\rangle = |1\rangle$ .

More interestingly, if both  $E_p$  and  $E_z$  beams are present together, the steady state solution then is not a statistical mixture of the two trap states  $|\psi_0\rangle$  and  $|\psi_-\rangle$ , but a three component CPT states [7]

$$|\Psi\rangle = \frac{(\Omega_p/\Omega_z)|g_o\rangle - |g_-\rangle + |g_+\rangle}{\sqrt{2 + |(\Omega_p/\Omega_z)|^2}},\tag{1}$$

which can be rewritten as

 $<sup>{}^{1}|3</sup>S_{1/2}, F = 1\rangle$  and  $|3P_{1/2}, F = 1\rangle$  of Na is an equally valid setup with equivalent configuration. We use the dipole-dipole interactions between one Na and one <sup>87</sup>Rb atom also in a later part of this paper.

$$|\psi\rangle = \sin(\theta/2)|\psi_0\rangle + \exp(i\phi)\cos(\theta/2)|\psi_-\rangle.$$
 (2)  
Or

$$\Psi \rangle = \sin(\theta/2)|0\rangle + \exp(i\phi)\cos(\theta/2)|1\rangle, \tag{3}$$

where

5

$$\sin(\theta/2) = \frac{\Omega_p}{\sqrt{2|\Omega_z|^2 + |\Omega_p|^2}} \quad \text{and} \quad \cos(\theta/2) = \frac{(\sqrt{2}\Omega_z)}{\sqrt{2|\Omega_z|^2 + |\Omega_p|^2}} \tag{4}$$

and  $(\theta/2) = \tan^{-1} \left( \Omega_p / \sqrt{2} \Omega_z \right)$ . Any desired value of  $\theta$  can be obtained by varying the ratio of  $\left( \Omega_p / \sqrt{2} \Omega_z \right)$ , where  $\Omega_{p,z} = d \cdot E_{p,z} / 2\hbar$ . The phase factor  $\phi$  in (3) can also be obtained by controlling the phase between the two beams  $E_{p,z} = \mathcal{E}_{p,z} \exp[i(\omega t - k \cdot x - \phi_{p,z})]$ . If the setup is as in figure 1a, then rotating the HWP will distribute the intensity between  $E_p$  and  $E_z$  and will positioning it appropriately will produce any desired  $\theta$ . Keeping a variable retarder at one of the output ports of PBS will also control the phase  $\phi$ .

The operation then can be mathematically expressed by

$$H(\theta) = \begin{pmatrix} \sin(\theta/2) & e^{i\phi}\cos(\theta/2) \\ -e^{-i\phi}\cos(\theta/2) & \sin(\theta/2) \end{pmatrix},$$
(5)

which, acting on the basis vectors

$$|0\rangle \equiv \begin{pmatrix} 1\\0 \end{pmatrix}$$
 and  $|1\rangle \equiv \begin{pmatrix} 0\\1 \end{pmatrix}$ , (6)

leads to dressed state vectors

$$|\Phi_{-}\rangle \equiv \begin{pmatrix} \sin(\theta/2) \\ -e^{-i\phi}\cos(\theta/2) \end{pmatrix} \text{ and } |\Phi_{+}\rangle \equiv \begin{pmatrix} e^{i\phi}\cos(\theta/2) \\ \sin(\theta/2) \end{pmatrix}.$$
(7)

Operator  $H(\theta)$ , reduces to a Hadamard when  $\theta/2$  is set to  $45^0$  and  $\phi = 1$ , which is equivalent of setting the halfwaveplate of fig. 1a to at  $45^0$ . The state  $\Phi_+$  is is a CPT state, as given in equation (3)

This scheme for state preparation has certain distinct advantages. (i) The qubit states (6) as well as state  $\Phi_+$  are CPT states. CPT states are end points of atom-laser interaction and the atoms eventually reach CPT states via non-CPT states as shown by Cohen-Tannoudji and Reynaud [4]. This means that the state preparation is reliable and the desired state is always prepared. (ii) Once the states are prepared, the atoms in these state no longer interact with the laser that prepares them. This eliminates need for precise time-control of the lasers. The state preparation is therefore robust and certain. (iii) The state preparation involves only cw beams and does not require any complex pulse shaping schemes. (v) Since it does not involve single photon processes, lasers with nominally high intensity can be used. This would allow very precise control of phase  $\phi$  while allowing fluctuations in the intensity. (vi) Any desired superposition corresponding to any desired Bloch vector can be prepared by simply varying the intensity ratio between two laser beams. Due to all of above, the configuration allows a robust and reliable preparation of two qubit states and its superposition and also the method of state preparation is very easy. In the following sections, methods of performing single qubit and two-qubit operations are discussed.

# 2. Operations of logic gates

## 2.1 Single Qubit Operations

Setting  $\theta/2 = 0$  in (5) will result in a rotation, which is the NOT operation

$$H(\theta = 0) = \begin{pmatrix} 0 & e^{i\phi} \\ -e^{-i\phi} & 0 \end{pmatrix},$$
(8)

which converts  $|\psi\rangle = \sin(\theta/2)|0\rangle + \exp(i\phi)\cos(\theta/2)|1\rangle$  to  $|\psi\rangle = \cos(\theta/2)|0\rangle + \exp(i\phi)\sin(\theta/2)|1\rangle$ , for any value of existing  $\theta/2$ .

This is an intriguing situation since setting  $\theta/2 = 0$  in equation (5) is equivalent to setting  $\Omega_p = 0$  in (4), which is equivalent to switching off  $E_p$  beam and thus always creating the atoms in state  $|1\rangle$ , no matter what is the original state. This discrepancy can be understood in the manner that the NOT operation always operates on the full dressed state  $|\Phi_+\rangle$  and hence valid.

# 2.2 Two-qubit operation; C-Phase gate

Two-qubit opeations can be obtained in a manner similar to the earlier works that exploited the dipole-dipole interaction [11, 13], except using magnetic dipole-dipole interaction between spin states instead of electric dipoles.

In an external magnetic field  $\vec{B}$ , the spin vectors align at an angle that depends on their m<sub>F</sub>value and also makes a Larmor precision about  $\vec{B}$ , with a frequency  $\omega_L = \gamma_L |B|$ .  $\gamma_L$  is the gyromagnetic ratio of the atom and |B| is the value of the magnetic field. The atom can now be flipped from one m<sub>F</sub>state to the other by applying an oscillatory magnetic field perpendicular to  $\vec{B}$ , and at a frequency equal to difference between the two corresponding Larmor frequencies. The dipole-dipole interaction between the spins now manifest as a shift in the Larmor frequencies and hence the resonance frequency for the oscillatory magnetic field also shifts as shown in figure 2. [9, 10]

As in case of electric dipoles, the spin dipole interaction is also inversely dependent on the cube of the distance between them, given by

$$V_{dd} = \frac{\mu_0}{4\pi} \frac{\gamma_L^2}{r^3} \left[ S_1 . S_2 - 3(S_1 . n)(S_2 . n) \right]. \tag{9}$$

Which gets reduced to  $V_{dd} = \frac{\mu_0}{4\pi} \frac{\gamma_L^r}{r^3} (3\cos^2\theta_s - 1)$ , for two degenerate  $m_F$  levels[9]. Here *r* is the normal distance between the two atoms,  $\theta_s$  the angle between the spin

Here *r* is the normal distance between the two atoms,  $\theta_s$  the angle between the spin directions and  $\vec{r}$ ,  $\mu_0$  the permittivity of free space and the ratio  $\mu_0/4\pi$  is a scaling factor for MKS units. The energy levels of the states atom pairs can be shown as in figure 2a. This interaction *V* causes a mixing of the pair states  $|01\rangle$  and  $|10\rangle$  as well as cause a shift in the energies as shown in figure 2 b. The energy for the transition  $|00\rangle \leftrightarrow |10\rangle + |01\rangle$  is shifted by  $\Omega_m = 2(\mu_0/(4\pi)) \cdot (\gamma_L^2/r^3)$ 

If a pulse of oscillatory magnetic field at frequency  $\omega_L + 2\Omega$  the atoms will absorb its energy only if they are in the state  $|10\rangle + |01\rangle$ , and travel to state  $|11\rangle$ . If such a pulse has a McCall-Hahn pulse area of  $2\pi$ , the the atom will return to state  $|10\rangle + |01\rangle$ , but now with a phase factor of  $\pi$ . Atoms in any other state will not be affected. If the atom pair is in state  $|10\rangle - |01\rangle$  instead, the phase factor already exists for  $|01\rangle$  state. Therefore, if the atoms are brought together, the rf pulse applied and then taken apart, only the atoms in state  $|01\rangle$  will return to  $-|01\rangle$ .



**Figure 2.** (a) Energy diagram of two-atom system (b) shows effect of the dipole-dipole interaction. The state  $|10\rangle - |01\rangle$  is not coupled by radio frequency transition either to  $|00\rangle$  or to  $|11\rangle$ . The energy difference in case of non-interaction situation is equal to  $\hbar\omega_L$  where  $\omega_L$  is the Larmor frequency. See text for amount of shift in case of (b).



**Figure 3.** Energy diagram of two-atom system for two different atoms. (a) without the spin dipole-dipole interaction and (b) Energy shifts due to dipole interaction. Note that there is no mixing. dashed lines in (b) are position of unshifted energy levels.

Another option, following Ryabstev and co-workers [11] is to bring the atoms together and hold them close for a specific period. Since the dipole-dipole interaction causes a mixing of the states  $|01\rangle$  and  $|10\rangle$  for form a time dependent superposition

$$|\Psi_{dd}(t)\rangle = \cos(V_{dd}t/\hbar)|10\rangle - i \sin(V_{dd}t/\hbar)|01\rangle.$$

, the atom pair oscillates between  $|01\rangle$  and  $|10\rangle$  with a half-period  $T = 1/2(\hbar\pi)/V_{dd}$ . The scheme then involves holding the atoms together for *T* and taking apart, which gives a control-swap gate or for 2*T*, which will result in a controlled swap gate.

The serious drawback of this scheme is that one can not distinguish a-priori between control atom and the logic atom since they are both identical. An option then is to use two different atoms, with same level configuration.

## 2.3 heterogeneous atoms

Sodium, with  $3S_{1/2}$ , F = 1 and  $3P_{1/2}$ , F = 1 triplets, shows an identical behaviour of state preparation and qubit operations, but the corresponding Larmor frequency is different. The dipole-dipole interaction between spins of Sodium atom and <sup>87</sup>Rb will cause only a level shift instead of a mixing states  $|01\rangle$  and  $|10\rangle$  as showin figure 3 [9]. The amount of shift is  $\Omega = (\mu_0/4\pi)(\gamma_1\gamma_2/r^3)(3\cos^2\theta_s - 1)$ , where  $\gamma_1$  and  $\gamma_2$  are gyromagnetic ratios of the Sodium and <sup>87</sup>Rb atoms respectively.

Now a controlled NOT gate can be obtained by using a rf pulse with a frequency  $\omega_2 + \Omega$  with a McCall-Hahn area of  $2\pi$ , or a pair of pulses with frequencies  $\omega_2 + \Omega$  and  $\omega_1 + \Omega$ 

times in a STIRAP like fashion to obtain a controlled swap gate.

## 3. conclusion

It is shown that a system that exhibits two trapping states can be obtained in <sup>87</sup>Rb and Sodium atom interacting with two lasers that couple its F=1  $\leftrightarrow$  F=1 transition. They can be mapped to the qubit states  $|1\rangle$  and  $|0\rangle$  and can be used for quantum computation. Any required superposition state  $|\psi\rangle = \sin(\theta/2)|0\rangle + \exp(i\phi)\cos(\theta/2)|1\rangle$  can be prepared. Since this involves CPT states and also ground levels, it is very robust against decoherences. Single qubit and two-qubit operations are described with these states. However, the dipole-dipole interaction between spin states is weaker than that between electric dipole states and hence the shift is small. But the typical value of Larmor frquency for most alkali atoms are about a 100 kHz and measuring a small shift in the Radio frequency domain is technologically feasible. The major technical difficulty with this scheme may be with having to move the atoms nearer and apart as and when required for the two qubit operation.

## 4. acknowledgment

Several useful discussions led to fine tuning of this work. In particular, I thank Hema Ramachandran of Raman Institute and R. Srikanth of Poorna Prajna Institute, Bangalore, Prasanta Panigrahi of IISER, Kolkota and Surya P. Tewari of University of Hyderabad, for helpful discussions.

## References

- Neilson M. A. and Chuang. I.L. *Quantum Computation and Quantum Information*, (Cambridge, 2000); Joachim Stolze & Dieter Suter, *Quantum Computing*, (Wiley-VCH, Weinheim 2004)
- [2] Khitrin, A. K. and B. M. Fung, *Phys. Rev.* A 64, 32306 (2001), Ranbir Das and Anil Kumar, *Phys. rev.* A 68, 032304 (2003) and references therein. J. I. Cirac and P. Zoller, *Phys. Rev. Lett.* 74, 4091 (1995), Ferdinand Schmidt-Kaler, Hartmut Häffner, Mark Riebe, Stephan Gulde, Gavin P. T. Lancaster, Thomas Deuschle, Christoph Becher, Christian F. Roos, Jürgen Eschner & Rainer Blatt, *Nature*, 422, 408 (2003), Daniel Loss and David diVincenzo, *Phys. Rev.* A 57, 120 (1998)
- [3] E. Arimondo Progress in Optics, XXXV, 257 ed: E.Wolf, (North Holland 1996)
- [4] C. Cohen-Tannoudji and S. Reynaud, J.Phys. B10 345 . (1977) ibid, 2311.
- [5] Alan Corney, Atomic and Laser Spectroscopy, (Clarendon, Oxford, 1977)
- [6] Dmitry Budker, Derek F. Kimball and David P. DeMille, *Atomic Physics*, (Oxford University Press, 2004), section 2.5
- [7] Ashok Vudayagiri and Surya P. Tewari, J. Phys. B 39, 3919(2006)
- [8] J. R. Kuklinski, U. Gaubatz, F. T. Hioe, and K. Bergmann, Phys. Rev. A 40 (1989),6741
- [9] C. Cohen-Tannoudji, B. Diu and F. Laloë, *Quantum Mechanics*, (John Wiley, 1977), (see section 'Complement B<sub>X1</sub>' on page 1120)
- [10] Jackson, J. D. Classical Electrodynamics, (Wiley Eastern, New Delhi, 1989)
- [11] I. I. Ryabtsev et.al. J. Phys. B 38 (2005), 421
- [12] S. L. McCall and E. L. Hahn, *Phys. Rev. Lett.* 18 908, (1967); L. Allen and J. H. Eberly, *Optical Resonance and two level atoms*, (Dover, New York, 1987)
- [13] Brennen G. K., Caves C. M., Poul S. Jessen and Ivan H. Deutsch, *Phys. Rev. Lett.* 82(1999), 1060