Analysis of the role of von Neumann's projection postulate in the canonical scheme of quantum teleportation and main quantum algorithms

Andrei Khrennikov

School of Mathematics and Systems Engineering International Center of Mathematical Modeling in Physics and Cognitive Sciences Växjö University, S-35195, Sweden

November 26, 2018

Abstract

Modern development of quantum technologies based on quantum information theory stimulated analysis of proposed computational, cryptographic and teleportational schemes from the viewpoint of quantum foundations. It is evident that not all mathematical calculations performed in complex Hilbert space can be directly realized in physical space. Recently by analyzing the original EPR paper we found that they argument was based on the misuse of the von Neumann's projection postulate. Opposite to von Neumann, Einstein, Podolsky and Rosen (EPR) applied this postulate to observables represented by operators with degenerate spectra. It was completely forbidden by von Neumann's axiomatics of QM. It is impossible to repeat the EPR considerations in the von Neumann's framework. In this note we analyze quantum teleportation by taking into account von Neumann's projection postulate. Our analysis shows that so called quantum teleportation is impossible in von Neumann's framework. On the other hand, our analysis implies that the main quantum algorithms are totally consistent with von Neumann's projection postulate.

1 Introduction

As a consequence of tremendous development of theoretical basis of quantum information theory, quantum technologies became an established domain of experimental research (in particular in laser physics) which is directed toward realization of market-oriented projects in future . Such a situation stimulated analysis of proposed computational, cryptographic and delectation schemes from the viewpoint of quantum foundations. It is evident that not all mathematical calculations performed in complex Hilbert space can be directly realized in physical space, cf. e.g. [1] - [6]. In particular, quantum information theory (if its is not considered as a purely mathematical formalism) should be coupled to quantum measurement theory. The most extensive analysis of foundations of this theory was performed by von Neumann [7]. His book became really the Bible of the so called Copenhagen interpretation of QM.¹

Therefore by considering different constructions which arise in quantum information theory we should check their matching with (in particular) von Neumann's theory of quantum measurement. One of the most intriguing modern constructions of quantum information theory is the quantum teleportation scheme [15]. In this note we shall analyze it from the viewpoint of the quantum measurement theory. As it might be already expected from our analysis of the EPR paper [16], the quantum teleportation scheme does not match von Neumann's theory

¹We recall that by this interpretation any state of an *individual physical system* is described by a wave function ψ . This interpretation was created by Bohr, Heisenberg, Pauli, von Neumann, Fock, Landau. It is commonly used in quantum experimental research. It is important for our further considerations to point out that: "The state of a system after measurement is determined by the von Neumann projection postulate." This interpretation is typically confronted with so called statistical (or ensemble) interpretation. By the latter a wave function ψ is not an attribute of a single physical system (e.g. electron). A wave function ψ (as well as a density matrix ρ) describes an ensemble of identically prepared physical system. Here the projection postulate determines not the state of a system (after the corresponding measurement), but the probability distribution of an ensemble of (output-)systems. This interpretation is due to Einstein, Podolsky, Rosen, De Broglie, Schrödinger, Bohm,..., Ballentine, De Baere, Manko, Khrennikov,... Its modern version is often called the Växjö interpretation of QM, see [8], [9]: contextual statistical realist interpretation of QM. The latter interpretation combines the ensemble interpretation of quantum state (not only mixed, but even pure!) with Bohr's principle of complementarity. We remark that, although the the ensemble interpretation of quantum state is widely used in theoretical research, e.g. quantum tomography [10]–[14], it is still not common for quantum experimental physics.

of measurement. In principle, one can proceed up to the very last step of the quantum teleportation scheme. However, to finalize the quantum teleportation procedure, Bob should perform a measurement in the basis which is unknown for him.

On the other hand, our analysis implies that the main quantum algorithms are totally consistent with von Neumann's projection postulate.

2 Von Neumann's projection postulate

Von Neumann performed a very deep analysis of the measurement process described by QM. One of the fundamental questions which he studied was determination of the output state after measurement. Suppose that an observable A was measured and the value A = awas obtained. This measurement was performed for some initially prepared state, for simplicity we assume that it was a pure state $|\psi\rangle$. What is a post-measurement state corresponding to this result? The answer is given by the projection postulate [7]:

Part 1: If the spectrum of the operator A (we shall use the same symbol for an observable and the corresponding operator) is nondegenerate, then the post-measurement state is given by the eigenvector corresponding to the eigenvalue a.

This part of the projection postulate is well known. But, unfortunately, the second (not less fundamental) part of the projection postulate is missing in the majority of books and papers:

Part 2: If the spectrum of the operator A is degenerate, then the post-measurement *state is not determined*. To determine it, a refinement measurement should be performed.

By a refinement measurement von Neumann understood measurement of any observable C which is compatible with A and represented by the operator with nondegenerate spectrum such that results of the A-measurement can be obtained as a function of results of the C-measurement, i.e., A = f(C).

The crucial point is that in all experiments with **entangled sys**tems $S_1 + S_2$ partial measurements on such systems (e.g. on one of particles) are described by operators with degenerate spectra, see [16] for details. Acting in the tensor product $H_1 \otimes H_2$ induces degeneration (which was absent in e.g. S_1 -Hilbert space H_1). Thus such a measurement does not determine state. To determine the out-put state, one should perform a refinement measurement. But such a refinement measurement is always performed on both particles. The latter fact destroys all dreams about quantum nonlocality and its possible technological applications.

3 Analysis (due to von Neumann) of the teleportation scheme

We shall proceed across the quantum teleportation scheme, see [15] or simply the corresponding article in wikipedia, and point to applications of the projection postulate. There are Alice (A) and Bob (B), and Alice has a qubit in some arbitrary quantum state $|\psi\rangle$. Assume that this quantum state is not known to Alice and she would like to send this state to Bob. Suppose Alice has a qubit that she wants to teleport to Bob. This qubit can be written generally as: $|\psi\rangle = \alpha |0\rangle + \beta |1\rangle$.

The quantum teleportation scheme requires Alice and Bob to share a maximally entangled state before, for instance one of the four Bell states: $|\Phi^+ \rangle = \frac{1}{\sqrt{2}}(|0 \rangle_A \otimes |0 \rangle_B + |1 \rangle_A \otimes |1 \rangle_B), |\Phi^- \rangle = \frac{1}{\sqrt{2}}(|0 \rangle_A \otimes |0 \rangle_B - |1 \rangle_A \otimes |1 \rangle_B), |\Psi^+ \rangle = \frac{1}{\sqrt{2}}(|0 \rangle_A \otimes |1 \rangle_B + |1 \rangle_A \otimes |0 \rangle_B)$ $), |\Psi^- \rangle = \frac{1}{\sqrt{2}}(|0 \rangle_A \otimes |1 \rangle_B - |1 \rangle_A \otimes |0 \rangle_B)$. Alice takes one of the particles in the pair, and Bob keeps the other one. We will assume that Alice and Bob share the entangled state $|\Phi^+ \rangle$. So, Alice has two particles (the one she wants to teleport, and A, one of the entangled pair), and Bob has one particle, B. In the total system, the state of these three particles is given by

$$|\psi \rangle \otimes |\Phi^+ \rangle = (\alpha |0 \rangle + \beta |1 \rangle) \otimes \frac{1}{\sqrt{2}} (|0 \rangle \otimes |0 \rangle + |1 \otimes |1 \rangle)$$

Alice will then make a partial measurement in the Bell basis on the two qubits in her possession. To make the result of her measurement clear, we will rewrite the two qubits of Alice in the Bell basis via the following general identities (these can be easily verified): $|0 > \otimes |0 > = \frac{1}{\sqrt{2}}(|\Phi^+ > + |\Phi^- >), |0 > \otimes |1 > = \frac{1}{\sqrt{2}}(|\Psi^+ > + |\Psi^- >), |1 > \otimes |0 > = \frac{1}{\sqrt{2}}(|\Psi^+ > - |\Psi^- >), |1 > \otimes |1 > = \frac{1}{\sqrt{2}}(|\Phi^+ > - |\Phi^- >),$ Evidently the results of her (local) measurement is that the three-particle state would collapse to one of the following four states (with equal probability of obtaining each): $|\Phi^+ > \otimes (\alpha |0 > +\beta |1 >), |\Phi^- > \otimes (\alpha |0 > \beta |1 >$

 $-\beta|1\rangle$, $|\Psi^+\rangle \otimes (\beta|0\rangle + \alpha|1\rangle$), $|\Psi^-\rangle \otimes (-\beta|0\rangle + \alpha|1\rangle$). The four possible states for Bob's qubit are unitary images of the state to be teleported. The crucial step, the local measurement done by Alice on the Bell basis, is done. It is clear how to proceed further. Alice now has complete knowledge of the state of the three particles; the result of her Bell measurement tells her which of the four states the system is in. She simply has to send her results to Bob through a classical channel. Two classical bits can communicate which of the four results she obtained. After Bob receives the message from Alice, he will know which of the four states his particle is in. Using this information, he performs a unitary operation on his particle to transform it to the desired state $\alpha|0\rangle + \beta|1\rangle$:

If Alice indicates her result is $|\Phi^+\rangle$, Bob knows his qubit is already in the desired state and does nothing. This amounts to the trivial unitary operation, the identity operator.

If the message indicates $|\Phi^-\rangle$, Bob would send his qubit through the unitary gate given by the Pauli matrix $\sigma_3 = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}$ to recover the state. If Alice's message corresponds to $|\Psi^+\rangle$, Bob applies the gate $\sigma_1 = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$ to his qubit. Finally, for the remaining case, the appropriate gate is given by $\sigma_3\sigma_1 = i\sigma_2 = \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}$. Teleportation is therefore achieved.

The main problem is that Alice's measurement is represented by a degenerate operator in the 3-qubit space. It is nondegenerate with respect to her 2-quibits, but not in the total space. Thus the standard conclusion that by obtaining e.g. A = 1, Alice can be sure that Bob obtained the right state $|\psi\rangle$, does not match the quantum measurement theory. According to von Neumann, to get this state Bob should perform a refinement measurement. To to perform it Bob, should know the state $|\psi\rangle$. It seems that it is the end of the story about quantum teleportation. We remark that quantum teleportation does trivially not applicable if one uses so called statistical [18] (or ensemble) interpretation of QM. However, in this paper we used not it, but the conventional Copenhagen interpretation due to von Neumann.

A number of people (e.g. Richard Gill and Marcus Appleby) who commented my reanalysis of the role of projection postulate in the modern version of the Copenhagen interpretation and especially in quantum information pointed out that, even if Einstein, Podolsky and Rosen used the von Neumann projection postulate in the improper way, the EPR-version of the projection postulate is nowadays commonly used. I agree with them that, although EPR did not use the real von Neumann postulate, but they used the version of the projection postulate which is nowadays known as Luders' postulate [17] (by which it is possible to consider operators with degenerate spectra in the same way as nondegenerate), see [16] for details. Here the out-put state is given by the projection of the original state onto the corresponding eigenspace. But the price of this use is quantum nonlocality which also became canonical in modern QM. It was completely absent in the original Copenhagen interpretation, since if one follows J. von Neumann no trace of quantum nonlocality would be found. I recall that so called EPR states were studied in details in von Neumann's book, but without EPRs paradoxical consequences: either incompleteness or nonlocality. One could not completely exclude the possibility that the original von Neumann's analysis of quantum measurements was wrong and that, in spite of his demand for sharp distinguishing of measurements for observables with degenerate and nondegenerate spectra, it is always possible to apply Lüders' postulate. We also remark that conclusion of an ultimate von Neumann's mathematical analysis of quantum measurement procedures totally coincide with views of Niels Bohr, see his reply to Einstein [19]. In such a case it should be openly pointed out that the interpretation of QM which is commonly used in quantum information theory is not the conventional Copenhagen interpretation of Bohr-Heisenberg-von Neumann-Fock-Landau-Pauli, ..., but a new "quantum information interpretation of QM." Opposite to the conventional Copenhagen interpretation, its QI-version is based on Lüders' postulate and hence quantum nonlocality. We recall that, although EPR had also used Lüders' postulate, they still considered nonlocality as totally unphysical. Their output from misuse of the von Neumann projection postulate was "naive realism" – assigning values of two incompatible physical variables to the same physical system. By the QI-interpretation of QM such a possibility is excluded, because of Bell's theorem. Although Bell's theorem does not provide an ultimate proof of this statement, see e.g. [20] and [6] and literature hereby, I would agree with rejection of naive realism. However, I point out that, opposite to the QI-interpretation, the conventional Copenhagen interpretation could peacefully escape both nonlocality and naive realism via the application of the proper version (namely,

von Neumann's one) of the projection postulate.²

Conclusion. If one proceed in the Copenhagen framework in the proper way, the quantum teleportation scheme would not work.

4 Quantum algorithms and von Neumann's projection postulate

4.1 Deutsch-Jozsa algorithm

Let us start with the Deutsch-Jozsa algorithm. Since we are interested only in the final measurement (unitary evolutions given by quantum gates are not important for our analysis), we just write the out-put state of quantum computation:

$$|\psi\rangle = \frac{1}{\sqrt{2^n}} \sum_{z=0}^{2^n-1} \sum_{x=0}^{2^n-1} (-1)^{xz+f(x)} |z\rangle \otimes \left[\frac{|0\rangle - |1\rangle}{\sqrt{2}}\right].$$
(1)

Then we perform measurement of "what is written" in the argument register z. We consider the operator

$$\widehat{P} = \sum_{z=0}^{2^n - 1} z | z > \otimes | z > .$$
(2)

This is the operator with nodegenerate spectrum in the "argument-space". Since the final state $|\psi\rangle$ is factorized into the "argument state",

$$\frac{1}{\sqrt{2^n}} \sum_{z=0}^{2^n-1} \sum_{x=0}^{2^n-1} (-1)^{xz+f(x)} |z| >$$

and the "function state" $\left[\frac{|0\rangle-|1\rangle}{\sqrt{2}}\right]$, we can forget about the last one.

4.2 Simon's algorithm

We now move to Simon's algorithm. Here the output state after a cicle of quantum computation is given by

$$|\psi\rangle = \frac{1}{\sqrt{2^n}} \sum_{k=0}^{2^n-1} \left(\frac{1}{\sqrt{2^n}} \sum_{j=0}^{2^n-1} (-1)^{jk} | j \rangle \otimes |f(k)\rangle \right).$$
(3)

²Thus it could be done without Bell's inequality.

Although this state is not factorized, we can proceed as it will be descfribed in the following subsection.

4.3 Application of von Neumann's quantum formalism to partial measurements

First, we recall von Neumann's formalization of Born's probability postulate (the probabilistic interpretation of the wave function):

(PI) The probability that in the state ψ the quantity with operator \widehat{R} take on values from an interval Δ_1 is

$$P_{\psi}(R \in \Delta_1) = ||E(\Delta_1)\psi||^2, \tag{4}$$

where $E(\lambda)$ is the resolution of the identity belonging to \widehat{R} .

We point out that \widehat{R} need not have a nondegenerate spectrum! Thus the PI-postulate and the projection postulate are not identical in their structures. One might say that PI is closer to Lüders' postulate (which is simply von Neumann's postulate without taking into account the structure of spectrum).

Let H_1 and H_2 be two complex finite dimensional Hilbert spaces, dim $H_i \ge 2$. Let $\hat{a}_1 : H_1 \to H_1$ be a self-adjoint operator. The Hilbert space H_i represents (quantum) states of the system $s_i, i = 1, 2$. The operator \hat{a}_1 represents an observable a_1 corresponding to measurements on s_1 . The composite system $s = (s_1, s_2)$ is described by the tensor product space $H = H_1 \otimes H_2$.

Suppose that \hat{a}_1 has purely discrete nondegenerate spectrum: $\alpha_j, j = 1, ..., N = \dim H_1$.

We remark that measurement of a_1 on s_1 can be considered as a measurement on $s = (s_1, s_2)$. It is represented by the operator $\widehat{A}_1 = \widehat{a}_1 \otimes I$. Its spectrum coincides with spectrum of \widehat{a}_1 .

Suppose that $s = (s_1, s_2)$ is described by a state $\psi \in H$. We performed the a_1 -measurement. Let $a_1 = \alpha_j$.

PROB). On the one hand, it is a measurement on $s = (s_1, s_2)$. It is described by \widehat{A}_1 . Thus by PI the probability to get $a_1 = \alpha_j$ is given by

$$P(a_1 = \alpha_j) = \|E_{\alpha_j} \otimes I\psi\|^2.$$

Thus the probability of the result of a partial measurement (i.e. on one system) is given by the PI-postulate for the state space of the composite system s. PROJ). On the other hand, the same measurement can be considered as simply a measurement on s_1 (if systems are isolated at the moment of measurement!). We recall that in H_1 the spectrum of \hat{a}_1 is nondegenerate. Hence, by the von Neumann's projection postulate the resulting state in H_1 would always be the eigenstate $|\alpha_j \rangle$ of \hat{a}_1 – independently of the initial state of s_1 . The latter independence from the initial state is important, because by determining the state $|\psi \rangle$ of the composite system s, we do not (in general) determine states of subsystems. Thus, in spite of this difficulty, we are able to determine the resulting state in H_1 after a partial measurement.

We now apply von Neumann's formalism to the state (3) which is produced at the end of a cycle of Simon's algorithm and the operator (2) which plays the role of \hat{a}_1 in previous considerations. By PROB we shall get the probability of the result z; by PROJ we really get a state $|j\rangle$ which is orthogonal to the vector $|x\rangle$ determining period.

In the same way we can consider Grover's algorithm and Shor's algorithm.

This paper was written during my visit to Danish Technical University (Copenhagen) which was supported by Informatics and Mathematical Modelling grant of this university. Introductory lectures on quantum information which I gave to students and teachers of the department of Computer Science and Engineering stimulated my analysis of quantum information schemes. I would like to thank Paul Fischer for hospitality and discussions on possibility of practical realization of quantum information schemes from the point of view of (classical) computer science.

References

- I. V. Volovich, Quantum Cryptography in space and Bell's theorem, in : Foundations of Probability and Physics, Ed. A. Khrennikov, World Sci. 2001, pp.364-372.
- [2] I.V. Volovich, Towards Quantum Information Theory in Space and Time, in: Quantum Theory: Reconsideration of Foundations, Ed. A.Khrennikov, Vaxjo University Press, 2002, pp.423-440.
- [3] Y. Ozhigov, Classical collective behavior instead of quantum dynamics, arXiv:quant-ph/0702237 (2007).

- Y. Ozhigov, Simulation of quantum dynamics via classical collective behavior arXiv:quant-ph/0602155 (2006).
- [5] Y. Ozhigov, Amplitude quanta in multiparticle system simulation, *Russian Microelectronics*, 35, 1, 53-65 (2006).
- [6] K. Hess and W. Philipp, "Bell's theorem: critique of proofs with and without inequalities", in *Foundations of Probability and Physics*-3, edited by A. Yu. Khrennikov, AIP Conference Proceedings Ser. **750**, Melville, New York, 2006, pp. 150-157.
- [7] J. von Neumann, Mathematical foundations of quantum mechanics, Princeton Univ. Press, Princeton, N.J., 1955.
- [8] A. Khrennikov. Vxj Interpretation of Quantum Mechanics. http://xxx.lanl.gov/abs/quant-ph/0202107.
- [9] A. Khrennikov, On foundations of quantum theory. In: Quantum Theory: Reconsideration of Foundations. Ser. Math. Modelling, 2, 163-196, Växjö Univ. Press, Växjö, 2002.
- [10] V. I. Manko, J. of Russian Laser Research, 17, 579-584 (1996).
- [11] V. I. Manko and E. V. Shchukin, J. Russian Laser Research, 22, 545-560 (2001).
- [12] M. A. Manko, V. I. Manko, R. V. Mendes, J. Russian Laser Research, 27, 507-532.
- [13] S. De Nicola, R. Fedele, M. A. Man'ko and V. I. Man'ko, Quantum tomography, wave packets, and solitons. J. of Russian Laser Research, 25, 1071-2836, 2004.
- [14] O. V. Manko and V. I. Manko, J. Russian Laser Research, 25, 477-492 (2004).
- [15] C. H. Bennett, G. Brassard, C. Crpeau, R. Jozsa, A. Peres, W. K. Wootters, Teleporting an Unknown Quantum State via Dual Classical and Einstein-Podolsky-Rosen Channels, *Phys. Rev. Lett.*, **70**, 1895-1899 (1993).
- [16] A. Khrennikov, On the problem of completeness of QM: von Neumann against Einstein, Podolsky, and Rosen. arXiv:0805.1511.
- [17] G. Lüders, Ann. Phys., Lpz, 8, 322 (1951).
- [18] L. E. Ballentine, *Quantum mechanics*, Englewood Cliffs, New Jersey, 1989.
- [19] N. Bohr, Phys. Rev. 48, 696-700 (1935).

[20] A. Khrennikov, Analysis of explicit and implicit assumptions in theorems of J. von Neumann and J. Bell. J. of Russian Laser Research, 28, 244-254 (2007).