

Cosmological Kaluza-Klein branes in black brane spacetimes

Masato Minamitsuji*

*Arnold-Sommerfeld-Center for Theoretical Physics, Department für Physik,
Ludwig-Maximilians-Universität, Theresienstr. 37, D-80333, Munich, Germany*

We discuss cosmological evolutions of a Kaluza-Klein (KK) brane in the $D(> 6)$ -dimensional bulk spacetime. By “KK brane”, we mean the brane which involves KK compactified spatial dimensions as well as ordinary four dimensional spacetime dimensions into the worldvolume. The bulk spacetime is composed of two copies of a patch of D -dimensional black three-brane solution. The near-horizon geometry of the black brane spacetime is given by $AdS_5 \times S^{(D-5)}$ and in the asymptotic infinity the spacetime approaches D -dimensional Minkowski. We excise the region $r > r_b$ (r_b is a certain radial position) from the original black brane spacetime and glue two copies of the remaining spacetime at $r = r_b$. The KK brane is located at $r = r_b$. We identify these two copies by imposing the reflection (Z_2) symmetry with respect to the KK brane. In order to realize cosmology on the brane, we consider the motion of the KK brane into the above bulk spacetime toward the asymptotic infinity. We find that the brane Universe cannot reach the asymptotic infinity, irrespectively of the components of matter on the brane. The early time cosmology coincides with that in the second Randall-Sundrum (RS II) model. The deviation from the extremal condition gives rise to the dark radiation type contribution on the brane. We also discuss several experimental and observational bounds.

Since the stimulating proposals by Randall and Sundrum (RS) [1, 2], the cosmological aspects of braneworld models have been studied in various literature [3]. In the second RS (RS II) model [1], at the low energy scales the four-dimensional cosmology is recovered on the brane due to the strong warping of the extra dimension. Then, six and higher-dimensional models have started to attract growing interests. In higher dimensions, there will be various types of braneworld models, for example, those with flux-stabilized compact extra dimensions, with brane intersections. Codimension-two brane models with flux-stabilized extra dimensions have been originally focused on as the simple realization of large extra dimensions [4] and as a way of resolution of the cosmological constant problem [5] (see, however, e.g., [6]). Recently, these six dimensional models have been discussed from the different point of views. It is well-known that a four-dimensional defect in six or higher dimensions generically has a problem on localization of ordinary matter on the brane due to the stronger self-gravity. Ways to circumvent this problem should be developed. In six dimensional models, the ways of brane regularization to study gravity and cosmology on the brane have been studied, e.g., in [7, 8, 9, 10]. Here, we focus on the scheme employed in [7, 8, 10], in which the original three-brane is replaced with a four-brane with a compactified Kaluza-Klein (KK) circle. Such a way can be generalized to the cases of higher dimensions and called *Kaluza-Klein braneworld*. General properties of KK braneworlds have been investigated, e.g., in Ref. [11].¹ One possible way

to construct the cosmological models from the more fundamental higher-dimensional theory is to start from the dimensionally reduced theory after a compactification. However, the dynamics of compactified extra-dimensions could generically affect the brane dynamics non-trivially and then the lower-dimensional effective theory would break down. In order to take the dynamics of extra spatial dimensions, the scheme of KK braneworld is the one of the most rigorous approaches.

Cosmological Kaluza-Klein branes:

We briefly review the cosmology on a $(D - 1)$ -dimensional KK brane in D -dimensional spacetime. We apply the well-established method developed in Ref. [16]. We start from the following static $D(> 6)$ -dimensional metric ansatz:

$$ds^2 = -F(r)^2 dt^2 + A(r)^2 \delta_{ij} dx^i dx^j + B(r)^2 dr^2 + C(r)^2 h_{ab} dy^a dy^b. \quad (1)$$

We assume the Z_2 symmetry with respect to the brane. h_{ab} represents the metric of $(D - 5)$ -dimensional compact space. The indices $i(= 1, 2, 3)$ and a run over the ordinary three-dimensional space and $(D - 5)$ -dimensional compact space, respectively. The brane proper time τ and induced metric are given by

$$-F(r(\tau))^2 \dot{t}^2 + B(r(\tau))^2 \dot{r}^2 = -1, \\ ds_{\text{ind}}^2 = -d\tau^2 + a^2 \delta_{ij} dx^i dx^j + C(a)^2 h_{ab} dy^a dy^b. \quad (2)$$

where the cosmological scale factor is given by $a = A|_{r=r_b}$ (r_b is the radial position of the brane) and $\dot{}$ means the derivative with respect to the brane proper time τ .

*Email: masato.minamitsuji@physik.uni-muenchen.de

¹ See for several attempts to realize the localization of ordinary matter and gravity on the brane in higher dimensions, e.g., by constructing regular solitonic solutions [12], by replacing the brane with the core region where matter is distributed smoothly

[13], by adding higher curvature terms to the theory [14] and by considering brane intersections [15].

Then, we derive the equations of motion of the brane in the static bulk. The (t, t) , (i, j) (more precisely $(i, j) - (t, t)$) and (a, b) (more precisely $(a, b) - (t, t)$) components of Israel junction condition are given by

$$\left(\frac{3}{a} + \frac{(D-5)C_{,a}}{C}\right)\sqrt{\tilde{B}^{-2} + \dot{a}^2} = \frac{\rho}{2M_D^{D-2}}, \quad (3)$$

$$\begin{aligned} & \frac{\ddot{a} + \frac{\tilde{B}_{,a}}{\tilde{B}}\dot{a}^2}{\sqrt{\tilde{B}^{-2} + \dot{a}^2}} + \left(\frac{F_{,a}}{F} - \frac{1}{a}\right)\sqrt{\tilde{B}^{-2} + \dot{a}^2} \\ &= -\frac{1}{2M_D^{D-2}}(\rho + p), \end{aligned} \quad (4)$$

$$\begin{aligned} & \frac{\ddot{a} + \frac{\tilde{B}_{,a}}{\tilde{B}}\dot{a}^2}{\sqrt{\tilde{B}^{-2} + \dot{a}^2}} + \left(\frac{F_{,a}}{F} - \frac{C_{,a}}{C}\right)\sqrt{\tilde{B}^{-2} + \dot{a}^2} \\ &= -\frac{1}{2M_D^{D-2}}(\rho + q), \end{aligned} \quad (5)$$

where $\tilde{B}da = Bdr$ and M_D represents the D -dimensional gravitational scale. ρ , q and p are $((D-5)$ -dimensional) energy density, pressure along the internal dimensions and that along the external dimensions, respectively. If there are dynamical bulk degrees of freedom than the metric, we need to take their jump conditions into account. The (t, t) -component of the Israel junction conditions Eq. (3) gives the effective Friedmann equation:

$$\left(\frac{\dot{a}}{a}\right)^2 = \frac{\rho^2}{4M_D^{2(D-2)}\left(3 + \frac{(D-5)a}{C}\frac{dC}{da}\right)^2} - \frac{1}{a^2\tilde{B}^2}. \quad (6)$$

Black three-brane solutions:

As one of the simplest higher-dimensional solutions, we consider the D -dimensional black three-brane solutions (see e.g., [17]). We start from the following D -dimensional theory:

$$S = \frac{M_D^{D-2}}{2} \int d^D X \sqrt{-G} \left(R[G] - \frac{1}{2(D-5)!} F_{[D-5]}^2 \right), \quad (7)$$

where G_{AB} and $R[G]$ are the D -dimensional metric and the Ricci scalar associated with the metric G_{AB} , respectively. $F_{[D-5]}$ represents $(D-5)$ -form field strength. This theory contains a series of the black three-brane solutions, whose metric is given by

$$\begin{aligned} ds^2 &= H^{-1/2} \left(-f dt^2 + \delta_{ij} dx^i dx^j \right) \\ &+ H^{2/(D-6)} \left(f^{-1} dr^2 + r^2 h_{ab} dy^a dy^b \right), \end{aligned} \quad (8)$$

where δ_{ij} and h_{ab} are metric of flat three-dimensional space and $(D-5)$ -sphere, respectively. We also define

$$H(r) := 1 + \frac{Q}{r^{D-6}}, \quad f(r) = 1 - 2\frac{\mu}{r^{D-6}}, \quad (9)$$

and h_{ab} is the metric of $(D-5)$ -sphere. μ and Q are parameters of the solutions. We assume that the function f is positive. Thus, the radial coordinate r has the minimal

value at the horizon position $r = r_{\min} := (2\mu)^{1/(D-6)}$, whereas there is no finite maximal value of the coordinate r . The case that $\mu = 0$ corresponds to the extremal solutions. The near horizon geometry is $\text{AdS}_5 \times S^{(D-5)}$. The effective curvature radius of AdS spacetime is given by the parameter Q

$$\ell := \frac{4}{D-6} Q^{1/(D-6)}. \quad (10)$$

The $(D-5)$ -form field dynamically acts on $S^{(D-5)}$. The magnetic $(D-5)$ -form field strength is given by

$$F^{y^1 \dots y^{(D-5)}} = \frac{1}{\sqrt{-G}} \epsilon^{y^1 \dots y^{D-5}} E'(r) \quad (11)$$

where

$$E(r) = \sqrt{\frac{Q}{Q + 2\mu} \frac{D-2}{2(D-6)} \frac{f(r)}{H(r)}}. \quad (12)$$

At the spatial infinity $r \rightarrow \infty$, the spacetime approaches D -dimensional Minkowski. In the case $D = 10$, the solution corresponds to a stack of coincident D3-branes at the low energy scales.

The reason to consider the black brane spacetime is as follows: Firstly, the spacetime contains the warped spatial dimension. The well-behaved cosmological feature of the RS II model is due to the realization of the warped structure. Such a warped structure of extra dimensions may also help to construct the realistic brane cosmological models in higher dimensions. Also, in string compactifications, the highly warped structure is induced gravitationally by the branes localized at a certain place in the internal space. The black brane spacetime is one of the simplest examples of such spacetimes. Secondly, how to make the compact extra-dimensions invisible is also non-trivial problem. For this purpose, the flux-stabilization of extra dimensions would be useful. In the black brane solutions, the magnetic $(D-5)$ -form makes the size of the spherical dimensions invisible in the near-horizon region.

We apply the previously mentioned KK braneworld scheme to the black brane spacetime. The bulk geometry is constructed by the standard cut-and-paste procedure. We excise the region $r > r_b$ from the original black brane spacetime, where r_b is a certain radial position, and glue two copies of the remaining spacetime at $r = r_b$. Then, two copies of the patch $r_{\min} < r < r_b$ are bounded by a codimension-one brane at $r = r_b$. The compact spatial dimensions along $(D-5)$ -sphere (as well as the ordinary four-dimensional spacetime dimensions) are involved into the brane worldvolume as KK compactified dimensions. We identify two copies of the bulk by imposing the reflection (Z_2 -)symmetry with respect to the KK brane. An expanding cosmology could be realized as a brane motion from the spatial black brane horizon toward the spatial infinity.

Cosmology on the KK brane:

The bulk metric functions A , B , C and F in Eq. (1) are now

$$\begin{aligned} A &= \frac{1}{H^{1/4}}, \quad B = \left(\frac{f^{1/2}}{H^{1/(D-6)}} \right)^{-1}, \\ C &= rH^{1/(D-6)}, \quad F = \frac{f^{1/2}}{H^{1/4}}. \end{aligned} \quad (13)$$

The cosmic scale factor is defined by $a(\tau) := A(r_b)$. For the nonzero μ , the cosmic scale factor has non-zero minimal value:

$$a_{\min} = \left(\frac{2\mu/Q}{1 + 2\mu/Q} \right)^{1/4}, \quad (14)$$

which corresponds to $r = r_{\min}$. On the other hand, in any case, the cosmic scale factor also has the maximal size $a = 1$ for $r \rightarrow \infty$. However, as we will see the brane Universe cannot reach $a = 1$. In order to have enough cosmological expansions, we assume the near-extremal condition $\mu/Q \ll 1$. The physical size of KK directions on the brane is given by $C = Q^{1/(D-6)}/(1 - a^4)^{1/(D-6)}$

and thus in the near horizon region $a \ll 1$ is as small as the effective AdS radius ℓ .

After integrating over the KK dimensions, the effective four-dimensional energy density is given by

$$\Sigma = \rho \frac{Q^{(D-5)/(D-6)} \Omega_{(D-5)}}{(1 - a^4)^{(D-5)/(D-6)}}. \quad (15)$$

We decompose this into the constant and time-dependent parts: $\Sigma = \Sigma_0 + \Sigma_1$, where Σ_0 is the constant part of the effective brane energy density and Σ_1 is the remaining time-dependent part. The constant part of the brane energy density can also be decomposed into two parts $\Sigma_0 = \Sigma_{\text{RS}} + \delta$, where

$$\Sigma_{\text{RS}} := 6 \left(\frac{D-6}{4} \right) M_D^{D-2} Q \Omega_{(D-5)} \left(1 + \frac{2\mu}{Q} \right)^{1/2}. \quad (16)$$

In such a decomposition, from Eq. (6), we obtain the effective Friedmann equation:

$$\begin{aligned} \left(\frac{\dot{a}}{a} \right)^2 &= -\frac{8(D-5)}{Q^{2/(D-6)}} \left(\frac{D-6}{4} \right)^2 \left(1 + \frac{2\mu}{Q} \right) \frac{a^4(1-a^4)^{2+2/(D-6)} [3(D-6) - a^4(D-8)]}{[3(D-6) + (D-2)a^4]^2} + \frac{1}{3} \lambda_{\text{eff}} \frac{(1-a^4)^{4+2/(D-6)}}{\left(1 + \frac{D-2}{3(D-6)} a^4 \right)^2} \\ &+ \frac{1}{3M_4^2} \frac{(1-a^4)^{4+2/(D-6)}}{\left(1 + \frac{D-2}{3(D-6)} a^4 \right)^2} \Sigma_1 + \left(\frac{D-6}{4} \right)^2 \frac{\left(1 + \frac{2\mu}{Q} \right)}{Q^{2/(D-6)}} \left(\frac{\Sigma_1}{\Sigma_{\text{RS}}} \right)^2 \frac{(1-a^4)^{4+2/(D-6)}}{\left(1 + \frac{D-2}{3(D-6)} a^4 \right)^2} \\ &+ \left(\frac{D-6}{4} \right)^2 (1-a^4)^{2+2/(D-6)} \frac{2\mu}{a^4 Q^{1+2/(D-6)}}. \end{aligned} \quad (17)$$

Here,

$$M_4^2 := \frac{4}{D-6} \frac{M_D^{D-2} Q^{(D-4)/(D-6)} \Omega_{D-5}}{\left(1 + \frac{2\mu}{Q} \right)^{1/2} \left(1 + \frac{\delta}{\Sigma_{\text{RS}}} \right)}. \quad (18)$$

and

$$\frac{\lambda_{\text{eff}}}{3} := \ell^{-2} \left(1 + \frac{2\mu}{Q} \right) \left(\frac{2\delta}{\Sigma_{\text{RS}}} + \frac{\delta^2}{\Sigma_{\text{RS}}^2} \right), \quad (19)$$

are the effective four-dimensional gravitational coupling and cosmological constant, respectively, since in the early times $a \ll 1$, the Friedmann equation is reduced to the form

$$\left(\frac{\dot{a}}{a} \right)^2 \approx \frac{1}{3} \lambda_{\text{eff}} + \frac{1}{3M_4^2} (\Sigma_1 + \Sigma_{\text{DR}}) + \frac{1 + \frac{2\mu}{Q}}{\ell^2} \left(\frac{\Sigma_1}{\Sigma_0} \right)^2 \quad (20)$$

The length scale ℓ is given by Eq. (10). We also define the energy density:

$$\Sigma_{\text{DR}} := 3M_4^2 \left(\frac{D-6}{4} \right)^2 \frac{2\mu}{Q^{1+2/(D-6)} a^4}. \quad (21)$$

The result is nothing but the one in the RS II model [3]. Especially, in the low energy regime $\Sigma_1 \ll \Sigma_{\text{RS}}$, the standard cosmology is recovered except for the term Σ_{DR} . By combining Eq. (16) with (18), we find

$$\Sigma_{\text{RS}} \simeq \frac{6M_4^2}{\ell^2} \left(1 + \frac{2\mu}{Q} \right). \quad (22)$$

In the case that $\delta = 0$, as in the original model of RS [1], the effective cosmological constant λ_{eff} vanishes. The deviation from the extremal condition $\mu \neq 0$ gives rise to the so-called *dark radiation* (see Ref.[3]), given by Σ_{DR} . In the RS II cosmology, the dark radiation arises due to the presence of a black hole in the bulk. In the current case, similarly the near-horizon geometry is approximately given by the product of the five-dimensional Schwarzschild-AdS with $S^{(D-5)}$. As we will see later, the pressure equation in the corresponding bulk region also has the same structure as in the RS II cosmology.

In the later times, cosmology could deviate from the RS one. To see this, it is useful to rewrite the effective Friedmann equation into the form in an analogy with the classical mechanics: $\dot{a}^2 + U(a) = 0$, where for instance,




FIG. 1: The function $U(a)$ is shown as a function of a for $D = 10$, $\delta/\Sigma_{\text{RS}} = 0.5$ and $\Sigma_1 = 0$.

in the extremal case $\mu = 0$, the potential term is given by

$$U(a) := -\ell^{-2}a^2(1-a^4)^{2+2/(D-6)} \times \left[\frac{(1-a^4)^2 \left(1 + \frac{\delta}{\Sigma_{\text{RS}}} + \frac{\Sigma_1}{\Sigma_{\text{RS}}}\right)^2}{\left(1 + \frac{(D-2)}{3(D-6)}a^4\right)^2} - 1 \right]. \quad (23)$$

As easily seen, the potential term becomes positive before arriving the maximal value $a = a_{\text{max}}$. In the case of the vacuum brane $\Sigma_1 = 0$, this maximal value of cosmic scale factor is given by

$$a_{\text{max}} = \left(\frac{\delta}{\delta + \frac{4(D-5)}{3(D-6)}\Sigma_{\text{RS}}} \right)^{1/4} \quad (24)$$

In Fig. 1 and 2, the typical behaviors of the potential $U(a)$ without the ordinary matter $\Sigma_1 = 0$ and with dust fluid on the brane are shown, respectively. The function $U(a)$ vanishes at $a = a_{\text{max}}$. After the brane reaches the maximal size, the brane goes back toward $a \rightarrow 0$. Similar behaviors can be seen in the near-extremal case. Thus, we find that *the brane Universe cannot reach the asymptotic infinity, irrespectively of the components of the matter on the brane*. In other words, observers on the brane never reach the KK compactified extra dimensions. The result implies that the cosmological brane cannot escape from the gravitational potential induced by the black brane horizon.

As previously mentioned, in the special case that $D = 10$, the black three-brane solution corresponds to a low energy description of a stack of N coincident (BPS for $\mu = 0$) D3-branes in type IIB string theory. These parameters in the solution can be expressed as $M_{10}^8 = 2/(2\pi)^7 g_s^2 \alpha'^4$, $Q = \ell^4 = 4\pi N g_s \alpha'^2$ where $\ell_s = (\alpha')^{1/2}$ and g_s represent string length scale and string coupling constant, respectively. Note that the supergravity description of D3-branes is valid in the case that the effective AdS curvature scale $\ell = Q^{1/4}$ is much larger than ℓ_s (hence $N g_s \gg 1$) and quantum gravitational corrections are negligible, i.e., $\ell \gg M_{10}^{-1}$ (hence $N \gg 1$). Thus,




FIG. 2: The function $U(a)$ is shown as a function of a for $D = 10$, $\delta/\Sigma_{\text{RS}} = 0.5$. $\Sigma_1/\Sigma_0 = 0.005/a^3$ is chosen as dust fluid.

assuming the near-extremal condition $\mu/Q \ll 1$ and the optimistic choice of the brane tension $\delta/\Sigma_{\text{RS}} = O(1)$, the effective four-dimensional quantities are also written as

$$M_4^2 \simeq \frac{N^{3/2}}{8\pi^{5/2} g_s^{1/2} \alpha'}, \quad \Sigma_{\text{RS}} \simeq \frac{3N}{8\pi^3 g_s \alpha'^2}. \quad (25)$$

Although our approach is higher-dimensional, in the sense that we start from the original higher-dimensional theory and then secondly integrate over the internal dimensions after obtaining the junction conditions, we now confirm that in the early time the brane cosmology obtained in our approach is equivalent to the one in the approach in the effective theory approach in which the internal dimensions are integrated over firstly.

Experimental and observational bounds:

As in the cases of the RS II cosmology (see e.g., [18]), several experimental and observational bounds are imposed. A lower bound on the RS brane tension is given by the condition that the density square term in the effective Friedmann equation must be suppressed before the Big Bang nucleosynthesis (BBN): $\Sigma_{\text{RS}} > (1\text{MeV})^4$. But a more stringent bound is given from the results of the table top tests on Newton's law, implying $\ell < 0.1\text{mm}$: $\Sigma_{\text{RS}} > (10^3\text{GeV})^4$ in the assumption that $\mu/Q \ll 1$, since $M_4 \approx 10^{18}\text{GeV}$.

In the ten-dimensional ($D = 10$) case, from the Eq. (25), we find

$$N^2 \simeq \frac{24\pi^2 M_4^4}{\Sigma_{\text{RS}}}, \quad g_s \alpha'^2 \simeq \left(\frac{3}{2}\right)^{3/2} \frac{M_4^2}{\pi^2 \Sigma_{\text{RS}}^{3/2}}. \quad (26)$$

The condition that $\Sigma_{\text{RS}} > (10^3\text{GeV})^4$ (hence $\ell < 0.1\text{mm}$) implies $N < 10^{30}$ and $M_{10} \sim (g_s \alpha'^2)^{-1/8} > 10^{-5}\text{GeV}$. Thus, there is a large parameter space which is consistent with the conditions that $N g_s \gg 1$ and $N \gg 1$ for which the classical description of the solution is valid. Note that $(\ell/0.1\text{mm}) \simeq (N/10^{30})$, since $N \simeq 2\pi M_4 \ell$.

As we have seen, in the near-horizon region the effective cosmology on the brane is approximated well by the

one in the RS II model. According to the *AdS/conformal field theory (CFT)*, more precisely *AdS₅/CFT₄, correspondence* [19], which states that in type IIB string theory the (super)gravity theory in the five-dimensional AdS spacetime is equivalent to the $\mathcal{N} = 4$ super-conformal $U(N)$ Yang-Mills theory at the boundary, gravity in RS II braneworld seems to be equivalent to the four-dimensional gravity coupled to CFT on the brane. There are several examples in which the equivalence (up to $O(\ell^2)$) has been confirmed [20], including the case of cosmological branes. Such an equivalence should be valid for our cosmological brane. The five-dimensional effective gravitational scale is related to the four-dimensional one $M_5^3 \approx M_4^2/\ell$. Then, the CFT parameters are related to the five-dimensional gravitational ones: for instance, the CFT degrees of freedom is given by $N^2 \approx (2\pi)^2 M_5^3 \ell^3$. Since ℓ^{-1} and N^2 are the cut-off energy scale and the degrees of freedom of CFT, respectively, the relation $M_4^2 \simeq N^2 \ell^{-2}$ can be seen as a typical example that the bound $M_4^2 \gtrsim n\Lambda^2$ (discussed in Ref. [22]) is saturated. Here, n is the number of species of quantum fields with mass scale Λ .

Inspired by the AdS/CFT correspondence, it has been conjectured that a black hole on a RS II brane is classically unstable [21]. Such a conjecture is based on the argument that the a five dimensional black hole localized on the brane should be equivalent to the four-dimensional quantum-corrected black hole of the gravity theory coupled to CFT. From the CFT point of

view, such a brane black hole could decay into large ($\sim N^2$) CFT degrees of freedom and its lifetime τ_{BH} should be much shorter than that of standard one: $\tau_{\text{BH}} \simeq 10^4 (M_{\text{BH}}/M_\odot)^3 (0.1\text{mm}/\ell)^2 \text{year}$ (or equivalently, $\tau_{\text{BH}} \simeq 10^4 (M_{\text{BH}}/M_\odot)^3 (10^{30}/N)^2 \text{year}$) [21], where M_{BH} and M_\odot are the masses of a black hole and the Sun, respectively. If the conjecture in RS II model is true, of course this must be true in our construction (as long as the brane stays in the near horizon region). This could give a stronger bound on ℓ . For example, in order for a black hole with solar mass $M_{\text{BH}} \simeq M_\odot$, to survive today, τ_{BH} must be longer than the age of the Universe $\sim 10^{10} \text{year}$: this condition requires $\ell < 10^{-4} \text{mm}$ (and hence $N < 10^{27}$).

Effective pressure equations:

The other non-trivial components of Israel junction condition dominate the acceleration/deceleration of our brane Universe. As well as the effective energy density Eq. (15), by integrating over the KK directions, the effective four-dimensional pressure along the ordinary three-space P and that along the KK directions \tilde{Q} are introduced from the $((D-1)$ -dimensional) pressures p and q , respectively. As for the effective energy density Σ , those pressures are also decomposed into the constant and time-dependent parts in general: $P = P_0 + P_1$, $\tilde{Q} = \tilde{Q}_0 + \tilde{Q}_1$. Note that the constant part is given by $P_0 = \tilde{Q}_0 = -\Sigma_0$. By substituting the explicit expressions for $F_{,a}/F$, $\tilde{B}_{,a}/\tilde{B}$ and $C_{,a}/C$ into Eqs. (4) and (5), we obtain

$$\begin{aligned} & \frac{\ddot{a}}{a} - \frac{(D-6) - a^4(5D-26)}{(D-6)(1-a^4)} \frac{\dot{a}^2}{a^2} + \left(\frac{D-6}{4}\right)^2 \frac{4\mu(1-a^4)^{2+2/(D-6)}}{a^4 Q^{1+2/(D-6)}} = -\frac{1}{2M_4^2} \frac{(1-a^4)^{3+2/(D-6)}}{1 + \frac{D-2}{3(D-6)} a^4} \left(1 + \frac{\Sigma_1}{\Sigma_0}\right) (\Sigma_1 + P_1) \\ & \frac{\ddot{a}}{a} + \frac{\dot{a}^2}{a^2} \frac{4a^4}{1-a^4} + \left[1 - \frac{4a^4}{(1-a^4)(D-6)} + \frac{\frac{4\mu}{\tilde{Q}}}{a^4(1+\frac{2\mu}{\tilde{Q}}) - \frac{2\mu}{\tilde{Q}}}\right] \left(\frac{D-6}{4}\right)^2 \frac{(1-a^4)^{2+2/(D-6)}}{a^4 Q^{2/(D-6)}} (a^4(1+\frac{2\mu}{\tilde{Q}}) - \frac{2\mu}{\tilde{Q}}) \\ & = -\frac{1}{2M_4^2} \frac{(1-a^4)^{3+2/(D-6)}}{1 + \frac{D-2}{3(D-6)} a^4} \left(1 + \frac{\Sigma_1}{\Sigma_0}\right) (\Sigma_1 + \tilde{Q}_1), \end{aligned} \quad (27)$$

where M_4 is defined in Eq. (18). In the early times $a \ll 1$ and low energy scales $\Sigma_1/\Sigma_0 \ll 1$, the external pressure equation can be reduced to the form

$$\frac{\ddot{a}}{a} - \frac{\dot{a}^2}{a^2} \approx -\frac{1}{2M_4^2} (\Sigma_1 + P_1 + \Sigma_{\text{DR}} + P_{\text{DR}}), \quad (28)$$

where the dark energy density and pressure are given by Eq. (21) $P_{\text{DR}} = \Sigma_{\text{DR}}/3$. This is nothing but the result in the standard cosmology except for the dark radiation type contribution in the non-extremal case.

Hereafter, we focus on the extremal case $\mu = 0$ and discuss the behavior of the pressure in the KK direction

\tilde{Q}_1 . In the early times $a \ll 1$, we find

$$\frac{1}{Q^{1/2}} \left(1 + \frac{\delta}{\Sigma_{\text{RS}}}\right)^2 \approx \frac{1}{6M_4^2} (3P_1 - 2\Sigma_1 - 3\tilde{Q}_1). \quad (29)$$

Note that in the above Eq. (29), the left-hand-side is constant. Thus, in the right-hand-side, the combination $3P_1 - 2\Sigma_1 - 2\tilde{Q}_1$ must be totally time-independent. Also, in the above derivation, it is assumed that $\Sigma_1/\Sigma_0 \ll 1$. For the realistic brane matter $P_1/\Sigma_0 \ll 1$ should also be satisfied. In order for Eq. (29) to be compatible, \tilde{Q}_1 should be as large as $|\tilde{Q}_1| \sim \Sigma_0$ in contrast to Σ_1 and P_1 .

One might think that the junction conditions for the $(D-5)$ -form field might restrict the motion of the brane. But now, it is not the case. Only the non-trivial com-

ponent of the $(D - 5)$ -form field strength is $Fy^1 \cdots y^{D-5}$, where all the coordinates y^a represent the dimensions of $S^{(D-5)}$, and the product of the form field with the normal vector $n_A F^{A_1 \cdots A_{D-5}}$ is vanishing. Thus all the junction conditions of the form become trivial. In other words, the $(D - 5)$ -form field cannot couple with the brane matter, if it is magnetic.

Summary:

We discussed cosmology in the brane Universe in $D(> 6)$ -dimensional bulk spacetime. The scheme called *Kaluza-Klein braneworld* is one of the well-motivated ways of brane regularization, in discussing the behavior of a self-gravitating brane in higher dimensions. A KK brane involves the compact spatial dimensions as well as the ordinary four-dimensional spacetime into the brane worldvolume.

We consider the D -dimensional black three-brane solutions. In the near horizon region the spacetime structure is approximately given by $\text{AdS}_5 \times S^{(D-5)}$ and in the asymptotic infinity it approaches the D -dimensional Minkowski. We excise the region $r > r_b$ from the original black brane spacetime, and glue two copies of the remaining spacetime at $r = r_b$. Then, two copies of the patch $r_{\min} < r < r_b$ are bounded by a codimension-one KK brane at $r = r_b$. The KK brane now involves the compact spatial dimensions along $(D - 5)$ -sphere into the worldvolume. We identify two copies of the bulk by imposing the reflection (Z_2 -)symmetry with respect to the KK brane. We derive all the components of Israel junction conditions, by assuming exact cosmological symmetry on the (KK) brane. The junction conditions for the $(D - 5)$ form field do not restrict the brane motion. In other words, the brane matter does not couple

to the bulk $(D - 5)$ -form field. After integrating over the compact KK dimensions, we have derived the effective cosmological equations on the brane. We find that irrespectively of components of the matter on the brane, the brane cannot reach the asymptotic infinity of the bulk. Thus, observers on the (KK) brane never see these $(D - 5)$ extra dimensions.

In the early times, when the KK brane moves in the near-horizon region, the brane cosmology coincides with that in the five-dimensional Randall-Sundrum (RS II) model. If the original black three-brane solution is near-extremal $\mu/Q \ll 1$, the dark radiation type contribution arises. It is natural since the near horizon geometry is the product of five-dimensional AdS-Schwarzschild with $S^{(D-5)}$, which is similar to the RS model. We also discussed experimental and observational bounds.

There would be various extensions of our considerations. As one of the possible extensions, it would be important to consider the cases without Z_2 -symmetry across the brane, which are rather generic in the higher dimensions. In the present set-up, the physical size of our Universe approaches the maximal size. This would be the disadvantage of our simplest model. In order to avoid this problem and to obtain ever expanding brane Universe, it would be useful to consider the time-dependence of the extra dimensions.

Acknowledgement

This work was supported in part by the Transregional Research Centre TRR33 ‘‘The Dark Universe’’.

-
- [1] Phys. Rev. Lett. **83**, 4690 (1999) [arXiv:hep-th/9906064].
 - [2] L. Randall and R. Sundrum, Phys. Rev. Lett. **83**, 3370 (1999) [arXiv:hep-ph/9905221].
 - [3] V. A. Rubakov, Phys. Usp. **44**, 871 (2001) [Usp. Fiz. Nauk **171**, 913 (2001)] [arXiv:hep-ph/0104152]; D. Langlois, Prog. Theor. Phys. Suppl. **148**, 181 (2003) [arXiv:hep-th/0209261]; R. Maartens, Living Rev. Rel. **7**, 7 (2004) [arXiv:gr-qc/0312059].
 - [4] N. Arkani-Hamed, S. Dimopoulos and G. R. Dvali, Phys. Lett. B **429**, 263 (1998) [arXiv:hep-ph/9803315]; I. Antoniadis, N. Arkani-Hamed, S. Dimopoulos and G. R. Dvali, Phys. Lett. B **436**, 257 (1998) [arXiv:hep-ph/9804398].
 - [5] J. W. Chen, M. A. Luty and E. Ponton, JHEP **0009**, 012 (2000) [arXiv:hep-th/0003067]; S. M. Carroll and M. M. Guica, arXiv:hep-th/0302067; I. Navarro, JCAP **0309**, 004 (2003) [arXiv:hep-th/0302129]; Y. Aghababaie, C. P. Burgess, S. L. Parameswaran and F. Quevedo, Nucl. Phys. B **680**, 389 (2004) [arXiv:hep-th/0304256].
 - [6] I. Navarro, Class. Quant. Grav. **20**, 3603 (2003) [arXiv:hep-th/0305014]; H. P. Nilles, A. Papazoglou and G. Tasinato, Nucl. Phys. B **677**, 405 (2004) [arXiv:hep-th/0309042]; H. M. Lee, Phys. Lett. B **587**, 117 (2004) [arXiv:hep-th/0309050]; J. Vinet and J. M. Cline, Phys. Rev. D **70**, 083514 (2004) [arXiv:hep-th/0406141]; J. Garriga and M. Porrati, JHEP **0408**, 028 (2004) [arXiv:hep-th/0406158].
 - [7] J. Louko and D. L. Wiltshire, JHEP **0202**, 007 (2002) [arXiv:hep-th/0109099]. B. M. N. Carter, A. B. Nielsen and D. L. Wiltshire, JHEP **0607**, 034 (2006) [arXiv:hep-th/0602086].
 - [8] M. Peloso, L. Sorbo and G. Tasinato, Phys. Rev. D **73**, 104025 (2006) [arXiv:hep-th/0603026]; E. Papantonopoulos, A. Papazoglou and V. Zamarias, JHEP **0703**, 002 (2007) [arXiv:hep-th/0611311]; T. Kobayashi and M. Minamitsuji, Phys. Rev. D **75**, 104013 (2007) [arXiv:hep-th/0703029]; F. Arroja, T. Kobayashi, K. Koyama and T. Shiromizu, arXiv:0710.2539 [hep-th].
 - [9] J. Vinet and J. M. Cline, Phys. Rev. D **70**, 083514 (2004) [arXiv:hep-th/0406141]; J. Vinet and J. M. Cline, Phys. Rev. D **71**, 064011 (2005) [arXiv:hep-th/0501098];
 - [10] E. Papantonopoulos, A. Papazoglou and V. Zamarias, arXiv:0707.1396 [hep-th]; M. Minamitsuji and D. Langlois, Phys. Rev. D **76**, 084031 (2007) [arXiv:0707.1426 [hep-th]].
 - [11] N. Chatillon, C. Macesanu and M. Trodden, Phys.

- Rev. D **74**, 124004 (2006) [arXiv:gr-qc/0609093]; D. Yamauchi and M. Sasaki, Prog. Theor. Phys. **118N2**, 245 (2007) [arXiv:0705.2443 [gr-qc]]; S. Kanno, D. Langlois, M. Sasaki and J. Soda, Prog. Theor. Phys. **118N4**, 701 (2007) [arXiv:0707.4510 [hep-th]]
- [12] T. Gherghetta, E. Roessl and M. E. Shaposhnikov, Phys. Lett. B **491**, 353 (2000) [arXiv:hep-th/0006251]; I. Oda, Phys. Rev. D **64**, 026002 (2001) [arXiv:hep-th/0102147]; E. Roessl and M. Shaposhnikov, Phys. Rev. D **66**, 084008 (2002) [arXiv:hep-th/0205320].
- [13] B. Carter, R. A. Battye and J. P. Uzan, Commun. Math. Phys. **235**, 289 (2003) [arXiv:hep-th/0204042]; R. A. Battye, B. Carter and A. Mennim, Phys. Rev. D **71**, 104026 (2005) [arXiv:hep-th/0412053]; I. Navarro and J. Santiago, JHEP **0502**, 007 (2005) [arXiv:hep-th/0411250]; S. A. Appleby and R. A. Battye, Phys. Rev. D **76**, 124009 (2007) [arXiv:0707.4238 [hep-ph]].
- [14] C. Charmousis and R. Zegers, JHEP **0508**, 075 (2005) [arXiv:hep-th/0502170]; C. Charmousis and R. Zegers, Phys. Rev. D **72**, 064005 (2005) [arXiv:hep-th/0502171].
- [15] I. Navarro and J. Santiago, JHEP **0404**, 062 (2004) [arXiv:hep-th/0402204]; E. Gravanis and S. Willison, J. Math. Phys. **47**, 092503 (2006) [arXiv:hep-th/0412273];
- [16] H. A. Chamblin and H. S. Reall, Nucl. Phys. B **562**, 133 (1999) [arXiv:hep-th/9903225]; P. Kraus, JHEP **9912**, 011 (1999) [arXiv:hep-th/9910149]; A. Kehagias and E. Kiritsis, JHEP **9911**, 022 (1999) [arXiv:hep-th/9910174]; D. Ida, JHEP **0009**, 014 (2000) [arXiv:gr-qc/9912002].
- [17] G. T. Horowitz and A. Strominger, Nucl. Phys. B **360**, 197 (1991); M. J. Duff, R. R. Khuri and J. X. Lu, Phys. Rept. **259**, 213 (1995) [arXiv:hep-th/9412184]; K. S. Stelle, arXiv:hep-th/9701088;
- [18] J. M. Cline, C. Grojean and G. Servant, Phys. Rev. Lett. **83**, 4245 (1999) [arXiv:hep-ph/9906523]; R. Maartens, D. Wands, B. A. Bassett and I. Heard, Phys. Rev. D **62**, 041301 (2000) [arXiv:hep-ph/9912464]; D. Langlois, R. Maartens, M. Sasaki and D. Wands, Phys. Rev. D **63**, 084009 (2001) [arXiv:hep-th/0012044].
- [19] J. M. Maldacena, Adv. Theor. Math. Phys. **2**, 231 (1998) [Int. J. Theor. Phys. **38**, 1113 (1999)] [arXiv:hep-th/9711200]; S. S. Gubser, I. R. Klebanov and A. M. Polyakov, Phys. Lett. B **428**, 105 (1998) [arXiv:hep-th/9802109]; E. Witten, Adv. Theor. Math. Phys. **2**, 253 (1998) [arXiv:hep-th/9802150].
- [20] S. W. Hawking, T. Hertog and H. S. Reall, Phys. Rev. D **62**, 043501 (2000) [arXiv:hep-th/0003052]; M. J. Duff and J. T. Liu, Class. Quant. Grav. **18**, 3207 (2001) [Phys. Rev. Lett. **85**, 2052 (2000)] [arXiv:hep-th/0003237]; T. Shiromizu and D. Ida, Phys. Rev. D **64**, 044015 (2001) [arXiv:hep-th/0102035]; T. Tanaka, arXiv:gr-qc/0402068.
- [21] T. Tanaka, Prog. Theor. Phys. Suppl. **148**, 307 (2003) [arXiv:gr-qc/0203082]; R. Emparan, A. Fabri and N. Kaloper, JHEP **0208**, 043 (2002) [arXiv:hep-th/0206155]; R. Emparan, J. Garcia-Bellido and N. Kaloper, JHEP **0301**, 079 (2003) [arXiv:hep-th/0212132].
- [22] G. Dvali, arXiv:0706.2050 [hep-th]; G. Dvali and M. Redi, Phys. Rev. D **77**, 045027 (2008) [arXiv:0710.4344 [hep-th]].