

Universal slow fall-off to the unique AdS infinity in Einstein-Gauss-Bonnet gravity

Hideki Maeda*

Centro de Estudios Científicos (CECS), Arturo Prat 514, Valdivia, Chile

(Dated: January 26, 2023)

In this paper, the following two propositions are proven under the dominant energy condition for the matter field in the higher-dimensional spherically symmetric spacetime in Einstein-Gauss-Bonnet gravity in the presence of a cosmological constant Λ . First, for $\Lambda \leq 0$ and $\alpha \geq 0$ without a fine-tuning to give a unique anti-de Sitter vacuum, where α is the Gauss-Bonnet coupling constant, vanishing generalized Misner-Sharp mass is equivalent to the maximally symmetric spacetime. Under the fine-tuning, it is equivalent to the vacuum class I spacetime. Second, under the fine-tuning with $\alpha > 0$, the asymptotically anti-de Sitter spacetime in the higher-dimensional Henneaux-Teitelboim sense is only a special class of the vacuum class I spacetime. The latter means the universal slow fall-off to the unique anti-de Sitter infinity in the presence of physically reasonable matter.

PACS numbers: 04.20.Cv, 04.20.Ha, 04.50.-h.

1. Introduction: Gravitation physics in higher dimensions is a prevalent subject of current research motivated mainly by string theory. In this context, it is well known that the most natural extension of general relativity in higher dimensions as a theory of quasi-linear second-order partial differential equations is not general relativity itself but Lovelock gravity [1].

The Lovelock Lagrangian comprises the dimensionally extended Euler densities. In n dimensions, the first $[n/2]$ curvature terms appear in the field equations, where $[x]$ denotes the integer part of x . In the even-dimensional case, however, the last $((n/2)$ -th) term becomes a topological invariant and does not contribute to the field equations. The Lovelock tensor $\mathcal{G}_{\mu\nu}$ derived from the Lovelock Lagrangian has the following properties: (1) $\mathcal{G}_{\mu\nu}$ is symmetric, (2) $\mathcal{G}_{\mu\nu}$ contains up to the second derivative of the metric, (3) $\nabla_\nu \mathcal{G}^{\mu\nu} \equiv 0$, and (4) $\mathcal{G}_{\mu\nu}$ is linear in the second derivative of the metric. The fourth condition is derived by other three in four dimensions.

Lovelock gravity including general relativity as a special case is obviously a gauge theory for the (local) Lorentz group because its Lagrangian comprises the Lorentz scalars. However, it is not a gauge theory for the Poincaré group in general, which is the standard symmetry group in particle physics [2]. From the viewpoint of the gauge principle for the unified theory, it is expected that the gravitation theory is represented as a gauge theory for the Poincaré group or some group which contains the Lorentz group and the symmetry group analogous to translations in a flat spacetime. Miraculously, under the fine-tuning between the coupling constants, Lovelock gravity can be a gauge theory for the Poincaré, de Sitter (dS), or anti-de Sitter (AdS) group. The last two groups are the smallest nontrivial choices of such required groups containing the translation symmetry group on a pseudosphere. Unfortunately, this miracle happens only in *odd* dimensions, nevertheless this so-called Chern-Simons gravity has been of particular interest as an aesthetic way to the unified theory [2].

On the other hand, among all the classes of Lovelock gravity, the second-order Lovelock theory so-called Einstein-Gauss-Bonnet gravity has been intensively investigated because the renormalizable Gauss-Bonnet term, a special combination of the curvature-squared terms, appears in the low-energy limit of the heterotic string theory [3]. The history of the black-hole physics in Einstein-Gauss-Bonnet gravity

began from the well-known Boulware-Deser-Wheeler solution corresponding to the Schwarzschild-Tangherlini solution in general relativity [4]. This solution has been generalized to the topological case with a cosmological constant and the Maxwell field [5] and has occupied the central position in the research of the Gauss-Bonnet black holes. Based on this generalized Boulware-Deser-Wheeler solution, the effects of the Gauss-Bonnet term on the stability [6] and the black-hole thermodynamics [7] have been investigated. (We refer [8] for the recent review.) Intriguingly, the solutions in Einstein-Gauss-Bonnet gravity are classified into two branches, one of which admits the general relativistic limit, while the other not. As a result, the theory generically admits two distinct (A)dS vacua stemming from its quadratic nature.

Recently, asymptotically AdS black holes with a scalar hair have attracted much attention, which was first discovered numerically as a counterexample of the black-hole no-hair conjecture [9]. In particular, theories of AdS gravity coupled to a scalar field with mass at or slightly above the Breitenlohner-Freedman bound [10] are called *designer gravity* [11]. Designer gravity admits a large class of asymptotically AdS spacetimes with slower fall-off conditions than the standard ones, of which boundary conditions are defined by an essentially arbitrary function prescribing the relation between the asymptotic value of the scalar field and its normal derivative at infinity. In the context of the AdS/CFT correspondence [12], asymptotically AdS black-hole solutions in designer gravity have been applied to the study of the cosmic censorship conjecture [13] or big-bang singularities in cosmology [14].

In the present paper, we show that such remarkable slow fall-off to the AdS infinity is universal in spherically symmetric spacetimes containing *any* matter satisfying the dominant energy condition in Einstein-Gauss-Bonnet gravity with a fine-tuning of the coupling constants to give a unique AdS vacuum, where the theory becomes Chern-Simons gravity in five dimensions. We adopt the units in which only the n -dimensional gravitational constant G_n is retained.

2. Preliminaries: The action for Einstein-Gauss-Bonnet

gravity in the $n(\geq 5)$ -dimensional spacetime is given by

$$S = \int d^n x \sqrt{-g} \left[\frac{1}{2\kappa_n^2} (R - 2\Lambda + \alpha L_{\text{GB}}) \right] + S_{\text{matter}}, \quad (1)$$

$$L_{\text{GB}} := R^2 - 4R_{\mu\nu}R^{\mu\nu} + R_{\mu\nu\rho\sigma}R^{\mu\nu\rho\sigma}, \quad (2)$$

where $\kappa_n := \sqrt{8\pi G_n}$ and Λ is a cosmological constant. L_{GB} is called the Gauss-Bonnet term, which does not contribute to the field equations in four dimensions. α is the coupling constant of the Gauss-Bonnet term. In the low-energy limit of heterotic string theory, α is regarded as the inverse string tension and positive-definite [3], however we don't fix the sign of α at this stage.

The gravitational equation of the action (1) is

$$G^\mu{}_\nu + \alpha H^\mu{}_\nu + \Lambda \delta^\mu{}_\nu = \kappa_n^2 T^\mu{}_\nu, \quad (3)$$

where

$$G_{\mu\nu} := R_{\mu\nu} - \frac{1}{2}g_{\mu\nu}R, \quad (4)$$

$$H_{\mu\nu} := 2 \left[RR_{\mu\nu} - 2R_{\mu\alpha}R^\alpha{}_\nu - 2R^{\alpha\beta}R_{\mu\alpha\nu\beta} + R_\mu{}^{\alpha\beta\gamma}R_{\nu\alpha\beta\gamma} \right] - \frac{1}{2}g_{\mu\nu}L_{\text{GB}} \quad (5)$$

and $T^\mu{}_\nu$ is the energy-momentum tensor for matter fields obtained from S_{matter} in Eq. (1).

Suppose the n -dimensional spacetime $(\mathcal{M}^n, g_{\mu\nu})$ to be a warped product of an $(n-2)$ -dimensional constant curvature space (K^{n-2}, γ_{ij}) and a two-dimensional orbit spacetime (M^2, g_{ab}) under the isometries of (K^{n-2}, γ_{ij}) . Namely, the line element is given by

$$g_{\mu\nu}dx^\mu dx^\nu = g_{ab}(y)dy^a dy^b + r^2(y)\gamma_{ij}(z)dz^i dz^j, \quad (6)$$

where $a, b = 0, 1$; $i, j = 2, \dots, n-1$. Here r is a scalar on (M^2, g_{ab}) with $r = 0$ defining its boundary and γ_{ij} is the unit metric on (K^{n-2}, γ_{ij}) with its sectional curvature $k = \pm 1, 0$. We assume that (K^{n-2}, γ_{ij}) is compact. The generalized Misner-Sharp mass in Einstein-Gauss-Bonnet gravity is a scalar function on (M^2, g_{ab}) defined by

$$m := \frac{(n-2)V_{n-2}^k}{2\kappa_n^2} \left\{ -\tilde{\Lambda}r^{n-1} + r^{n-3}[k - (Dr)^2] + \tilde{\alpha}r^{n-5}[k - (Dr)^2]^2 \right\}, \quad (7)$$

where $\tilde{\alpha} := (n-3)(n-4)\alpha$, $\tilde{\Lambda} := 2\Lambda/[(n-1)(n-2)]$, D_a is a metric compatible linear connection on (M^2, g_{ab}) and $(Dr)^2 := g^{ab}(D_a r)(D_b r)$ [15, 16]. V_{n-2}^k denotes the area of K^{n-2} . For $1 + 4\tilde{\alpha}\tilde{\Lambda} = 0$, Eq. (7) gives

$$m = \frac{(n-2)V_{n-2}^k}{8\tilde{\alpha}\kappa_n^2} r^{n-5} \{ r^2 + 2\tilde{\alpha}[k - (Dr)^2] \}^2, \quad (8)$$

which is non-negative (non-positive) for $\alpha > (<)0$.

The line element may be written locally in the double-null coordinates as

$$ds^2 = -2e^{-f(u,v)} du dv + r^2(u,v)\gamma_{ij}dz^i dz^j. \quad (9)$$

The metric functions e^{-f} and r^2 are non-zero and finite to avoid the coordinate singularities. Null vectors $(\partial/\partial u)$ and $(\partial/\partial v)$ are taken to be future-pointing. The expansions of two independent future-directed radial null vectors¹ are defined as

$$\theta_+ := (n-2)r^{-1}r_{,v}, \quad (10)$$

$$\theta_- := (n-2)r^{-1}r_{,u}. \quad (11)$$

Then, the quasi-local mass m is expressed as

$$m = \frac{(n-2)V_{n-2}^k}{2\kappa_n^2} r^{n-3} \left[-\tilde{\Lambda}r^2 + \left(k + \frac{2}{(n-2)^2} r^2 e^f \theta_+ \theta_- \right) + \tilde{\alpha}r^{-2} \left(k + \frac{2}{(n-2)^2} r^2 e^f \theta_+ \theta_- \right)^2 \right]. \quad (12)$$

An $(n-2)$ -surface with $\theta_+ \theta_- > (<)0$ is called a *trapped* (*untrapped*) *surface*. We fix the orientation of the untrapped surface by $\theta_+ > 0$ and $\theta_- < 0$, i.e., $\partial/\partial u$ and $\partial/\partial v$ are ingoing and outgoing null vectors, respectively.

The most general material stress-energy tensor $T_{\mu\nu}$ in this symmetric spacetime is given by

$$T_{\mu\nu} dx^\mu dx^\nu = T_{uu}(u,v) du^2 + 2T_{uv}(u,v) du dv + T_{vv}(u,v) dv^2 + p(u,v) r^2 \gamma_{ij} dz^i dz^j. \quad (13)$$

The variation of m is determined by the field equations as

$$m_{,v} = \frac{1}{n-2} V_{n-2}^k e^f r^{n-1} (T_{uv} \theta_+ - T_{vv} \theta_-), \quad (14)$$

$$m_{,u} = \frac{1}{n-2} V_{n-2}^k e^f r^{n-1} (T_{uv} \theta_- - T_{uu} \theta_+). \quad (15)$$

In the present paper, we assume the dominant energy condition for the matter field, which implies

$$T_{uu} \geq 0, \quad T_{vv} \geq 0, \quad T_{uv} \geq 0 \quad (16)$$

and assures that a causal observer measures the non-negative energy density and the energy flux is a future-directed causal vector.

For the proof of our main results, we review the generalized Birkhoff's theorem in Einstein-Gauss-Bonnet gravity [16, 18]. In the vacuum case, Eqs. (14) and (15) give $m = M$, where M is a constant. The vacuum spacetime can be completely classified by the following theorem. (See Proposition 1 in [16] for the proof.)

Theorem 1 (*The generalized Birkhoff's theorem.*) An n -dimensional vacuum spacetime in Einstein-Gauss-Bonnet gravity with the metric form (6) is isometric to one of the followings: (i) the generalized Boulware-Deser-Wheeler solution if $(Dr)^2 \neq 0$, (ii) the Nariai-type solution if r is constant, and (iii) the class I solution if $(Dr)^2 = k + r^2/(2\tilde{\alpha})$.

¹ In [16, 17], they are erroneously mentioned as radial null geodesics.

The generalized Boulware-Deser-Wheeler solution [4, 5] is given as

$$ds^2 = -f(r)dt^2 + f^{-1}(r)dr^2 + r^2\gamma_{ij}dz^i dz^j, \quad (17)$$

$$f(r) := k + \frac{r^2}{2\tilde{\alpha}} \left[1 \mp \sqrt{1 + \frac{8\kappa_n^2 \tilde{\alpha} M}{(n-2)V_{n-2}^k r^{n-1}} + 4\tilde{\alpha}\tilde{\Lambda}} \right]. \quad (18)$$

The Nariai-type solution [19, 20] is given as

$$ds^2 = -(1 - \sigma\rho^2)dt^2 + \frac{d\rho^2}{1 - \sigma\rho^2} + r_0^2\gamma_{ij}dz^i dz^j, \quad (19)$$

$$\sigma := \left[\frac{2(n-3) + 2\tilde{\alpha}(n-5)kr_0^{-2}}{r_0^2 + 2\tilde{\alpha}k} \right] k, \quad (20)$$

where r_0^2 is the real and positive root of the following algebraic equation (see [16] for the existence condition):

$$(n-1)\tilde{\Lambda} = \frac{(n-3)k}{r_0^2} + \frac{(n-5)\tilde{\alpha}k^2}{r_0^4}. \quad (21)$$

The relation $r_0^2 + 2\tilde{\alpha}k = 0$ does not hold since it gives a contradiction. The quasi-local mass of the Nariai-type spacetime is given by

$$m = -\frac{(n-2)V_{n-2}^k r_0^{n-5}}{2\kappa_n^2} (\tilde{\Lambda}r_0^4 - kr_0^2 - k^2\tilde{\alpha}), \quad (22)$$

$$= \frac{(n-2)kV_{n-2}^k r_0^{n-5}}{(n-1)\kappa_n^2} (r_0^2 + 2k\tilde{\alpha}), \quad (23)$$

where we used Eq. (21) for eliminating Λ . Thus, m is non-zero because of $r_0^2 + 2k\tilde{\alpha} \neq 0$.

The class I solution [16, 18] exists only for $1 + 4\tilde{\alpha}\tilde{\Lambda} = 0$ as

$$ds^2 = -g(r)e^{2\delta(t,r)}dt^2 + \frac{dr^2}{g(r)} + r^2\gamma_{ij}dz^i dz^j, \quad (24)$$

$$g(r) := k + \frac{r^2}{2\tilde{\alpha}}, \quad (25)$$

where $\delta(t, r)$ is an *arbitrary* function. We note that the class I solution is not static in general and the quasi-local mass is identically zero ($m \equiv 0$) for this solution.

Next, we also review the vanishing mass theorem in the asymptotically AdS spacetime for $1 + 4\tilde{\alpha}\tilde{\Lambda} = 0$. Equation (8) shows that the quasi-local mass is non-negative (non-positive) for $\alpha > (<)0$. Then, by the combination of the asymptotic analysis and the monotonic property of m on untrapped surfaces under the dominant energy condition, the following theorem is shown for positive α . (See Proposition 7 in [16] for the proof.)

Theorem 2 (*Vanishing mass in asymptotically AdS spacetime with $1 + 4\tilde{\alpha}\tilde{\Lambda} = 0$.)* Suppose $1 + 4\tilde{\alpha}\tilde{\Lambda} = 0$ with $\alpha > 0$ and the dominant energy condition in an n -dimensional asymptotically AdS spacetime. Then, $m \equiv 0$ holds on the untrapped spacelike hypersurface.

In the above theorem, we employed the higher-dimensional generalization of the Henneaux-Teitelboim asymptotically AdS boundary conditions for the metric components adopting the coordinates $x^\mu = \{t, \rho, z^i\}$ [21]. The metric under consideration can be written as $g_{\mu\nu} = g_{\mu\nu}^{(0)} + h_{\mu\nu}$, where $g_{\mu\nu}^{(0)}$ is the metric of the AdS spacetime, from which deviation is represented by $h_{\mu\nu}$. In the global coordinates, we have

$$g_{\mu\nu}^{(0)} dx^\mu dx^\nu = -(1 + \ell_{\text{eff}}^{-2} \rho^2) dt^2 + \frac{d\rho^2}{(1 + \ell_{\text{eff}}^{-2} \rho^2)} + \rho^2 d\Omega_{n-2}^2, \quad (26)$$

$$\ell_{\text{eff}}^2 := -\frac{1}{2\tilde{\Lambda}} \left(1 \pm \sqrt{1 + 4\tilde{\alpha}\tilde{\Lambda}} \right), \quad (27)$$

where $d\Omega_{n-2}^2$ is the line element of a unit $(n-2)$ -sphere. ℓ_{eff} is the effective curvature radius in the spacetime, so we assume $1 + 4\tilde{\alpha}\tilde{\Lambda} \geq 0$ hereafter for ℓ_{eff}^2 to be real. The AdS spacetime (26) corresponds to the generalized Boulware-Deser-Wheeler solution (17) with $k = 1$ and $M = 0$. The fall-off condition is such that

$$h_{tt} = c_{tt}\rho^{-n+3} + O(\rho^{-n+2}), \quad (28a)$$

$$h_{\rho\rho} = c_{\rho\rho}\rho^{-n-1} + O(\rho^{-n-2}), \quad (28b)$$

$$h_{t\rho} = c_{t\rho}\rho^{-n} + O(\rho^{-n-1}), \quad (28c)$$

$$h_{\rho i} = c_{\rho i}\rho^{-n} + O(\rho^{-n-1}), \quad (28d)$$

$$h_{ti} = c_{ti}\rho^{-n+3} + O(\rho^{-n+2}), \quad (28e)$$

$$h_{ij} = c_{ij}\rho^{-n+3} + O(\rho^{-n+2}), \quad (28f)$$

where c_{tt}, \dots, c_{ij} are functions independent of ρ . In the spherically symmetric spacetime, which is of our interest here, c_{tt}, \dots, c_{ij} are independent of z^i and $c_{\rho i} = c_{ti} \equiv 0$.

3. Main results: Now we show our main results. As seen before, if the spacetime is maximally symmetric or class I spacetime, the quasi-local mass m is identically zero. Indeed, for $k = 1$, $\Lambda \leq 0$, and $\alpha \geq 0$, its inverse also holds under the dominant energy condition.

Proposition 1 (*Vanishing mass spacetime.*) Under the dominant energy condition for $k = 1$, $\Lambda \leq 0$, and $\alpha \geq 0$, $m \equiv 0$ is equivalent to the maximally symmetric spacetime for $1 + 4\tilde{\alpha}\tilde{\Lambda} \neq 0$ and the class I spacetime (24) for $1 + 4\tilde{\alpha}\tilde{\Lambda} = 0$.

As seen in Eq. (27), the special tuning between the coupling constants $1 + 4\tilde{\alpha}\tilde{\Lambda} = 0$ allows the theory to have a unique (A)dS vacuum and become Chern-Simons gravity in five dimensions [22]. Proposition 1 is proven by the combination of the following two lemmas together with Theorem 1.

Lemma 1 If $m \equiv 0$ for $k = 1$, $\Lambda \leq 0$, and $\alpha \geq 0$, then $\theta_+ \theta_- < 0$, i.e., the spacetime consists of the untrapped surfaces.

Proof. Trivial from Eq. (12). ■

Lemma 2 Under the dominant energy condition, if $m = 0$ on the untrapped surface, then $T_{\mu\nu} = 0$.

Proof. By the variation formulae (14) and (15), $T_{uu} = T_{vv} = T_{uv} = 0$ on the untrapped surface. Then, the energy-momentum conservation equation $T^{A\nu}{}_{;\nu} = 0$, where $A = u$ or v , gives

$$pr_{,A}e^f = 0. \quad (29)$$

We have $r_{,A} \neq 0$ on the untrapped surface, so that $p = 0$ there, which completes the proof. ■

Then, by the combination of Proposition 1 and Theorem 2, it is easy to show the following proposition about the asymptotically AdS spacetime for $1 + 4\tilde{\alpha}\tilde{\Lambda} = 0$ with $\alpha > 0$.

Proposition 2 (*Asymptotically AdS spacetime with $1 + 4\tilde{\alpha}\tilde{\Lambda} = 0$.)* Suppose $1 + 4\tilde{\alpha}\tilde{\Lambda} = 0$ with $\alpha > 0$ and the dominant energy condition in an n -dimensional asymptotically AdS spacetime. Then, the spacetime is represented by the class I solution (24) with $k = 1$ and $\delta(t, r)$ satisfying the fall-off condition (28).

4. *Discussions:* Properties of the generalized Misner-Sharp mass (7) have been fully investigated in [16]. It inherits the characteristics such as monotonicity or positivity from the Misner-Sharp mass in general relativity and is its natural counterpart in Einstein-Gauss-Bonnet gravity. As an application, this quasi-local mass played an essential role to reveal the dynamical properties of the Gauss-Bonnet black holes [17]. Proposition 1 obtained in the present paper is another remarkable property in the spherically symmetric case in addition to the results in [16], which claims the equivalence between the vanishing quasi-local mass and the maximally symmetric spacetime for $\alpha \geq 0$ and $\Lambda \leq 0$ with $1 + 4\tilde{\alpha}\tilde{\Lambda} \neq 0$.

The case with $1 + 4\tilde{\alpha}\tilde{\Lambda} = 0$ is exceptional in Proposition 1, which admits the theory to have a unique (A)dS vacuum as well as the non-maximally symmetric vacuum solution with vanishing quasi-local mass. Proposition 2 concerns with this exceptional case and claims that, in the presence of physically reasonable matter satisfying the dominant energy condition, even if the metric of some spherically symmetric solution of the Einstein-Gauss-Bonnet equations with $1 + 4\tilde{\alpha}\tilde{\Lambda} = 0$ and $\alpha > 0$ reduces to the AdS metric at infinity, the fall-off rate is necessarily slower than the condition (28). The generalized Boulware-Deser-Wheeler solution (17) for $n \geq 6$ with positive M is a vacuum example with such slow fall-off. It is seen that the fall-off rate to the AdS metric changes in the case of $1 + 4\tilde{\alpha}\tilde{\Lambda} = 0$. This phenomenon was first pointed out in the study of the static black holes with and without the Maxwell field in the class of Lovelock gravity admitting a unique (A)dS vacuum [22]. (The case with $k = 2$ in [22] corresponds to ours.) Our proposition claims that it is universal under the dominant energy condition even in the highly dynamical situation.

In the n -dimensional Kerr-Myers-Perry-AdS spacetime, the fall-off condition (28) is certainly satisfied. Although its counterpart in Einstein-Gauss-Bonnet gravity has not been found yet, it would also exhibit the slow fall-off to the unique AdS infinity for $1 + 4\tilde{\alpha}\tilde{\Lambda} = 0$. Under the standard fall-off condition (28), several definitions of the global mass in the asymptotically

AdS spacetime have been given in Einstein-Gauss-Bonnet gravity [23]. However, the universal slow fall-off means that they are diverging in the case of $1 + 4\tilde{\alpha}\tilde{\Lambda} = 0$ with $\alpha > 0$. This fact forces us to reformulate the global mass to give a finite value under the slower fall-off condition in this special case. This problem has been investigated in Chern-Simons gravity in [24].

Since the slow fall-off to the unique AdS vacuum has been confirmed in the vacuum case and in the presence of the Maxwell field [22], it is naturally expected to be a universal property under the dominant energy condition also in the class of Lovelock gravity admitting a single AdS vacuum. Our main results have been obtained essentially by using the mass variation formulae (14) and (15) and the energy-momentum conservation equation. These variation formulae are exactly the same as those in general relativity, which enable us to prove the propositions in parallel with the general relativistic case. In our recent paper [16], a further generalization of the Misner-Sharp quasi-local mass in general Lovelock gravity was proposed, with which the mass variation formulae were conjectured to hold. We expect that a large set of the results obtained in the present paper and in [16] is generalized in a very straightforward manner. They will provide us a firm ground in the research of Lovelock gravity.

The author thanks Cristián Martínez, Masato Nozawa, Ricardo Troncoso, and Jorge Zanelli for discussions and comments. The author was supported by Fondecyt grant 1071125. The Centro de Estudios Científicos (CECS) is funded by the Chilean Government through the Millennium Science Initiative and the Centers of Excellence Base Financing Program of Conicyt. CECS is also supported by a group of private companies which at present includes Antofagasta Minerals, Arauco, Empresas CMPC, Indura, Naviera Ultragas, and Telefónica del Sur.

* Electronic address: hideki@cecs.cl

- [1] D. Lovelock, *J. Math. Phys.* **12**, 498 (1971).
- [2] J. Zanelli, arXiv:hep-th/0502193.
- [3] B. Zwiebach, *Phys. Lett.* **B156**, 315 (1985); D.J. Gross and E. Witten, *Nucl. Phys.* **B277**, 1 (1986); D.J. Gross and J.H. Sloan, *Nucl. Phys.* **B291**, 41 (1987); R.R. Metsaev and A.A. Tseytlin, *Phys. Lett.* **B191**, 354 (1987); R.R. Metsaev and A.A. Tseytlin, *Nucl. Phys.* **B293**, 385 (1987).
- [4] D.G. Boulware and S. Deser, *Phys. Rev. Lett.* **55**, 2656 (1985); J.T. Wheeler, *Nucl. Phys.* **B268**, 737 (1986).
- [5] J.T. Wheeler, *Nucl. Phys. B* **273**, 732 (1986); D.L. Wiltshire, *Phys. Lett.* **B169**, 36 (1986); R.G. Cai, *Phys. Rev. D* **65**, 084014 (2002); M. Cvetič, S. Nojiri, and S.D. Odintsov, *Nucl. Phys.* **B628**, 295 (2002).
- [6] S. Deser and Z. Yang, *Classical Quantum Gravity* **6**, L83 (1989); G. Dotti and R.J. Gleiser, *Classical Quantum Gravity* **22**, L1 (2005); R. Konoplya, *Phys. Rev. D* **71**, 024038 (2005); G. Dotti and R.J. Gleiser, *Phys. Rev. D* **72**, 044018 (2005); R.J. Gleiser and G. Dotti, *Phys. Rev. D* **72**, 124002 (2005); S.K. Chakrabarti and K.S. Gupta, *Int. J. Mod. Phys. A* **21**, 3565 (2006); S.K. Chakrabarti, *Gen. Relativ. Gravit.* **39**, 567 (2007); R.G. Daghighi, G. Kunstatter, and J. Ziprick, *Classical Quantum Gravity* **24**, 1981 (2007); M. Beroiz, G. Dotti, and R.J. Gleiser,

- Phys. Rev. D **76**, 024012 (2007); J. Liu, Commun. Theor. Phys. **47**, 647 (2007); R.A. Konoplya and A. Zhidenko, Phys. Rev. D **77**, 104004 (2008); T. Hirayama, arXiv:0804.3694 [gr-qc].
- [7] R.C. Myers and J.Z. Simon, Phys. Rev. D **38**, 2434 (1988); S. Nojiri, S.D. Odintsov, and S. Ogushi, Phys. Rev. D **65**, 023521 (2001); M. Cvetič, S. Nojiri, and S.D. Odintsov, Nucl. Phys. **B628**, 295 (2002); Y.M. Cho and I.P. Neupane, Phys. Rev. D **66**, 024044 (2002); I.P. Neupane, Phys. Rev. D **67**, 061501(R) (2003); I.P. Neupane, Phys. Rev. D **69**, 084011 (2004); R.-G. Cai and Q. Guo, Phys. Rev. D **69**, 104025 (2004); T. Clunan, S.F. Ross, and D.J. Smith, Classical Quantum Gravity **21**, 3447 (2004); T. Torii and H. Maeda, Phys. Rev. D **71**, 124002 (2005); J. Grain, A. Barrau, and P. Kanti, Phys. Rev. D **72**, 104016 (2005); G. Kofinas and R. Olea, Phys. Rev. D **74**, 084035 (2006).
- [8] C. Charmousis, arXiv:0805.0568 [gr-qc]; C. Garraffo and G. Giribet, arXiv:0805.3575 [gr-qc].
- [9] T. Torii, K. Maeda, and M. Narita, Phys. Rev. D **64**, 044007 (2001).
- [10] P. Breitenlohner and D.Z. Freedman, Annals Phys. **144**, 249 (1982).
- [11] M. Henneaux, C. Martínez, R. Troncoso, and J. Zanelli, Phys. Rev. D **65**, 104007 (2002); M. Henneaux, C. Martínez, R. Troncoso, and J. Zanelli, Phys. Rev. D **70**, 044034 (2004); T. Hertog and K. Maeda, JHEP **0407**, 051 (2004); T. Hertog and K. Maeda, Phys. Rev. D **71**, 024001 (2005); T. Hertog and G.T. Horowitz, Phys. Rev. Lett. **94**, 221301 (2005); T. Hertog and S. Hollands, Classical Quantum Gravity **22**, 5323 (2005); M. Henneaux, C. Martínez, R. Troncoso, and J. Zanelli, Annals Phys. **322**, 824 (2007); A.J. Amsel and D. Marolf, Phys. Rev. D **74**, 064006 (2006) [Erratum-ibid. D **75**, 029901 (2007)]; D. Marolf and S.F. Ross, JHEP **0611**, 085 (2006); S. de Haro and A.C. Petkou, JHEP **0612**, 076 (2006); T. Hertog, Classical Quantum Gravity **24**, 141 (2007); T. Hertog, Phys. Rev. D **74**, 084008 (2006); S. de Haro, I. Papadimitriou, and A.C. Petkou, Phys. Rev. Lett. **98**, 231601 (2007); A.J. Amsel, T. Hertog, S. Hollands, and D. Marolf, Phys. Rev. D **75**, 084008 (2007) [Erratum-ibid. D **77**, 049903 (2008)]; D. Marolf and S.F. Ross, JHEP **0805**, 055 (2008).
- [12] J.M. Maldacena, Adv. Theor. Math. Phys. **2**, 231 (1998) [Int. J. Theor. Phys. **38**, 1113 (1999)].
- [13] T. Hertog, G.T. Horowitz, and K. Maeda, Phys. Rev. Lett. **92**, 131101 (2004); T. Hertog, G.T. Horowitz, and K. Maeda, Phys. Rev. D **69**, 105001 (2004); M. Alcubierre, J.A. Gonzalez, M. Salgado, and D. Sudarsky, M. Gutperle and P. Kraus, JHEP **0404**, 024 (2004); M. Dafermos, Adv. Theor. Math. Phys. **9**, 575 (2005); D. Garfinkle, Phys. Rev. D **69**, 124017 (2004); V.E. Hubeny, X. Liu, M. Rangamani, and S. Shenker, JHEP **0412**, 067 (2004); T. Hertog, G.T. Horowitz, and K. Maeda, arXiv:gr-qc/0405050; D. Garfinkle, Phys. Rev. D **70**, 104015 (2004); A.V. Frolov, Phys. Rev. D **70**, 104023 (2004).
- [14] T. Hertog and G.T. Horowitz, JHEP **0407**, 073 (2004); T. Hertog and G.T. Horowitz, JHEP **0504**, 005 (2005); N. Turok, B. Craps, and T. Hertog, arXiv:0711.1824 [hep-th]; B. Craps, T. Hertog, and N. Turok, arXiv:0712.4180 [hep-th].
- [15] H. Maeda, Phys. Rev. D **73**, 104004 (2006).
- [16] H. Maeda and M. Nozawa, Phys. Rev. D **77**, 064031 (2008).
- [17] M. Nozawa and H. Maeda, Classical Quantum Gravity **25**, 055009 (2008).
- [18] C. Charmousis, J-F. Dufaux, Classical Quantum Gravity **19**, 4671 (2002); S. Deser and B. Tekin, Classical Quantum Gravity **20**, 4877 (2003); R. Zegers, J. Math. Phys. **46**, 072502 (2005); S. Deser and J. Franklin, Classical Quantum Gravity **22**, L103 (2005).
- [19] D. Lorenz-Petzold, Prog. Theor. Phys. **78**, 969 (1987).
- [20] H. Maeda and N. Dadhich, Phys. Rev. D **75**, 044007 (2007).
- [21] M. Henneaux and C. Teitelboim, Commun. Math. Phys. **98**, 391 (1985); S. Hollands, A. Ishibashi, and D. Marolf, Classical Quantum Gravity **22**, 2881 (2005).
- [22] J. Crisostomo, R. Troncoso, and J. Zanelli, Phys. Rev. D **62**, 084013 (2000).
- [23] S. Deser and B. Tekin, Phys. Rev. Lett. **89**, 101101 (2002); S. Deser and B. Tekin, Phys. Rev. D **67**, 084009 (2003); A. Padilla, Classical Quantum Gravity **20**, 3129 (2003).
- [24] P. Mora, R. Olea, R. Troncoso, and J. Zanelli, JHEP **0406**, 036 (2004); P. Mora, R. Olea, R. Troncoso, and J. Zanelli, JHEP **0602**, 067 (2006).