

# On the classical equivalence of superstring field theories

---

**Ehud Fuchs, Michael Kroyter**

*Max-Planck-Institut für Gravitationsphysik  
Albert-Einstein-Institut  
14476 Golm, Germany  
udif@aei.mpg.de, mikroyt@aei.mpg.de*

**ABSTRACT:** We construct mappings that send solutions of the cubic and non-polynomial open superstring field theories to each other. We prove that the action is invariant under the maps and that gauge orbits are mapped into gauge orbits. It follows that the perturbative spectrum around solutions is the same in both theories. The mappings also preserve the string field reality condition.

We generalize to the cases of a non-BPS D-brane and of multi-D-brane systems. We analyze the recently found analytic solutions of the cubic action, both in the BPS sector and the non-BPS sector and show that they span a one parameter family of solutions with empty cohomology and identical action, which suggests that they are gauge equivalent. We write the gauge transformations relating these solutions explicitly. This seems to suggest that open superstring field theory is able to describe a vacuum solution even around a BPS D-brane.

**KEYWORDS:** String Field Theory.

---

## Contents

<b>1. Introduction</b>	<b>1</b>
<b>2. Mapping the cubic theory to the non-polynomial one</b>	<b>4</b>
2.1 The map	5
2.2 Equations of motion	6
2.3 Gauge equivalence of solutions and cohomology	7
2.4 The action	8
<b>3. Formal gauge form of solutions in the cubic theories</b>	<b>10</b>
3.1 Gauge form of the solutions	10
3.2 Formal gauge representation for the bosonic solutions	12
<b>4. Incorporating the GSO(−) sector</b>	<b>13</b>
<b>5. Vacuum solutions</b>	<b>13</b>
5.1 On the BPS D-brane	14
5.2 On the Non-BPS D-brane	15
<b>6. Conclusions</b>	<b>18</b>

---

## 1. Introduction

Covariant open superstring field theory was first constructed by Witten [1], following his construction of the bosonic theory [2]. The formal structure of the theories is very similar. They both rely on the algebraic structures obeyed by the star-product, the BRST operator  $Q$  and the integration over the space of string fields. There are two new features that arise in the case of the superstring. One is the existence of several sectors, namely GSO( $\pm$ ) and NS/R, that should be separately handled. The other is the notion of picture number [3].

Picture number is the expression of the existence of zero modes for the superghosts of the RNS formalism. Just as the zero modes of the fermionic  $bc$  ghost system imply a two-fold degeneracy in the description of vertex operators, namely integrated/unintegrated vertex, the zero modes of the bosonic  $\beta\gamma$  system imply the existence of an infinite number of equivalent vertices, distinguished by a new quantum number, called picture number. Picture number is most easily dealt with using fermionization [3], where the  $\beta, \gamma$  system is expressed in terms of the  $\eta, \xi, \phi$  variables,

$$\beta = e^{-\phi} \partial \xi, \quad \gamma = \eta e^{\phi}. \quad (1.1)$$

The operation of integration over the space of string fields is realized using a CFT expectation value, which is zero for fields of zero picture number. This implies that one has either to assign non-zero picture number to the string field or to append it to the operations of star-product/derivation/integration.

Witten chose to define the string field in the natural  $(-1)$  picture and to append a picture changing operator to the cubic term in the action. This picture changing operator had to be inserted at the interaction (mid-string) point  $z = i$  in order to preserve the associativity of the star product. Soon thereafter it was noticed that in calculations of scattering amplitudes and in iteration of gauge transformations picture changing operators formally collide [4, 5]. This is an immediate consequence of inserting the operator at  $z = i$ , which is invariant under the star-product. The collision of picture changing operators produces singularities, since the OPE of the picture changing operator with itself is divergent. It was shown that the theory can be regularized to leading order by adding counter terms (already at the classical level), but it was also shown that the problems persist at higher order and it was not understood if and how it could be systematically solved.

Resolutions of these problems can be achieved by leaving the string fields at ghost number one, while modifying their picture number to zero [6, 7, 8]. The modified action (in the NS+ sector) reads,

$$S = -\frac{1}{g_o^2} \int Y_{-2} \left( \frac{1}{2} \Psi Q \Psi + \frac{1}{2} \Psi^3 \right), \quad (1.2)$$

where,  $g_o$  is the open string coupling constant,  $Y_{-2}$  is the double-step inverse picture changing operator and the string fields are multiplied in the action using Witten's star product, which we leave implicit throughout the paper.

There are two possible definitions for  $Y_{-2}$ , namely the chiral the non-chiral operators. The chiral is a local operator, whose exact form will not be of interest for us, while the non-chiral is a non-local operator, defined using the doubling trick as

$$Y_{-2}(z) = Y(z)Y(\bar{z}). \quad (1.3)$$

This definition cannot be used on the real line, where picture changing operators and vertex operators are usually inserted, due to the singular OPE of  $Y$  with itself. However, it makes perfect sense when inserted in the string mid-point,  $z = i$ . In this formalism there are no problems of collisions, since while the interaction vertex comes with a  $Y_{-2}$  factor, the propagator comes with a factor of its inverse, which is well defined. These operators are canceled in pairs and scattering amplitudes can be calculated and give (at least on-shell and for tree diagrams) the expected results [9].

The two different choices define two a-priori inequivalent theories. The theory that uses the chiral operator was shown to have some strange features in level truncation [10]. Also, being chiral it does not respect the twist symmetry that one expects it to obey. For these reasons the non-chiral theory was used in most studies of cubic superstring field theory and is considered the more reliable one. This is the version of the theory we shall be using here. The non-chiral  $Y_{-2}$  was recently used also in the construction of boundary superstring field theory [11].

From the action (1.2) one derives the equation of motion

$$Y_{-2}(i)(Q\Psi + \Psi^2) = 0. \quad (1.4)$$

The operator  $Y_{-2}(i)$  has a non-trivial kernel. However, this kernel consists of states of a somewhat singular nature that are localized at  $z = \pm i$ . It is not clear if it has a non-trivial intersection with the correct space of open string fields, especially since it is not clear what should this space be. Assuming that this kernel poses no problem, the equation of motion gets the desired form

$$Q\Psi + \Psi^2 = 0. \quad (1.5)$$

The action is invariant under the gauge transformation

$$\Psi \rightarrow e^{-\Lambda}(\Psi + Q)e^{\Lambda}. \quad (1.6)$$

Another resolution of the problems, not relying on picture changing operators, was obtained by Berkovits [12], who developed another form of superstring field theory. This theory is non-polynomial, resembling a WZW-theory, although (unlike closed/heterotic string field theory [13, 14, 15]) string fields are still multiplied using only Witten's star-product. The non-polynomial theory uses ghost number zero, picture number zero string fields that reside in the large Hilbert space. The large Hilbert space consists of two copies of the small (standard) Hilbert space, the usual one and another one, which is multiplied by  $\xi_0$ . This redundancy is taken care of by a novel gauge symmetry, which implies that only the copy proportional to  $\xi_0$  is physical. The action reads,

$$S = \frac{1}{2g_0^2} \oint \left( e^{-\Phi} Q(e^{\Phi}) e^{-\Phi} \eta_0(e^{\Phi}) - \int_0^1 dt e^{-t\Phi} \partial_t e^{t\Phi} [e^{-t\Phi} \eta_0 e^{t\Phi}, e^{-t\Phi} Q e^{t\Phi}] \right), \quad (1.7)$$

and the gauge symmetry is

$$e^{\Phi} \rightarrow e^{-Q\Lambda_Q} e^{\Phi} e^{\eta_0\Lambda_\eta}. \quad (1.8)$$

In (1.7) and elsewhere we use the square brackets to denote the graded commutator. That is,  $[A, B]$  is an anti-commutator if  $A$  and  $B$  are odd (as it is here) and a commutator otherwise. The integral over string fields corresponds to a CFT expectation value. For the non-polynomial theory this should be calculated in the large Hilbert space. To distinguish this case we define

$$\oint A \equiv \langle A \rangle_{Large}, \quad \int A \equiv \langle A \rangle_{Small}. \quad (1.9)$$

The integrals are related in a simple way

$$\oint \xi_0 A = \int A. \quad (1.10)$$

From the action (1.7) the following equation of motion is derived,

$$\eta_0(G^{-1}QG) = 0, \quad (1.11)$$

where we define

$$G = e^{\Phi}. \quad (1.12)$$

Equation (1.11) states that  $G^{-1}QG$  lives in the small Hilbert space, where the string fields of the cubic theory reside. It also has the correct quantum numbers to be identified with  $\Psi$ . This expression takes exactly the form of a gauge solution of the cubic theory (1.6). It is not a genuine gauge solution, since the “gauge field”  $G$  lives in the large Hilbert space. It is nevertheless a solution of the cubic theory. Thus, there exists a simple map of solutions of the non-polynomial theory to solutions of the cubic theory [16, 17],

$$\Psi = G^{-1}QG. \tag{1.13}$$

The purpose of this paper is to show that (at least classically and in the NS sector) the cubic superstring field theory and Berkovits’ non-polynomial superstring field theory are equivalent. We prove this in section 2 by adding to the map (1.13) also a map that works in the other direction. This map also sends solutions to solutions and leaves the action invariant. We further show that both maps respect the gauge symmetries, so we obtain a one to one mapping of gauge orbits of solutions in both directions. In section 3 we show how to write the solutions of the cubic theory as formal gauge solutions. In this form our map takes the form of a mapping of gauge fields. We also comment on bosonic string field theory. In section 4 we describe how the GSO(−) sector is added to the theory and to our mapping. This enables us to generalize the mapping to non-BPS D-branes and to multi D-brane systems.

In section 5 we describe analytic solutions in both theories on BPS and non-BPS D-branes that seem to describe the vacuum without the original D-brane. These solutions are based on the recent solutions of [18, 19, 20]. While such solutions are to be expected for the non-BPS D-brane, where a tachyon is present, it may seem strange to have them on the BPS D-brane, where a physical tachyon is absent. Moreover, a BPS D-brane carries a RR charge, which is absent in the vacuum without the D-brane. Thus, such solutions describe a change in closed string moduli, which is usually not expected to be possible to describe within open string field theory. We claim, that it seems to be the case that such solutions are nevertheless realized. We comment on this issue and offer some other conclusive remarks in section 6.

## 2. Mapping the cubic theory to the non-polynomial one

In this section we construct a map of string fields from the cubic superstring field theory to string fields in the non-polynomial theory. We construct the map in 2.1 and show that it sends solutions to solution in 2.2. We compose this map and the map (1.13), which goes in the opposite direction and is defined only for solutions. The composition in one direction gives the identity transformation on solutions of the cubic theory and in the other direction it gives a gauge transformation on the space of solutions of the non-polynomial theory. Further, we prove in 2.3 that both maps send gauge equivalent solutions to gauge equivalent solutions and that all gauge orbits are accessible. Thus, the two maps define a one to one correspondence of gauge orbits of solutions. A direct consequence of this fact is that the cohomologies around solutions in both theories agree. Finally, in 2.4 we study the action of solutions and prove that it has the same value in both theories.

## 2.1 The map

A map of  $\Psi$  to  $\Phi$  should contain multiplication by a  $\xi_0 = \xi(0)$  factor, which reduces the ghost number by one, while increasing the picture number by one. In order to get the correct quantum numbers, we need also an operator that decreases the picture number by one, without modifying the other quantum numbers, such as ghost number and conformal weight. Such an operator exists, namely, the inverse picture changing operator,

$$Y(z) = (c\partial\xi e^{-2\phi})(z). \quad (2.1)$$

Combining the two operators we get the operator

$$P(z) = (\xi Y)(z) = -(c\xi\partial\xi e^{-2\phi})(z), \quad (2.2)$$

that can be used to define the desired mapping. So far we did not fix  $z$ . In fact, we can define a map

$$\Phi = P\Psi, \quad (2.3)$$

using a linear combination

$$P = \sum_i k_i P(z_i). \quad (2.4)$$

The operator  $P$  contains  $\xi_0$  as long as  $\sum k_i \neq 0$ . It is also possible to define  $P$  using different insertion points for  $Y(z)$  and  $\xi(z)$ . This will become useful in 2.4.

An important property of  $P$  is that it is nilpotent,

$$P^2 = 0. \quad (2.5)$$

This follows from the OPE

$$P(z)P(w) = -\frac{z-w}{12}(cc'\xi\xi'\xi''\xi'''\xi''''e^{-4\phi})(w) + \mathcal{O}((z-w)^2). \quad (2.6)$$

This is in contrast to the OPE of the inverse picture changing operator  $Y$  with itself,

$$Y(z)Y(w) = -\frac{(cc'\xi'\xi''e^{-4\phi})(w)}{(z-w)^2} + \mathcal{O}\left(\frac{1}{z-w}\right). \quad (2.7)$$

The picture changing operator

$$X(z) = [Q, \xi(z)] \quad (2.8)$$

suffers from a similar singularity. The appearance of these singularities in Witten's theory were the source of a need for its modification.

We demand that the  $P$  operator be nilpotent also under the star-product

$$(P\Phi_1)(P\Phi_2) = 0. \quad (2.9)$$

We achieve this by inserting  $P(z)$  at the string mid-point

$$P = k_1 P(i) + k_2 P(-i), \quad (2.10)$$

which is invariant under the star-product.

The operator  $P(z)$  is the inverse of the BRST charge  $Q$  in the sense that

$$[Q, P(z)] = 1, \quad \forall z. \quad (2.11)$$

We can define a string field

$$A = P(z) |1\rangle, \quad (2.12)$$

where  $|1\rangle$  is the identity string field. This state obeys,

$$QA = |1\rangle. \quad (2.13)$$

From the above it follows that the cohomology of  $Q$  is trivial [21, 22] (in the large Hilbert space). The same also holds for  $\eta_0$  that obeys

$$[\eta_0, \xi(z)] = 1, \quad \forall z. \quad (2.14)$$

The property (2.11) is crucial for our construction. Thus, we shall demand that the coefficients in (2.10) sum to unity. Specifically, we choose to work with the combination

$$P = \frac{e^{i\frac{\pi}{4}}P(i) + e^{-i\frac{\pi}{4}}P(-i)}{\sqrt{2}}, \quad (2.15)$$

since it would prove useful in the calculation of the action. Similarly we define

$$Y = \frac{e^{i\frac{\pi}{4}}Y(i) + e^{-i\frac{\pi}{4}}Y(-i)}{\sqrt{2}}. \quad (2.16)$$

## 2.2 Equations of motion

Suppose that  $\Phi$  is given by (2.3), (2.10). The nilpotency of  $P$  under the star-product (2.9) implies,

$$G = 1 + P\Psi, \quad G^{-1} = 1 - P\Psi. \quad (2.17)$$

Let us map  $G$  back to  $\Psi'$  using (1.13). For  $G$  of the form (2.17), we find,

$$\Psi' = G^{-1}QG = (1 - P\Psi)(\Psi - PQ\Psi) = (1 - P\Psi)(\Psi + P\Psi\Psi) = \Psi. \quad (2.18)$$

Here, we used (2.11), then we used the equation of motion for  $\Psi$  (1.5) and the nilpotency of  $P$  under the star-product (2.9). We conclude that the transformation (1.13) is the inverse of (2.3) in this direction. This is not true when the transformations are composed in the other order.

Proving that  $G$  defined by our map is a solution is now straightforward, as

$$\eta_0(G^{-1}QG) = \eta_0\Psi = 0, \quad (2.19)$$

since  $\Psi$  resides in the small Hilbert space and so is annihilated by  $\eta_0$ . We conclude that given our map, the equation of motion for  $\Phi$  follows from that of  $\Psi$ .

### 2.3 Gauge equivalence of solutions and cohomology

Not all solutions of the non-polynomial theory can be written in the form (2.3). Here, we prove that all of them are gauge equivalent to ones that can be so written. Then, we prove that the two maps (2.3) and (1.13) map gauge equivalent solutions to gauge equivalent solutions. Thus, the maps define a one to one and onto mapping of gauge orbits of solutions of both theories.

Parametrize an arbitrary solution  $G$  as

$$G = \frac{1}{1 - \Phi}. \quad (2.20)$$

In this scheme (1.13) takes the form,

$$\Psi = G^{-1}QG = -QG^{-1}G = Q\Phi \frac{1}{1 - \Phi}. \quad (2.21)$$

Map it now back to the non-polynomial theory using (2.3),

$$\tilde{G} = 1 + PQ\Phi \frac{1}{1 - \Phi} = 1 + ((1 - QP)\Phi) \frac{1}{1 - \Phi} = (1 - Q(P\Phi))G. \quad (2.22)$$

We conclude that the state to which  $\Psi$  is mapped is gauge equivalent to the original  $G$ . Important immediate corollaries are the existence of a solution of the form (2.3) in all gauge orbits, the gauge equivalence of any two solutions that produce the same  $\Psi$  via (1.13) and the fact that choosing different coefficients in the definition (2.10) results in gauge equivalent solutions, as long as the coefficients sum up to unity.

It may seem strange that the gauge transformation relating two solutions, which give rise to the same solution in the cubic theory, is the one based on  $Q$ , present also in the cubic theory. We should remember, though, that in this equation,  $Q$  acts on string fields residing in the large Hilbert space, which includes more degrees of freedom. This extra amount of degrees of freedom is fixed by the other,  $\eta_0$ -based gauge transformation, which we did not fix.

Let there be two gauge equivalent solutions of the non-polynomial theory,

$$\tilde{G} = e^{-Q\Lambda_Q} G e^{\eta_0\Lambda_\eta}. \quad (2.23)$$

It follows that

$$\begin{aligned} \tilde{G}^{-1}Q\tilde{G} &= e^{-\eta_0\Lambda_\eta} G^{-1} e^{Q\Lambda_Q} Q (e^{-Q\Lambda_Q} G e^{\eta_0\Lambda_\eta}) \\ &= e^{-\eta_0\Lambda_\eta} G^{-1} (QG e^{\eta_0\Lambda_\eta} + GQ e^{\eta_0\Lambda_\eta}) = e^{-\Lambda} (\Psi + Q) e^\Lambda, \end{aligned} \quad (2.24)$$

where we defined

$$\Lambda = \eta_0\Lambda_\eta. \quad (2.25)$$

We see that gauge equivalent states are mapped under (1.13) to gauge equivalent states.

Now, let there be two gauge equivalent string fields of the cubic theory,

$$\tilde{\Psi} = e^{-\Lambda} (\Psi + Q) e^\Lambda. \quad (2.26)$$



We get,

$$\begin{aligned} 1 + P\tilde{\Psi} &= 1 + P(-Qe^{-\Lambda} + e^{-\Lambda}\Psi)e^\Lambda = (Q(Pe^{-\Lambda}) + Pe^{-\Lambda}\Psi)e^\Lambda = Q(Pe^{-\Lambda})(1 + P\Psi)e^\Lambda \\ &= Q(Pe^{-Q\Lambda})(1 + P\Psi)e^\Lambda = e^{-Q\Lambda}Q(1 + P\Psi)e^{\Lambda_Q}, \end{aligned} \quad (2.27)$$

where we used (2.11), (2.14) and defined

$$\Lambda_Q = P\Lambda, \quad \Lambda_\eta = \xi\Lambda, \quad (2.28)$$

and the location of the  $\xi$  insertion is arbitrary. The result (2.27) takes the form of a gauge transformation (1.8) of  $(1 + P\Psi)$ . We conclude that in this direction gauge equivalent states are mapped to gauge equivalent states even off-shell.

Let us now consider linearizing the theories around two equivalent solutions. The linearized equation of motion of both theories coincide under the maps, since the exact equation coincides and similarly for the linearized gauge transformations. Solutions of the linearized equation of motion modulo linearized gauge transformations define the cohomology around the solution. We conclude that the linearized map around a solution can be interpreted as a map between the cohomology spaces of the solutions, which is one to one and onto. This suggests that solutions are mapped to solutions with the same physical content.

## 2.4 The action

Another physical property of a solution is its action. The action of the non-polynomial theory (1.7) can be put in the form,

$$S = \frac{1}{2g_o^2} \oint \left( -Qe^\Phi \eta_0 e^{-\Phi} + \int_0^1 dt \Phi[\eta_0 e^{-t\Phi}, Qe^{t\Phi}] \right), \quad (2.29)$$

and we can explicitly write down,

$$e^{t\Phi} = 1 + tP\Psi, \quad e^{-t\Phi} = 1 - tP\Psi. \quad (2.30)$$

We use the map to write the first term as

$$S_1 = -\frac{1}{2g_o^2} \oint QG\eta_0 G^{-1} = \frac{1}{2g_o^2} \oint (\Psi - PQ\Psi)Y\Psi = -\frac{1}{2g_o^2} \int (YQ\Psi)(Y\Psi). \quad (2.31)$$

Here we used the fact that in the resulting expression  $\xi_0$  appears only in  $P$ , the relation between  $P$  and  $Y$  and the relation between the integration in the large and small Hilbert spaces (1.10).

We now face a problem. In the above equation we have a collision of  $Y$ 's, which is singular. In order to evaluate the action one has to employ some sort of regularization. We propose to replace  $P$  by a regularized  $P_\epsilon$ , in which the  $\xi$  and  $Y$  insertions approach the potentially singular points at a different pace,

$$P_\epsilon = \frac{e^{i\frac{\pi}{4}}\xi((1 - \epsilon_1)i)Y((1 - \epsilon_2)i) + e^{-i\frac{\pi}{4}}\xi(-(1 - \epsilon_1)i)Y(-(1 - \epsilon_2)i)}{\sqrt{2}}. \quad (2.32)$$

There are several ways to think of such a regularization. One could consider it simply as a trick in order to evaluate a singular expression in the action. Another option exists if one considers the case  $\epsilon_1 = \epsilon_2$ . Then, it can represent the regularization of the star product itself, in the spirit of Witten's original construction. This interpretation was used in various contexts in recent developments [23, 18, 24], where it was represented using

$$\mathcal{O} \rightarrow s^{-L_0} \mathcal{O} s^{L_0}. \quad (2.33)$$

Alternatively, we can think of the regularization as defining a family of string fields, of which our choice is the limiting case. Now, there are corrections to (2.17). However, if we let  $\epsilon_1$  approach zero in a much faster pace than  $\epsilon_2$ , then we only have to add one more term in this expansion, since higher order terms will be subleading even when contracted with  $Y$ 's. One can see that the contributions to the action from including this term is zero, so it can be safely neglected. Plugging the regularization we obtain

$$S_1 = -\frac{1}{2g_o^2} \int \frac{i(Y(z_1)Y(z_2) - Y(z_1^*)Y(z_2^*)) + (Y(z_1)Y(z_2^*) + Y(z_2)Y(z_1^*))}{2} Q\Psi\Psi, \quad (2.34)$$

where we defined

$$z_1 = (1 - \epsilon_2)i, \quad z_2 = \frac{i}{1 - \epsilon_2}. \quad (2.35)$$

The two terms in the first parentheses give exactly the results one would obtain from choosing either  $k_1 = 1, k_2 = 0$  or  $k_1 = 0, k_2 = 1$  in (2.10). Below (2.22) we explained that solutions based on different choices of  $k_i$  are gauge equivalent. Hence, these terms give the difference of the quadratic part of the action evaluated on gauge equivalent solutions. We shall see below that in  $S_2$  one gets also the difference of the cubic terms, so together these terms cancel out. Thus, effectively we get the relation

$$YY \simeq Y_{-2}. \quad (2.36)$$

We can now take the limit  $\epsilon_2 \rightarrow 0$  and get

$$S_1 = -\frac{1}{2g_o^2} \int Y_{-2} Q\Psi\Psi. \quad (2.37)$$


For the second term in the action we write

$$S_2 = \frac{1}{2g_o^2} \oint \int_0^1 dt P\Psi[-tY\Psi, t\Psi - tPQ\Psi] = -\frac{1}{6g_o^2} \oint P\Psi[Y\Psi, \Psi] = -\frac{1}{3g_o^2} \int Y_{-2}\Psi\Psi\Psi. \quad (2.38)$$

Here we used the fact that in our regularization  $P^2Y$  goes to zero, then we evaluated the simple  $t$  integral and moved to the small Hilbert space, where we wrote the result using (2.36). Summing up the two terms  $S_1 + S_2$  gives exactly the action of the cubic theory.

Of course, by using (2.36) in the last step of (2.38) we neglected the contributions that should cancel the first parentheses of (2.34). We can represent these terms in a manifestly symmetric way as in fig. 1. Here, we use a coordinate system in which the three  $\Psi$ 's

are inserted at  $\pm\sqrt{3}$  and at zero and the  $Y$ 's are inserted at the locations derived from using  $\epsilon_2 = 0.005$ . The mirror image of this figure appears with a minus sign, so these terms correspond to the difference of the cubic term in the action of the gauge equivalent solutions based on  $P(\pm i)$ . The coefficients are easily seen to agree with those of (2.34), so these terms cancel out as promised.



**Figure 1:** The 3-vertex with two regularized  $Y(i)$  insertions in the upper half plane. The operators generating  $\Psi$  are inserted at the green points using appropriate conformal transformations and the (zero-weight)  $Y$ 's are inserted at the red points. Coordinate axes and half a unit disk are drawn for perspective.

By using the specific choice of  $P$  (2.15), we showed that the action of the non-polynomial solution is exactly the same as that of the cubic theory with the non-chiral choice of  $Y_{-2}$ . One is free to work with any choice of  $P$  in the non-polynomial theory, since they all give gauge equivalent solutions. However, the specific choice of  $P$  (2.15) has the advantage of sending real string fields to real string fields. Since (1.13) sends real solutions to real solutions, we conclude that the mappings we use are consistent with the string field reality conditions.

### 3. Formal gauge form of solutions in the cubic theories

In this section we discuss the formal gauge representation of solutions. In 3.1 we give the formal gauge form of solutions of the cubic superstring field theory and describe our map as a map of gauge fields. Then, in 3.2 we discuss the problems in obtaining the same formal representation for solutions of the bosonic theory and give some directions towards this end.

#### 3.1 Gauge form of the solutions

In [18] a mapping of bosonic solutions, given in a formal gauge form, to the non-polynomial theory, was considered. That mapping could also be used to map superstring fields from the cubic theory, as we are doing here. Let  $\Lambda$  be a gauge field defining a solution to a cubic (bosonic or supersymmetric) string field theory via

$$\Psi = e^{-\Lambda} Q e^{\Lambda}. \tag{3.1}$$

Define a corresponding solution in the non-polynomial theory by defining the gauge fields,

$$\Lambda_\eta = \xi(z)\Lambda, \quad \Lambda_Q = P(z)\Lambda, \quad (3.2)$$

for some  $z$ . A pure gauge solution of the non-polynomial theory is given by

$$e^\Phi = e^{-Q\Lambda_Q} e^{\eta_0\Lambda_\eta} = e^{P(z)Q\Lambda - \Lambda} e^\Lambda, \quad (3.3)$$

and we see that the choice of  $z$  for  $\xi$  drops out of the expression for the solution, but remains for  $P$ . Expansion in powers of  $\Lambda$  gives a simple leading order behaviour,

$$\Phi_1 = P(z)Q\Lambda = P(z)\Psi_1, \quad (3.4)$$

which has a canonical form (up to the choice of  $z$ ) and agrees with the first order of (2.3) (with the same choice of  $z$ ).

We can change our scheme [25, 18] and redefine

$$1 + \Phi = (1 - Q\Lambda_Q) \frac{1}{1 - \eta_0\Lambda_\eta} = (1 + P(z)Q\Lambda - \Lambda) \frac{1}{1 - \Lambda} = 1 + (P(z)Q\Lambda) \frac{1}{1 - \Lambda}. \quad (3.5)$$

If we choose to work with a  $P$  of the form (2.10), we can write that as

$$1 + \Phi = 1 + P(Q\Lambda \frac{1}{1 - \Lambda}) = 1 + P\Psi, \quad (3.6)$$

which is exactly the map (2.3).

An important question raised in [25, 18] is how general is the representation of solutions as formal gauge solutions. We can now give a simple answer for the case of the cubic superstring field theory: all its solutions can be formally represented as gauge solutions via (1.13), where  $G$  is defined by (2.17).

One may wonder what happens when this gauge representation is used in order to obtain a general formal gauge form of solutions of the non-polynomial theory via (3.2). Now, the gauge field of the cubic theory is given by

$$\Lambda = \log G = P\Psi. \quad (3.7)$$

Plugging this  $\Lambda$  into (3.2) results in an expression, which generally differs from what one would like to obtain and in particular is trivial,

$$\tilde{G} = 1, \quad (3.8)$$

if the same  $z$  is used in the definition of  $P$  and in (3.2). This implies that we cannot use our construction to obtain a general formal gauge representation for the non-polynomial theory. This should have been expected, since we did not extend the space in any way that would allow the introduction of formal gauge fields.

### 3.2 Formal gauge representation for the bosonic solutions

The map of gauge fields to the non-polynomial theory (3.2), (3.3) can also be used to map bosonic string fields to the non-polynomial theory<sup>1</sup>. On the other hand, the mapping (2.3) cannot be generally used to map bosonic solutions to analogous ones in the non-polynomial theory. The reason being that the supersymmetric BRST charge contains terms that do not exist in the bosonic theory,

$$Q = Q_0 + Q_1 + Q_2 . \quad (3.9)$$

Here, the subscript of  $Q$  counts its  $\phi$  charge. The bosonic BRST charge is contained in  $Q_0$ . The difference  $Q_0 - Q_{bos}$  is zero when acting upon bosonic states not containing factors of the  $b$  ghost. The  $Q_2$  term on the other hand gives zero upon terms not containing the  $c$  ghost. The  $Q_1$  term is zero on terms not containing the  $\partial X^\mu$  oscillators. Thus, bosonic solutions are generally modified when translated to the analogous supersymmetric ones. This implies that (2.18) does not simply generalize to the bosonic case, since the use there of the supersymmetric  $Q$  is important for its derivation. We do not have a general form for a formal gauge representation for bosonic solutions.

One can still map bosonic solutions into the supersymmetric ones, in those cases where a formal gauge representation of the solution exists,

$$\Psi_{bos} = e^{-\Lambda_{bos}} Q_{bos} e^{\Lambda_{bos}} . \quad (3.10)$$

One can seek a formal gauge representation for bosonic solutions by enlarging the bosonic Hilbert space using a couple of canonically conjugated anti-commuting variables  $\tilde{b}_0, \tilde{c}_0$ . Define their conformal weight to be zero and their ghost number to be  $-1$  and  $1$  respectively. Also, define

$$\tilde{c}_0 |0\rangle = 0 . \quad (3.11)$$

Then, the redefinition

$$\tilde{Q}_{bos} = Q_{bos} + \tilde{c}_0 , \quad (3.12)$$

does not change the cohomology problem when restricted to the subspace without the  $\tilde{b}_0$  mode. In the enlarged space, on the other hand,  $\tilde{Q}_{bos}$  has a trivial cohomology,

$$\tilde{Q}_{bos} \Psi = 0 \quad \Rightarrow \quad \Psi = Q(\tilde{b}_0 \Psi) . \quad (3.13)$$

We also define that these operators simply multiply with respect to the star product. Then, we can write all bosonic solutions as formal gauge solutions,

$$\Psi = G^{-1} \tilde{Q}_{bos} G , \quad G = 1 + \tilde{b}_0 \Psi . \quad (3.14)$$

---

<sup>1</sup>The idea of mapping bosonic solutions to supersymmetric ones may seem strange, since the two theories are obviously different. However, it seems natural to assume that a bosonic string field, without an explicit dependence on more than ten space-time directions, can be mapped to some sort of a supersymmetric string field, whose physical meaning is similar. Using such a map we would be able to learn about the superstring from its simpler bosonic counterpart. For another relation of the two theories see [26].

While the above may be useful for the bosonic theory, it does not help us with mapping bosonic solutions to supersymmetric ones. A redefinition of the supersymmetric  $Q$  according to

$$\tilde{Q} = Q + \tilde{c}_0, \quad (3.15)$$

may be used to define a supersymmetric string field  $G^{-1}\tilde{Q}G$ . Unfortunately, this string field contains in general a  $\tilde{b}_0$  piece, due to the  $Q - Q_{bos}$  term. So it cannot be used for deriving supersymmetric solutions from the bosonic ones.

#### 4. Incorporating the GSO(−) sector

Our construction can be generalized to the cases of non-BPS D-branes and D-brane systems. Recall first that in order to describe the GSO(−) sector one has to introduce into the theory the so called “internal Chan-Paton indices” [27, 28]. For the non-polynomial theory one has to tensor the GSO(+) sector with the two dimensional unit matrix  $\mathbf{1}$  and the GSO(−) sector with the Pauli matrix  $\sigma_1$ . The two gauge fields of a GSO(+) sector are to be tensored with  $\sigma_3$ , while the gauge fields of the GSO(−) sectors get a factor of  $i\sigma_2$ . The operators  $Q, \eta_0$  are also tensored with  $\sigma_3$  and in the action a trace over the new matrix space should be performed and the result should be normalized by  $2 = \text{tr}(\mathbf{1})$ . With these simple rules all the algebraic properties hold just as in the case of a single GSO(+) sector. The non-BPS D-brane can be described by introducing one GSO(+) and one GSO(−) sector, while for more complicated brane systems one has to introduce in addition to the internal space also the genuine Chan-Paton space, with each factor tensored also in the internal space according to its GSO parity. The same story works for the cubic theory. Now, however, the gauge fields are the ones that get the factors of  $\mathbf{1}, \sigma_1$  and the string fields are tensored with  $\sigma_3, i\sigma_2$ . The only other modification is the inclusion of a factor of  $\sigma_3$  also for  $Y_{-2}$ .

The algebraic properties that are important for our construction are the commutation relations (2.11, 2.14). In order that they will continue to hold we have to assign a factor of  $\sigma_3$  to  $P(z)$  and  $\xi(z)$ , which implies that  $Y(z)$  carries a factor of  $\mathbf{1}$ . Then, upon moving from the large to the small Hilbert space by integrating the  $\xi_0$ , the leftover  $\sigma_3$  factor combines with the two  $Y$ ’s and the relation (2.36) also remains intact. With these simple assignments all the proofs of section 2 carry over to the general brane system without any further modification. We conclude that there is no obstruction for generalizing our map to an arbitrary D-brane system.

#### 5. Vacuum solutions

In this section we discuss vacuum solutions in superstring field theory. In 5.1 we describe the preliminary proposal for a vacuum solution in the non-polynomial theory on a BPS D-brane given in [18]. We explain the problem of the proposal and correct it. Then, we discuss the analogous construction, by Erler, in the cubic theory [19] and show that the two are related by (2.3). This proves that the solution of the non-polynomial theory shares the properties of correct action and zero cohomology derived in [19] for the cubic solution.

The solution on the BPS D-brane was criticized in [20], where a solution living on a non-BPS D-brane was constructed. In 5.2 we describe the criticism and the new solution and prove that the cohomology of the solution is trivial as expected. We further study the one parameter family of solutions interpolating Erler’s solution on the non-BPS D-brane and the new solution and show that all the members in this family are gauge equivalent.

### 5.1 On the BPS D-brane

In [18], it was suggested that the map of gauge solutions (3.2),(3.3) from the bosonic to the non-polynomial theory can be used to find a “tachyon vacuum” in the supersymmetric theory. Since there is no tachyon on the BPS D-brane the new vacuum cannot represent tachyon condensation. It can still, potentially, represent a state where the D-brane is absent.

During most of that work the operator  $P(0)$  was used. This results in

$$\Lambda_Q = (P B c)(0) |0\rangle = P(0) |0\rangle, \quad (5.1)$$

implying that

$$Q \Lambda_Q = |0\rangle, \quad (5.2)$$

which lives in the small Hilbert space. The gauge transformation  $\eta_0 \Lambda_\eta$  always leads, by definition, to a state in the small Hilbert space. The solution resides in the small Hilbert space and is therefore pure gauge. Since the source of illness here is the OPE of  $P(z)$  and  $Bc(z)$ , an immediate remedy would be to move  $P(z)$  to another point. Here, we suggest to move it to  $\pm i$ , that is, to use  $P$  of (2.15), so as to get

$$\Lambda_Q = P \Lambda. \quad (5.3)$$

This expression lives in the large Hilbert space.

In order to calculate the coefficients of the various fields we have to expand  $P$  around  $z = 0$ . This is done in a derivative, i.e., conformal weight expansion, by noting that

$$P = (\cos \partial - \sin \partial) P(0) = P(0) - P'(0) - \frac{1}{2} P''(0) + \dots \quad (5.4)$$

To find the lowest non-trivial field we can use (3.6) and recall that for  $\Psi$  the lowest level field with non-zero value is  $c(0) |0\rangle$ . Acting with (5.4) on  $c(0) |0\rangle$  we get

$$P c(0) |0\rangle \approx -P'(0) c(0) |0\rangle = -c c' \xi \xi' e^{-2\phi} |0\rangle. \quad (5.5)$$

The field  $c c' \xi \xi' e^{-2\phi}$  lives in the large Hilbert space, so it may give rise to some physical content. It does not obey the Siegel gauge, so it was not studied much in the level-truncation literature. It may seem to describe an unintegrated vertex operator because of the  $c$  factor. However, a direct evaluation reveals that it does not correspond to a vertex operator. This field is the only field other than the identity field with all quantum numbers being zero. It follows that there is no way to correct it in order to obtain a vertex operator so it is an auxiliary field.

In [19], Erler suggested to use the trivial mapping of Schnabl’s gauge field to the cubic superstring field theory, in order to obtain a non-perturbative vacuum in this theory. He further showed that it gives the expected action of a solution describing the absence of the D-brane and that the cohomology around the solution is empty. The cohomology proof is in fact straightforward. The resulting string field is almost identical to the bosonic one and differs only due to the new terms in  $Q$ . This gives,

$$\Psi = \Psi_{bos} + B\gamma^2 |0\rangle = \Psi_{bos} + B\eta\partial\eta e^{2\phi} |0\rangle. \quad (5.6)$$

The cohomology proof in the bosonic case [22] used the operator

$$A = B \int_0^1 dt \Omega^t, \quad (5.7)$$

where  $\Omega = |0\rangle$  is the vacuum and its powers are defined to be the appropriate wedge states. This field obeys the relation

$$Q_\Psi A = QA + \Psi A + A\Psi = 1, \quad (5.8)$$

which implies that the cohomology is empty. Since the only new term in (5.6) is simply proportional to  $B$  and contain no  $c$  factors, the above equation holds without modifying  $A$  for the supersymmetric case as well.

The calculation of the action relies on the fact that in order to get a non-trivial result after the integration, the integrand has to be proportional to  $e^{-2\phi}$ . The picture changing operator  $Y_{-2}$  supplies a  $-4$  factor to the  $\phi$  momentum in the action, so the string field should supply a factor of two. This can come either from the  $\gamma^2$  term in (5.6) or from  $Q$  acting on  $c$ . Finally, all expectation values give linear functions rather than the trigonometric functions one gets in the bosonic case. This fact enables a much simpler evaluation, which results in the correct D-brane tension.

Mapping Erler’s solution using (2.3) is equivalent to using the modified gauge representation (5.3). From the discussion of section 2 we conclude that the solution of the non-polynomial theory also has the correct action and cohomology.

## 5.2 On the Non-BPS D-brane

In [20], Aref’eva, Gorbachev and Medvedev (hereafter AGM), criticized Erler’s work. First, it was claimed that a vacuum solution on a non-BPS D-brane should non exist. Then, it was claimed that the condensation of the string field  $c(0) |0\rangle$ , which is the leading order in Erler’s solution, should not correspond to the tachyon vacuum anyway, since this field, called “the BPS tachyon” is an auxiliary field and not a vertex operator and is an artifact of working in the zero picture. It is easy to see that if one tries to map it to the “natural”  $-1$  picture, it is mapped to zero. In fact, a solution based on this vertex operator (in the theory with chiral  $Y_{-2}$ ) was found using lowest-level truncation in [29], where it was interpreted as a supersymmetry breaking solution. In AGM it was suggested that the correct resolution of these problems is to define the string field on the non-BPS D-brane



using a gauge field with Erler's gauge field in the GSO(+) sector<sup>2</sup> and with

$$\Lambda_- = B\gamma(0)|0\rangle, \quad (5.9)$$

in the GSO(-) sector.

It is immediate that the action is the same as in Erler's case. We think of the AGM solution as being expanded around Erler's one and write the difference

$$\tilde{\Psi} = \Psi_{AGM} - \Psi_E. \quad (5.10)$$

From general considerations it follows that

$$S_{AGM} = S_E + \frac{1}{6} \int \frac{1}{2} \text{tr}(Y_{-2} \tilde{\Psi}^3). \quad (5.11)$$

The form of the gauge field (5.9) implies that

$$\tilde{\Psi}_+ \approx \gamma B \gamma, \quad (5.12)$$

carries momentum two, while the GSO(-) part given by (split-string notations [30, 31, 32])

$$\tilde{\Psi}_- = Fc \frac{KB}{1-\Omega} \gamma F + F\gamma \frac{KB}{1-\Omega} cF, \quad (5.13)$$

carries one unit of  $\phi$  momentum. It follows that the minimal  $\phi$ -momentum power that one can get in the above action is four, which implies

$$S_{AGM} = S_E. \quad (5.14)$$

The proof (not given in AGM) that the cohomology of the AGM solution is empty relies again on the same  $A$  (5.7) as in the previous cases, only now it should be tensored with  $\sigma_3$ . For (5.8) to hold in this case, we only have to prove

$$[A, \tilde{\Psi}] = 0. \quad (5.15)$$

This is trivial for the GSO(+) part (5.12), which is proportional to  $B$ , without any  $c$  insertions. The GSO(-) part (5.13) gives

$$[\tilde{\Psi}, A] = \left( -F\gamma \frac{KB}{1-\Omega} cFB \int_0^1 \Omega^t + \int_0^1 \Omega^t BFc \frac{KB}{1-\Omega} \gamma F \right) \otimes \sigma_1. \quad (5.16)$$

Expanding the denominator we can write the terms in parentheses as

$$\begin{aligned} KB \int_0^\infty (\Omega^t(\gamma(0)|0\rangle) - (\gamma(0)|0\rangle)\Omega^t) &= B \int_0^\infty \partial_t(\Omega^t(\gamma(0)|0\rangle) - (\gamma(0)|0\rangle)\Omega^t) = \\ B(\Omega^\infty(\gamma(0)|0\rangle) - (\gamma(0)|0\rangle)\Omega^\infty) &= 0, \end{aligned}$$

where we used the fact that  $K$  acts as a derivation with respect to the strip length and the final identity is correct to order  $N^{-3}$ , just as the original proof of the vanishing of the cohomology [22].

---

<sup>2</sup>A GSO(+) component is indispensable, since the action of a purely GSO(-) solution is trivially zero.

The calculation of the action, the triviality of the cohomology and the existence of  $\gamma$  in the gauge field seem to be in favour of the interpretation of the AGM solution as representing tachyon condensation on the non-BPS D-brane. Nevertheless, one may wonder whether this solution is indeed different from Erler's one, or are they gauge equivalent. Furthermore, the reasoning regarding the form of the solution as well as the evaluation of the action and the cohomology remains unchanged if one multiplies the GSO(-) gauge field  $\Lambda_-$  by an arbitrary parameter  $\epsilon$ ,

$$\Lambda_+ = Bc(0)|0\rangle, \quad \Lambda_- = \epsilon B\gamma(0)|0\rangle. \quad (5.17)$$

In this manner one gets a one parameter family  $\Psi_\epsilon$  of solutions generated by (5.17). This family interpolates between Erler's solution and the AGM solution. All these solutions have the same action and trivial cohomology so it is natural to expect them to be gauge equivalent. If this is indeed the case, one is to conclude that the  $\epsilon = 0$  solution can also be used to describe tachyon condensation on the non-BPS D-brane. It is then very natural to assume that Erler's solution on the BPS D-brane and its counterpart in the non-polynomial theory indeed manage to describe the state without the BPS D-brane despite the fact that there are no tachyons living on it.

Writing the gauge transformation between two solutions in the  $\epsilon$ -family is (at least formally) immediate. Since these solutions are formally given as gauge solutions one only has to compose these two gauge transformation to get a new one. Then, the question is whether the resulting transformation is a genuine gauge transformation or whether it is singular in any sense.

We call the gauge transformation from Erler's solution, with  $\epsilon = 0$  to the  $\epsilon$ -solution  $\Psi_\epsilon$ ,  $e^{\Lambda_E^\epsilon}$ . Recall however that the gauge form of the solutions is given in the left scheme, so we can write

$$\begin{aligned} e^{\Lambda_E^\epsilon} &= (1 - \Lambda_E) \frac{1}{1 - \Lambda_\epsilon} = 1 \otimes \mathbf{1} + \epsilon B\gamma(0)|0\rangle \otimes \sigma_1, \\ e^{-\Lambda_E^\epsilon} &= (1 - \Lambda_\epsilon) \frac{1}{1 - \Lambda_E} = 1 \otimes \mathbf{1} - \epsilon B\gamma(0)|0\rangle \otimes \sigma_1, \end{aligned} \quad (5.18)$$

where we used the relation

$$\Lambda_-^2 = \Lambda_- \Lambda_+ = 0, \quad (5.19)$$

to get the final expressions. These transformations form an abelian group with the simple multiplication rule,

$$e^{\Lambda_E^\epsilon} e^{\Lambda_E^{\tilde{\epsilon}}} = e^{\Lambda_E^{\epsilon+\tilde{\epsilon}}}, \quad (5.20)$$

where we again used (5.19). We see no sign of a singularity at any value of  $\epsilon$ .

We can also check that this gauge transformation acts correctly on  $A$

$$A_\epsilon = e^{-\Lambda_E^\epsilon} A e^{\Lambda_E^\epsilon} = A. \quad (5.21)$$

A singularity in the gauge transformation (5.18) would probably express itself as a singularity in the above equation. This does not happen. Finally, one may want to calculate some invariants of the  $\Psi_\epsilon$  solution as in [33, 34] and verify that they are  $\epsilon$ -independent. We

did not try this method, but we believe that if tried it would indeed result in  $\epsilon$ -independent expressions.

We can map the  $\epsilon$ -family to the non-polynomial theory. Explicitly, the solution

$$\begin{aligned} \Psi_\epsilon = & \left( Fc \frac{KB}{1-\Omega} cF + F\gamma \left( \frac{\epsilon^2 K}{1-\Omega} + (1-\epsilon^2) \right) B\gamma F \right) \otimes \sigma_3 \\ & + \epsilon \left( Fc \frac{KB}{1-\Omega} \gamma F + F\gamma \frac{KB}{1-\Omega} cF \right) \otimes (i\sigma_2), \end{aligned} \quad (5.22)$$

is mapped to

$$\begin{aligned} \Phi_\epsilon = & P \left( Fc \frac{KB}{1-\Omega} cF + F\gamma \left( \frac{\epsilon^2 K}{1-\Omega} + (1-\epsilon^2) \right) B\gamma F \right) \otimes \mathbf{1} \\ & + \epsilon P \left( Fc \frac{KB}{1-\Omega} \gamma F + F\gamma \frac{KB}{1-\Omega} cF \right) \otimes \sigma_1. \end{aligned} \quad (5.23)$$

While this mapping seems to be trivial, it serves us to address one last important point. The  $B$  line integral runs from  $-i$  to  $i$  and the  $B$  operator has non-trivial commutation relation with  $P$ . It follows that again we have to regularize in order to make sense of the expression. We let  $B$  pass to the right of the regularized  $P$  insertion, as is implicit in (5.23). Then, one may wonder whether the commutation relation of  $B$  and  $P$  would invalidate the nilpotency of  $\Phi$ , which was inherited from that of  $P$ . Again, we can establish the nilpotency by defining a much faster pace in the  $\epsilon_1$  direction compared with that of the  $\epsilon_2$  direction in (2.32). All our treatment remains consistent when such a regularization is used.

When using wedge state algebra in the evaluation of expressions involving  $\Phi$ , one has to include the so called phantom-terms. The form and origin of these terms is quite clear by now [19]. They originate from replacing terms of the form  $(1-\Omega)^{-1}$  with their series form. The phantom-terms should be added to (5.22) as described in [19, 20]. This will result in their presence also in (5.23). There is no need for the phantom piece, neither in (5.22) nor in (5.23) when level-truncation is being used for the evaluation of expressions.

## 6. Conclusions

In this work we presented mappings between gauge orbits of the cubic and the non-polynomial superstring field theories. An obvious question is whether it is possible to extend our construction to the quantum theory. Two important challenges stand in the way: the extension of the maps to general (off-shell) string fields and the inclusion of the Ramond sector.

The space of string fields that one has to map for the off-shell theory is larger than the space of string fields we considered in this work. This stems from the fact that both theories are infinitely reducible gauge theories. Moreover, the reducibility of the gauge symmetry is realized only on-shell. This suggests that quantization should be addressed using the BV formalism [35, 36, 37] (see [38, 39] for reviews), as was done for the bosonic theory in [40, 41, 42]<sup>3</sup>. The enlargement of the space of fields comes from the inclusion of

---

<sup>3</sup>A BRST study of the bosonic string was performed in [43, 44].

the (second quantized) ghosts and antifields. It can happen that the maps can be defined in this enlarged space despite not being defined on the smaller space. These hypothetical maps would have to commute with the superbrackets and would therefore have the interpretation of canonical transformations.

The fact that the gauge symmetries of the two theories are different may be an obstacle for this ambitious plan. A possible resolution is to gauge fix one of the gauge symmetries of the non-polynomial theory. One can find in the literature the gauge fixing

$$\xi_0 \Phi = 0. \tag{6.1}$$

This is not invariant under the star product. A more adequate gauge choice seems to be

$$\xi(i)\Phi = 0, \tag{6.2}$$

or the analogous form in the physical case of a theory with the reality condition included. This gauge fixing is immediately solved by writing

$$\Phi = \xi(i)\tilde{\Phi}, \tag{6.3}$$

and our map reduces in this case to

$$\tilde{\Phi} = Y(i)\Psi. \tag{6.4}$$

Now, the spaces and gauge symmetries of both theories are quite similar. For the theory with the reality condition imposed, this construction does not work in this way, since  $P$  of (2.15) cannot be written as a product of  $\xi$  and  $Y$ . One possible resolution of this problem is to impose the gauge condition directly with  $P$ , with the immediate solution being (2.3). The fact that the transformation is linear implies that at least formally no problematic measure factors would emerge. The question is how to generalize the inverse map (1.13) so as to include also off-shell states and anti-fields.

The inclusion of the Ramond sector is essential, since Ramond states appear in loops even in scattering processes of NS states. This raises the familiar problems related to defining an action for the Ramond sector in superstring field theory [16, 45]. We hope that it will be possible to extend this work to the Ramond sector. Presumably it could also help to resolve some of the problems related to the definition of the Ramond sector action.

Understanding the Ramond sector is desirable already at the classical level. For one thing, one may want to be able to study classical Ramond sector solutions. Another important issue is the evaluating of the complete spectrum of the theory around a classical solution. As we mentioned in 5.2, a preliminary numerical study of a solution analogous to Erler's one performed in [29] resulted in the interpretation of the new vacuum as a supersymmetry breaking one. Understanding the Ramond sector will enable us to study this question analytically. We believe that this solution represents the closed string vacuum and so should have a vanishing cohomology also in the Ramond sector. Not only that the supersymmetry should not be broken around it, but also the solution should be left invariant under a larger supersymmetry. Other aspects of supersymmetry are also of importance.

A common objection to the use of the cubic theory is its usage of the (inverse) picture changing operators  $Y(\pm i)$ . The problem being that these operators have non-trivial kernels, which may result in the equations of motion (1.4) and (1.5) being inequivalent. It is not clear if this is really a problem, since the problematic states are of a somewhat exotic form, i.e., they have a factor that is localized at the string mid-point and it is not clear if such states belong to the space of string fields that one has to consider. We used just these states in our map to the non-polynomial theory. Hence, we may conclude that the cubic theory on a space not including these states is equivalent to the non-polynomial one on a space where they are included. One may still wonder whether it is possible to define an adequate space of string fields for the cubic theory. It is known for example that the space of finite combinations of local insertions on the vacuum is not closed under the star product. Closeness under the star product is certainly a very desirable property of the looked-for space. Can one define a space closed under the star product, that is large enough for including the expected classical solutions, yet not containing states with insertions at  $\pm i$ ?

In order to get some intuition about the needed space of string fields we may assist the remarkable developments in the field during the last couple of years, following Schnabl's analytical solution [46]. Subsequent work [30, 47, 22] enabled the proof of Sen's conjectures [48, 49], analytical solutions describing marginal deformations were found [50, 51, 52, 53, 54, 25, 18, 55, 56, 57], scattering amplitudes and gauge conditions were studied [58, 59, 24, 60] and much more [61, 62, 63, 64, 65, 66, 67, 68, 69, 70]. These developments used to a large extent wedge states [71, 72] with local insertions [73] and with insertions of the line integrals  $B, K$ . It seems plausible that the space of string fields should be based on such states, which do form a closed algebra under the star product (similar but extended subalgebras may include general surface states [74, 75] with local insertions and the subalgebra of wedge and butterfly states [76, 77] with local insertions). Of course, infinite linear combinations are still needed. This may result in no restriction at all on the space, since without restricting the coefficients all possible states can be formed from local insertions, even without the introduction of wedge states. Thus, we have to assume that the coefficients should be restricted not to grow, in some sense, faster than in the known cases. If this is the case, the formal problems in our construction can be solved, as was explained at the end of 5.2. Obviously, more research in this direction is needed.

We showed that the vacuum solution for non-BPS D-brane is gauge equivalent to a solution that can be defined also on the BPS D-brane. This convinces us that the solution for the BPS D-brane should be taken seriously. The main criticism on this solution is that it contradicts our understanding that the BPS D-brane is stable. In other words, a condensation of a BPS D-brane would correspond to a non-continuous change in the Ramond-Ramond charge. Yet, we can think of a continuous dynamical process in which a D-brane disappears. For example, we can start with a system of two D5-branes, send one D5-brane to spatial infinity and remain with only one D5-brane. One might worry that hiding one D-brane at infinity is only a trick and try considering a compact space. However, on a compact space the total R-R charge has to vanish. This means that the possibility of hiding the D-brane at infinity is correlated with the existence of a non-vanishing R-R charge. We could also consider space-time filling D9-branes. In this case

tadpole cancellation would constrain the total charge to zero. All the above arguments are related to the close-string sector, which we did not consider in this paper. Still, it is important to show that they work out correctly, otherwise inconsistencies would probably show up at 1-loop calculations. A related issue is the common expectation that open string field theory should describe only changes of open string moduli. It seems that open string field theory can influence the close-string sector, at least to some extent. It would be very desirable to understand better the realm of open string field theory.

## Acknowledgments

We would like to thank the organizers of the workshop “String field theory and related aspects” held at the Arnold Sommerfeld Center, Munich, for hospitality and for providing such a stimulating research environment. We thank the participants of this conference for many discussions. We would also like to thank Ido Adam, Guillaume Bossard, Stefan Fredenhagen and Stefan Theisen for useful discussions. The work of M. K. was supported by a Minerva fellowship during the initiation of this project. This research was supported by the German-Israeli Project cooperation (DIP H.52).

## References

- [1] E. Witten, *Interacting field theory of open superstrings*, *Nucl. Phys.* **B276** (1986) 291.
- [2] E. Witten, *Noncommutative geometry and string field theory*, *Nucl. Phys.* **B268** (1986) 253.
- [3] D. Friedan, E. J. Martinec, and S. H. Shenker, *Conformal invariance, supersymmetry and string theory*, *Nucl. Phys.* **B271** (1986) 93.
- [4] C. Wendt, *Scattering amplitudes and contact interactions in Witten’s superstring field theory*, *Nucl. Phys.* **B314** (1989) 209.
- [5] I. Y. Arefeva and P. B. Medvedev, *Anomalies in Witten’s field theory of the NSR string*, *Phys. Lett.* **B212** (1988) 299.
- [6] C. R. Preitschopf, C. B. Thorn, and S. A. Yost, *Superstring field theory*, *Nucl. Phys.* **B337** (1990) 363–433.
- [7] I. Y. Arefeva, P. B. Medvedev, and A. P. Zubarev, *New representation for string field solves the consistence problem for open superstring field*, *Nucl. Phys.* **B341** (1990) 464–498.
- [8] I. Y. Arefeva, P. B. Medvedev, and A. P. Zubarev, *Background formalism for superstring field theory*, *Phys. Lett.* **B240** (1990) 356–362.
- [9] I. Y. Arefeva, D. M. Belov, A. A. Giryavets, A. S. Koshelev, and P. B. Medvedev, *Noncommutative field theories and (super)string field theories*, [hep-th/0111208](#).
- [10] B. V. Urosevic and A. P. Zubarev, *On the component analysis of modified superstring field theory actions*, *Phys. Lett.* **B246** (1990) 391–398.
- [11] A. Ishida and S. Teraguchi, *On the general action of boundary (super)string field theory*, [0805.1826](#).
- [12] N. Berkovits, *Super-Poincaré invariant superstring field theory*, *Nucl. Phys.* **B450** (1995) 90–102, [[hep-th/9503099](#)].

- [13] B. Zwiebach, *Closed string field theory: Quantum action and the B-V master equation*, *Nucl. Phys.* **B390** (1993) 33–152, [[hep-th/9206084](#)].
- [14] Y. Okawa and B. Zwiebach, *Heterotic string field theory*, *JHEP* **07** (2004) 042, [[hep-th/0406212](#)].
- [15] N. Berkovits, Y. Okawa, and B. Zwiebach, *WZW-like action for heterotic string field theory*, *JHEP* **11** (2004) 038, [[hep-th/0409018](#)].
- [16] N. Berkovits, *The Ramond sector of open superstring field theory*, *JHEP* **11** (2001) 047, [[hep-th/0109100](#)].
- [17] A. Kling, O. Lechtenfeld, A. D. Popov, and S. Uhlmann, *Solving string field equations: New uses for old tools*, *Fortsch. Phys.* **51** (2003) 775–780, [[hep-th/0212335](#)].
- [18] E. Fuchs and M. Kroyter, *Marginal deformation for the photon in superstring field theory*, *JHEP* **11** (2007) 005, [[arXiv:0706.0717](#) [[hep-th](#)]].
- [19] T. Erler, *Tachyon vacuum in cubic superstring field theory*, *JHEP* **01** (2008) 013, [[0707.4591](#)].
- [20] I. Y. Aref’eva, R. V. Gorbachev, and P. B. Medvedev, *Tachyon solution in cubic Neveu-Schwarz string field theory*, [0804.2017](#).
- [21] I. Ellwood, B. Feng, Y.-H. He, and N. Moeller, *The identity string field and the tachyon vacuum*, *JHEP* **07** (2001) 016, [[hep-th/0105024](#)].
- [22] I. Ellwood and M. Schnabl, *Proof of vanishing cohomology at the tachyon vacuum*, *JHEP* **02** (2007) 096, [[hep-th/0606142](#)].
- [23] E. Fuchs and M. Kroyter, *Universal regularization for string field theory*, *JHEP* **02** (2007) 038, [[hep-th/0610298](#)].
- [24] M. Kiermaier, A. Sen, and B. Zwiebach, *Linear b-gauges for open string fields*, *JHEP* **03** (2008) 050–050, [[0712.0627](#)].
- [25] E. Fuchs, M. Kroyter, and R. Potting, *Marginal deformations in string field theory*, *JHEP* **09** (2007) 101, [[arXiv:0704.2222](#) [[hep-th](#)]].
- [26] N. Berkovits and C. Vafa, *On the uniqueness of string theory*, *Mod. Phys. Lett.* **A9** (1994) 653–664, [[hep-th/9310170](#)].
- [27] N. Berkovits, A. Sen, and B. Zwiebach, *Tachyon condensation in superstring field theory*, *Nucl. Phys.* **B587** (2000) 147–178, [[hep-th/0002211](#)].
- [28] I. Y. Arefeva, D. M. Belov, and A. A. Giryavets, *Construction of the vacuum string field theory on a non-BPS brane*, *JHEP* **09** (2002) 050, [[hep-th/0201197](#)].
- [29] I. Y. Arefeva, P. B. Medvedev, and A. P. Zubarev, *Nonperturbative vacuum for superstring field theory and supersymmetry breaking*, *Mod. Phys. Lett.* **A6** (1991) 949–958.
- [30] Y. Okawa, *Comments on Schnabl’s analytic solution for tachyon condensation in Witten’s open string field theory*, *JHEP* **04** (2006) 055, [[hep-th/0603159](#)].
- [31] T. Erler, *Split string formalism and the closed string vacuum*, *JHEP* **05** (2007) 083, [[hep-th/0611200](#)].
- [32] T. Erler, *Split string formalism and the closed string vacuum. II*, *JHEP* **05** (2007) 084, [[hep-th/0612050](#)].

- [33] I. Ellwood, *The closed string tadpole in open string field theory*, 0804.1131.
- [34] T. Kawano, I. Kishimoto, and T. Takahashi, *Gauge invariant overlaps for classical solutions in open string field theory*, 0804.1541.
- [35] J. Zinn-Justin, *Renormalization of gauge theories*, . Lectures given at Int. Summer Inst. for Theoretical Physics, Jul 29 - Aug 9, 1974, Bonn, West Germany.
- [36] I. A. Batalin and G. A. Vilkovisky, *Gauge algebra and quantization*, *Phys. Lett.* **B102** (1981) 27–31.
- [37] I. A. Batalin and G. A. Vilkovisky, *Quantization of gauge theories with linearly dependent generators*, *Phys. Rev.* **D28** (1983) 2567–2582.
- [38] M. Henneaux and C. Teitelboim, *Quantization of gauge systems*, . Princeton, USA: Univ. Pr. (1992) 520 p.
- [39] J. Gomis, J. Paris, and S. Samuel, *Antibracket, antifields and gauge theory quantization*, *Phys. Rept.* **259** (1995) 1–145, [hep-th/9412228].
- [40] M. Bochicchio, *Gauge fixing for the field theory of the bosonic string*, *Phys. Lett.* **B193** (1987) 31.
- [41] M. Bochicchio, *String field theory in the Siegel gauge*, *Phys. Lett.* **B188** (1987) 330.
- [42] C. B. Thorn, *Perturbation theory for quantized string fields*, *Nucl. Phys.* **B287** (1987) 61.
- [43] T. G. Erler and D. J. Gross, *Locality, causality, and an initial value formulation for open string field theory*, hep-th/0406199.
- [44] T. G. Erler, *Level truncation and rolling the tachyon in the lightcone basis for open string field theory*, hep-th/0409179.
- [45] Y. Michishita, *A covariant action with a constraint and Feynman rules for fermions in open superstring field theory*, *JHEP* **01** (2005) 012, [hep-th/0412215].
- [46] M. Schnabl, *Analytic solution for tachyon condensation in open string field theory*, *Adv. Theor. Math. Phys.* **10** (2006) 433–501, [hep-th/0511286].
- [47] E. Fuchs and M. Kroyter, *On the validity of the solution of string field theory*, *JHEP* **05** (2006) 006, [hep-th/0603195].
- [48] A. Sen, *Descent relations among bosonic D-branes*, *Int. J. Mod. Phys.* **A14** (1999) 4061–4078, [hep-th/9902105].
- [49] A. Sen, *Universality of the tachyon potential*, *JHEP* **12** (1999) 027, [hep-th/9911116].
- [50] M. Schnabl, *Comments on marginal deformations in open string field theory*, *Phys. Lett.* **B654** (2007) 194–199, [hep-th/0701248].
- [51] M. Kiermaier, Y. Okawa, L. Rastelli, and B. Zwiebach, *Analytic solutions for marginal deformations in open string field theory*, *JHEP* **01** (2008) 028, [hep-th/0701249].
- [52] T. Erler, *Marginal solutions for the superstring*, *JHEP* **07** (2007) 050, [arXiv:0704.0930 [hep-th]].
- [53] Y. Okawa, *Analytic solutions for marginal deformations in open superstring field theory*, *JHEP* **09** (2007) 084, [arXiv:0704.0936 [hep-th]].



- [54] Y. Okawa, *Real analytic solutions for marginal deformations in open superstring field theory*, *JHEP* **09** (2007) 082, [[arXiv:0704.3612](#) [[hep-th](#)]].
- [55] I. Kishimoto and Y. Michishita, *Comments on solutions for nonsingular currents in open string field theories*, *Prog. Theor. Phys.* **118** (2007) 347–369, [[0706.0409](#)].
- [56] M. Kiermaier and Y. Okawa, *Exact marginality in open string field theory: a general framework*, [arXiv:0707.4472](#) [[hep-th](#)].
- [57] M. Kiermaier and Y. Okawa, *General marginal deformations in open superstring field theory*, [arXiv:0708.3394](#) [[hep-th](#)].
- [58] H. Fuji, S. Nakayama, and H. Suzuki, *Open string amplitudes in various gauges*, *JHEP* **01** (2007) 011, [[hep-th/0609047](#)].
- [59] L. Rastelli and B. Zwiebach, *The off-shell Veneziano amplitude in Schnabl gauge*, *JHEP* **01** (2008) 018, [[0708.2591](#)].
- [60] M. Kiermaier and B. Zwiebach, *One-loop Riemann surfaces in Schnabl gauge*, [0805.3701](#).
- [61] E. Fuchs and M. Kroyter, *Schnabl's  $\mathcal{L}_0$  operator in the continuous basis*, *JHEP* **10** (2006) 067, [[hep-th/0605254](#)].
- [62] L. Rastelli and B. Zwiebach, *Solving open string field theory with special projectors*, *JHEP* **01** (2008) 020, [[hep-th/0606131](#)].
- [63] Y. Okawa, L. Rastelli, and B. Zwiebach, *Analytic solutions for tachyon condensation with general projectors*, [hep-th/0611110](#).
- [64] L. Bonora, N. Bouatta, and C. Maccaferri, *Towards open-closed string duality: Closed strings as open string fields*, [hep-th/0609182](#).
- [65] I. Ellwood, *Rolling to the tachyon vacuum in string field theory*, *JHEP* **12** (2007) 028, [[0705.0013](#)].
- [66] O.-K. Kwon, *Marginally deformed rolling tachyon around the tachyon vacuum in open string field theory*, [arXiv:0801.0573](#) [[hep-th](#)].
- [67] A. Ishida, C. Kim, Y. Kim, O.-K. Kwon, and D. D. Tolla, *Tachyon vacuum solution in open string field theory with constant B field*, [0804.4380](#).
- [68] T. Kawano, I. Kishimoto, and T. Takahashi, *Schnabl's solution and boundary states in open string field theory*, [0804.4414](#).
- [69] A. Bagchi and A. Sen, *Tachyon condensation on separated brane-antibrane system*, *JHEP* **05** (2008) 010, [[0801.3498](#)].
- [70] S. Hellerman and M. Schnabl, *Light-like tachyon condensation in open string field theory*, [arXiv:0803.1184](#) [[hep-th](#)].
- [71] L. Rastelli and B. Zwiebach, *Tachyon potentials, star products and universality*, *JHEP* **09** (2001) 038, [[hep-th/0006240](#)].
- [72] L. Rastelli, A. Sen, and B. Zwiebach, *Boundary CFT construction of D-branes in vacuum string field theory*, *JHEP* **11** (2001) 045, [[hep-th/0105168](#)].
- [73] M. Schnabl, *Wedge states in string field theory*, *JHEP* **01** (2003) 004, [[hep-th/0201095](#)].
- [74] A. LeClair, M. E. Peskin, and C. R. Preitschopf, *String field theory on the conformal plane. 1. Kinematical principles*, *Nucl. Phys.* **B317** (1989) 411.

- [75] A. LeClair, M. E. Peskin, and C. R. Preitschopf, *String field theory on the conformal plane. 2. Generalized gluing*, *Nucl. Phys.* **B317** (1989) 464.
- [76] E. Fuchs and M. Kroyter, *On surface states and star-subalgebras in string field theory*, *JHEP* **10** (2004) 004, [[hep-th/0409020](#)].
- [77] S. Uhlmann, *A note on kappa-diagonal surface states*, *JHEP* **11** (2004) 003, [[hep-th/0408245](#)].