Non-Abelian Strings in Hot or Dense QCD

Eiji NAKANO^{a*}, Muneto NITTA^{b**} and Taeko MATSUURA^{c***}

^aGSI, Planckstr.1, D-64291 Darmstadt, Germany

^bDepartment of Physics, Keio University, Hiyoshi, Yokohama,

Kanagawa 223-8521, Japan

^cParticle Therapy Division, Research Center for Innovative Oncology,

National Cancer Center Hospital East, Japan

Different types of non-Abelian vortex-strings appear in dense or hot QCD, both of which possess non-Abelian internal orientation zero modes. We calculate the interaction between them and find the universal repulsion for dense QCD (color superconductivity) and the dependence on the orientations for hot QCD (chirally broken phase). This is a review article based on our papers.^{1),2)}

§1. Introduction and Summary

It is one of the most important subjects to determine the phase structure of QCD in study of the strong interaction. QCD is expected to exhibit color superconductivity at high baryon density region, where the color $SU(3)_{\rm C}$, the flavor $SU(3)_{\rm F}$ and the baryon $U(1)_{\rm B}$ symmetries are spontaneously broken down to the the color-flavor locked (CFL) symmetry $SU(3)_{\rm C+F}$.³⁾ According to the spontaneously broken $U(1)_{\rm B}$ there appear topological (vortex-)strings which wind around it.^{4),5)} These are global vortices⁶⁾ or superfluid vortices in a rotating media. The existence of vortices in a core of a neutron star may become a proof of realization of the color super-conductivity. It has been shown⁷⁾ that there also exist non-Abelian strings which wind around both $U(1)_{\rm B}$ and $SU(3)_{\rm C-F}$ simultaneously because of the non-trivial homotopy $\pi_1[U(3)] \simeq \mathbb{Z}^{\dagger}$.[†] The interesting is that these vortices carry a color gauge flux but behave as superfluid vortices in the energetic point of view. Moreover they have normalizable orientational zero modes in the internal space, associated with further breaking of the residual symmetry $SU(3)_{\rm C+F}$ in the presence of the strings.²

On the other hand, topological strings are also expected to form during the chiral phase transition⁸⁾ where approximate axial symmetry $U(1)_{\rm A}$ is spontaneously broken as well as chiral symmetry $SU(3)_{\rm L} \times SU(3)_{\rm R}$ broken down to the vector-like symmetry $SU(3)_{\rm L+R}$. These strings wind around the spontaneously broken $U(1)_{\rm A}$ which is explicitly broken by anomaly at low temperature but recovers at high temperature. Non-Abelian strings also exist due to $\pi_1[U(3)_{\rm A}] \simeq \mathbb{Z}^{(9),10}$ These strings also have orientational zero modes in the internal space associated with breakdown

typeset using $\mathcal{PTPTEX.cls}$ (Ver.0.9)

^{*)} e.nakano(at)gsi.de

^{**)} nitta(at)phys-h.keio.ac.jp, speaker at the conference

^{***)} tamatsuu(at)east.ncc.go.jp

^{†)} Similar but different non-Abelian vortex-strings have been extensively studied in the context of supersymmetric gauge theories and superstring theory.^{11), 12)} Apart from supersymmetry, only but crucial difference is that $U(1)_{\rm B}$ is gauged in those context.

of $SU(3)_{L+R}$ in the presence of strings. They are however non-normalizable¹⁾ unlike those appearing in the color superconductor.

Let us recall that, in the case of usual superconductors, their stability in the presence of external magnetic fields depends on whether topological (vortex-)strings attract or repel each other; the former is unstable (Type I) whereas the latter is stable (Type II). Therefore the purpose of our study is to calculate the interaction between two parallel non-Abelian strings with general relative orientational zero modes in the both cases of 1) dense and 2) hot QCD. We find followings:

1) In the case of dense QCD, i.e. in the CFL phase, we find the universal repulsion between strings which does not depend on their orientations. This shows that the superfluid $U(1)_{\rm B}$ vortices^{4),5)} are unstable to decay, and that the most fundamental objects in the color superconductor is non-Abelian strings. Moreover it implies the stability of color superconductors in the presence of external color gauge fields.

2) In the case of hot QCD i.e. in the chirally broken phase, we find that the force between strings depends on their relative orientations: when the orientations are the same, the repulsive force reaches the maximum, whereas when the relative orientation becomes the maximum, no force exists. This implies the marginal instability of $U(1)_{\rm A}$ strings.⁸⁾ For more detail see the original papers.^{1),2)}

§2. Non-Abelian Strings in Dense QCD

We start from the most general Landau-Ginzburg(LG) Lagrangian

$$\mathcal{L} = \operatorname{tr}(D\Phi)^{\dagger}(D\Phi) - m^{2}\operatorname{tr}(\Phi^{\dagger}\Phi) - \lambda_{1}(\operatorname{tr}\Phi^{\dagger}\Phi)^{2} - \lambda_{2}\operatorname{tr}\left[(\Phi^{\dagger}\Phi)^{2}\right] - \frac{1}{4}F_{ij}^{a}F^{aij}, (2\cdot1)$$

where Φ is a 3 by 3 matrix of scalar fields on which the symmetries act as

$$\Phi \to e^{i\alpha} U_{\mathcal{C}} \Phi U_{\mathcal{F}}^t, \quad e^{i\alpha} \in U(1)_{\mathcal{B}}, \quad U_{\mathcal{C}} \in SU(3)_{\mathcal{C}}, \quad U_{\mathcal{F}} \in SU(3)_{\mathcal{F}}.$$
(2.2)

In the vacuum $\langle \Phi \rangle = v\mathbf{1}$ the symmetries are spontaneously broken down to the CFL symmetry $SU(3)_{L+R}$. Hereafter we take v = 1 for simplicity by a suitable rescaling. In the polar coordinates (θ, ρ) , a single non-Abelian string is given by⁷

$$\Phi(\theta, \rho) = \operatorname{diag}(e^{i\theta} f(\rho), g(\rho), g(\rho)) \sim \operatorname{diag}(e^{i\theta}, 1, 1) \quad \text{for} \quad \rho \gg \lambda, \tag{2.3}$$

$$A_z = a(\rho) \operatorname{diag}(2, -1, -1) \sim (1/\rho) \operatorname{diag}(2, -1, -1) \quad \text{for} \quad \rho \gg \lambda,$$
 (2.4)

where f and g are profile functions of ρ behaving as $f \sim g \sim 1$ far from the core of a string $(\rho \gg \lambda)$ and $f \sim 0$ $(g \neq 0)$ near the core of a string. Here the size λ of the core is either the penetration depth or the coherent length. A numerical solution with approximation g = 1 can be found.⁷ Note that the solution carries a color flux $(2\cdot 4)$ taking a value in the Lie algebra of $SU(3)_{\rm C}$.

The solution (2.3) breaks the CFL symmetry $SU(3)_{C+F}$. Therefore acting it on the solution (2.3) we obtain a continuous family of solutions. Those solutions with different color flux one-to-one correspond to a complex projective space²⁾

$$\frac{SU(3)_{\mathrm{C+F}}}{SU(2)_{\mathrm{C+F}} \times U(1)_{\mathrm{C+F}}} \simeq \mathbf{C}P^2.$$
(2.5)




Fig. 1. Configuration of two strings with interval d = 2a in polar coordinate (ρ, θ) .

Let us calculate the force between two strings with general orientations of $\mathbb{C}P^2$. First we put one string with a fixed orientation, given by the solution $(2\cdot3)$ with $\theta \to \theta_1$ at one side of Fig. 1. The other string put at the other side of Fig. 1 should have generic orientation, which can be obtained in general by acting $SU(3)_F$ on the form of the solution $(2\cdot3)$ with $\theta \to \theta_2$. However in focusing on relative orientation between two strings, we can consider an $SU(2)_F$ subgroup (in the upper-left 2 by 2 block) without loss of generality: $\begin{pmatrix} e^{i\theta_2}f & 0\\ 0 & g \end{pmatrix} \begin{pmatrix} a^* & -b\\ b^* & a \end{pmatrix} = \begin{pmatrix} e^{i\theta_2}af & e^{i\theta_2}bf\\ -b^*g & a^*g \end{pmatrix}$. The point here is that this flavor rotated configuration reverts back to original one at long distance by a color gauge rotation $\begin{pmatrix} a^* & -be^{i\theta_2}F(\rho_2)\\ b^*e^{-i\theta_2}F(\rho_2) & a \end{pmatrix} \in SU(3)_C$, with an arbitrary regular function F with conditions F(0) = 0 and $F(\infty) = 1$, to

$$\begin{pmatrix} |a|^2 f e^{i\theta_2} + |b|^2 g e^{i\theta_2 F} & a^* b \left[-e^{i\theta_2 F} + f e^{i\theta_2} \right] \\ ab^* \left[-1 + f e^{i(1-F)\theta_2} \right] & |a|^2 g + |b|^2 f e^{i(1-F)\theta_2} \end{pmatrix} \sim \begin{pmatrix} e^{i\theta_2} & 0 \\ 0 & 1 \end{pmatrix} \text{ for } \rho_2 \gg \lambda.(2.6)$$

With this expression now the situation has become much simpler in evaluation of long range force. The interaction energy of two strings, $E = \int F$, is given by subtracting energy of two isolated string systems from that of two-body system. The Abrikosov's product ansatz with (2.6) results in the energy density F and the energy E:

$$F = \pm \frac{2}{3} \left[\frac{-a^2 + \rho^2}{a^4 + \rho^4 - 2a^2 \rho^2 \cos(2\theta)} \right], \quad E = \pm \frac{2\pi}{3} \left[-\ln 4 + \ln\{\left(a^2 + L^2\right)/a^2\} \right], (2.7)$$

with the infrared cutoff L. The upper(lower) sign denotes a string-(anti-)string configuration. This gives the force between strings:²⁾

$$f(a,L) = \mp \frac{\partial E}{2\partial a} = \pm \frac{2\pi}{3} \left(\frac{1}{a} - \frac{a}{a^2 + L^2} \right) \sim \pm \frac{2\pi}{3a}, \qquad (2.8)$$

where the most-right side denotes the large volume limit $L \to \infty$. Magnitude of the force looks shrunk by factor 1/3 from the known result of Abelian global strings.⁶⁾

§3. Non-Abelian Strings in Hot QCD

The LG Lagrangian (linear sigma model) for chiral symmetry breaking is

$$\mathcal{L} = \operatorname{tr}(\vec{\partial} \Phi^{\dagger} \vec{\partial} \Phi) - m^2 \operatorname{tr}(\Phi^{\dagger} \Phi) - \lambda_1 (\operatorname{tr} \Phi^{\dagger} \Phi)^2 - \lambda_2 \operatorname{tr}[(\Phi^{\dagger} \Phi)^2]$$
(3.1)

where Φ is a 3 by 3 matrix of scalar fields on which the symmetries act as

$$\Phi \to e^{i\alpha} U_{\rm L} \Phi U_{\rm R}^{\dagger}, \quad e^{i\alpha} \in U(1)_{\rm A}, \quad U_{\rm L} \in SU(3)_{\rm L}, \quad U_{\rm R} \in SU(3)_{\rm R}.$$
(3.2)

These symmetries are spontaneously broken to the vector-like symmetry $SU(3)_{L+R}$ by $\langle \Phi \rangle = v\mathbf{1}$. A single non-Abelian string with vanishing anomaly has the form $\Phi(\theta, \rho) = \text{diag}(e^{i\theta}f(\rho), g(\rho), g(\rho))$. A numerical solution has been obtained.¹⁰⁾ This solution has orientation $\mathbb{C}P^2$ by replacing C + F in (2.5) by L + R.

We now calculate the inter-vortex force in the situation of Fig. 1. The second string is of the form $g\Phi g^{\dagger}$ with $g = \cos\left(\frac{\alpha}{2}\right)\mathbf{1}_2 + i\vec{n}\cdot\vec{\sigma}\sin\left(\frac{\alpha}{2}\right)$ with $\vec{\sigma} = (\sigma_1, \sigma_2, \sigma_3)$ the Pauli matrices and \vec{n} a unit three vector. Defining $e^{i\beta} = n_x - in_y$, we have

$$g\Phi(\theta_2,\rho)g^{\dagger} = \begin{pmatrix} e^{i\theta_2}\cos^2\left(\frac{\alpha}{2}\right) + \sin^2\left(\frac{\alpha}{2}\right) & \frac{i}{2}(1-e^{i\theta_2})e^{i\beta}\sin\alpha\\ -\frac{i}{2}(1-e^{i\theta_2})e^{-i\beta}\sin\alpha & \cos^2\left(\frac{\alpha}{2}\right) + e^{i\theta_2}\sin^2\left(\frac{\alpha}{2}\right) \end{pmatrix}.$$
 (3.3)

By using the Abrikosov's product ansatz again, we obtain the interaction energy density \tilde{F} and integrated energy \tilde{E} : $\tilde{F} = \pm 3(1 + \cos \alpha)F/2$, $\tilde{E} = \pm 3(1 + \cos \alpha)E/2$, respectively. Then the force between two strings is given by¹

$$\tilde{f}(a,\alpha,L) = \mp \frac{\partial E}{2\partial a} = \pm (1+\cos\alpha) \left(\frac{\pi}{a} - \frac{\pi a}{a^2 + L^2}\right) \sim \pm (1+\cos\alpha)\frac{\pi}{a}, \quad (3.4)$$

where the most-right side denotes the large volume limit $L \to \infty$. Unlike the previous case (2.8), this depends on the relative orientation α . For parallel orientations ($\alpha = 0$) this reaches the maximal coinciding with the Abelian case,⁶) whereas for orthogonal orientations ($\alpha = \pi$) this vanishes.

Acknowledgements

The work of E.N. is supported by Center for Theoretical Sciences, National Taiwan University under grant NO. NSC96-2811-M-002-024. The work of M.N. is supported in part by Grant-in-Aid for Scientific Research (No. 20740141) from the Ministry of Education, Culture, Sports, Science and Technology.

References

- 1) E. Nakano, M. Nitta and T. Matsuura, arXiv:0708.4092 [hep-ph].
- 2) E. Nakano, M. Nitta and T. Matsuura, arXiv:0708.4096 [hep-ph].
- For a review see M. G. Alford, A. Schmitt, K. Rajagopal and T. Schafer, "Color superconductivity in dense quark matter," arXiv:0709.4635 [hep-ph].
- 4) M. M. Forbes and A. R. Zhitnitsky, Phys. Rev. D 65 (2002), 085009.
- 5) K. Iida and G. Baym, Phys. Rev. D 66 (2002), 014015; K. Iida, Phys. Rev. D 71 (2005), 054011.
- E. P. S. Shellard, Nucl. Phys. B 283 (1987), 624; L. Perivolaropoulos, Nucl. Phys. B 375 (1992), 665.
- 7) A. P. Balachandran, S. Digal and T. Matsuura, Phys. Rev. D 73 (2006), 074009.
- X. Zhang, T. Huang and R. H. Brandenberger, Phys. Rev. D 58 (1998), 027702; R. H. Brandenberger and X. Zhang, Phys. Rev. D 59 (1999), 081301; A. P. Balachandran and S. Digal, Int. J. Mod. Phys. A 17 (2002), 1149.
- 9) A. P. Balachandran and S. Digal, Phys. Rev. D 66 (2002), 034018.
- 10) M. Nitta and N. Shiiki, Phys. Lett. B 658 (2008), 143.
- A. Hanany and D. Tong, JHEP 0307 (2003), 037; JHEP 0404 (2004), 066; R. Auzzi,
 S. Bolognesi, J. Evslin, K. Konishi and A. Yung, Nucl. Phys. B 673 (2003), 187; M. Shifman and A. Yung, Phys. Rev. D 70 (2004), 045004; Rev. Mod. Phys. 79 (2007), 1139.
- M. Eto, Y. Isozumi, M. Nitta, K. Ohashi and N. Sakai, Phys. Rev. Lett. 96 (2006), 161601;
 J. Phys. A 39 (2006), R315; Phys. Rev. D 72 (2005), 025011; M. Eto et al., Phys. Rev. D 74 (2006), 065021; Phys. Rev. Lett. 98 (2007), 091602; arXiv:0802.1020 [hep-th].