Exact Scale Invariance in Mixing of Binary Candidates in Voting Model

Shintaro Mori*

Department of Physics, School of Science, Kitasato University, Kitasato 1-15-1, Sagamihara, Kanagawa 228-8555, Japan

Masato Hisakado[†]

Standard and Poor's,

Marunouchi 1-6-5, Chiyoda-ku,

Tokyo 100-0005, Japan

(Dated: February 6, 2020)

Abstract

We introduce a voting model and discuss the scale invariance in the mixing of candidates. The Candidates are classified into two categories $\mu \in \{0,1\}$ and are called as 'binary' candidates. There are in total $N = N_0 + N_1$ candidates, and voters vote for them one by one. The probability that a candidate gets a vote is proportional to the number of votes. The initial number of votes ('seed') of a candidate μ is set to be s_{μ} . After infinite counts of voting, the probability function of the share of votes of the candidate μ obeys gamma distributions with the shape exponent s_{μ} in the thermodynamic limit $Z_0 = N_1 s_1 + N_0 s_0 \to \infty$. Between the cumulative functions $\{x_{\mu}\}$ of binary candidates, the power-law relation $1 - x_1 \sim (1 - x_0)^{\alpha}$ with the critical exponent $\alpha = s_1/s_0$ holds in the region $1 - x_0, 1 - x_1 << 1$. In the double scaling limit $(s_1, s_0) \to (0, 0)$ and $Z_0 \to \infty$ with $s_1/s_0 = \alpha$ fixed, the relation $1 - x_1 = (1 - x_0)^{\alpha}$ holds exactly over the entire range $0 \le x_0, x_1 \le 1$. We study the data on horse races obtained from the Japan Racing Association for the period 1986 to 2006 and confirm scale invariance.

PACS numbers: 02.50.Ga,89.65.-s

^{*}Electronic address: mori@sci.kitasato-u.ac.jp

[†]Electronic address: masato hisakado@standardandpoors.com

I. INTRODUCTION

Scale-invariant behaviour has attracted considerable attention on account of its ubiquity in natural and man-made phenomena [1]. Many possible candidate mechanisms that gives rise to power-law distributions have been proposed thus far. The Yule process is a widely applicable mechanism for generating power-law distributions [2]. Originally, it has been proposed to explain why the distribution of the number of species in a genus, a family, or any other taxonomic group follows a power law [3]. Now, it has found wide applications in other areas [1, 4].

Consider the distribution of the number of species in a genus. Suppose first that new species appear but they never die; species are only ever added to genera and never removed. Species are added to genera by speciation, the splitting of one species into two. If we assume that this happens at some stochastically constant rate, then it follows that a genus with kspecies will gain new species at a rate proportional to k, since each of the k species has the same chance per unit time of dividing into two. In addition, suppose that a new species that belongs to a new genus is added once every m speciation events. So m+1 new species appear for each new genus and there are m+1 species per genus. Thus the number of general goes up steadily as does the number of species within each genus. We denote the fraction of genera that has k species by $p_{k,n}$, where n denotes the total number of genera and n measures the passage of time in the model. At each time-step one new species founds a new genus, thereby increasing n by 1, and m other species are added to various pre-existing general which are selected in proportion to the number of species they already have. By solving the master equation for $p_{k,n}$ in the limit $n \to \infty$, $p_k \equiv \lim_{n \to \infty} p_{k,n}$ behaves as $p_k \sim k^{2+\frac{1}{m}}$. The Yule process has been adopted and generalized to explain power laws in many other systems. An important feature of this process is that the probability that a genus with kspecies will gain new species is proportional to k. This 'rich-get-richer' process is the most important factor in exhibiting power-law behaviour. The feature that n increases infinitely is also important in generating power-law behaviour.

In this study, we introduce a voting model and discuss the scale invariance in the mixing of candidates. The candidates are classified into two categories $\mu \in \{0,1\}$ and are called as 'binary' candidates. The probability that a candidate get a vote is proportional to the number of votes, which is the same as the relation in the Yule process. The main difference

between our model and the Yule process is that the number of candidates is fixed in our model. In the Yule process, n increases and in the limit $n \to \infty$, power-law behaviour is observed. In our model, the distribution of the number of votes does not show power-law behaviour. However, our model exhibits scale-invariant behaviour. This behaviour is observed in the mixing of the binary candidates. Furthermore, the power law holds over the entire range in a double scaling limit.

The organization of this paper is as follows. In section II, we introduce the voting model. We select a candidate (initial number of votes s_{μ}) and show that the probability density function of the share of votes u, of the candidate obeys a gamma distribution function with the shape exponent s_{μ} in the thermodynamic limit $Z_0 = N_1 s_1 + N_0 s_0 \to \infty$. We also show that the joint probability density function of u for any k candidates is given by the direct product of the gamma distributions in the same limit. We discuss the scale invariance in the mixing of the binary candidates in section III. The cumulative function $1-x_{\mu}$ of candidates μ is given by the incomplete gamma function. The power-law relation $1-x_1 \sim (1-x_0)^{\alpha}$ with the exponent $\alpha = s_1/s_0$ holds in the region $1 - x_0, 1 - x_1 << 1$. Furthermore, in the double scaling limit $\{s_{\mu}\}\to 0$ and $Z_0\to\infty$ with $\alpha=s_1/s_0$ fixed, the relation $1-x_1=(1-x_0)^{\alpha}$ holds exactly over the entire range $0 \le x_0, x_1 \le 1$. Using the data on horse races, we verify these results in section IV. We show that scale invariance holds over the wide range of cumulative functions. In addition, we show that the probability distribution functions of u are well described by gamma distributions. Section V is dedicated to the summary and concluding remarks. Appendix A is devoted to the derivation of the joint probability distribution function of u for any k candidates. In Appendix B, we map the voting model to a branching process and easily derive the gamma distribution function.

II. VOTING MODEL FOR BINARY CANDIDATES

Consider a voting model for N candidates. Voters vote for them one by one, and the result of each voting is announced promptly. The time variable $t \in \{0, 1, 2 \cdots, T\}$ counts the number of the votes. The candidates are classified into two categories $\mu \in \{0, 1\}$ and are called as binary candidates. There are N_{μ} candidates in each category and $N_0 + N_1 = N$. The main result of this section is that the scaled share of votes u_i^{μ} of a candidate μ obeys a gamma distribution with the shape exponent s_{μ} in the thermodynamic limit $N_0, N_1 \to \infty$.

We denote the number of votes of *i*th candidate $\mu \in \{0, 1\}$ at time t as $\{X_{i,t}^{\mu}\}_{i \in \{1, \dots, N_{\mu}\}}$. At t = 0, $X_{i,t}^{\mu}$ takes the initial value $X_{i,0}^{\mu} = s_{\mu} > 0$. If the *i*th candidate μ gets a vote at t, $X_{i,t}^{\mu}$ increases by one unit.

$$X_{i,t+1}^{\mu} = X_{i,t}^{\mu} + 1.$$

A voter casts a vote for the total N candidates at a rate proportional to $X_{i,t}^{\mu}$. The probability $P_{i,t}^{\mu}$ that the ith candidate μ gets a vote at t is

$$P_{i,t}^{\mu} = \frac{X_{i,t}^{\mu}}{Z_t} \tag{1}$$

$$Z_t = \sum_{\mu=0}^{1} \sum_{i=1}^{N_{\mu}} X_{i,t}^{\mu} = N_1 s_1 + N_0 s_0 + t.$$
 (2)

The problem of determining the probability of the *i*th candidate μ getting n votes up to T is equivalent to the famous Pólya's urn problem [5, 6, 7]. If the change in $X_{i,t}^{\mu}$ is given by

$$\Delta X_{i,t}^{\mu} \equiv X_{i,t}^{\mu} - X_{i,t-1}^{\mu},$$

the sequence $(\Delta X_{i,1}^{\mu}, \cdots, \Delta X_{i,T}^{\mu})$ is called Pólya's urn sequence. This sequence is an exchangeable stochastic process, and the joint distribution of $(X_{i,1}^{\mu}, \cdots, X_{i,T}^{\mu})$ is given by

$$\operatorname{Prob}(\Delta X_{i,1}^{\mu} = x_1, \cdots, \Delta X_{i,T}^{\mu} = x_T) = \frac{(s_{\mu})_k (Z_0 - s_{\mu})_{T-k}}{(Z_0)_T}.$$

Here, $k = \sum_{t=1}^{T} x_t$ and $(a)_n \equiv a \cdot (a+1) \cdot (a+2) \cdot \cdot \cdot (a+n-1)$ is the rising factorial. This distribution depends only on k, and not on the particular order of (x_1, \dots, x_T) . This distribution is invariant under the permutations of the entries and, hence, it is called exchangeable.

Furthermore, the expectation value of $\Delta X_{i,t}^{\mu}$, denoted by p_{μ} , does not depend on t.

$$p_{\mu} \equiv <\Delta X_{i,t}^{\mu} > = \frac{s_{\mu}}{Z_0}.\tag{3}$$

The correlation function ρ_{μ} between $\Delta X_{i,t}^{\mu}$ and $\Delta X_{i,t'}^{\mu}$ $(t' \neq t)$ is also constant [7] as ρ_{μ} .

$$\rho_{\mu} \equiv \text{Corr}(\Delta X_{i,t}^{\mu}, \Delta X_{i,t'}^{\mu}) \equiv \frac{\langle \Delta X_{i,t}^{\mu} \Delta X_{i,t'}^{\mu} \rangle - p^2}{p(1-p)} = \frac{1}{Z_0 + 1}, \quad t \neq t'.$$
 (4)

The probability that the ith candidate μ gets n votes up to T is given by the beta binomial distribution

$$\operatorname{Prob}(X_{i,T}^{\mu} - s_{\mu} = n) = {}_{T}C_{n} \cdot \frac{(s_{\mu})_{n}(Z_{0} - s_{\mu})_{T-n}}{(Z_{0})_{T}}.$$
(5)

 $(a)_n$ is written as $(a)_n = \frac{\Gamma(a+n)}{\Gamma(a)}$ and this relation can also be written as

$$Prob(X_{i,T}^{\mu} - s_{\mu} = n) = {}_{T}C_{n} \cdot \frac{\Gamma(s_{\mu} + n)}{\Gamma(s_{\mu})} \frac{\Gamma(Z_{0} - s_{\mu} + T - n)}{\Gamma(Z_{0} - s_{\mu})} \frac{\Gamma(Z_{0})}{\Gamma(Z_{0} + T)}.$$
 (6)

Using a definition of beta function $B(a,b) \equiv \frac{\Gamma(a)\Gamma(b)}{\Gamma(a+b)}$, we can rewrite the expression as

$$Prob(X_{i,T}^{\mu} - s_{\mu} = n) = {}_{T}C_{n} \cdot \frac{B(s_{\mu} + n, Z_{0} - s_{\mu} + T - n)}{B(s_{\mu}, Z_{0} - s_{\mu})}.$$
 (7)

B(a,b) is also written as $B(a,b) = \int_0^1 p^{a-1} (1-p)^{b-1} dp$, we get the next expression

$$\operatorname{Prob}(X_{i,T}^{\mu} - s_{\mu} = n) = {}_{T}C_{n} \cdot \int_{0}^{1} p^{n} (1 - p)^{T - n} \frac{p^{s_{\mu} - 1} (1 - p)^{Z_{0} - s_{\mu}}}{B(s_{\mu}, Z_{0} - s_{\mu})} dp. \tag{8}$$

After infinite counts of voting, i.e. $T \to \infty$, the share of votes $x_i^{\mu} \equiv \lim_{T \to \infty} \frac{X_{i,T}^{\mu} - s_{\mu}}{T}$, becomes the beta distributed random variable beta $(s_{\mu}, Z_0 - s_{\mu})$ on [0, 1].

$$p(x) \equiv \lim_{T \to \infty} \text{Prob}(X_{i,T}^{\mu} - s_{\mu} = Tx) \cdot T = \frac{x^{s_{\mu} - 1} (1 - x)^{Z_0 - s_{\mu} - 1}}{B(s_{\mu}, Z_0 - s_{\mu})}.$$
 (9)

Here, we use the identity $\lim_{T\to\infty} {}_T C_{Tx} p^{Tx} (1-p)^{T(1-x)} \cdot T = \delta(x-p)$. This result has been derived by Pólya [5].

Next, we focus on the thermodynamic limit $N_0, N_1 \to \infty$ and $Z_0 = N_0 s_0 + N_1 s_1 \to \infty$. The expectation value of x_i^{μ} is $\langle x_i^{\mu} \rangle = p_{\mu} = \frac{s_{\mu}}{Z_0}$. We introduce a variable $u_i^{\mu} \equiv (Z_0 - s_{\mu} - 1)x_i^{\mu}$. The distribution function $p_{s_{\mu}}(u)$ in the thermodynamic limit is given as

$$p_{s_{\mu}}(u) \equiv \lim_{Z_0 \to \infty} p(x_i^{\mu} = \frac{u}{Z_0 - s_{\mu} - 1}) = \frac{1}{\Gamma(s_{\mu})} e^{-u} u^{s_{\mu} - 1}.$$
 (10)

The share of votes, u, of a candidate μ obeys a gamma distribution function with s_{μ} .

In general, the joint probability distribution function of the scaled share of votes of k different candidates becomes the direct product of k gamma distribution functions in the limit $Z_0 \to \infty$. We denote the k candidates as $\{(\mu_j, i_j)\}_{j=1,\dots,k}$ and denote the scaled share of votes as $\{u_j\}_{j=1,\dots,k}$. The joint probability distribution function is given as

$$p(u_1, \dots, u_k) = \prod_{j=1}^k p_{s_{\mu_j}}(u_j).$$
 (11)

The derivation of the result is given in Appendix A. It should be noted that in the thermodynamic limit, the correlation among $\{u_j\}_{j=1,\dots,k}$ vanishes. Hence, by mapping the voting problem to a continuous time branching process, we can derive the gamma distribution function $p_{s_{\mu}}(u)$ easily (refer Appendix B). In the branching process, the stochastic processes of the increase in $\{X_{i,t}^{\mu}\}$ are independent of each other.

III. SCALE INVARIANCE IN MIXING OF BINARY CANDIDATES

In this section, we discuss the mixing of the binary candidates. After many counts of voting $T \to \infty$, the binary candidates are distributed in the space of u according to the gamma distribution in the thermodynamic limit $Z_0 \to \infty$. If $s_1 > s_0$, a candidate belonging to category $\mu = 1$ has a higher probability of getting many votes than a candidate belonging to category $\mu = 0$. Even the latter can obtain many votes. It is also possible that the former may get few votes. Thus, there is a mixing of the binary candidates. We see a scale invariant behaviour appears in the mixing. Between the cumulative functions of the binary candidates $1 - x_{\mu}$, the power-law relation $1 - x_1 \sim (1 - x_0)^{\alpha}$ with the exponent $\alpha = s_1/s_0$ holds.

In order to study the mixing configuration, we arrange the N candidates according to the size of u_i^{μ} as

$$u_{i_1}^{\mu_1} > u_{i_2}^{\mu_2} > \dots > u_{i_N}^{\mu_N}, \quad \mu_k \in \{0, 1\}.$$
 (12)

Using the ranking information $\{\mu_k\}_{k=1,\dots,N}$, we draw a path $\{(x_{0,k},x_{1,k})\}_{k=0,\dots,N}$ in two-dimensional space (x_0,x_1) from $(x_{0,0},x_{1,0})=(0,0)$ to $(x_{0,N},x_{1,N})=(1,1)$ as

$$x_{\mu,k} = \frac{1}{N_{\mu}} \sum_{j=1}^{k} \delta_{\mu_j,\mu}.$$
 (13)

If $\mu_k = \mu$, the path extends in x_μ direction. The pictorial representation of the mixing of binary objects is known as a receiver operating characteristic (ROC) curve [8]. If $s_1 >> s_0$, the binary candidates are well separated on the axis of u, and the first N_1 candidates belong to category $\mu = 1$ and the last N_0 candidates belong to category $\mu = 0$. The path goes straight from (0,0) to (0,1) and then turns right to the end point (1,1). If $s_1 = s_0$, the path almost runs diagonally to the end point. If $s_1 > s_0$ holds, the path resembles a upward convex curve from (0,0) to (1,1).

The distribution function of the candidate μ on the axis of u is given by the gamma distribution with the shape exponent s_{μ} . The ROC curve $(x_0(t), x_1(t))$ of the parameter $t \in [0, \infty]$ is given by its cumulative function as

$$x_{\mu}(t) = \int_{1}^{\infty} p_{s_{\mu}}(u) du. \tag{14}$$

Using the incomplete gamma function of the first kind $\gamma(s,t) \equiv \int_0^t e^{-u} \cdot u^{s-1} du$ [9], the ROC




FIG. 1: ROC curve of mixing configuration. \bigcirc represents candidate belonging to category $\mu = 1$. \times represents candidate belonging to category $\mu = 0$. At the top of the figure, the order of three candidates from category $\mu = 1$ and five candidates from category $\mu = 0$ is shown.

curve is given as

$$1 - x_{\mu}(t) = \frac{1}{\Gamma(s_{\mu})} \cdot \gamma(s_{\mu}, t). \tag{15}$$

Near the end point, $(x_0, x_1) \simeq (1, 1)$, in other words, in the small u region $(t \simeq 0)$, the incomplete gamma function $\gamma(s_{\mu}, t)$ behaves as

$$\gamma(s_{\mu}, t) \sim t^{s_{\mu}}.\tag{16}$$

As $1 - x_{s_{\mu}}(t) \propto t^{s_{\mu}}$, the following relation holds:

$$1 - x_1 \sim (1 - x_0)^{\alpha}$$
 with $\alpha = \frac{s_1}{s_0}$. (17)

The density of good candidates, ρ_1 , in terms of the cumulative function of bad candidates, $1-x_0$, is given as

$$\rho_1 = \frac{d(1-x_1)}{d(1-x_0)} \propto (1-x_0)^{\alpha-1}.$$
(18)

 ρ_1 obeys the power law with the exponent $\alpha - 1$.

Furthermore, in the limit $(s_1, s_0) \to (0, 0)$ with $\alpha = s_1/s_0$ fixed, the relation $1 - x_1 = (1 - x_0)^{\alpha}$ holds. The proof is given as follows. $\gamma(s, t)$ is expressed using Kummer's confluent

hypergeometric function M(a, b, t) [9] as

$$\gamma(s,t) = \frac{1}{s}t^s \cdot M(s,s+1,-t). \tag{19}$$

The cumulative function $1 - x_{\mu}(t)$ is then given as

$$1 - x_{\mu}(t) = \frac{t^{s_{\mu}}}{\Gamma(s_{\mu} + 1)} \cdot M(s_{\mu}, s_{\mu} + 1, -t). \tag{20}$$

Thus, we obtain

$$(1 - x_0)^{\alpha} = (1 - x_1) \frac{\Gamma(s_1 + 1)}{M(s_1, s_1 + 1, -t)} \left(\frac{M(s_0, s_0 + 1, -t)}{\Gamma(s_0 + 1)} \right)^{\alpha}.$$
 (21)

In the limit $s_{\mu} \to 0$, both $\Gamma(s_{\mu}+1)$ and $M(s_{\mu}, s_{\mu}+1, -t)$ become equal to 1 and the following relation holds.

$$1 - x_1 = (1 - x_0)^{\alpha}, \ 0 \le x_0, x_1 \le 1.$$
 (22)

Thus, the scale-invariant relation holds over the entire range $0 \le x_0, x_1 \le 1$. The feature is remarkable from the viewpoint of statistical physics. Usually, the power-law relation does hold only in the tail.

The relative probability that a candidate gets the first vote (t = 0) is given by s_{μ} . If the candidate get the first vote, his/her score increases by 1 and the relative probability becomes $s_{\mu} + 1$. In the limit $s_{\mu} \to 0$, the additional score +1 or the weight of a single vote becomes crucially important. The probability that the candidate gets the next vote becomes equal to 1, which is exemplified by the behaviour of ρ_{μ} , given by (4).

$$\rho_{\mu} = \frac{1}{Z_0 + 1} = \frac{1}{N_0 s_0 + N_1 s_1 + 1} \to 1 \quad \text{if } \{s_{\mu}\} \to 0.$$
 (23)

After infinite counts of voting, the candidate occupies the first position in the order of candidates according to the number of votes. Then, we neglect this candidates in the voting problem and consider the remaining N-1 candidates. Similarly, if a candidate is selected randomly with the relative probability s_{μ} , he/she occupies the second position. Thus, the voting problem reduces to a random choice problem with the relative probability s_{μ} in the limit $\{s_{\mu}\} \to 0$. At (x_0, x_1) on the ROC curve, the probability that the next candidate belongs category μ is proportional to $(1-x_{\mu})s_{\mu}$. The coordinates of the ROC curve (x_0, x_1) grow according to the following relation:

$$dx_{\mu} \propto (1 - x_{\mu}) \cdot s_{\mu}.$$

Solving this relation, we get (22).

Finally, we discuss the limit in the derivation of the exact scale invariance. In the derivation of the gamma distribution, we take the thermodynamic limit $Z_0 = N_1 s_1 + N_0 s_0 \to \infty$. With the gamma distribution, (22) holds in the limit $\{s_{\mu}\} \to 0$. In order that (22) holds, these two limits, $Z_0 \to \infty$ and $\{s_{\mu}\} \to 0$, should go together. $\{s_{\mu}\}$ approaches zero more slowly than $\{N_{\mu}\}$ approaches infinity. In other words, in the double scaling limit $Z_0 \to \infty$ and $\{s_{\mu}\} \to 0$ with $\alpha = s_1/s_0$ fixed, (22) holds. So the above intuitive explanation of the exact scale invariance may be too naive. If we take the limit $\{s_{\mu}\} \to 0$ without the limit $Z_0 \to \infty$, ρ_{μ} becomes 1. The firstly chosen candidate get all the remaining votes and there does not occur the mixing of the binary candidates. The double scaling limit is crucial in the emergence of the exact scale invariance.

IV. DATA ANALYSIS OF HORSE RACES

We verify the results of the voting model, particularly the scale invariance in the mixing of binary candidates. We study all the data on horse race betting obtained from the Japan Racing Association (JRA) for the period 1986 to 2006. There have been 71549 races and in which a total of 901366 horses have participated. We select the winning horses as candidate belonging to category $\mu = 1$. For candidate belonging to category $\mu = 0$, we consider two cases; losing horses and horses finishing second. In a race, no one knows which horse will win. Betters only have partial information on the horses, which is embedded in the initial values $\{s_{\mu}\}$. The results of betting are announced at short intervals. Betters usually presume that the horses which get many votes are strong. They come to know which horses are considered to be strong by other betters. These features are incorporated in the voting model. Betters do not always bet to strong horses. Some betters may prefer betting to a horse that can coin more money even if it is considered to be 'weaker' than a horse that can coin less money. However, in the bet to win, only the better who bets to the winning horse coin the bet. Hence, the assumption is not so unrealistic. We also note the reason why we can treat multiple categories, 2nd finishing horse and losing horse, as the category $\mu = 0$. For the betters, the only difference between the losing horses and finishing second ones is their confidence. By tuning parameter s_0 , we can treat the two categories on the same footing.

Next, we explain the meaning of the initial values $\{s_{\mu}\}$. The probability that a candidate μ is selected is proportional to s_{μ} as $<\Delta X_{i,t}^{\mu}>=s_{\mu}/Z_{0}$. The ratio s_{1}/s_{0} is a measure of the accuracy of the knowledge of betters. On the other hand, ρ_{μ} is given by (4). If the scale of $\{s_{\mu}\}$ is small, the decisions of betters are crucially affected by the choices of other betters. In the limit $\{s_{\mu}\}\to\infty$, their decisions are not affected by the choices of other betters. The scale of $\{s_{\mu}\}$ is a measure of the degree of similarity ('copycat') of the choices of betters.

In the early stage of voting, $\{s_{\mu}\}$ is the only available information. Voters decide on horses on the basis of $\{s_{\mu}\}$ and they are 'intelligent', because their decisions are not affected by the choices of other betters. As the voting process proceeds, the importance of the cumulative number of votes exceeds that of the initial scores, and voters become 'copycat'. If one control the scale of $\{s_{\mu}\}$ (or the weight of a single vote), the passage timing from the initial 'intelligent' stage to the late 'copycat' stage should change.

TABLE I: Data on horse race betting obtained from the Japan Racing Association (JRA) for the period 1986 to 2006. There are 71549 races and 71650 winning (finishing first) horses. 71590 horses are finishing second. The difference between N^{Win} and N^{2nd} indicates that there occurs a tie in the race. In the third column, we show the average value of the share of the votes in each category. The fourth column shows the values v^{ν}/c , here c is the estimated value of the scale parameter in (24). About the estimation of c, please the main text and Figure 2.

Category ν	$N^{ u}$	v^{ν} [%]	v^{ν}/c
Win	71650	21.23	1.769
2nd	71590	15.40	1.283
Lose	829716	6.80	0.567

We denote the three categories of horses as $\nu \in \{\text{Win, 2nd, Lose}\}$ and the number of horses in each category as N^{ν} . v_i^{ν} denotes the share of votes of the *i*th horse in the category ν , and v^{ν} denotes the average value of v_i^{ν} . In Table I, we summarize the data on horse races. A difference between N^{Win} and N^{2nd} indicates that there is a tie in the race.

We have shown that the share of votes, u, obeys a gamma distribution function with s_{μ} . In order to check whether v_i^{ν} obeys a gamma distribution function, we have to set the scale c between v_i^{ν} and u as follows:

$$v_i^{\nu} = c \cdot u. \tag{24}$$




FIG. 2: Logarithmic plot of probability distribution functions of shares of votes. The curves from the top to bottom indicate the data for ν =Win, 2nd and Lose. The gamma distribution functions with c = 0.12 and s_{ν} are also plotted.

The same c should be used for all categories. Assuming that u obeys the gamma probability distribution with s_{ν} , v_{i}^{ν} obeys the following probability distribution function:

$$p(v_i^{\nu} = v) = p_{s_{\nu}}(v) = \frac{1}{c \cdot \Gamma(s_{\nu})} (\frac{v}{c})^{s_{\nu}-1} \exp(-\frac{v}{c}).$$

The expectation value of v_i^{ν} is

$$< v_i^{\nu} >_{\mu} = \int_0^{\infty} p_{s_{\nu}}(v) v dv = c \cdot s_{\nu}.$$

If we set c, it is possible to estimate s_{ν} of the horses in category ν as $s_{\nu} = v^{\nu}/c$.

Figure 2 shows the probability distribution functions of v_i^{μ} . In the same figure, we show the result of fitting with the gamma probability functions. Using the least square method in the range $v \in [0.01, 1.0]$, we set c = 0.12 and $s_{Win} = 1.659$, $s_{2nd} = 1.258$ and $s_{Lose} = 0.529$. Comparing with the values in the fourth column in Table I, it is observed the values of s_{ν} and v^{ν}/c are close to each other in all categories, implying that the bulk shapes of the probability functions of v_i^{ν} are well described by the gamma distributions.

We study the mixing properties of the binary horses by employing the method explained in the text. We adopt the Win-Lose pair and Win-2nd pair as the binary pairs. Figure




FIG. 3: Double-logarithmic plot of ROC curves $(1 - x_0, 1 - x_1)$. The curves of the Win-Lose pair (solid line) and the Win-2nd pair (dashed line) are plotted. The fitting curves given by $1 - x_1 = a \cdot (1 - x_0)^{\alpha}$ (dash-dotted line) are also plotted.

TABLE II: The initial value s_{μ} in each category and the critical exponent α are plotted. In the last column, we show the predicted values of α by the voting model.

Pair	s_1	s_0	α	s_1/s_0
Win vs Lose	1.659	0.529	1.81	3.134
Win vs 2nd	1.659	1.258	1.12	1.318

3 shows the double-logarithmic plot of the ROC curve $(1 - x_0, 1 - x_1)$ for the two pairs. The plots show scale-invariant behaviour over the wide range of $1 - x_1$. In the case of the Win-2nd pair, scale invariance holds over the range $10^{-5} < 1 - x_1 < 10^{-1}$. Using the least square method in the range $0 \le 1 - x_0 \le 0.1$, we estimate the critical exponent α . The values of the parameters and other data are summarized in Table II. The estimated values of α are considerably different from those predicted by the model; i.e. s_1/s_0 .

V. CONCLUDING REMARKS

In this study, we have introduced a simple voting model in order to discuss the mixing of binary candidates with initial number of votes s_0 and s_1 . As the voting process proceeds, the candidates are mixed in the space of the share of votes, u. We have shown that the probability distribution of u of a candidate μ obeys a gamma distribution function with the shape exponent s_{μ} in the thermodynamic limit $Z_0 \to 0$. The joint probability distribution of k different candidates is given as the direct product of the gamma distributions. The mixing configuration of the binary candidates exhibits scale invariance in the small u region. In particular, in the double scaling limit $Z_0 \to \infty$ and $\{s_{\mu}\} \to 0$ with $\alpha = s_1/s_0$ fixed, the scale invariance holds over the entire range. The cumulative function of the binary candidates obeys $1 - x_1 = (1 - x_0)^{\alpha}$ for $0 \le x_0, x_1 \le 1$.




FIG. 4: Voting model and Random Young diagram model. As the voting process proceeds, the order of the binary candidates and the Young diagram change. The complementary space of the ROC curve corresponds the Young diagram.

The data on horse races obtained from JRA also show that scale invariance holds over the wide range of cumulative functions. The distribution functions of the share of votes, u, are well described by the gamma distribution functions, implying that the behaviour of betters is well described by the voting model. However a clear discrepancy is observed in the critical behaviour. Although our voting model describes the mechanism of scale invariance in the mixing of binary candidates, it may be too simple to describe the behaviour of betters in real cases. Thus far, dividends have been reported to exhibit power-law behaviour. Another betting model has been proposed in [10]. A detailed study of real data, in particular the time

series of the number of votes, should clarify the mechanism of scale invariance in betting systems.

We also note that our model is related to the random Young diagram problem [11]. This problem pertains to the probabilistic growth of a Young diagram. A parabolic shape [12] and a quadrant shape [13] have been obtained for the asymptotic shape. The complementary part of the ROC curve, which is embedded in the fourth quadrant, corresponds the Young diagram. In our model, the ROC curve $(x_0(t), x_1(t))$ given by (15) describes the asymptotic shape of the Young diagram. In particular, it is described by the relation $1 - x_1 = (1 - x_0)^{\alpha}$ in the double scaling limit. Figure 4 shows the correspondence between the voting model and the random Young diagram problem. As the voting process proceeds, the order of the binary candidates and the Young diagram change.




FIG. 5: Triple-logarithmic plot of ROC curve $(1-x_{1st}, 1-x_{2nd}, 1-x_{3rd})$. x_{ν} denotes the cumulative function of the horses finishing 1st, 2nd and 3rd.

It is also possible to study the voting model with many categories of candidates with the usage of many different initial values $\{s_{\mu}\}$. u of the candidates in each category becomes a gamma distributed random variable. Scale invariance does hold between any pair of categories. Figure 5 shows the triple-logarithmic plot of the cumulative functions of the winning (x_{1st}) , finishing second (x_{2nd}) and finishing third (x_{3rd}) horses. In the linear part of the curve, scale invariance holds between any pair of categories.

Acknowledgments

We thank Dr. Emmanuel Guitter for his useful discussions on the branching process. We also thank Takumi Nakaso for helping us with the data analysis of horse races.

- [1] M.E.J.Newman, Contem. Phys. 46,323(2005).
- [2] G.U. Yule, Philos. Trans. R.Soc.London B213,21(1925).
- [3] J.C.Willis and G.U.Yule, Nature **109**,177(1922).
- [4] A.-L.Barabasi and R. Albert, Science **286**,509 (1999).
- [5] G. Pólya, Sur quelques points de la théorie des probabilités, Ann. Inst. Henri Poincaré 1,117(1931).
- [6] T. Hiillet, J.Phys.A41,505005(2008).
- [7] M. Hisakado, K. Kitsukawa and S. Mori, J.Phys.A39,15365(2006).
- [8] B.Enleman, E.Hayden and D.Tasche, Testing rating accuracy, www.risk.net (2003).
- [9] M.Abramowitz and I.A. Stegun, eds. *Handbook of Mathematical Functions with Formulas, Graphs and Mathematical Tables*, Dover, New York(1972).
- [10] K.Park and E.Dommany, Europhys.Lett.**53**,419(2001).
- [11] G.W.Andrews and Kimmo Eriksson, *Integer Partitions*, Cambridge University Press, Cambridge (2004).
- [12] J. Rost, Z.Wahrsch. Verw. Gebiete 58,41(1981).
- [13] W. Jockusch, J.Propp and P.Shor, Random domino titling and the arctic circle theorem, available at http://www.math.wisc.edu/~propp/articles.html.
- [14] P. Haccou, P.Jagers and V.A.Vatutin *Branching Processes*, Cambridge University Press, Cambridge (2005).
- [15] F.David, P. Di Francesco, E.Guitter and T.Jonsson, J.Stat.Mech., 02011(2007).

APPENDIX A: JOINT PROBABILITY DISTRIBUTION FUNCTION

We start from the expression of the joint probability function given by

$$\operatorname{Prob}(\{X_{i_j,T}^{\mu_j} - s_{\mu_j} = n_j\}_{j=1,\dots,k}) = \frac{T!}{(Z_0)_T} \prod_{i=1}^{k+1} \left[\frac{(s_{\mu_j})_{n_j}}{n_j!} \right]. \tag{A1}$$

Here, $s_{\mu_{k+1}} \equiv Z_0 - \sum_{j=1}^k s_{\mu_j}$ and $n_{k+1} \equiv T - \sum_{j=1}^k n_j$. Using the Dirichlet distribution function, we can rewrite the expression as

$$\operatorname{Prob}(\{X_{i_{j},T}^{\mu_{j}} - s_{\mu_{j}} = n_{j}\}_{j=1,\dots,k}) = \frac{T!}{\prod_{i=1}^{k+1} n_{i}!} \prod_{i=1}^{k} \left[\int_{0}^{1 - \sum_{j=1}^{i-1} p_{j}} dp_{i} \right] \prod_{i=1}^{k+1} \left[\frac{p^{n_{i} + s_{\mu_{i}} - 1}}{\Gamma(s_{\mu_{i}})} \right] \Gamma(Z_{0}). \tag{A2}$$

The expectation value of $\Delta X_{i_j,t}^{\mu_j} = X_{i_j,t+1}^{\mu_j} - X_{i_j,t}^{\mu_j}$ is given by

$$p_{\mu_j} = <\Delta X_{i_j,t}^{\mu_j}> = \frac{s_{\mu_j}}{Z_0}.$$
 (A3)

The correlation between $\Delta X_{i_j,t}^{\mu_j}$ and $\Delta X_{i_k,t}^{\mu_k}$ $(k \neq j)$ is given as

$$\rho_{\mu_j,\mu_k} = -\sqrt{\frac{s_{\mu_j}s_{\mu_k}}{(1 - \frac{s_{\mu_j}}{Z_0})(1 - \frac{s_{\mu_k}}{Z_0})}} \frac{1}{Z_0(1 + Z_0)}.$$
(A4)

By changing the integral variables from $\{p_i\}_{i=1,\dots,k}$ to $\{h_i\}_{i=1,\dots,k}$ as $p_i = (1-\sum_{j=1}^{i-1}p_j)h_i = \prod_{j=1}^{i-1}(1-h_j)h_i$, we obtain

$$\operatorname{Prob}(\{X_{i_{j},T}^{\mu_{j}} - s_{\mu_{j}} = n_{j}\}_{j=1,\dots,k}) = \frac{\Gamma(Z_{0})}{\Gamma(s_{\mu_{k+1}})} \prod \left[\frac{1}{\Gamma(s_{\mu_{i}})} \left({}_{T-\sum_{j=1}^{i-1} n_{j}} C_{n_{i}} \cdot \int_{0}^{1} dh_{i} h_{i}^{n_{i}+s_{\mu_{i}}-1} (1-h)^{T-\sum_{j=1}^{i} n_{j}+Z_{0}-\sum_{j=1}^{i} s_{\mu_{j}}-1} \right) \right]. \tag{A5}$$

We focus on the share of votes of candidates in the limit $T \to \infty$. We introduce y_i as $n_i = (T - \sum_{j=1}^{i-1} n_j)y_i = T \prod_{j=1}^{i-1} (1 - y_j)y_i$ and define the joint distribution function as

$$P(\{y_j\}_{j=1,\cdots k}) \equiv \lim_{T \to \infty} \text{Prob.}(\{X_{i_j,T}^{\mu_j} - s_{\mu_j} = T \prod_{l=1}^{j-1} (1 - y_l) y_j\}_{j=1\cdots k}) \cdot \prod_{i=1}^k (T - \sum_{j=1}^{i-1} n_j). \quad (A6)$$

The joint function $P(\{y_j\}_{j=1,\dots,k})$ is given by

$$P(\{y_j\}_{j=1,\dots,k}) = \frac{\Gamma(Z_0)}{\Gamma(s_{\mu_{k+1}})} \prod_{i=1}^k \left[\frac{1}{\Gamma(s_{\mu_i})} y_i^{s_{\mu_i}-1} (1-y_i)^{Z_0 - \sum_{j=1}^i s_{\mu_j} - 1} \right]. \tag{A7}$$

We introduce the variable x_i as $x_i = (1 - \sum_{j=1}^{i-1} x_j)y_i$, which is related to n_i as $n_i = T \cdot x_i$. The joint probability function $P(\{x_j\}_{j=1,\dots,k})$ is then given as

$$P(\lbrace x_j \rbrace_{j=1,\dots,k}) = \frac{\Gamma(Z_0)}{\Gamma(s_{\mu_{k+1}})} \prod_{i=1}^k \left[\frac{1}{\Gamma(s_{\mu_i})} x_i^{s_{\mu_i}-1} \right] (1 - \sum_{j=1}^k x_j)^{s_{\mu_{k+1}}-1}.$$
 (A8)

Finally, we introduce the variable $\{u_i\}$ as $u_i \equiv (s_{\mu_{k+1}} - 1)x_i$. In the thermodynamic limit $Z_0, s_{\mu_{k+1}} \to \infty$, we obtain

$$P(\{u_j\}_{j=1,\dots,k}) = \prod_{j=1}^k \left[\frac{e^{-u_j}}{\Gamma(s_{\mu_j})} u_j^{s_{\mu_j}-1} \right] = \prod_{j=1}^k p_{s_{\mu_j}}(u_j).$$
 (A9)

APPENDIX B: CONTINUOUS TIME BRANCHING PROCESS

We translate the discrete time voting problem $\{X_{i,t}^{\mu}\}_{i=1,\dots,N_{\mu}}$ to a continuous time branching process $\{X_{i}^{\mu}(t)\}_{i=1,\dots,N_{\mu}}$ [14], because the latter is more tractable than the former [15]. Figure 6 shows the mapping process. Let $X_{i}^{\mu}(t)$ denote the number of offspring of s_{μ} individuals. Each individual is substituted by two offspring at its death (branching) and the probability that an individual dies during time dt is given by dt. The number of offspring of each individual is denoted as $\{x_{i,k}^{\mu}(t)\}_{k=1,\dots,s_{\mu}}$.

$$X_i^{\mu}(t) = \sum_{k=1}^{s_{\mu}} x_{i,k}^{\mu}(t) \quad , \quad x_{i,k}^{\mu}(0) = 1.$$
 (B1)

The substitution of the individuals by two offspring corresponds to the process of getting a vote. The frequency of deaths or the probability of getting another vote is proportional to $X_i^{\mu}(t)$. This relation is the same as that in the discrete time voting model. The counts of voting, t, corresponds to the counts of branchings. If branching takes place t times up to t', the following relation holds.

$$X_i^{\mu}(t') = X_{i,t}^{\mu}$$

The expectation values $< x_{i,k}^{\mu}(t) >$ and $< X_i^{\mu}(t) >$ increase with e^t . Next, we introduce the scaled variables $U_i^{\mu}(t)$ and $u_{i,k}^{\mu}(t)$ as

$$U_i^{\mu}(t) \equiv e^{-t} X_i^{\mu}(t) \text{ and } u_{i,k}^{\mu}(t) \equiv e^{-t} x_{i,k}^{\mu}(t).$$
 (B2)

We focus on the following probability distributions:

$$p_{s_{\mu}}(u)du \equiv \lim_{t \to \infty} \text{Prob}(u \le U_i^{\mu}(t) \le u + du)$$
 (B3)

$$p(u)du \equiv \lim_{t \to \infty} \text{Prob}(u \le u_{i,k}^{\mu}(t) \le u + du).$$
 (B4)




FIG. 6: Mapping voting model to branching process. The left-hand side figure shows a voting process with $N_1 = N_0 = 2$. \bigcirc represents candidate belonging to category $\mu = 1, s_1 = 2$ and \times represents candidate belonging to category $\mu = 0, s_0 = 1$. The right-hand side figure shows the corresponding branching process. \bullet represents the initial individual and offspring. Candidate belonging to category $\mu = 1(\mathbf{0})$ is composed of two individuals (one individual).




FIG. 7: Pictorial representation of self-consistent relation among u, u_1 and u_2 . An individual splits at $t = \tau$ for the first time producing two offspring appears. Because of the time lag τ , the relation $u = (u_1 + u_2)e^{-\tau}$ holds.

In order to obtain p(u), we consider the situation in which an individual splits at $t = \tau$ for the first time. The resulting two offspring continue the branching process. The scaled number of offspring of the individual is denoted as u. Those of the two offspring are denoted as u_1 and u_2 . Figure 7 gives a pictorial representation of the relation among u, u_1 and u_2 . We observe that these variables satisfy the following relation:

$$u = (u_1 + u_2)e^{-\tau}$$
.

Furthermore, u_1 and u_2 obey the same probability distribution as that obeyed by u, and the probability that an individual splits for the first time during $\tau \leq t \leq \tau + d\tau$ is $e^{-\tau}d\tau$. Thus, we obtain

$$p(u) = \int_0^\infty e^{-\tau} d\tau \int_0^\infty du_1 \int_0^\infty du_2 p(u_1) p(u_2) \delta(u - (u_1 + u_2)e^{-\tau}).$$
 (B5)

Introducing $X = e^{-\tau}$, the relation is rewritten as

$$p(u) = \int_0^1 dX \int_0^\infty du_1 \int_0^\infty du_2 p(u_1) p(u_2) \delta(u - (u_1 + u_2)X).$$
 (B6)

Using the Laplace transform of p(u), $\hat{p}(s) \equiv \int_0^\infty p(u)e^{-su}du$, it can be shown that $\hat{p}(s)$ satisfies the following integral equation:

$$\hat{p}(s) = \frac{1}{s} \int_0^s \hat{p}(v) dv. \tag{B7}$$

Differentiating (B7) with respect to s, we obtain the following differential equation.

$$s\frac{d\hat{p}(s)}{ds} = \hat{p}^2(s) - \hat{p}(s). \tag{B8}$$

(B8) can be solved easily to obtain

$$\hat{p}(s) = \frac{1}{1+as}. ag{B9}$$

Using the normalization condition $\langle u \rangle = 1$ and the inverse Laplace transform, we get

$$p(u) = e^{-u}. (B10)$$

We obtain $p_{s_{\mu}}(u)$ by convolution as

$$p_{s_{\mu}} = \prod_{i=1}^{s_{\mu}} \left[\int_{0}^{\infty} du_{i} p(u_{i}) \right] \delta(u - \sum_{i=1}^{s_{\mu}} u_{i})$$

$$= \frac{1}{\Gamma(\mu)} u^{\mu-1} e^{-u}.$$
(B11)

 U_i^{μ} obeys a gamma distribution with the shape exponent s_{μ} given by (10). We note that the result (10) is derived in the thermodynamic limit, where the correlation among $\{u_j\}_{j=1,\dots k}$ vanishes. On the other hand, in the continuous time branching process, the splitting processes of each individual and offspring are independent of each other. As a result, we obtain the gamma distribution which appears in the voting model in the thermodynamic limit.