# From Black-Scholes and Dupire formulae to last passage times of local martingales

Part A : The infinite time horizon

Amel Bentata∗and Marc Yor†

June,  $2^{nd}$ , 2008

- 1. These notes are the first half of the contents of the course given by the second author at the Bachelier Seminar (February 8-15-22 2008) at IHP. They also correspond to topics studied by the first author for her Ph.D.thesis.
- 2. Comments are welcome and may be addressed to : bentata@clipper.ens.fr.

<sup>\*</sup>Université Paris 6, Laboratoire de Probabilités et Modèles Aléatoires, CNRS-UMR 7599, 16, rue Clisson, 75013 Paris Cedex, France

<sup>&</sup>lt;sup>†</sup>Université Paris 6, Laboratoire de Probabilités et Modèles Aléatoires, CNRS-UMR 7599, 16, rue Clisson, 75013 Paris Cedex, France-Institut Universitaire de France

# Contents



## References [42](#page-41-0)

## A rough description of Part A of the course

The starting point of the course has been the elementary remark that the following holds :

<span id="page-3-0"></span>
$$
\mathbb{E}\left[\left(\mathcal{E}_t-1\right)^{\pm}\right] = \mathbb{P}\left(4\mathbf{N}^2 \leq t\right),\tag{1}
$$

where on the  $LHS$ ,  $\mathcal{E}_t = \exp\left(B_t - \frac{t}{2}\right)$  $\left(\frac{t}{2}\right)$ , for  $(B_t)$  a standard Brownian motion, and  $N \stackrel{\text{law}}{=} B_1$  is the standard Gaussian variable. The identity [\(1\)](#page-3-0) may follow from inspection of the Black-Scholes formula, but seemed to deserve further explanation.

The full course consists in ten notes, the contents of the first five are :

In Note 1, it is shown that a wide extension of [\(1\)](#page-3-0) holds with  $\mathcal{E}_t$  being replaced by a continuous local martingale  $M_t \geq 0$ , converging to 0, as  $t \to \infty$ , and with  $4N^2$  being replaced by the last passage time at 1 by M.

This motivates the study, in Note 2, of the law of  $\mathcal{G}_K = \sup\{t, M_t = K\}.$ In this note, we recover the computation of the laws of the last passage times for transient diffusions, as obtained by Pitman-Yor in [\[22\]](#page-42-0), and we extend these results in a natural manner, when  $(M_t, t \ge 0)$  is only assumed to be a positive local martingale, converging to 0, as  $t \to \infty$ .

In Note 3, a connection is made with some representation of Azema supermartingales associated with ends  $L$  of previsible random time sets; it turns out that  $L = \mathcal{G}_K$  is a particular case of such random times; hence, the obtained supermartingales are particular cases of Azema's supermartingales. This Note 3 also leads us to present the progressive enlargement of filtration formulae in this setup.

In Note 4, the main formula :

$$
\mathbb{P}\left(\mathcal{G}_K \le t | \mathcal{F}_t\right) = \left(1 - \frac{M_t}{K}\right)^+, \tag{2}
$$

on which most of our previous discussion has been based is shown to generalize in the form :

$$
\mathbb{E}\left[1_{\{\mathcal{G}_K\leq t\}}\left(K-M_{\infty}\right)^{+}|\mathcal{F}_t\right] = \left(K-M_t\right)^{+},\tag{3}
$$

in the case where  $(M_t, t \geq 0)$  is only assumed to take values in  $\mathbb{R}^+$ , but  $M_{\infty}$ is not necessarily equal to 0. We then explain how to obtain a formula for  $\mathbb{P}(\mathcal{G}_K \leq t | \mathcal{F}_t).$ 

In Note 5, we integrate the previous results with respect to  $K$ , in a similar manner as one may recover Itô's formula from Tanaka 's formula. This note bears quite some similarity with the paper by Az $\acute{e}$ ma-Yor [\[1\]](#page-41-1) on local times.

## <span id="page-4-0"></span>1 Note 1 : Option Prices as Probabilities

### <span id="page-4-1"></span>1.1 A first question

One of the pillars of modern mathematical finance has been the computation (and the understanding !) of the quantities :

$$
\mathbb{E}\left[ \left( \mathcal{E}_t - K \right)^{\pm} \right],
$$

where :

$$
\mathcal{E}_t = \exp\bigg(B_t - \frac{t}{2}\bigg),\,
$$

with  $(B_t)$  a Brownian motion starting from 0.

In an explicit form<sup>[1](#page-4-2)</sup>, the Black-Scholes formula writes :

<span id="page-4-5"></span>
$$
\mathbb{E}\left[\left(\mathcal{E}_t - K\right)^+\right] = \left(1 - K\right) + \mathbb{E}[(K - \mathcal{E}_t)^+\right] \tag{4}
$$

<span id="page-4-4"></span>
$$
= \mathcal{N}\left(-\frac{\log K}{\sqrt{t}} + \frac{\sqrt{t}}{2}\right) - K\mathcal{N}\left(-\frac{\log K}{\sqrt{t}} - \frac{\sqrt{t}}{2}\right) \tag{5}
$$

where :

$$
\mathcal{N}(x) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{x} dy e^{-y^2/2}.
$$

Since  $(\mathcal{E}_t, t \ge 0)$  is a martingale, both  $(\mathcal{E}_t - K)^+$  and  $(K - \mathcal{E}_t)^+$  are submartingales; hence :

$$
t \to C^{\pm}(t, K) = \mathbb{E}\left[ (\mathcal{E}_t - K)^{\pm} \right]
$$

are increasing functions of t.

They are also continuous, and<sup>[2](#page-4-3)</sup>:

i. 
$$
C^+(0, K) = (1 - K)^+; C^+(\infty, K) = 1;
$$

ii. 
$$
C^-(0, K) = (K - 1)^+
$$
;  $C^-(\infty, K) = K$ .

Consequently :

i'. if  $K \geq 1$ ,  $C^+(t, K)$  increases from 0 (for  $t = 0$ ), to 1 (for  $t = \infty$ );

ii'. if  $K \leq 1$ ,  $\frac{1}{K}C^{-}(t, K)$  increases from 0 (for  $t = 0$ ), to 1 (for  $t = \infty$ ).

<sup>&</sup>lt;sup>1</sup>Formula [\(5\)](#page-4-4) extends easily when we replace  $\mathcal{E}_t$  by  $\exp(\sigma B_t + \nu t)$ , so there is no loss of generality to take :  $\sigma = 1$ ,  $\nu = -1/2$ .

<span id="page-4-3"></span><span id="page-4-2"></span><sup>&</sup>lt;sup>2</sup>That  $C^+(\infty, K) = 1$  is most easily seen using [\(4\)](#page-4-5), and the fact that  $\mathcal{E}_t \to_{t \to \infty} 0$ .

Therefore, in both cases,  $C^+(\bullet, K)$ , and  $C^-(\bullet, K)$  are distribution functions of a certain random variable  $X^{\pm}$  taking values in  $\mathbb{R}^{+}$ .

Can we identify the corresponding distribution?

Or, even better, can we find, in our Brownian (Black-Scholes) framework, a random variable whose distribution function is  $C^{+/-}(\bullet, K)$ ?

To motivate the reader's interest, we assert, right away, taking  $K = 1$ , that there is the formula :

<span id="page-5-0"></span>
$$
\mathbb{E}\left[\left(\mathcal{E}_t-1\right)^+\right]=\mathbb{E}\left[\left(\mathcal{E}_t-1\right)^-\right]=\mathbb{P}\left(4B_1^2\leq t\right).
$$
\n(6)

We think of this formula as "an alternative Black-Scholes formula". Furthermore, formula [\(6\)](#page-5-0) has been very helpful to answer M.Qian's question : given a probability measure  $\mu(dt)$  on  $\mathbb{R}^+$ , can one compute :

$$
\int_0^\infty \mu(dt) \mathbb{E}\left[ (\mathcal{E}_t - 1)^{\pm} \right] ? \tag{7}
$$

Indeed from [\(6\)](#page-5-0), the previous quantity equals :

$$
\mathbb{E}\left[\overline{\mu}(4B_1^2)\right],\tag{8}
$$

where :  $\overline{\mu}(x) = \mu([x,\infty))$ . For example, if :

$$
\mu(dt) = \lambda e^{-\lambda t} dt,
$$

then :

$$
\int_0^\infty \lambda \, dt \, e^{-\lambda t} \mathbb{E} \left[ \left( \mathcal{E}_t - 1 \right)^{\pm} \right] = \frac{1}{\sqrt{1 + 8\lambda}}. \tag{9}
$$

Question 1.1. It also seems of interest to ask the following extension of M.Qian's question : what is the law of :

$$
\mathcal{E}_{\mu}^{\pm} \equiv \int_0^{\infty} \mu(dt) \left(\mathcal{E}_t - 1\right)^{\pm},
$$

in particular in the case  $\mu(dt) = \lambda e^{-\lambda t} dt$ ? We may start by computing moments of this variable  $\mathcal{E}_{\mu}^{\pm}$ .

In fact, in July 1997, Prof. Miura asked the second author for the law of  $\int_0^t ds \, (\mathcal{E}_s - 1)^+$ , in order to obtain the price of "Area options", that is :  $\mathbb{E}\left[\left(\int_0^t ds\, \left(\mathcal{E}_s-1\right)^+ - K\right)^+\right]$ .

Below, we give a clear probabilistic explanation of formula [\(6\)](#page-5-0), and even more generally of the extended alternative Black-Scholes formula :

<span id="page-5-1"></span>
$$
\mathbb{E}\left[\left(\mathcal{E}_t - K\right)^{\pm}\right] = \left(1 - K\right)^{\pm} + \sqrt{K} \mathbb{E}\left[1_{\{4B_1^2 \le t\}} \exp\left(-\frac{(\log K)^2}{8B_1^2}\right)\right].\tag{10}
$$

### <span id="page-6-0"></span>1.2 A first answer

In fact, the previous question admits a general answer, which does not require to work within a Brownian framework.

Let  $(M_t, t \geq 0)$  denote a continuous local martingale, defined on  $(\Omega, \mathcal{F}, \mathcal{F}_t, \mathbb{P});$ we assume that  $M_t \geq 0$  and  $M_t \to 0$  when  $t \to \infty$ . Let  $\mathcal{M}_0^+$  denote the set of these particular local martingales, we insist that we allow local martingales...

<span id="page-6-3"></span>**Theorem 1.1.** Let  $\mathcal{G}_K = \sup\{t, M_t = K\}$  with the convention  $\sup\{\emptyset\} = 0$ . Then :

<span id="page-6-5"></span>
$$
\left(1 - \frac{M_t}{K}\right)^+ = \mathbb{P}\left(\mathcal{G}_K \le t | \mathcal{F}_t\right). \tag{11}
$$

(*t* maybe replaced by any stopping time  $T$ ). Consequently :

<span id="page-6-2"></span>
$$
\mathbb{E}\left[\left(1-\frac{M_t}{K}\right)^+\right] = \mathbb{P}\left(\mathcal{G}_K \le t\right). \tag{12}
$$

Proof. a) Note that :

$$
(\mathcal{G}_K \le t) = \left(\sup_{s \ge t} M_s \le K\right). \tag{13}
$$

b) From the next lemma, we have conditionally on  $\mathcal{F}_t$ :

<span id="page-6-1"></span>
$$
\sup_{s \ge t} M_s \stackrel{\text{law}}{=} \frac{M_t}{\mathbf{U}},\tag{14}
$$

where **U** is uniform on [0, 1] and independent from  $\mathcal{F}_t$ . Consequently,

$$
\mathbb{P}\left(\sup_{s\geq t} M_s \leq K|\mathcal{F}_t\right) = \mathbb{P}\left(\frac{M_t}{\mathbf{U}} < K|\mathcal{F}_t\right) = \left(1 - \frac{M_t}{K}\right)^+.\tag{15}
$$

Now formula [\(14\)](#page-6-1) follows from the elementary, but very useful lemma :

<span id="page-6-4"></span>**Lemma 1.1** (Doob's maximal identity). If  $(N_t, t \ge 0) \in \mathcal{M}_0^+$ , then :

$$
\sup_{t\geq 0} N_t \stackrel{\text{law}}{=} \frac{N_0}{\mathbf{U}},\tag{16}
$$

where U is uniform on [0, 1] and independent from  $\mathcal{F}_0$ .

*Proof.* We use Doob's optional stopping theorem : if  $T_a = \inf\{t, N_t = a\}$ , (with the convention inf $\{\emptyset\} = \infty$ ), then, if  $a > N_0$ :

$$
\mathbb{E}\left[N_{T_a}|\mathcal{F}_0\right] = N_0
$$
  

$$
a \mathbb{P}\left(T_a < \infty|\mathcal{F}_0\right) = N_0,
$$

that is  $\mathbb{P}\left(\sup_{t\geq 0} N_t > a | \mathcal{F}_0\right) = \frac{N_0}{a}$  $\frac{v_0}{a}$ . This yields to the result.

Exercise 1.1. Denote :

$$
\overline{\mathcal{E}}_{(t,\infty)} \equiv \sup_{s \geq t} \mathcal{E}_s.
$$

Prove that the process :  $(\lambda_t \equiv \mathcal{E}_t/\mathcal{E}_{(t,\infty)}, t \geq 0)$  is strictly stationary, with common law U.

More generally, prove that the process  $(\lambda_t^{(p)} \equiv \mathcal{E}_t / \tilde{\mathcal{E}}_t^{(p)})$  $t_t^{(p)}, t \ge 0$ ) is strictly stationary, where :

$$
\tilde{\mathcal{E}}_t^{(p)} = \left( \int_t^\infty du \, \exp p \left( B_u - \frac{u}{2} \right) \right)^{1/p}.
$$

Show that :

$$
\lambda_t^{(p)} \to_{p \to \infty} \lambda_t.
$$

What is the common law  $U_p$  of the  $\lambda_t^{(p)}$  $\stackrel{(p)}{t} \stackrel{\prime}{s}$ ?

#### <span id="page-7-0"></span>1.3 The law of  $G_a^{(\nu)}$  $\dot{a}$

Coming back to our original question in Section 1.1, we observe that formula [\(12\)](#page-6-2), in the Brownian framework, gives :

$$
\mathbb{E}\left[\left(1-\frac{\mathcal{E}_t}{K}\right)^+\right] = \mathbb{P}\left(\mathcal{G}_K \le t\right). \tag{17}
$$

Hence, taking  $K = 1$ , it suffices to obtain the identity:

<span id="page-7-1"></span>
$$
\mathcal{G}_1 \stackrel{\text{law}}{=} 4B_1^2,\tag{18}
$$

to recover formula [\(6\)](#page-5-0); this identity [\(18\)](#page-7-1) may be simply obtained by time inversion, since :

$$
G_1 = \sup\{t, \mathcal{E}_t = 1\} = \sup\{t, B_t - \frac{t}{2} = 0\}
$$

hence :

$$
\mathcal{G}_1 \stackrel{\text{law}}{=} \frac{1}{T_{1/2}} \stackrel{\text{law}}{=} \frac{4}{T_1} \stackrel{\text{law}}{=} 4B_1^2. \tag{19}
$$

 $\Box$ 

We are now bound to describe the law of  $G_a^{(\nu)} = \sup\{t, B_t + \nu t = a\}$ , for all  $a, \nu \in \mathbb{R}$ . These laws are well-known, thanks again to the stability by time inversion for Brownian motion : if  $(B_u)$  is a Brownian motion, then :

$$
\hat{B}_u = uB_{\frac{1}{u}}, \quad u > 0
$$

is also a Brownian motion.

As a consequence :

$$
\left(T_a^{(\nu)}, G_a^{(\nu)}\right) \stackrel{\text{law}}{=} \left(\frac{1}{G_\nu^{(a)}}, \frac{1}{T_\nu^{(a)}}\right). \tag{20}
$$

The (separate) laws of  $T_a^{(\nu)}$  and  $G_a^{(\nu)}$  are (for  $a > 0, \nu \in \mathbb{R}$ ):

$$
\mathbb{P}\left(T_a^{(\nu)} \in dt\right) = \frac{dt \, a}{\sqrt{2\pi t^3}} \exp\left(-\frac{(a - \nu t)^2}{2t}\right),\tag{21}
$$

and

$$
\mathbb{P}\left(G_a^{(\nu)} \in dt\right) = |\nu| \frac{dt}{\sqrt{2\pi t}} \exp\left(-\frac{(a - \nu t)^2}{2t}\right). \tag{22}
$$

We refer to [\[22\]](#page-42-0) for some further discussion about time inversion.

**Exercise 1.2.** Give the expression of the joint law of  $(T_a^{(\nu)}, G_a^{(\nu)})$ .

Exercise 1.3. Recover the Black-Scholes formula thanks to the knowledge of the laws of  $G_a^{(\pm 1/2)}$ .

Exercise 1.4. Establish formula [\(10\)](#page-5-1).

#### <span id="page-8-0"></span>1.4 Other universal laws

We now come back to the setup of Section 2; we would like to understand better why a "universal law", such as the uniform, occurs in the framework of Theorem [1.1.](#page-6-3)

Recall that :

<span id="page-8-1"></span>
$$
N_t = \beta_{\langle N \rangle_t}, \quad t \ge 0,
$$
\n(23)

where  $(\beta_u)$  is a Brownian motion starting from  $N_0$ .

Since  $N_t \to 0$  when  $t \to \infty$ , one has :

$$
\langle N \rangle_{\infty} = T_0(\beta). \tag{24}
$$

Now, we see that :

$$
\sup_{t\geq 0} N_t = \sup_{u\leq T_0(\beta)} \beta_u.
$$
\n(25)

Hence, taking for simplicity  $N_0 = 1$ , we see why the law of  $\sup_{t \geq 0} N_t$  is universal, i.e : it is the law of  $\sup_{u \leq T_0(\beta)} \beta_u$ , which, as we have already shown, is the law of  $\frac{1}{U}$ .

Now, it may be natural to see whether some other functionals of  $N$ , say  $F(N)$ , maybe reduced to the corresponding functionals of  $\beta$ , killed at  $T_0(\beta)$ , i.e :  $F(N) = F(\beta_{\Lambda T_0(\beta)})$ . In this case  $F(N)$  will have "the universal law" of  $F(\beta_{\cdot\wedge T_0(\beta)})$ .

#### Question 1.2. Characterize the universal functionals F.

To identify at least some such functionals, let us recall the definition of the local times of  $N$ , via the occupation measure :

$$
f \to \int_0^t d\langle N \rangle_s f(N_s), \quad f : \mathbb{R}^+ \to \mathbb{R}^+, \text{Borel.}
$$

which is absolutely continuous with respect to the Lebesgue measure; indeed :

<span id="page-9-1"></span>
$$
\int_0^t d\langle N \rangle_s f(N_s) = \int_0^\infty dx f(x) \mathcal{L}_t^x(N), \tag{26}
$$

where  $(\mathcal{L}_t^x(N); x \geq 0, t \geq 0)$  is the jointly continuous family of local times of N.

From the Dubins-Schwarz relation (see [\(23\)](#page-8-1)), we obtain :

$$
\mathcal{L}_t^x(N) = \mathcal{L}_{\langle N \rangle_t}^x(\beta). \tag{27}
$$

Consequently :

$$
\mathcal{L}_{\infty}^{x}(N) = \mathcal{L}_{T_{0}(\beta)}^{x}(\beta). \tag{28}
$$

Hence, the local time process  $(\mathcal{L}^x_\infty(N), x \geq 0)$  is a universal functional, whose law, that is the law of the process  $(\mathcal{L}^x_{T_0(\beta)}(\beta), x \ge 0)$  is well-known and is the subject of the following Ray-Knight theorem.

<span id="page-9-0"></span>**Theorem 1.2** (Ray-Knight). Let  $(\beta_u, u \leq T_0(\beta))$  be a Brownian motion starting at 1, considered up to time  $T_0(\beta)$ , its first time when it hits 0. Then :  $(Z_x = \mathcal{L}^x_{T_0(\beta)}(\beta), x \ge 0)$  satisfies :

$$
Z_x = 2 \int_0^x \sqrt{Z_y} \, d\gamma_y + 2 \left( x \wedge 1 \right), \tag{29}
$$

where  $(\gamma_y, y \geq 0)$  is a Brownian motion. In other words,

i.  $(Z_r, x \leq 1)$  is a  $BESQ_0(2)$ ;

- ii.  $Z_1$  is distributed as 2e, where e is a standard exponential variable;
- iii. Conditionally on  $Z_1 = z$ ,  $(Z_{1+x}, x \ge 0)$  is a  $BESQ_z(0)$ .

Exercise 1.5. Recover the universal result :

$$
\sup_{t \leq T_0(\beta)} \beta_t \stackrel{\text{law}}{=} \frac{1}{\mathbf{U}},
$$

from Theorem [1.2.](#page-9-0)

Proof. (A possible one!)

Call  $\Sigma = \sup_{t \leq T_0(\beta)} \beta_t$ , and note that :

$$
\Sigma = 1 + \inf \{ x \ge 1, Z_x = 0 \}.
$$

By time reversal ,

$$
\Sigma = 1 + \sup\{t, \hat{Z}_t = Z_1\},\
$$

where  $(\hat{Z}_t)$  is a  $BESQ_0(4)$ . Hence :

$$
\Sigma \stackrel{\text{law}}{=} 1 + \frac{Z_1}{2\gamma_1} \stackrel{\text{law}}{=} 1 + \frac{\mathbf{e}}{\mathbf{e}'} \stackrel{\text{law}}{=} \frac{\mathbf{e'} + \mathbf{e}}{\mathbf{e}'} \stackrel{\text{law}}{=} \frac{1}{\mathbf{U}}.
$$
 (30)

It may be of interest to give the general Laplace transform of :

<span id="page-10-0"></span>
$$
\int_0^\infty d\langle N \rangle_s f(N_s) = \int_0^{T_0(\beta)} du f(\beta_u). \tag{31}
$$

We refer to [\[5\]](#page-41-2).

However, we may identify directly the law of the  $RHS$  of [\(31\)](#page-10-0) when f is a power function :  $f(x) = x^{\alpha}$   $\alpha > 0$ . Indeed, applying It At''s formula to  $(\beta_u^{\alpha}, u \leq T_0(\beta))$ , it is easily shown that :

<span id="page-10-1"></span>
$$
\beta_u^{\alpha} = \rho_{\alpha^2 \int_0^u ds \beta_s^{2(\alpha - 1)}}, \quad u \le T_0(\beta), \tag{32}
$$

where  $(\rho_h, h \ge 0)$  is a BES process, with dimension  $d_{\alpha} = 2 - \frac{1}{\alpha}$  $\frac{1}{\alpha}$ . Consequently, formula [\(32\)](#page-10-1) yields :

$$
\alpha^2 \int_0^{T_0(\beta)} ds \,\beta_s^{2(\alpha-1)} \stackrel{\text{law}}{=} T_0(\rho) \stackrel{\text{law}}{=} \mathcal{G}_1(\rho'),\tag{33}
$$

where  $(\rho'_u, u \ge 0)$  is the *BES* process starting from 0, with dimension :  $d'_{\alpha} = 2 + \frac{1}{\alpha}$ . Then, elementary arguments using Lemma [\(1.1\)](#page-6-4) lead to :

$$
\mathcal{G}_1 \stackrel{\text{law}}{=} \frac{1}{2\gamma_{1/2\alpha}},
$$

where  $\gamma_{\nu}$  indicates a gamma variable with parameter  $\nu$ . For this proof, we refer to [\[28\]](#page-43-0), p16–17.

## <span id="page-11-0"></span>2 Note 2 : Computing the law of  $\mathcal{G}_K$

In Note 1, we have shown (under our current hypotheses) :

<span id="page-11-2"></span>
$$
\mathbb{E}\left[\left(1-\frac{M_t}{K}\right)^+\right] = \mathbb{P}\left(\mathcal{G}_K \le t\right). \tag{34}
$$

As a motivation for this note, remark that when  $M_t = \mathcal{E}_t$ , the LHS of [\(34\)](#page-11-2) is known : this is the Black-Scholes formula ! Consequently, we can recover from the Black-Scholes formulae (see [\(4\)](#page-4-5) and [\(5\)](#page-4-4)) the law of  $\mathcal{G}_K$ .

#### <span id="page-11-1"></span>2.1 A general result

Here, we aim to give a formula for the law of  $\mathcal{G}_K$  associated to our general local martingale  $(M_t)$ , and its local times  $\mathcal{L}^x_t(M)$  as defined via  $(26)$ :

<span id="page-11-5"></span>
$$
\int_0^t d\langle M \rangle_s f(M_s) = \int_0^\infty dx f(x) \mathcal{L}_t^x(M). \tag{35}
$$

To proceed, we need to make some further hypotheses on M :

- $(H_1)$  for every  $t > 0$ , the law<sup>[3](#page-11-3)</sup> of the r.v.  $M_t$  admits a density  $(m_t(x), x \ge 0)$ , and :  $(t, x) \to m_t(x)$  may be chosen continuous on  $(0, \infty)^2$ ;
- $(H_2) d\langle M \rangle_t = \sigma_t^2 dt$ , and there exists a jointly continuous function :

$$
(t, x) \to \theta_t(x) = \mathbb{E}\left[\sigma_t^2 | M_t = x\right] \quad \text{on} \quad (0, \infty)^2.
$$

Then, the following holds :

<span id="page-11-7"></span>**Theorem 2.1.** The law of  $\mathcal{G}_K$  is given by :

<span id="page-11-4"></span>
$$
\mathbb{P}\left(\mathcal{G}_K \in dt\right) = \left(1 - \frac{a}{K}\right)^+ \epsilon_0(dt) + \frac{1_{\{t > 0\}}}{2K} \theta_t(K) m_t(K) dt, \tag{36}
$$

where  $a = M_0$ .

Proof. a) Using Tanaka's formula, one obtains:

<span id="page-11-6"></span>
$$
\mathbb{E}\left[\left(K - M_t\right)^+\right] = \left(K - a\right)^+ + \frac{1}{2} \mathbb{E}\left[\mathcal{L}_t^K(M)\right].\tag{37}
$$

<span id="page-11-3"></span> $3$ We are grateful to F. Delarue for pointing out the exercise in [\[20\]](#page-42-1) p.97 which gives a sufficient condition.

Thus, from [\(34\)](#page-11-2), there is the relationship :

$$
\mathbb{P}\left(\mathcal{G}_K \in dt\right) = \left(1 - \frac{a}{K}\right)^+ \epsilon_0(dt) + \frac{1_{\{t > 0\}}}{2K} d_t \left(\mathbb{E}\left[\mathcal{L}_t^K(M)\right]\right),\tag{38}
$$

and formula [\(36\)](#page-11-4) is now equivalent to the following expression for  $d_t\left(\mathbb{E}\left[\mathcal{L}_t^K(M)\right]\right)$  :

<span id="page-12-1"></span>
$$
d_t\left(\mathbb{E}\left[\mathcal{L}_t^K(M)\right]\right) = dt\,\theta_t(K)\,m_t(K)\quad (t>0). \tag{39}
$$

b) We now prove [\(39\)](#page-12-1). The density of occupation formula [\(35\)](#page-11-5) for the local martingale  $(M_t)$  writes : for every  $f : \mathbb{R}^+ \to \mathbb{R}^+$ , Borel,

<span id="page-12-2"></span>
$$
\int_0^t ds \,\sigma_s^2 f(M_s) = \int_0^\infty dK f(K) \mathcal{L}_t^K(M). \tag{40}
$$

Thus, taking expectations on both sides of [\(40\)](#page-12-2), we obtain :

<span id="page-12-3"></span>
$$
\mathbb{E}\left[\int_0^t ds \,\sigma_s^2 f(M_s)\right] = \int_0^\infty dK f(K) \mathbb{E}\left[\mathcal{L}_t^K(M)\right].\tag{41}
$$

The  $LHS$  of [\(41\)](#page-12-3) equals :

$$
\int_0^t ds \mathbb{E} \left[ \mathbb{E} \left[ \sigma_s^2 | M_s \right] f(M_s) \right] = \int_0^\infty dK f(K) \int_0^t ds \, m_s(K) \, \theta_s(K) \tag{42}
$$

and formula [\(39\)](#page-12-1) now follows easily from [\(41\)](#page-12-3).

 $\Box$ 

**Exercise 2.1.** Give the particular case of formula [\(36\)](#page-11-4) when  $M_t = \mathcal{E}_t$ , thus recovering again the law of  $\mathcal{G}_K$  in the Brownian framework.

### <span id="page-12-0"></span>2.2 Some connection with the Dupire formula

We recall our original notation :

$$
C^{\pm}(t,K) = \mathbb{E}\left[ \left( \mathcal{E}_t - K \right)^{\pm} \right],
$$

which we now extend to our general martingale case, i.e :

$$
C^{\pm}(t, K) = \mathbb{E}\left[\left(M_t - K\right)^{\pm}\right].
$$

Theorem 2.2. The following identities hold :

$$
\frac{\partial}{\partial T}(C^-(T,K)) \stackrel{\text{(a)}}{=} \theta_T(K) \frac{\partial^2}{\partial K^2} C^-(T,K) \stackrel{\text{(b)}}{=} 2K\gamma_K(T),\tag{43}
$$

where  $(\gamma_K(T), T > 0)$  is the density of  $\mathcal{G}_K$ .

Comment : The identity (a) is also found, up to minor differences, in Klebaner [\[8\]](#page-41-3). In general, connections between local times and the Black-Scholes and Dupire formulae had been noticed for quite some time by several authors. However, the identity (b) seems, to the best of our knowledge, to be new.

Proof. Thanks to [\(37\)](#page-11-6), one has :

$$
\frac{\partial}{\partial T}(C^-(T,K)) = \frac{1}{2} \frac{\partial}{\partial T} \mathbb{E}[\mathcal{L}_T^K],\tag{44}
$$

and, clearly :

$$
\frac{\partial^2}{\partial K^2} C^-(T, K) = m_T(K). \tag{45}
$$

 $\Box$ 

From [\(39\)](#page-12-1), we obtain :

$$
\frac{\partial}{\partial T}\mathbb{E}[\mathcal{L}_T^K] = \theta_T(K)m_T(K) = 2K\gamma_K(T).
$$

We refer to [\[6\]](#page-41-4) and [\[7\]](#page-41-5) for the "true" Dupire formula.

### <span id="page-13-1"></span><span id="page-13-0"></span>2.3 Specialising to transient diffusions

#### 2.3.1 General framework

We present here some results which can be found in [\[22\]](#page-42-0), chapter 6.

We consider the canonical realisation of a transient diffusion

$$
(R_t, t \geq 0; \mathbb{P}_x, x \in \mathbb{R}^+) \text{ on } C(\mathbb{R}^+, \mathbb{R}^+).
$$

For simplicity, we suppose that :

i. 
$$
\mathbb{P}_x(T_0 < \infty) = 0, x > 0;
$$

ii.  $\mathbb{P}_x$  (lim<sub>t→∞</sub>  $R_t = \infty$ ) = 1,  $x > 0$ .

As a consequence of (i) and (ii), there exists a scale function s for this diffusion which satisfies  $s(0^+) = -\infty$  and  $s(\infty) = 0$ . Let  $\Gamma$  be the infinitesimal generator of the diffusion<sup>[4](#page-13-2)</sup>, and take the speed measure  $m$  to be such that :

$$
\Gamma = \frac{1}{2} \frac{d}{dm} \frac{d}{ds} \, .
$$

<span id="page-13-2"></span><sup>&</sup>lt;sup>4</sup>This is the classical Itô-Mc Kean presentation; see also Borodin-Salminen [\[4\]](#page-41-6); for "practical" cases, see 2.3.2

Let

$$
g_y = \sup\{t > 0, R_t = y\}.
$$

Then, by applying the results of the previous section to  $M_t = -s(R_t)$ , we may obtain the following theorem :

**Theorem 2.3** (Pitman-Yor, [\[22\]](#page-42-0), section 6). For all  $x, y > 0$ ,

<span id="page-14-1"></span>
$$
\mathbb{P}_x(g_y \in dt) = \frac{-1}{2s(y)} p_t^{\bullet}(x, y) dt,
$$
\n(46)

where  $p_t^{\bullet}(x, y) (= p_t^{\bullet}(y, x))$  is the density of the semigroup  $P_t(x, dy)$  with respect to  $m(dy)$ .

Proof. a) Previous arguments show that :

<span id="page-14-0"></span>
$$
\mathbb{P}_x\left(g_y \le t\right) = \mathbb{E}_x\left[\left(1 - \frac{M_t}{(-s(y))}\right)^+\right],\tag{47}
$$

by changing the space variable :  $\mu = s(x)$ , which corresponds to putting the diffusion R in its natural scale, i.e : replacing it by  $M_t = -s(R_t)$ .

b) Tanaka's formula now yields, from [\(47\)](#page-14-0) :

$$
\mathbb{P}_x\left(g_y \le t\right) = \left(1 - \frac{s(x)}{s(y)}\right)^+ - \frac{1}{2s(y)} \mathbb{E}\left[\mathcal{L}_t^{-s(y)}(M)\right].\tag{48}
$$

Formula [\(46\)](#page-14-1) will now follow from :

$$
\frac{\partial}{\partial t} \left( \mathbb{E}_x \left[ \mathcal{L}_t^{-s(y)}(M) \right] \right) = p_t^{\bullet}(x, y). \tag{49}
$$

In turn, this formula follows from the density of occupation formula for our diffusion  $R :$  for any  $f : \mathbb{R}^+ \to \mathbb{R}^+$ , Borel :

<span id="page-14-2"></span>
$$
\int_0^t ds f(R_s) = \int m(dy) f(y) l_t^y,
$$
\n(50)

where  $(l_t^y)$  $t_i^y$ ) is the family of diffusion local times (see, e.g., [\[4\]](#page-41-6), II.13 and V.). On the  $LHS$ , we obtain:

$$
\mathbb{E}_x \left[ \int_0^t ds f(R_s) \right] = \int m(dy) \int_0^t ds \, p_s^{\bullet}(x, y) \, f(y). \tag{51}
$$

Thus, [\(50\)](#page-14-2) implies that :

<span id="page-14-3"></span>
$$
\mathbb{E}_x[l_t^y] = \int_0^t ds \, p_s^\bullet(x, y) \tag{52}
$$

On the other hand, there is the following relationship between the diffusion and martingale local times :

<span id="page-15-1"></span>
$$
l_t^y = \mathcal{L}_t^{-s(y)}(M). \tag{53}
$$

 $\Box$ 

Finally, formula [\(46\)](#page-14-1) follows from [\(52\)](#page-14-3) and [\(53\)](#page-15-1).

#### <span id="page-15-0"></span>2.3.2 In practice . . .

In practice, it may be useful to write formula [\(46\)](#page-14-1) in terms of the density  $p_t(x, y)$  of the semigroup  $P_t(x, dy)$  with respect to the Lebesgue measure dy (and not  $m(dy)$ , which may not be so "natural" as a reference measure).

We assume that the infinitesimal generator is of the form :

$$
\Gamma = \frac{1}{2}a(x)\frac{d^2}{dx} + b(x)\frac{d}{dx}
$$
\n(54)

Consequently :

$$
\frac{dm}{dy} = \frac{1}{s'(y)a(y)},\tag{55}
$$

and

$$
p_t^{\bullet}(x, y) = p_t(x, y)s'(y)a(y), \qquad (56)
$$

so that formula [\(46\)](#page-14-1) becomes :

<span id="page-15-2"></span>
$$
\mathbb{P}_x(g_y \in dt) = -\left(\frac{s'(y)a(y)}{2s(y)}\right) p_t(x, y)dt. \tag{57}
$$

**Exercise 2.2.** Recover the law of  $G_a^{(\nu)}$  from formula [\(57\)](#page-15-2)

**Exercise 2.3.** Write explicitly formula [\(57\)](#page-15-2) for  $(R_t)$  a transient BES process, i.e : the  $\mathbb{R}^+$ -valued diffusion with infinitesimal generator :

$$
\frac{1}{2}\frac{d^2}{dx} + \frac{\delta - 1}{2}\frac{d}{dx}, \quad \delta > 2.
$$

Answer :

$$
g_a(R) \stackrel{\text{law}}{=} \frac{a^2}{2\gamma_\nu},
$$

when  $R_0 = 0$ . See [\[28\]](#page-43-0).

## <span id="page-16-0"></span>2.4 Other examples of explicit computations of the law of  $\mathcal{G}_K$

We present here the following examples : the killed Brownian motion, the inverse of a 3-dimensional Bessel process, and an example of an inhomogeneous Markov process for which we can compute  $m_t(x)$ . For more details, see [\[12\]](#page-41-7). These examples will be detailed in the appendix of Part B, in section 11.

**Example 2.1.**  $M_t = B_{t \wedge T_0}$ , where  $(B_t, t \geq 0)$  is a Brownian motion starting from 1 and  $T_0 = \inf\{t \geq 0, B_t = 0\}$ . Then for every  $K \leq 1$ ,

$$
\mathcal{G}_K(M) \stackrel{\text{law}}{=} \frac{\mathbf{U}_K^2}{\mathbf{N}^2},\tag{58}
$$

where  $U_K$  is a uniform r.v. on  $[1 - K, 1 + K]$  and independent from N a standard gaussian r.v.

Example 2.2.  $M_t = \frac{1}{R}$  $\frac{1}{R_t}$  where  $(R_t, t \geq 0)$  is a 3-dimensional Bessel process starting from 1. Then for every  $K < 1$ ,

$$
\mathcal{G}_K(M) \stackrel{\text{law}}{=} \frac{\tilde{\mathbf{U}}_K^2}{\mathbf{N}^2},\tag{59}
$$

where  $\tilde{\mathbf{U}}_K$  is a uniform r.v. on  $[\frac{1}{K} - 1, \frac{1}{K} + 1]$ , assumed to be independent from  $N$  a standard gaussian r.v.

Exercise 2.4.  $M_t = \cosh(B_t) \exp \left(-\frac{t}{2}\right)$  $\left(\frac{t}{2}\right)$  where  $(B_t, t \geq 0)$  is a Brownian motion starting from 0. Use Theorem [2.1](#page-11-7) to compute the law of  $\mathcal{G}_K$ .

Exercise 2.5. Draw a Black-Scholes-last time ( :BS-LT) Table as follows :



In this Table,  $\beta_{a,b}$  denotes a beta variable with parameters  $(a, b)$ ,  $\gamma_a$  a gamma variable with parameter a.  $\exp\left(-\frac{2B_tB_1}{1-t}\right)$  $\frac{B_t B_1}{1-t}$ , t < 1, is a martingale with respect to  $\mathcal{F}_t \vee \sigma(B_1)$ .

# <span id="page-18-0"></span>3 Note 3 : Representation of some particular Azéma supermartingales

## <span id="page-18-1"></span>3.1 A general representation theorem and our particular case

Let  $L = \sup\{t, R_t \in \Gamma\}$ , where  $(R_t)$  is a transient diffusion, and  $\Gamma$  a compact set in  $\mathbb{R}^+$ . It is interesting to describe the pre-L process :  $(R_t, t \leq L)$ and the post-L process :  $(R_{L+t}, t \geq 0)$ ; this has been the subject of many studies in the Markovian literature([\[15\]](#page-42-2), [\[16\]](#page-42-3); [\[27\]](#page-43-1) for Brownian motion). The enlargement of filtration technique shows that these descriptions"follow"once the Azéma supermartingale :

$$
Z_t = Z_t^L = \mathbb{P}\left(L > t | \mathcal{F}_t\right)
$$

has been computed "explicitly".

For the moment, we give a general representation of  $(Z_t)$  in the following framework : let  $L$  be the end of a previsible set (on a given filtered probability space) such that :

$$
\begin{cases}\n(C) & \text{all } \mathcal{F}_t \text{ martingales are continuous;} \\
(A) & \text{for any stopping time } T, \mathbb{P}(L = T) = 0.\n\end{cases}
$$
\n(CA)

(C stands for continuous, and A for avoiding (stopping times)).

<span id="page-18-2"></span>**Theorem 3.1.** [[\[14\]](#page-42-4) or [\[19\]](#page-42-5)] Under  $(CA)$ , there exists a unique positive continuous local martingale  $(N_t, t \geq 0)$ , with  $N_0 = 1$ , such that :

<span id="page-18-3"></span>
$$
\mathbb{P}\left(L > t | \mathcal{F}_t\right) = \frac{N_t}{S_t},\tag{60}
$$

where  $S_t = \sup_{s \leq t} N_s, t \geq 0$ .

Exercise 3.1. a) Give the additive decomposition of the supermartingale  $\cdot \frac{N_t}{S_t}$  $\frac{N_t}{S_t}$  as :

$$
\mathbb{E}\left[\log(S_{\infty})|\mathcal{F}_t\right] - \log(S_t).
$$

 $Hint: from Itô's formula:$ 

$$
\frac{N_t}{S_t} = 1 + \int_0^t \frac{dN_s}{S_s} - \int_0^t \frac{N_s dS_s}{S_s^2} \n= 1 + \int_0^t \frac{dN_s}{S_s} - \int_0^t \frac{dS_s}{S_s},
$$

since  $dS_s$  only charges the set  $\{s, N_s = S_s\}$ . One obtains :

<span id="page-19-1"></span>
$$
\frac{N_t}{S_t} = 1 + \int_0^t \frac{dN_s}{S_s} - \log(S_t).
$$
 (61)

b) Prove that  $log(S_{\infty})$  is distributed exponentially. Answer :

$$
\log(S_{\infty}) \stackrel{\text{law}}{=} \log\left(\frac{1}{U}\right).
$$

c) We also note that the martingale  $\mathbb{E} [\log(S_{\infty}) | \mathcal{F}_t]$  belongs to BMO, since :

$$
\mathbb{E}\left[\log(S_{\infty}) - \log(S_t)|\mathcal{F}_t\right] \le 1.
$$

Rather than trying to prove Theorem [3.1,](#page-18-2) we now show how our previous formula  $(11)$ , i.e :

$$
\mathbb{P}\left(\mathcal{G}_K \le t | \mathcal{F}_t\right) = \left(1 - \frac{M_t}{K}\right)^+, \tag{62}
$$

or equivalently :

$$
\mathbb{P}\left(\mathcal{G}_K > t | \mathcal{F}_t\right) = \left(\frac{M_t}{K}\right) \wedge 1\tag{63}
$$

is a particular case of formula [\(60\)](#page-18-3).

<span id="page-19-2"></span>**Proposition 3.1.** Let  $M_0 \ge K$ , there is the representation :

<span id="page-19-3"></span>
$$
\left(\frac{M_t}{K}\right) \wedge 1 = \frac{N_t}{S_t},\tag{64}
$$

where

$$
\begin{cases}\nN_t &= \left(\frac{M_t}{K} \wedge 1\right) \exp\left(\frac{1}{2K} \mathcal{L}_t^K\right), \\
S_t &= \sup_{s \le t} N_s = \exp\left(\frac{1}{2K} \mathcal{L}_t^K\right).\n\end{cases} \tag{65}
$$

Proof. From Tanaka's formula :

<span id="page-19-0"></span>
$$
\frac{M_t}{K} \wedge 1 = 1 + \frac{1}{K} \int_0^t 1_{\{M_s \le K\}} dM_s - \frac{1}{2K} \mathcal{L}_t^K(M). \tag{66}
$$

The comparison of formulae [\(66\)](#page-19-0) and [\(61\)](#page-19-1) gives :

$$
\begin{cases}\n\int_0^t \frac{dN_s}{S_s} &= \frac{1}{K} \int_0^t 1_{\{M_s \le K\}} dM_s, \\
\frac{1}{2K} \mathcal{L}_t^K(M) &= \log(S_t).\n\end{cases} \tag{67}
$$

Hence :

$$
N_t = \left(\frac{M_t}{K} \wedge 1\right) S_t
$$
  
= 
$$
\left(\frac{M_t}{K} \wedge 1\right) \exp\left(\frac{1}{2K} \mathcal{L}_t^K\right).
$$

Since  $M_t \to 0$  when  $t \to \infty$ , it follows from the previous equality that :  $N_t \to 0$  when  $t \to \infty$ .

We now compare the results of Theorem [3.1](#page-18-2) and Proposition [3.1.](#page-19-2)

We remark that not every supermartingale of the form :  $(\frac{N_t}{S_t}, t \ge 0)$  can be written as  $(M_t \wedge 1)$  where  $M_0 \geq 1$  (there is no loss of generality in taking  $K = 1$ ).

Indeed, assuming [\(64\)](#page-19-3), with  $K = 1$ , we deduce that :

<span id="page-20-0"></span>
$$
d\langle N\rangle_s = \exp\left(\mathcal{L}_s^{(1)}\right)1_{\{M_s < 1\}} d\langle M\rangle_s. \tag{68}
$$

 $\Box$ 

Now, in a Brownian setting, we have  $d\langle N \rangle_s = n_s^2 ds$  and  $d\langle M \rangle_s = m_s^2 ds$ , for two  $(\mathcal{F}_s)$  previsible processes  $(m_s^2)$  and  $(n_s^2)$ . Note that [\(68\)](#page-20-0) implies :

$$
n_s^2 = \exp\left(\mathcal{L}_s^{(1)}\right) 1_{\{M_s < 1\}} m_s^2, \quad ds \, d\mathbb{P} \, \text{a.s.}
$$

Consequently,

$$
n_s^2 = 0
$$
,  $ds d\mathbb{P}$  a.s. on  $\{(s, \omega), M_s > 1\}$ .

However, this cannot be satisfied if we start from N such that  $n_s^2 > 0$ , for all  $s > 0$ .

Note that the random set  $\{s, M_s > 1\}$  is not empty; if it were, then the local time at 1 of M would be 0, and M would be identically equal to 1.

Question 3.1. It is now natural to ask the following : for which functions  $h: \mathbb{R}^+ \to [0,1],$  is it true that, for any  $(M_t, t \geq 0)$  in  $\mathcal{M}_0^+$ ,  $(h(M_t), t \geq 0)$  is an Azéma supermartingale? We shall call such a function an Azéma function.

Here is a partial answer to Question 3.1 :

**Proposition 3.2.** Assume that h is an Az $\acute{e}$ ma function such that :

- i.  $\{x : h(x) < 1\} = [0, K]$ , for some positive real K;
- ii.  $h''$  -in L.Schwartz'distribution sense- is a bounded measure;

Then :

$$
h(x) = \left(\frac{x}{K}\right) \wedge 1.
$$

*Proof.* a) From (*ii*), for any  $M \in \mathcal{M}_0^+$ , we may apply the Itô-Tanaka formula to write the canonical decomposition of  $(h(M_t), t \geq 0)$  as a semimartingale; we get :

<span id="page-21-1"></span>
$$
h(M_t) = h(M_0) + \int_0^t h'(M_s) dM_s + \frac{1}{2} \int h''(dx) \mathcal{L}_t^x(M). \tag{69}
$$

b) Since  $h(M_t)$  is an Azéma supermartingale, its increasing process in [\(69\)](#page-21-1) is carried by  $\{s : h(M_s) = 1\}$ . Therefore :

<span id="page-21-2"></span>
$$
\int h''(dx) \int_0^t 1_{\{h(M_s) < 1\}} d\mathcal{L}_s^x = 0. \tag{70}
$$

 $\Box$ 

Now, the LHS of [\(70\)](#page-21-2) equals :

$$
\int h''(dx) \; 1_{\{h(x) < 1\}} \mathcal{L}_t^x(M) = \int_{[0,K]} h''(dx) \mathcal{L}_t^x(M),
$$

as a consequence of (i). This is equivalent to :  $h''(dx) = 0$ , on  $[0, K],$  thus :  $h(x) = ax + b$ , on  $[0, K]$ ; furthermore,  $h(0) = 0$ , since :  $\lim_{t\to\infty} h(M_t) = 0$ , for any  $M \in \mathcal{M}_0^+$ . Thus :  $h(x) = ax$ , on  $[0, K[,$ and, applying (i) again yields to the result.

**Question 3.2.** Is it possible to relax further the hypotheses  $(i)$  and  $(ii)$ ?

## <span id="page-21-0"></span>3.2 Enlargement of filtration formulae

Under  $(CA)$ , there is a general expression for the transformation of a generic  $(\mathcal{F}_t)$ -martingale  $(\mu_t)$  into a  $(\mathcal{F}_t^L)$  semimartingale, where  $(\mathcal{F}_t^L)$  is the smallest filtration which contains  $(\mathcal{F}_t)$  and makes L a stopping time.

Then :

<span id="page-21-3"></span>
$$
\mu_t = \tilde{\mu}_t + \int_0^{t \wedge L} \frac{d\langle \mu, Z \rangle_s}{Z_s} + \int_L^t \frac{d\langle \mu, 1 - Z \rangle_s}{(1 - Z_s)}
$$
(71)

where  $(\tilde{\mu}_t)$  is a  $(\mathcal{F}_t^L)$  local martingale.

Now, since :

$$
Z_t = \frac{N_t}{S_t},
$$

(see formula [\(60\)](#page-18-3)), formula [\(71\)](#page-21-3) becomes :

<span id="page-22-1"></span>
$$
\mu_t = \tilde{\mu}_t + \int_0^{t \wedge L} \frac{d\langle \mu, N \rangle_s}{N_s} - \int_L^t \frac{d\langle \mu, N \rangle_s}{(S_s - N_s)}.
$$
\n(72)

Particularising again with  $L = \mathcal{G}_K$ , we have seen previously that :

$$
Z_t = \left(\frac{M_t}{K}\right) \wedge 1
$$

and

$$
N_t = \left( \left( \frac{M_t}{K} \right) \wedge 1 \right) \exp \left( \frac{\mathcal{L}_t^K}{2K} \right).
$$

Hence, applying  $(71)$  and  $(72)$ , we get :

<span id="page-22-2"></span>
$$
\mu_t = \tilde{\mu}_t + \int_0^{t \wedge \mathcal{G}_K} \frac{\mathbb{1}_{\{M_s < K\}} d\langle \mu, M \rangle_s}{M_s} - \int_{\mathcal{G}_K}^t \frac{d\langle \mu, M \rangle_s}{(K - M_s)}.
$$
\n(73)

It is of some interest to take  $\mu_s = M_s$ , formula [\(73\)](#page-22-2) then becomes :

<span id="page-22-3"></span>
$$
M_t = \tilde{M}_t + \int_0^{t \wedge \mathcal{G}_K} \frac{1_{\{M_s < K\}} d\langle M \rangle_s}{M_s} - \int_{\mathcal{G}_K}^t \frac{d\langle M \rangle_s}{(K - M_s)}.
$$
\n(74)

## <span id="page-22-0"></span>3.3 Study of the pre  $\mathcal{G}_K$ - and the post  $\mathcal{G}_K$ -processes

We now apply formula [\(74\)](#page-22-3) to give a description of the pre  $\mathcal{G}_K$ -process and the post  $\mathcal{G}_K$ -process.

a) The post  $\mathcal{G}_K$ -process : From [\(74\)](#page-22-3), we may write :

$$
M_{\mathcal{G}_K+t} = K + \hat{M}_t - \int_0^t \frac{d\langle M \rangle_{\mathcal{G}_K+t}}{(K - M_{\mathcal{G}_K+t})},\tag{75}
$$

where  $(\hat{M}_t, t \geq 0)$  is a  $\mathcal{F}_{\mathcal{G}_K+t}$  local martingale starting at 0. We introduce the notations :

<span id="page-22-5"></span>
$$
R_t = K - M_{\mathcal{G}_K + t};\tag{76}
$$

we have :

<span id="page-22-4"></span>
$$
R_t = -\hat{M}_t + \int_0^t \frac{d\langle M \rangle_{\mathcal{G}_K + u}}{R_u}.\tag{77}
$$

Since :  $\langle \hat{M} \rangle_t = \langle M \rangle_{\mathcal{G}_K+t} - \langle M \rangle_{\mathcal{G}_K}$ , we may write :  $\hat{M}_t = \beta_{\langle \hat{M} \rangle_t}$ , where  $(\beta_u)$  is a Brownian motion, we deduce from [\(77\)](#page-22-4) that :  $R_t = \rho_{\langle \hat{M} \rangle_t}$ , where  $(\rho_u, u \le \langle \hat{M} \rangle_{\infty})$  is a  $BES(3)$  process, considered up to :  $\langle \hat{M} \rangle_{\infty} =$  $T_K(\rho)$ , as deduced from [\(76\)](#page-22-5), and the fact that  $M_u \to 0$  when  $u \to \infty$ . We also note that :  $\langle M \rangle_{\mathcal{G}_K} = \mathcal{G}_K(\beta_{\cdot \wedge T_0}).$ 

b) The pre  $\mathcal{G}_K$ -process :

Here we take back the notations of subsection 2.1, but in order to see precisely the situation, we drop the continuity hypotheses  $(H_1)$  and  $(H_2)$  in that subsection. Theorem [2.1,](#page-11-7) which gives the law of  $\mathcal{G}_K$  (see [36\)](#page-11-4) is now completed by the following computation of the conditional law of the pre  $\mathcal{G}_K$ -process, given  $\mathcal{G}_K$ :

<span id="page-23-0"></span>**Theorem 3.2.** Let  $(\phi_u, u \geq 0)$  denote a positive,  $(\mathcal{F}_u)$  previsible process. Then :

a)

<span id="page-23-2"></span>
$$
\mathbb{E}\left[\phi_{\mathcal{G}_K}\right] = \mathbb{E}\left[\phi_0 \left(1 - \frac{M_0}{K}\right)^+\right] + \frac{1}{2K} \int_0^\infty ds \, m_s(K) \mathbb{E}\left[\phi_s \sigma_s^2 | M_s = K\right], \quad dK \, a.e,
$$
\n(78)

b) As a consequence of a), we recover :

<span id="page-23-4"></span>
$$
\mathbb{P}\left(\mathcal{G}_K \in ds\right) = \mathbb{E}\left[\left(1 - \frac{M_0}{K}\right)^+\right] \epsilon_0(ds) + \frac{ds}{2K} m_s(K) \mathbb{E}\left[\sigma_s^2 | M_s = K\right], \quad dK \ a.e,
$$
\n(79)

c) Furthermore :

<span id="page-23-5"></span>
$$
\mathbb{P}\left(\phi_{\mathcal{G}_K}|\mathcal{G}_K=s\right) = \frac{\mathbb{E}\left[\phi_s \sigma_s^2|M_s = K\right]}{\mathbb{E}\left[\sigma_s^2|M_s = K\right]}, \quad \mathbb{P}(\mathcal{G}_K \in ds) \text{ a.e.} \tag{80}
$$

The proof hinges on the balayage formula, which we first recall :

<span id="page-23-6"></span>**Lemma 3.1.** (see [\[23\]](#page-42-6)) Let  $(Y_t)$  be a continuous semimartingale, and  $g_Y(t) = \sup\{s \le t, Y_s = 0\}$ . Then, for any bounded previsible process  $(\phi_s, s \geq 0)$ , one has :

<span id="page-23-1"></span>
$$
\phi_{g_Y(t)} Y_t = \phi_0 Y_0 + \int_0^t \phi_{g_Y(s)} dY_s. \tag{81}
$$

Proof. (of Theorem [3.2\)](#page-23-0) :

We deduce from [\(81\)](#page-23-1), i.e : the balayage formula applied to  $(K - M_t)^+$ that :

<span id="page-23-3"></span>
$$
\mathbb{E}\left[\phi_{\mathcal{G}_K}\left(K - M_{\infty}\right)^+\right] = \mathbb{E}\left[\phi_0\left(K - M_0\right)^+\right] + \frac{1}{2}\mathbb{E}\left[\int_0^{\infty} \phi_s d\mathcal{L}_s^K\right],\tag{82}
$$

Now, [\(78\)](#page-23-2) is deduced from [\(82\)](#page-23-3) : under the present hypothesis :  $M_{\infty}$  = 0, [\(82\)](#page-23-3) writes :

$$
\mathbb{E}\left[\phi_{\mathcal{G}_K}\right] = \mathbb{E}\left[\phi_0 \left(1 - \frac{M_0}{K}\right)^+\right] + \frac{1}{2K} \mathbb{E}\left[\int_0^\infty \phi_s d\mathcal{L}_s^K\right].\tag{83}
$$

Now, [\(78\)](#page-23-2) will be proven if we show :

<span id="page-24-0"></span>
$$
\mathbb{E}\left[\int_0^\infty \phi_s d\mathcal{L}_s^K\right] = \int_0^\infty ds m_s(K) \mathbb{E}\left[\phi_s \sigma_s^2 | M_s = K\right], \quad dK \ a.e. \quad (84)
$$

In order to prove [\(84\)](#page-24-0), we use the density of occupation formula [\(40\)](#page-12-2) after integrating on both sides with respect to  $\phi_s$ , which yields, using  $(H_1)$  in Section 2.1 :

$$
\int_0^\infty ds \,\sigma_s^2 f(M_s)\phi_s = \int_0^\infty dK f(K) \int_0^\infty \phi_s d\mathcal{L}_s^K. \tag{85}
$$

Taking expectations of both sides, we obtain, with the help of  $(H_2)$  in Section 2.1 :

$$
\int_0^\infty dK f(K) \int_0^\infty ds m_s(K) \mathbb{E} \left[ \sigma_s^2 \phi_s | M_s = K \right]
$$

$$
= \int_0^\infty dK f(K) \mathbb{E} \left[ \int_0^\infty \phi_s d\mathcal{L}_s^K \right], \tag{86}
$$

which is easily shown to imply [\(84\)](#page-24-0). Then, replacing in [\(78\)](#page-23-2)  $\phi_s$  by  $\phi_s g(s)$ , for a generic, Borel,  $g : \mathbb{R}^+ \to \mathbb{R}^+$ , we deduce [\(79\)](#page-23-4) and [\(80\)](#page-23-5).

The particular case when  $(M_s)$  is Markovian, e.g : the Black-Scholes situation where  $M_s = \mathcal{E}_s$ , allows for some simplification of the above formula : in this case,  $\sigma_s = \sigma(s, M_s)$ , where  $(\sigma(s, x))$  is a deterministic function on  $([0, \infty))^2$ , and we obtain, from  $(80)$ :

$$
\mathbb{E}\left[\phi_{\mathcal{G}_K}|\mathcal{G}_K=s\right]=\mathbb{E}\left[\phi_s|M_s=K\right],\tag{87}
$$

i.e : conditionally on  $\mathcal{G}_K = s$ , the pre  $\mathcal{G}_K$ -process is the bridge (for M) on the time interval  $[0, s]$ , ending at K.

Exercise 3.2. Prove Lemma [3.1](#page-23-6) by applying the monotone class theorem , i.e :

a) show it for  $\phi_s = 1_{[0,T]}(s)$ , with T a stopping time,

b) apply the monotone class theorem.

Exercise 3.3. Prove, with the help of formula [\(81\)](#page-23-1), that the following processes are local martingales :

$$
f(S_t)(S_t - M_t) - \int_0^{S_t} dx f(x),
$$

<span id="page-25-0"></span>for any bounded Borel function  $f : \mathbb{R}^+ \to \mathbb{R}$ .

### 3.4 A larger framework

We refer to [\[17\]](#page-42-7). We now wish to explain how our basic formula [\(11\)](#page-6-5) which we now write as :

$$
\mathbb{E}_{\mathbb{P}}\left[F_t\left(1-\frac{M_t}{K}\right)^+\right] = \mathbb{E}\left[F_t\,1_{\{\mathcal{G}_K\leq t\}}\right],\tag{88}
$$

for every  $F_t \geq 0$ ,  $(\mathcal{F}_t)$  measurable, is a particular case of the following representation problem for certain (Skorokhod) submartingales.

Let us consider, on a filtered space  $(\Omega, \mathcal{F}, (\mathcal{F}_t))$ :

- a) a probability  $\mathbb{P}$ , and a positive process  $(X_t)$  which is adapted to  $(\mathcal{F}_t)$ , and integrable;
- b) a  $\sigma$ -finite measure  $\mathbb Q$  on  $(\Omega, \mathcal F)$  ( $\mathbb Q$  may be finite, even a probability, but we are also interested in the more general case where  $\mathbb Q$  is  $\sigma$ -finite);
- c) a positive  $\mathcal{F}\text{-measurable random variable } \mathcal{G}$  such that :

<span id="page-25-1"></span>
$$
\forall \Gamma_t \in \mathcal{F}_t, \quad \mathbb{E}_{\mathbb{P}}\left[\Gamma_t X_t\right] = \mathbb{Q}\left(\Gamma_t 1_{\{\mathcal{G} \le t\}}\right). \tag{89}
$$

Note that it follows immediately from [\(89\)](#page-25-1) that  $(X_t)$  is a  $(\mathbb{P}, \mathcal{F}_t)$  submartingale, since, for  $(s < t)$ , and  $\Gamma_s \in \mathcal{F}_s$ :

$$
\mathbb{E}_{\mathbb{P}}\left[\Gamma_s(X_t - X_s)\right] = \mathbb{Q}\left(\Gamma_s \, 1_{\{s \le \mathcal{G} \le t\}}\right) \ge 0. \tag{90}
$$

Conversely, we would like to find out which positive submartingales  $(X_t)$ , with respect to  $(\Omega, (\mathcal{F}_t), \mathbb{P})$  may be "represented" in the form [\(89\)](#page-25-1); that is, we seek a pair  $(\mathbb{Q}, \mathcal{G})$  such that [\(89\)](#page-25-1) is satisfied.

So far we have not solved this problem in its full generality, but we have three set-ups where the problem is solved. The next three subsections are devoted to the discussion of each of these cases.

However, the three cases are concerned with what we would like to call Skorokhod submartingales, i.e :  $(X_t)$  is a submartingale, such that :

$$
X_t = -\mathfrak{M}_t + L_t, \quad t \ge 0,
$$
\n<sup>(91)</sup>

with :

1.  $X_t \geq 0$ ;  $X_0 = 0$ ;

2.  $(L_t)$  is increasing, and  $(dL_t)$  is carried by the zeros of  $(X_t, t \ge 0)$ .

As is well known, this implies that :

$$
L_t = S_t(\mathfrak{M}) \equiv \sup_{s \leq t} \mathfrak{M}_s.
$$

We assume that,  $(\mathfrak{M}_t, t \geq 0)$  is a true martingale.

The three cases we shall consider are :

i.

$$
X_t = (1 - Y_t)^+, \quad t \ge 0,
$$
\n(92)

where  $(Y_t, t \geq 0)$  is a positive martingale, which converges to 0, as  $t \to \infty$  and with  $Y_0 = 1$ .

ii.

$$
X_t = S_t(\mathfrak{N}) - \mathfrak{N}_t \quad t \ge 0,
$$
\n<sup>(93)</sup>

where  $(\mathfrak{N}_t, t \geq 0)$  is a positive martingale, with  $\mathfrak{N}_0 = 1$ , and which converges to 0, as  $t \to \infty$ .

iii.

$$
X_t = |B_t|, \quad t \ge 0,\tag{94}
$$

where  $(B_t)$  is a standard Brownian motion.

#### <span id="page-26-0"></span>3.4.1 Case 1

Denote :

$$
G = \sup\{t, Y_t = 1\} = \sup\{t, X_t = 0\}.
$$

Then, we have shown that (see Theorem [1.1\)](#page-6-3) :

$$
\mathbb{P}\left(\mathcal{G}\leq t|\mathcal{F}_t\right)=\left(1-Y_t\right)^+\,. \tag{95}
$$

Therefore, in this case, we may write :

$$
\mathbb{E}\left[\Gamma_t X_t\right] = \mathbb{E}\left[\Gamma_t 1_{\{\mathcal{G}\leq t\}}\right],\tag{96}
$$

for every  $\Gamma_t \in \mathcal{F}_t$ . Thus,  $\mathbb{Q} = \mathbb{P}$  is convenient in this situation.

#### <span id="page-27-0"></span>3.4.2 Case 2

Again, we introduce :

$$
\mathcal{G} = \sup\{t, \mathfrak{N}_t = S_t(\mathfrak{N})\} = \sup\{t, X_t = 0\}.
$$

We have (see Theorem [3.1](#page-18-2) and Proposition [3.1\)](#page-19-2) :

$$
\mathbb{P}\left(\mathcal{G}\leq t|\mathcal{F}_t\right)=\frac{\mathfrak{N}_t}{S_t(\mathfrak{N})},\tag{97}
$$

and thus :

<span id="page-27-2"></span>
$$
\mathbb{E}\left[\Gamma_t\left(1-\frac{\mathfrak{N}_t}{S_t}\right)\right] = \mathbb{E}\left[\Gamma_t 1_{\{\mathcal{G}\leq t\}}\right].\tag{98}
$$

Since [\(98\)](#page-27-2) is valid for every  $\Gamma_t \in \mathcal{F}_t$ , and  $t \geq 0$ , we may write (98) in the equivalent form :

<span id="page-27-3"></span>
$$
\mathbb{E}\left[\Gamma_t\left(S_t - \mathfrak{N}_t\right)\right] = \mathbb{E}\left[\Gamma_t S_t \, 1_{\{\mathcal{G} \le t\}}\right].\tag{99}
$$

However, on  $(\mathcal{G} \leq t)$ , we have :  $S_t = S_{\infty}$ . Therefore, [\(99\)](#page-27-3) writes :

$$
\mathbb{E}\left[\Gamma_t\left(S_t - \mathfrak{N}_t\right)\right] = \mathbb{E}\left[\Gamma_t S_{\infty} \mathbf{1}_{\{\mathcal{G} \le t\}}\right],\tag{100}
$$

and a solution to [\(89\)](#page-25-1) is :

$$
\mathbb{Q} = S_{\infty} \cdot \mathbb{P}.\tag{101}
$$

However, we should note that  $\mathbb Q$  has infinite total mass, since :

$$
\mathbb{P}(S_{\infty} \in dt) = \frac{dt}{t^2} 1_{t \ge 1},
$$

i.e, from Lemma [1.1](#page-6-4) :

$$
S_{\infty} \stackrel{\text{law}}{=} \frac{1}{\mathbf{U}} \text{ with } \mathbf{U} \text{ uniform on } [0, 1].
$$

#### <span id="page-27-1"></span>3.4.3 Case 3

This study has been the subject of many considerations within the penalisation procedures of Brownian paths studied in [\[25\]](#page-42-8) and [\[18\]](#page-42-9).

In fact, on the canonical space  $\mathcal{C}(\mathbb{R}^+, \mathbb{R})$ , where we now denote  $(x_t, t \geq 0)$ as the coordinate process, and  $\mathcal{F}_t = \sigma\{x_s, s \leq t\}$ , then, if W denotes the Wiener measure, a  $\sigma$ -finite measure W has been contructed in [\[25\]](#page-42-8) and [\[18\]](#page-42-9) such that :

$$
\forall \Gamma_t \in \mathcal{F}_t, \quad \mathbb{W}\left(\Gamma_t | x_t\right) = \mathcal{W}\left(\Gamma_t 1_{\mathcal{G} \le t}\right),\tag{102}
$$

where  $\mathcal{G} = \sup\{s, x_s = 0\}$  is finite a.s under W. Thus, now a solution to [\(89\)](#page-25-1) is :

 $\mathbb{Q} = \mathcal{W}$ .

We note that  $W$  and  $W$  are naturally singular.

### <span id="page-28-0"></span>3.4.4 A comparative analysis of the three cases

We note that in case 1 and case 2,  $\{X_t, t \to \infty\}$  converges  $\mathbb P$  a.s. and that the solution to [\(89\)](#page-25-1) may be written, in both cases :

$$
\mathbb{E}\left[X_t\Gamma_t\right] = \mathbb{E}\left[X_{\infty}1_{\mathcal{G}\leq t}\right],\tag{103}
$$

where :  $\mathcal{G} = \sup\{t, X_t = 0\}$ . Is this the general case for Skorokhod submartingales which converge a.s.?

# <span id="page-29-0"></span>4 Note 4 : How are the previous results modified when  $M_{\infty} \neq 0$ ?

In this note, we work again with a continuous local martingale  $(M_t)$  taking values in  $\mathbb{R}^+$ , and starting from  $a > 0$ . We do not assume that  $M_\infty = 0$ ; thus :

$$
\mathbb{P}(M_{\infty} > 0) > 0.
$$

We ask a first question : can we describe the law of  $\sup_{t>0} M_t$ ? Also can we describe the law of  $\mathcal{G}_K = \sup\{t, M_t = K\}$ ?

## <span id="page-29-1"></span>4.1 On the law of  $S_{\infty} = \sup_{t>0} M_t$

Note that we cannot use the Dubins-Schwarz theorem :

$$
M_t = \beta_{\langle M \rangle_t}, \quad t \ge 0,
$$

in an efficient way, since in that generality,  $\langle M \rangle_{\infty}$  cannot be interpreted in terms of  $\beta$ .

Nonetheless, let us see how our argument involving Doob's optional stopping theorem (see lemma [1.1\)](#page-6-4) may be modified.

Let  $b > a = M_0$ , and  $T_b = \inf\{t, M_t = b\}$ . Then

$$
\mathbb{E}\left[M_{T_b}\right] = a,
$$

that is :

<span id="page-29-2"></span>
$$
b\,\mathbb{P}\left(S_{\infty}\geq b\right)+\mathbb{E}\left[M_{\infty}1_{\{S_{\infty}
$$

This leads us naturally to replace  $M_{\infty}$  by :

<span id="page-29-6"></span>
$$
\phi\left(S_{\infty}\right) = \mathbb{E}\left[M_{\infty}|S_{\infty}\right],\tag{105}
$$

with  $\phi(x) \leq x$ . Formula [\(104\)](#page-29-2) now becomes :

<span id="page-29-3"></span>
$$
b\,\mathbb{P}\left(S_{\infty}\geq b\right)+\mathbb{E}\left[\phi(S_{\infty})1_{\{S_{\infty}
$$

Assuming  $\phi$  as given, we consider [\(106\)](#page-29-3) as an equation for the distribution of  $S_{\infty}$ , and we obtain :

<span id="page-29-5"></span>**Proposition 4.1.** For simplicity, we assume that :  $\forall b > 0, \phi(b) < b$ . The law of  $S_{\infty}$  is given by :

<span id="page-29-4"></span>
$$
\mathbb{P}(S_{\infty} \ge b) = \exp\left(-\int_{a}^{b} \frac{dx}{x - \phi(x)}\right).
$$
 (107)

Comment : since  $S_{\infty} < \infty$  a.s., it follows from [\(107\)](#page-29-4) that :

$$
\int_{a}^{\infty} \frac{dx}{x - \phi(x)} = \infty.
$$
 (108)

*Proof.* of Proposition [4.1](#page-29-5) : from formula [\(106\)](#page-29-3), denoting  $\overline{\mu}(b) = \mathbb{P}(S_{\infty} \geq b)$ , we obtain :

$$
b\overline{\mu}(b) - \int_{a}^{b} d\overline{\mu}(x)\phi(x) = a.
$$
 (109)

Consequently :

$$
bd\overline{\mu}(b) - d\overline{\mu}(b)\phi(b) + db\overline{\mu}(b) = 0
$$

$$
(b - \phi(b))d\overline{\mu}(b) = -(db)\overline{\mu}(b).
$$

Then, the above equation yields :

$$
\overline{\mu}(b) = C \exp\left(-\int_{a}^{b} \frac{dx}{x - \phi(x)}\right),\tag{110}
$$

which implies  $C = 1$  by taking  $b = a$ .

<span id="page-30-0"></span>**Example 4.1.** We consider  $(B_t)$  issued from  $a > 0$ , and for  $\alpha < 1$ :

$$
T_a^{(\alpha)} = \inf\{t, B_t = \alpha S_t\},\tag{111}
$$

 $\Box$ 

to which we associate  $M_t = B_{t \wedge T_a^{(\alpha)}}$ . Then,  $\phi(x) = \alpha x$ ; consequently we have :

$$
\int_{a}^{b} \frac{dx}{(1-\alpha)x} = \frac{1}{1-\alpha} \log \left(\frac{b}{a}\right).
$$

Hence,

$$
\overline{\mu}(b) = \exp\left(-\frac{1}{1-\alpha}\log\left(\frac{b}{a}\right)\right)
$$

$$
= \left(\frac{a}{b}\right)^{1/(1-\alpha)}, \quad b \ge a,
$$

and :

$$
d\mu(b) = a^{1/(1-\alpha)} \left(\frac{\alpha}{1-\alpha}\right) \frac{db}{b^{\frac{2-\alpha}{1-\alpha}}} 1_{\{b \ge a\}}.
$$
 (112)

Question 4.1. Can we describe all the laws of  $(M_t, t \geq 0)$  which satisfy [\(105\)](#page-29-6) for a given  $\phi$ ? See Rogers [\[24\]](#page-42-10) where the law of  $(S_{\infty}, M_{\infty})$  is described in all generality...See also P.Vallois [\[26\]](#page-42-11). However, these authors assume that M is uniformly integrable...

**Question 4.2.** Under which condition(s) is  $(M_t, t \ge 0)$  uniformly integrable? (this question had a negative answer when  $M_{\infty} = 0$ , but now...?) A first answer :

We shall have

$$
\mathbb{E}[M_{\infty}] = a,
$$

which is satisfied if only if :

$$
\mathbb{E}\left[\phi(S_{\infty})\right] = a,
$$

i.e :

<span id="page-31-0"></span>
$$
\int_{a}^{\infty} dx \exp\left(-\int_{a}^{x} \frac{dy}{y - \phi(y)}\right) \frac{\phi(x)}{x - \phi(x)} = a.
$$
 (113)

In fact there is a more direct criterion which may be derived from  $(104)$ :

$$
\lim_{b \to \infty} b \, \mathbb{P}(S_{\infty} \ge b) = 0 \tag{114}
$$

and which amounts to :

<span id="page-31-1"></span>
$$
\lim_{b \to \infty} b \exp\left(-\int_a^b \frac{dx}{x - \phi(x)}\right) = 0. \tag{115}
$$

Note that, in all generality, it follows from  $(104)$  that :

$$
\lim_{b \to \infty} b \mathbb{P} \left( S_{\infty} \ge b \right) = a - \mathbb{E} \left[ M_{\infty} \right]
$$

Exercise 4.1. Prove that [\(113\)](#page-31-0) is equivalent to [\(115\)](#page-31-1).(Probably, integration by parts).

<span id="page-31-2"></span>**Example 4.2.** Going back to Example [4.1,](#page-30-0) when  $\phi(x) = \alpha x$ ,  $\alpha < 1$ , we get :

$$
\overline{\mu}(b) = C \frac{1}{b^{1/(1-\alpha)}}.
$$
\n(116)

Then, Example [4.2](#page-31-2) is a case of uniform integrability.

**Exercise 4.2.** Denote by  $\mathcal{M}^+$  the set of positive local martingales  $(M_t, t \geq 0)$ such that  $M_0 = 1$  and by :

$$
\mathcal{M}^{+,c} = \{ M \in \mathcal{M}^+, \lim_{b \to \infty} b \mathbb{P} \left( S_{\infty} \ge b \right) = 1 - c \}.
$$

a) Prove that :

$$
\mathcal{M}^+=\bigcup_{0\leq c\leq 1}\mathcal{M}^{+,c};
$$

(of course, this is a union of disjoint sets).

- b) Prove that  $c = 1$  iff  $(M_t)$  is ui.;
- c) Prove that  $c = 0$  iff  $M_t \rightarrow_{t \rightarrow \infty} 0$ ;
- d) Prove that for any  $c \in [0,1]$ ,  $M \in \mathcal{M}^{+,c}$  iff  $\mathbb{E}[M_\infty] = c$ .
- e) For any  $c \in [0,1]$ , give as many examples as possible of elements of  $\mathcal{M}^{+,c}$ .

Comments : A somewhat related discussion about the asymptotic behavior of  $\mathbb{P}(S_{\infty} \geq b)$  as well as that of  $\mathbb{P}(\langle M \rangle_{\infty} \geq b)$  is done in [\[2\]](#page-41-8).

Exercise 4.3. Give some examples of non uniform integrability obtained from the criterion [\(115\)](#page-31-1).

At this point, it is very natural to recall Azema-Yor's solution of Skorokhod's embedding as given in [\[3\]](#page-41-9) :

if  $\nu(dx)$  is a probability on  $\mathbb{R}$ , with  $\int \nu(dx) |x| < \infty$ , and  $\int \nu(dx)x = 0$ , then the stopping time :

$$
T_{\nu} = \inf\{t \ge 0, S_t \ge \psi_{\nu}(B_t)\},\
$$

where  $S_t = \sup_{s \le t} B_s$ ,  $B_0 = 0$ , and  $\psi_{\nu}(x) = \frac{1}{\nu(x,\infty)} \int_{[x,\infty)} \nu(dy) y$  solves Skorokhod's embedding problem, in that :  $B_{T_{\nu}} \sim \nu$ , and  $(B_{t \wedge T_{\nu}}, t \ge 0)$  is uniformly integrable. See Obloj [\[21\]](#page-42-12) for a thorough survey of Skorokhod's problem.

**Exercise 4.4.** Modify the Azema-Yor construction to obtain as many stopping times  $T'_\mu$  of Brownian motion  $(B_t, t \leq T_0)$ , where  $B_0 = 1$ , such that  $B_{T'_{\mu}} \sim \mu.$ 

More generally, one may ask :

**Question 4.3.** Given a stopping time T of  $(B_t, t \ge 0)$ , describe the set  $S_T$ of all the laws of B<sub>S</sub>, for all stopping times  $S \leq T$ , such that  $(B_{t \wedge S}, t \geq 0)$  is uniformly integrable.

## <span id="page-32-0"></span>4.2 Extension of our representation theorem in the case  $M_{\infty} \neq 0$

We now try to see how the formula (see  $(11)$ ):

$$
\mathbb{P}\left(\mathcal{G}_K \le t | \mathcal{F}_t\right) = \left(1 - \frac{M_t}{K}\right)^{+}
$$
\n(117)

is modified in the case  $M_{\infty} \neq 0$ .

Theorem 4.1. The following formula holds:

<span id="page-33-0"></span>
$$
\mathbb{E}\left[1_{\{\mathcal{G}_K\leq t\}}\left(K-M_{\infty}\right)^{+}|\mathcal{F}_t\right] = \left(K-M_t\right)^{+}.
$$
 (118)

Proof. We may prove formula [\(118\)](#page-33-0) in different ways.

First proof : It hinges on the balayage formula (see Lemma [3.1\)](#page-23-6) applied to  $Y_t = (K - M_t)^+$ ; we note :  $\mathcal{G}_K(s) = \sup\{u \leq s, M_u = K\}$ . The balayage formula [\(81\)](#page-23-1) now becomes :

<span id="page-33-2"></span>
$$
\phi_{\mathcal{G}_K(t)}(K - M_t)^+ = \phi_0(K - M_0)^+ - \int_0^t \phi_{\mathcal{G}_K(s)} 1_{\{M_s < K\}} dM_s + \frac{1}{2} \int_0^t \phi_s d\mathcal{L}_s^K,\tag{119}
$$

since  $d\mathcal{L}_s^K$  charges only the set of times for which  $M_s = K$ , i.e for which  $\mathcal{G}_K(s) = s.$ 

This formula applied between t and  $\infty$  yields :

<span id="page-33-1"></span>
$$
\mathbb{E}\left[\phi_{\mathcal{G}_K}\left(K - M_{\infty}\right)^+|\mathcal{F}_t\right] = \phi_{\mathcal{G}_K(t)}\left(K - M_t\right)^+ + \frac{1}{2}\mathbb{E}\left[\int_t^{\infty} \phi_s d\mathcal{L}_s^K|\mathcal{F}_t\right] \tag{120}
$$

Taking  $\phi_s = 1_{\{s \le t\}}$  and observing that  $\mathcal{G}_K(t) \le t$  and that  $\int_t^\infty 1_{\{s \le t\}} d\mathcal{L}_s^K =$ 0, we obtain :

$$
\mathbb{E}\left[1_{\{\mathcal{G}_K\leq t\}}\left(K-M_{\infty}\right)^{+}|\mathcal{F}_t\right] = \left(K-M_t\right)^{+}.
$$
 (121)

**Second proof :** We consider for  $T$  a stopping time :

$$
\mathbb{E}\left[1_{\{\mathcal{G}_K \leq T\}}\left(K - M_{\infty}\right)^{+}\right] = \mathbb{E}\left[1_{\{d_T = \infty\}}\left(K - M_{\infty}\right)^{+}\right],\tag{122}
$$

where  $d_T (\equiv d_T^K) = \inf\{t > T, M_t = K\}.$ Then,

$$
\mathbb{E}\left[1_{\{d_T=\infty\}}\left(K-M_{\infty}\right)^+\right] = \mathbb{E}\left[1_{\{d_T=\infty\}}\left(K-M_{d_T}\right)^+\right] \n= \mathbb{E}\left[\left(K-M_{d_T}\right)^+\right]
$$

We now note that, between T and  $d_T$ ,  $(\mathcal{L}_t^K)$  does not increase; hence, from Tanaka's formula, the previous quantity equals :

$$
\mathbb{E}\left[\left(K-M_T\right)^+\right].
$$

Therefore, we have obtained :

$$
\mathbb{E}\left[1_{\{\mathcal{G}_K \leq T\}}(K - M_{\infty})^+\right] = \mathbb{E}\left[(K - M_T)^+\right].\tag{123}
$$

This identity may be reinforced as :

$$
\mathbb{E}\left[1_{\{\mathcal{G}_K \leq T\}}\left(K - M_{\infty}\right)^{+}|\mathcal{F}_T\right] = \left(K - M_T\right)^{+}.
$$
 (124)

 $\Box$ 

## <span id="page-34-0"></span>4.3 On the law of  $\mathcal{G}_K$

It is quite natural in this section to introduce the conditional law :  $\nu_K(dm)$ of  $M_{\infty}$  given<sup>[5](#page-34-1)</sup>  $\mathcal{F}_{(\mathcal{G}_K)^{-}}$ , i.e :

$$
\mathbb{E}\left[f(M_{\infty})|\mathcal{F}_{(\mathcal{G}_K)^{-}}\right] = \int \nu_K(dm) f(m). \tag{125}
$$

In fact, it is the predictable process  $(\mu_u \equiv \mu_u^{(K)}, u \ge 0)$  defined via:

$$
\mathbb{E}\left[\left(K - M_{\infty}\right)^{+}|\mathcal{F}_{(\mathcal{G}_K)^{-}}\right] = \mu_{\mathcal{G}_K} = \int \nu_K(dm)\left(K - m\right)^{+},\tag{126}
$$

which will play an important role in the sequel.

**Theorem 4.2.** In the general case  $M_{\infty} \neq 0$ , the Azéma supermartingale :

$$
Z_t^K = \mathbb{P}\left(\mathcal{G}_K > t | \mathcal{F}_t\right)
$$

is given by :

<span id="page-34-3"></span>
$$
Z_t^K = \mathbb{E}\left[\frac{(K - M_{\infty})^+}{\mu_{\mathcal{G}_K}}|\mathcal{F}_t\right] - \frac{(K - M_t)^+}{\mu_{\mathcal{G}_K(t)}}.\tag{127}
$$

*Proof.* We start from [\(120\)](#page-33-1), which we write (for  $t = 0$ ) as :

$$
\mathbb{E}\left[\phi_{\mathcal{G}_K}\,\mu_{\mathcal{G}_K}\right] = \mathbb{E}\left[\phi_0\,(K - M_0)^+\right] + \frac{1}{2}\mathbb{E}\left[\int_0^\infty \phi_s d\mathcal{L}_s^K\right].\tag{128}
$$

Replacing  $(\phi_u \mu_u)$ , by  $(\phi_u)$ , this identity writes :

<span id="page-34-2"></span>
$$
\mathbb{E}\left[\phi_{\mathcal{G}_K}\right] = \mathbb{E}\left[\frac{\phi_0}{\mu_0}\left(K - M_0\right)^+\right] + \frac{1}{2}\mathbb{E}\left[\int_0^\infty \phi_s \frac{d\mathcal{L}_s^K}{\mu_s}\right].\tag{129}
$$

Then, applying formula [\(129\)](#page-34-2) to  $\phi_u \equiv 1_{[0,T]}(u)$ , with T a generic stopping time, we obtain :

$$
\mathbb{P}\left(\mathcal{G}_K \leq T\right) = \mathbb{E}\left[\frac{1}{\mu_0}\left(K - M_0\right)^+\right] + \frac{1}{2}\mathbb{E}\left[\int_0^T \frac{d\mathcal{L}_s^K}{\mu_s}\right]
$$
\n
$$
= \mathbb{E}\left[\frac{\left(K - M_T\right)^+\right]}{\mu_{\mathcal{G}_K(T)}}\right],\tag{130}
$$

<span id="page-34-4"></span>from the balayage formula. We shall now deduce formula [\(127\)](#page-34-3) from [\(130\)](#page-34-4) : to a set  $\Gamma_t \in \mathcal{F}_t$ , we associate the stopping time :

$$
T = \begin{cases} t, & \text{on } \Gamma_t \\ \infty, & \text{on } \Gamma_t^c. \end{cases}
$$
 (131)

<span id="page-34-1"></span> ${}^{5} \mathcal{F}_{(\mathcal{G}_K)^{-}} = \sigma\{H_{\mathcal{G}_K}; H \text{predictable}\}.$ 

Then, formula [\(130\)](#page-34-4) yields :

$$
\mathbb{E}\left[1_{\Gamma_{t}}1_{\{\mathcal{G}_{K}\leq t\}}\right]+\mathbb{E}\left[1_{\Gamma_{t}^{c}}\right]=\mathbb{E}\left[1_{\Gamma_{t}}\frac{\left(K-M_{t}\right)^{+}}{\mu_{\mathcal{G}_{K}\left(t\right)}}\right]+\mathbb{E}\left[1_{\Gamma_{t}^{c}}\frac{\left(K-M_{\infty}\right)^{+}}{\mu_{\mathcal{G}_{K}}}\right],\tag{132}
$$

which, by simply writing :  $1_{\Gamma_t^c} \equiv 1 - 1_{\Gamma_t}$ , we may write equivalently as :

$$
\mathbb{E}\left[1_{\Gamma_t}1_{\{\mathcal{G}_K>t\}}\right] = \mathbb{E}\left[1_{\Gamma_t}\left(\frac{\left(K-M_{\infty}\right)^+}{\mu_{\mathcal{G}_K}} - \frac{\left(K-M_t\right)^+}{\mu_{\mathcal{G}_K(t)}}\right)\right].\tag{133}
$$

This easily implies formula [\(127\)](#page-34-3).

It may be worth giving other expressions than [\(127\)](#page-34-3) for the supermartingale :

$$
Z_t^K \equiv \mathbb{P}\left(\mathcal{G}_K > t | \mathcal{F}_t\right).
$$

Note that, if we develop  $\left(\frac{(K-M_t)^+}{\mu} \right)$  $\frac{\Lambda - M_t}{\mu_{\mathcal{G}_K(t)}}$ ,  $t \geq 0$ , then, again due to the balayage formula, we obtain :

$$
Z_t^K = \frac{1}{2} \left( \mathbb{E} \left[ \int_0^\infty \frac{d\mathcal{L}_s^K}{\mu_s} | \mathcal{F}_t \right] - \int_0^t \frac{d\mathcal{L}_s^K}{\mu_s} \right). \tag{134}
$$

In a similar vein, in order to apply formula [\(127\)](#page-34-3), one needs to know how to compute the process  $(\mu_u, u \geq 0)$ . Now writing :

$$
(K - M_{\infty})^{+} = (K - M_{0})^{+} - \int_{0}^{\infty} 1_{\{M_{s} < K\}} dM_{s} + \frac{1}{2} \mathcal{L}_{\infty}^{K},
$$

we obtain :

<span id="page-35-0"></span>
$$
\mathbb{E}\left[\left(K - M_{\infty}\right)^{+}|\mathcal{F}_{(\mathcal{G}_K)^{-}}\right] = \left(K - M_0\right)^{+} - \mathbb{E}\left[\int_0^{\infty} \mathbf{1}_{\{M_s < K\}} dM_s|\mathcal{F}_{(\mathcal{G}_K)^{-}}\right] + \frac{1}{2}\mathcal{L}_{\infty}^K,\tag{135}
$$

since  $\mathcal{L}_{\infty}^K \equiv \mathcal{L}_{\mathcal{G}_K}^K$ . Thus, if we denote by  $(\gamma_u, u \ge 0)$  a previsible process such that :

<span id="page-35-1"></span>
$$
\mathbb{E}\left[\int_{\mathcal{G}_K}^{\infty} 1_{\{M_s < K\}} dM_s |\mathcal{F}_{(\mathcal{G}_K)^-}\right] = \gamma_{\mathcal{G}_K},\tag{136}
$$

we then deduce from [\(135\)](#page-35-0) that :

$$
\mu_{\mathcal{G}_K} = (K - M_0)^+ - \int_0^{\mathcal{G}_K} 1_{\{M_s < K\}} \, dM_s - \gamma_{\mathcal{G}_K} + \frac{1}{2} \mathcal{L}_{\infty}^K,\tag{137}
$$

that is :

$$
\mu_u = (K - M_0)^+ - \int_0^u 1_{\{M_s < K\}} dM_s - \gamma_u + \frac{1}{2} \mathcal{L}_u^K. \tag{138}
$$

Thus, we have shown that the computation of  $(\mu_u)$  is equivalent to that of  $(\gamma_u)$ , as defined implicitly in [\(136\)](#page-35-1).

 $\Box$ 

## <span id="page-36-0"></span>5 Note 5 : Let K vary...

In this note, we develop formulae to compute the dual predictable projections of certain raw (i.e : non adapted) increasing processes.

Precisely, if  $(R_t, t \geq 0)$  is a raw increasing process, there exists a unique predictable increasing process  $(A_t, t \ge 0)$  such that :  $\forall (\phi_t) \ge 0$ , predictable,

$$
\mathbb{E}\left[\int_0^\infty \phi_s \, dR_s\right] = \mathbb{E}\left[\int_0^\infty \phi_s \, dA_s\right].\tag{139}
$$

We shall always assume that :  $R_0 = 0$  and  $A_0 = 0$ . In the Strasbourg terminology,  $(A_t)$  is called the predictable dual projection  $(pdp)$  of  $(R_t)$ .

## <span id="page-36-1"></span>5.1 Some predictable dual projections under the hypothesis  $M_{\infty} = 0$

**Theorem 5.1.** a) For any  $K > 0$ ,  $1_{\{0 < \mathcal{G}_K \le t\}}$  admits as pdp  $\frac{1}{2K} \mathcal{L}_t^K$ .

b) Let  $S'_t = S_\infty - S_{(t,\infty)}$ , where  $S_{(t,\infty)} = \sup_{u \geq t} M_u$ , and  $S_\infty = S_{(0,\infty)} =$  $\sup_{u\geq 0} M_u$ . Then :  $(S'_t)$  admits as pdp  $\frac{1}{2} \int_0^t$  $d\langle M\rangle_s$  $\frac{M}{M_s}$  (with the convention that  $\frac{1}{M_s} = 0$ , for  $s \ge T_0(M)$ ).

*Proof.* 1. From formula [\(120\)](#page-33-1), we get, for  $\phi \geq 0$ , predictable, with  $\phi_0 = 0$ :

$$
\mathbb{E}\left[\phi_{\mathcal{G}_K} K\right] = \mathbb{E}\left[\frac{1}{2} \int_0^\infty d\mathcal{L}_s^K \phi_s\right].\tag{140}
$$

We have obtained a)

2. We now integrate both sides of this identity with respect to  $dK f(K)$ ,  $f \geq 0$ , Borel. Then :

<span id="page-36-2"></span>
$$
\mathbb{E}\left[\int_0^\infty dK f(K) \,\phi_{\mathcal{G}_K} K\right] = \mathbb{E}\left[\frac{1}{2} \int_0^\infty dK f(K) \int_0^\infty d\mathcal{L}_s^K \,\phi_s\right]. \tag{141}
$$

From the density of occupation formula [\(35\)](#page-11-5), the RHS is equal to :

<span id="page-36-3"></span>
$$
\frac{1}{2} \mathbb{E} \left[ \int_0^\infty d\langle M \rangle_s f(M_s) \phi_s \right]. \tag{142}
$$

We note that :

$$
(\mathcal{G}_K \leq t) = (S_{(t,\infty)} \leq K),
$$

i.e : the inverse of :  $K \to \mathcal{G}_K$  is :  $t \to S_{(t,\infty)}$ . As a consequence, we may express the  $LHS$  of [\(141\)](#page-36-2) as :

$$
\mathbb{E}\left[\int_0^\infty dS'_t\, f\left(S_{(t,\infty)}\right)S_{(t,\infty)}\,\phi_t\right],\tag{143}
$$

which we compare with [\(142\)](#page-36-3). Taking  $f(x) = 1/x$ , we obtain:

<span id="page-37-0"></span>
$$
\mathbb{E}\left[\int_0^\infty dS'_t \,\phi_t\right] = \frac{1}{2} \mathbb{E}\left[\int_0^\infty \frac{d\langle M\rangle_t}{M_t} \,\phi_t\right],\tag{144}
$$

which translates b).

As a check, we would like to show ( more directly, or in a different manner than above) that for a "good" martingale, with  $M_0 = 1$ , there is the identity :

$$
\mathbb{E}\left[S'_t\right] = \frac{1}{2} \mathbb{E}\left[\int_0^t \frac{d\langle M \rangle_s}{M_s}\right].\tag{145}
$$

First, we note that the  $RHS$  is equal, from Itô's formula, applied to :  $\phi(x) = x \log(x) - x$ , to :

$$
1 + \mathbb{E}\left[\phi(M_t)\right] = 1 + \mathbb{E}\left[M_t \log(M_t)\right] - 1\tag{146}
$$

Consequently, we wish to show :

$$
\mathbb{E}\left[S_t'\right] = \mathbb{E}\left[M_t \log(M_t)\right].\tag{147}
$$

The  $LHS$  is equal to:

$$
\mathbb{E}\left[S_{\infty} - S_{(t,\infty)}\right] = \mathbb{E}\left[S_t \vee \left(\frac{M_t}{\mathbf{U}}\right) - \frac{M_t}{\mathbf{U}}\right]
$$

$$
= \int_0^1 \frac{du}{u} \mathbb{E}\left[(uS_t - M_t)^+\right]
$$

$$
= \mathbb{E}\left[\int_0^{S_t} \frac{dv}{v}(v - M_t)^+\right].
$$

Now, we note that, for  $0 < a < b$ :

$$
\int_0^b \frac{dv}{v}(v-a)^+ = \int_a^b \frac{dv}{v}(v-a) = (b-a) - a\log\left(\frac{b}{a}\right).
$$
 (148)

Consequently, we need to show :

$$
\mathbb{E}\left[M_t \log(M_t)\right] = \mathbb{E}\left[\left(S_t - M_t\right) - M_t \log(S_t) + M_t \log(M_t)\right],\tag{149}
$$

 $\Box$ 

that is :

$$
0 = \mathbb{E}\left[ (S_t - M_t \log(S_t)) - M_t \right],\tag{150}
$$

which follows from the fact that  $\{S_t - M_t \log(S_t), t \geq 0\}$  is a martingale. Proof :

$$
M_t \log(S_t) = \int_0^t M_s \frac{dS_s}{S_s} + \int_0^t (\log(S_s)) dM_s,
$$

hence :

$$
S_t - M_t \log(S_t) = 1 - \int_0^t \log(S_s) dM_s.
$$

## <span id="page-38-0"></span>5.2 A comparison with the property :  $S_{\infty} \sim M_0/U$ .

We consider [\(144\)](#page-37-0), with  $\phi_t = f(M_t)$ ; we note that the LHS of (144) is :

$$
\mathbb{E}\left[\int_0^\infty d_t(S_{(t,\infty)})f(S_{(t,\infty)})\right] = 2\mathbb{E}\left[\int_0^{S_\infty} dx\,f(x)\right]
$$

$$
= 2\int_0^\infty dx\,f(x)\left(\frac{a}{x}\wedge 1\right).
$$
(151)

Going back to [\(144\)](#page-37-0), we now see that (using again our hypotheses  $(H_1)$  and  $(H_2)$  in Section 2.1):

$$
\mathbb{E}\left[\int_0^\infty dt \,\sigma_t^2 \,\frac{f(M_t)}{M_t}\right] = \int_0^\infty dt \int_0^\infty dK \, m_t(K) \,\frac{1}{K} \,\theta_t(K) \, f(K). \tag{152}
$$

Hence, we have obtained :

<span id="page-38-2"></span>
$$
\left(\frac{a}{K} \wedge 1\right) = \int_0^\infty dt \, \frac{m_t(K)}{2K} \theta_t(K). \tag{153}
$$

Now this identity agrees with the expression of the law of  $\mathcal{G}_K$ , given by  $(36)$ : the RHS of  $(153)$  is equal to:

$$
\mathbb{P}\left(\mathcal{G}_K > 0\right) = \left(\frac{a}{K}\right) \wedge 1. \tag{154}
$$

## <span id="page-38-1"></span>5.3 Some predictable dual projections in the general case  $M_{\infty} \neq 0$

Starting again from [\(119\)](#page-33-2), we obtain :

$$
\mathbb{E}\left[\phi_{\mathcal{G}_K}\left(K - M_{\infty}\right)^+\right] = \mathbb{E}\left[\frac{1}{2} \int_0^{\infty} d\mathcal{L}_s^K \phi_s\right].\tag{155}
$$

We now integrate both sides of this identity with respect to  $dK f(K)$ ,  $f \geq 0$ , Borel. Then :

<span id="page-39-1"></span>
$$
\mathbb{E}\left[\int_0^\infty dK f(K)\phi_{\mathcal{G}_k}(K - M_\infty)^+\right] = \mathbb{E}\left[\frac{1}{2}\int_0^\infty dK f(K)\int_0^\infty d\mathcal{L}_s^K\phi_s\right].\tag{156}
$$

The  $RHS$  of [\(156\)](#page-39-1) is, as seen before, still equal to :

$$
\frac{1}{2}\mathbb{E}\left[\int_0^\infty d\langle M\rangle_s f(M_s)\phi_s\right].
$$

We know that, for  $K > M_{\infty}$ ,

$$
(\mathcal{G}_K = u) \quad \text{iff} \quad (K = S_{(u,\infty)}) \,,
$$

and the  $LHS$  of [\(156\)](#page-39-1) equals :

$$
\mathbb{E}\left[\int_0^\infty dS'_t f(S_{(t,\infty)})\left(S_{(t,\infty)} - M_\infty\right)^+ \phi_t\right].\tag{157}
$$

We may now state the following :

**Theorem 5.2.** a) For any  $K > 0$ ,  $1_{\{0 \leq \mathcal{G}_K \leq t\}} (K - M_{\infty})^+$  admits  $\frac{1}{2} \mathcal{L}_t^K$  as pdp.

b) Let  $I_t = (S_\infty - M_\infty)^2 - (S_{(t,\infty)} - M_\infty)^2$ . Then,  $(I_t, t \ge 0)$  admits  $(\langle M \rangle_t)$ as pdp.

#### <span id="page-39-0"></span>5.4 A global approach

In this section, we provide a functional extension of :

<span id="page-39-2"></span>
$$
\mathbb{E}\left[\left(\mathcal{E}_t - K\right)^+\right] = \mathbb{P}\left(\mathcal{G}_K^{(1/2)} \le t\right),\tag{158}
$$

where :  $\mathcal{G}_K^{(1/2)} = \sup\{t, \exp\left(B_t + \frac{t}{2}\right)$  $(\frac{t}{2})=K$ .

In fact, we prove a general version of [\(158\)](#page-39-2), relative to a continuous, positive martingale  $(M_t, t \ge 0)$ , which converges to 0, a.s., as  $t \to \infty$ , and plays the role of  $(\mathcal{E}_t, t \ge 0)$  in the Brownian case. We assume that  $\mathbb{E}_{\mathbb{P}}[M_t] \equiv$ 1.

To state our result, we need to introduce a new probability  $\mathbb{P}^{(M)}$  such that :

$$
\mathbb{P}^{(M)}|_{\mathcal{F}_t}=M_t\cdot\mathbb{P}|_{\mathcal{F}_t}
$$

Theorem 5.3. The following holds :

For every absolutely continuous  $\Phi : \mathbb{R} \to \mathbb{R}^+$ , with  $\Phi(0) = 0$ , i.e.

$$
\Phi(x) = \int_0^x dy \, \phi(y), \text{ for } \phi \in L^1_{+,loc}(\mathbb{R}^+),
$$

there is the relation :

<span id="page-40-0"></span>
$$
\mathbb{E}_{\mathbb{P}}\left[\Phi(M_t)\right] = \mathbb{E}_{\mathbb{P}^{(M)}}\left[\phi\left(\inf_{s\geq t} M_s\right)\right].\tag{159}
$$

As a consequence, for any  $K > 0$ , one has :

<span id="page-40-1"></span>
$$
\mathbb{E}_{\mathbb{P}}\left[\left(M_t - K\right)^+\right] = \mathbb{P}^{(M)}\left(\mathcal{G}_K \le t\right),\tag{160}
$$

where  $\mathcal{G}_K = \sup\{u, M_u = K\}.$ 

Proof. We write :

$$
\mathbb{E}_{\mathbb{P}} \left[ \Phi(M_t) \right] = \mathbb{E}_{\mathbb{P}} \left[ \int_0^{M_t} dy \, \phi(y) \right] \n= \mathbb{E}_{\mathbb{P}} \left[ M_t \, \phi(\mathbf{U} M_t) \right],
$$

where U is uniform and independent from M. Thus, with the help of  $\mathbb{P}^{(M)}$ , we obtain, with  $N_u = \frac{1}{M_t}$  $\frac{1}{M_{t+u}}, u \geq 0$ :

$$
\mathbb{E}_{\mathbb{P}}\left[\Phi(M_{t})\right] = \mathbb{E}_{\mathbb{P}^{(M)}}\left[\phi\left(\frac{1}{\left(N_{t}/\mathbf{U}\right)}\right)\right]
$$

$$
= \mathbb{E}_{\mathbb{P}^{(M)}}\left[\phi\left(\frac{1}{\sup_{s\geq t}N_{s}}\right)\right]
$$
(161)
$$
= \mathbb{E}_{\mathbb{P}^{(M)}}\left[\phi\left(\inf_{s\geq t}M_{s}\right)\right],
$$

<span id="page-40-2"></span>which proves formula [\(159\)](#page-40-0).

Formula [\(160\)](#page-40-1) follows by taking  $\Phi(x) = (x - K)^+$ . Then,

$$
\phi(\inf_{s\geq t} M_s) = 1_{\{\inf_{s\geq t} M_s > K\}} = 1_{\{\mathcal{G}_K \leq t\}}.
$$

We also need to justify the equality :

$$
\mathbb{E}_{\mathbb{P}^{(M)}}\left[\phi\left(\frac{1}{(N_t/\mathbf{U})}\right)\right] = \mathbb{E}_{\mathbb{P}^{(M)}}\left[\phi\left(\frac{1}{\sup_{s\geq t} N_s}\right)\right]
$$

in [\(161\)](#page-40-2) by asserting that, under  $\mathbb{P}^{(M)}$ ,  $(N_u)$  belongs to  $\mathcal{M}_0^{(+)}$ , and then use Doob's maximal identity (see Lemma [1.1\)](#page-6-4).  $\Box$ 

## <span id="page-41-1"></span>References

- <span id="page-41-8"></span><span id="page-41-0"></span>[1] J.Az $\acute{e}$ ma and M.YOR(1978). En guise d'introduction. Astérisque. Local times,52–53.
- [2] J.Az $\acute{e}$ ma, R. F. GUNDY and M. YOR.(1978–1979). Sur l'intégrabilité uniforme des martingales continues. Séminaire de Probabilités XIV. Lecture Notes in Maths. 784, 53–61.
- <span id="page-41-9"></span>[3] J.Az $\acute{\text{EM}}$ A and M.YOR(1979). Une solution simple au problème de Skorokhod. Séminaire de Probabilités XIII. Lecture Notes in Maths. 721,90– 115 and 625–633.
- <span id="page-41-6"></span><span id="page-41-2"></span>[4] A. N. BORODIN and P. SALMINEN(2002). Handbook of Brownian Motion, Facts and Formulae. Birkhaüser. 2nd edition.
- <span id="page-41-4"></span>[5] C. Donati-Martin and M. Yor(1997). Some Brownian Functionals and their Laws. The Annals of Probability. 25, no. 3, 1011–1058.
- <span id="page-41-5"></span>[6] B. DUPIRE(1994). Pricing with a smile. *Risk Magazine* 7, 17–20.
- [7] B. Dupire(1997). Pricing and hedging with a smile. In M.A.H Dempster and S.R. Pliska. Mathematics of derivative securities, Isaac Newton Institute. Cambridge University Press, 103–111.
- <span id="page-41-3"></span>[8] F. KLEBANER(2002). Option price when the stock is a semimartingale. Elec. Comm. in Prob.7, 79–83
- [9] D. MADAN and M. YOR(2006). Itô's integrated formula for strict local martingales. Séminaire de Probabilités XXXIX. Lecture Notes in Maths. 1874,157–170.
- [10] D. MADAN, B. ROYNETTE and M. YOR(2008). Option Prices as Probabilities. Finance Research Letters.5,79–87.
- [11] D. MADAN, B. ROYNETTE and M. YOR(February 2008). An alternative expression for the Black-Scholes formula in terms of Brownian first and last passage times. *Preprint. IEC Nancy.*, no. 8. Not for publication.
- <span id="page-41-7"></span>[12] D. MADAN, B. ROYNETTE and M. YOR(March 2008). From Black-Scholes formula, to local times and last passage times for certain submartingales. *Preprint. IEC Nancy.*, no. 14. Not for publication.
- [13] D. MADAN, B. ROYNETTE and M. YOR(April 2008) Unifying Black-Scholes type formulae which involve last passage times up to finite horizon. To appear in Asia-Pacific Journal of Financial Markets.
- <span id="page-42-4"></span><span id="page-42-2"></span>[14] R. Mansuy and M. Yor(2006). Random times and enlargements of filtrations in a Brownian setting. Lecture Notes in Maths. 1873. Springer.
- <span id="page-42-3"></span>[15] P. W. MILLAR(1977). Random times and decomposition theorems. Proc. of Symposia in Pure Math. **31**, 91–103.
- <span id="page-42-7"></span>[16] P. W. Millar(1978). A path decomposition for Markov processes. Ann. of Prob. 6, 345–348.
- [17] J. Najnudel, B. Roynette and M. Yor(2007). On a remarkable σ-finite measure associated to Brownian penalisations. CRAS Note. 345,no. 8, 459–466 .
- <span id="page-42-9"></span><span id="page-42-5"></span>[18] J. NAJNUDEL, B. ROYNETTE and M. YOR(Submitted in February 2008). A global view of Brownian penalisations. Monograph.
- [19] A. Nikeghbali and M. Yor(2006). Doob's maximal identity, multiplicative decompositions and enlargements of filtrations. Ill. Journal of Maths. 50, 791-814.
- <span id="page-42-12"></span><span id="page-42-1"></span>[20] D. NUALART(2006). The Malliavin Calculus. Springer. 2nd edition.
- <span id="page-42-0"></span>[21] J. Obloj(2004) The Skorokhod embedding problem and its offspring. Prob. Surveys.1, 321–392.
- [22] J. Pitman and M. Yor(1981). Bessel processes and infinitely divisible laws. In : Stochastic integrals. ed : D. Williams. Lecture Notes in Maths. 851, 285–370.
- <span id="page-42-10"></span><span id="page-42-6"></span>[23] D. Revuz and M. Yor(2001). Continuous Martingales and Brownian Motion. Springer-Verlag. 3rd edition.
- [24] L. C. G. Rogers(1993). The joint law of the maximum and terminal value of a martingale. Prob. Th. Rel. F. 94, no. 4, 451–466.
- <span id="page-42-8"></span>[25] B. ROYNETTE and M. YOR(Submitted in July 2007). Penalising Brownian paths; Rigorous results and Meta-theorems. Monograph.
- <span id="page-42-11"></span>[26] P. Vallois(1994) Sur la loi du maximum et du temps local d'une martingale uniformément intégrable. Proc. London Math. Soc.3, no. 69, 399– 427.
- <span id="page-43-1"></span><span id="page-43-0"></span>[27] D. WILLIAMS(1970). Decomposing the brownian path. Bull. Amer. Math. Soc. 76, 871–873.
- [28] M. Yor(2001). Exponential functionals of Brownian motion and Related processes. Springer-Finance..

# Further reading

- a) A primer on the general theory of stochastic processes is :
	- A. NIKEGHBALI. (2006). An essay on the general theory of stochastic processes. Probab. Surv. 3, 345-412.
- b) Several computations of Azema supermartingales and related topics are also found in :
	- A. NIKEGHBALI.  $(2006)$ . Enlargements of filtrations and path decompositions at non stopping times. Probab. Theory Related  $Fields$  136, no. 4, 524-540.
	- A. NIKEGHBALI.  $(2006)$ . A class of remarkable submartingales. Stochastic Process. Appl. 116 , no. 6, 917-938.
	- A. NIKEGHBALI. (2006). Multiplicative decompositions and frequency of vanishing of nonnegative submartingales. J. Theoret. Probab. 19 , no. 4, 931-949.
	- A. NIKEGHBALI. (2008). A generalization of Doob's maximal identity. Submitted.

# Acknowledgments

What started it all is M. Qian's question.

M. Yor is also very grateful to D. Madan and B. Roynette for several attempts to solve various questions, rewriting, and summarizing...

Lecturing in Osaka and Ritsumeikan (October 2007), Melbourne and Sydney (December 2007), then finally at the Bachelier Séminaire (February 2008) has been a great help.

We gave further lectures in Oxford and at Imperial College, London, both in May 2008.