The parabolic Sturmian-function basis representation of the six-dimensional Coulomb Green's function

S. A. Zaytsev^{*}

Pacific National University, Khabarovsk, 680035, Russia

Abstract

The square integrable basis set representation of the resolvent of the asymptotic three-body Coulomb wave operator in parabolic coordinates is obtained. The resulting six-dimensional Green's function matrix is expressed as a convolution integral over separation constants.

PACS numbers: 03.65.Nk

arXiv:0806.0441v1 [math-ph] 3 Jun 2008

^{*}E-mail: zaytsev@fizika.khstu.ru

I. INTRODUCTION

It is well known that the Schrödinger equation for a three-body Coulomb system is asymptotically separable in terms of parabolic coordinates [1]

$$\xi_j = r_{ls} + \hat{\mathbf{k}}_{ls} \cdot \mathbf{r}_{ls}, \quad \eta_j = r_{ls} - \hat{\mathbf{k}}_{ls} \cdot \mathbf{r}_{ls}, \tag{1}$$

where \mathbf{r}_{ls} and \mathbf{k}_{ls} are the relative coordinate and momentum vectors between the particles l and s. Here j, l, s is a cyclic permutation of 1, 2, 3. The long-ranged six-dimensional operator, which provides a three-body continuum wave function, 3C-function [2, 3], that satisfies exact asymptotic boundary conditions for Coulomb systems in the region where the distances between the particles are large, reads as the sum of two-dimensional operators [1]:

$$\sum_{j=1}^{3} \frac{1}{\mu_{ls}(\xi_j + \eta_j)} \left[\hat{h}_{\xi_j} + \hat{h}_{\eta_j} + 2k_{ls} t_{ls} \right],$$
(2)

where $t_{ls} = \frac{Z_l Z_s \mu_{ls}}{k_{ls}}$, $\mu_{ls} = \frac{m_l m_s}{m_l + m_s}$. Here the one-dimensional operators \hat{h}_{ξ_j} and \hat{h}_{η_j} are defined by

$$\hat{h}_{\xi_j} = -2 \left(\frac{\partial}{\partial \xi_j} \xi_j \frac{\partial}{\partial \xi_j} + ik_{ls} \xi_j \frac{\partial}{\partial \xi_j} \right),
\hat{h}_{\eta_j} = -2 \left(\frac{\partial}{\partial \eta_j} \eta_j \frac{\partial}{\partial \eta_j} - ik_{ls} \eta_j \frac{\partial}{\partial \eta_j} \right).$$
(3)

In the previous paper [4] we introduce the six-dimensional operator \mathfrak{h} which is obtained by multiplying (2) on the left by $\prod_{j=1}^{3} \mu_{ls}(\xi_j + \eta_j)$:

$$\hat{\mathfrak{h}} = \mu_{13}(\xi_2 + \eta_2)\,\mu_{12}(\xi_3 + \eta_3)\hat{\mathfrak{h}}_1 + \mu_{23}(\xi_1 + \eta_1)\,\mu_{12}(\xi_3 + \eta_3)\hat{\mathfrak{h}}_2 + \mu_{23}(\xi_1 + \eta_1)\,\mu_{13}(\xi_2 + \eta_2)\hat{\mathfrak{h}}_3, \quad (4)$$

where

$$\hat{\mathfrak{h}}_j = \hat{h}_{\xi_j} + \hat{h}_{\eta_j} + 2k_{ls}t_{ls}.$$
(5)

The resolvent of the operator (4) can be used in the corresponding Lippmann-Schwinger equation for the three-body Coulomb wave function.

It has been suggested [4] to treat the operator (4) within the context of L^2 parabolic Sturmian basis set [5]

$$|\mathcal{N}\rangle = \prod_{j=1}^{3} \phi_{n_j \, m_j}(\xi_j, \, \eta_j),\tag{6}$$

$$\phi_{n_j \, m_j}(\xi_j, \, \eta_j) = \psi_{n_j}(\xi_j) \, \psi_{m_j}(\eta_j), \tag{7}$$

$$\psi_n(x) = \sqrt{2b}e^{-bx}L_n(2bx),\tag{8}$$

where b is the scaling parameter. In particular, a matrix representation \mathbf{G}_{j} of the resolvent for the two-dimensional operator

$$\hat{\mathfrak{h}}_j + \mu_{ls} C_j (\xi_j + \eta_j) \tag{9}$$

has been obtained which is formally the matrix inverse to the infinite matrix $[\mathbf{h}_j + \mu_{ls}C_j\mathbf{Q}_j]$ of the operator (9):

$$\left[\mathbf{h}_{j} + \mu_{ls}C_{j}\mathbf{Q}_{j}\right]\mathbf{G}_{j}(t_{ls};\ \mu_{ls}C_{j}) = \mathbf{I}_{j}.$$
(10)

Here

$$\mathbf{h}_{j} = \mathbf{h}_{\xi_{j}} \otimes \mathbf{I}_{\eta_{j}} + \mathbf{I}_{\xi_{j}} \otimes \mathbf{h}_{\eta_{j}} + 2k_{ls}t_{ls}\mathbf{I}_{j}$$
(11)

is the matrix of the operator $\hat{\mathbf{h}}_j$ (5) in the basis (7), \mathbf{I}_{ξ_j} , \mathbf{I}_{η_j} and $\mathbf{I}_j = \mathbf{I}_{\xi_j} \otimes \mathbf{I}_{\eta_j}$ are the unit matrices. $\mathbf{Q}_j = \mathbf{Q}_{\xi_j} \otimes \mathbf{I}_{\eta_j} + \mathbf{I}_{\xi_j} \otimes \mathbf{Q}_{\eta_j}$, where \mathbf{Q}_{ξ_j} and \mathbf{Q}_{η_j} are the matrices of ξ_j and η_j in basis (8), respectively.

In the previous paper [4] it has been suggested that the matrices \mathbf{G}_j of the twodimensional Green's functions be used to construct the matrix $\underline{\boldsymbol{\mathcal{G}}}$ which is inverse to the operator (4) matrix

$$\underline{\mathfrak{h}} = \mu_{13}\mu_{12}\mathbf{h}_1 \otimes \mathbf{Q}_2 \otimes \mathbf{Q}_3 + \mu_{23}\mu_{12}\mathbf{Q}_1 \otimes \mathbf{h}_2 \otimes \mathbf{Q}_3 + \mu_{23}\mu_{13}\mathbf{Q}_1 \otimes \mathbf{Q}_2 \otimes \mathbf{h}_3.$$
(12)

Namely, we proposed to express the six-dimensional Green's function matrix $\underline{\mathfrak{G}}$ as the convolution integral

$$\underline{\mathfrak{G}} = \aleph \int \int dC_1 dC_2 \,\mathbf{G}_1(t_{23}; \,\mu_{23}C_1) \otimes \mathbf{G}_2(t_{13}; \,\mu_{13}C_2) \otimes \mathbf{G}_3(t_{12}; \,-\mu_{12}(C_1 + C_2)), \quad (13)$$

where \aleph is a normalizing factor. Thus, our problem is now to determine the pathes of integration over the separation constants C_1 and C_2 in (13) and to find the corresponding normalizing factor \aleph such that the condition

$$\underline{\mathfrak{h}}\underline{\mathfrak{G}} = \mathbf{I}_1 \otimes \mathbf{I}_2 \otimes \mathbf{I}_3 \tag{14}$$

holds. For this purpose consider the product $\mathfrak{h} \underline{\mathfrak{G}}$. From the relation (10) we obtain

$$\underline{\mathfrak{h}} \, \underline{\mathfrak{G}} = \aleph \left\{ \int \int dC_1 dC_2 \left[\mathbf{I}_1 - \mu_{23} C_1 \mathbf{Q}_1 \mathbf{G}_1(t_{23}; \ \mu_{23} C_1) \right] \otimes \mu_{13} \mathbf{Q}_2 \mathbf{G}_2(t_{13}; \ \mu_{13} C_2) \right. \\ \left. \otimes \mu_{12} \mathbf{Q}_3 \mathbf{G}_3(t_{12}; \ -\mu_{12} (C_1 + C_2)) \right. \\ \left. \int \int dC_1 dC_2 \ \mu_{23} \mathbf{Q}_1 \mathbf{G}_1(t_{23}; \ \mu_{23} C_1) \otimes \left[\mathbf{I}_2 - \mu_{13} C_2 \mathbf{Q}_2 \mathbf{G}_2(t_{13}; \ \mu_{13} C_2) \right] \right. \\ \left. \otimes \mu_{12} \mathbf{Q}_3 \mathbf{G}_3(t_{12}; \ -\mu_{12} (C_1 + C_2)) \right.$$

$$\left. \left. \int \int dC_1 dC_2 \ \mu_{23} \mathbf{Q}_1 \mathbf{G}_1(t_{23}; \ \mu_{23} C_1) \otimes \mu_{13} \mathbf{Q}_2 \mathbf{G}_2(t_{13}; \ \mu_{13} C_2) \right. \\ \left. \otimes \left[\mathbf{I}_3 + \mu_{12} (C_1 + C_2) \mathbf{Q}_3 \mathbf{G}_3(t_{12}; \ -\mu_{12} (C_1 + C_2)) \right] \right\},$$

$$\left. \left. \left. \left. \left[\mathbf{I}_3 + \mu_{12} (C_1 + C_2) \mathbf{Q}_3 \mathbf{G}_3(t_{12}; \ -\mu_{12} (C_1 + C_2)) \right] \right\} \right\} \right\} \right\} \right\} \right\}$$

and hence

$$\underline{\mathfrak{h}} \, \underline{\mathfrak{G}} = \aleph \left\{ \int \int dC_1 dC_2 \left[\mathbf{I}_1 \otimes \mu_{13} \mathbf{Q}_2 \mathbf{G}_2(t_{13}; \ \mu_{13} C_2) \otimes \mu_{12} \mathbf{Q}_3 \mathbf{G}_3(t_{12}; \ -\mu_{12}(C_1 + C_2)) \right. \right. \\ \left. + \mu_{23} \mathbf{Q}_1 \mathbf{G}_1(t_{23}; \ \mu_{23} C_1) \otimes \mathbf{I}_2 \otimes \mu_{12} \mathbf{Q}_3 \mathbf{G}_3(t_{12}; \ -\mu_{12}(C_1 + C_2)) \right]$$

$$\left. + \left[\mu_{23} \mathbf{Q}_1 \int dC_1 \, \mathbf{G}_1(t_{23}; \ \mu_{23} C_1) \right] \otimes \left[\mu_{13} \mathbf{Q}_2 \int dC_2 \, \mathbf{G}_2(t_{13}; \ \mu_{13} C_2) \right] \otimes \mathbf{I}_3 \right\}.$$

$$(16)$$

As a first step towards our goal we consider the integrals

$$\int dC_j \mathbf{G}_j(t_{ls}; \ \mu_{ls}C_j) \tag{17}$$

inside the figure brackets on the right-hand side of (16).

In Sec. II completeness of the eigenfunctions of the two-dimensional operator (9) is considered. In particular, an integral representation of the matrix **A** which is inverse to the infinite matrix **Q** of the operator $(\xi + \eta)$ in the basis (7) is obtained. In Sec. III it is demonstrated that the integral (17) taken along an appropriate contour is proportional to the matrix **A** obtained in the previous section. Finally, Sec. IV presents a convolution integral representation of the six-dimensional Coulomb Green' function matrix.

II. COMPLETENESS RELATIONS

A. The continuous spectrum

Of particular interest are the regular solutions

$$f(\gamma, \tau, \xi, \eta) = u(\gamma, \tau, \xi) v(\gamma, \tau, \eta)$$
(18)

of the system

$$\left[\hat{h}_{\xi} + 2kt + \mu C\xi\right] u(\gamma, \tau, \xi) = 0, \qquad (19)$$

$$\left[\hat{h}_{\eta} + 2k(t_0 - t) + \mu C\eta\right] v(\gamma, \tau, \xi) = 0.$$
(20)

Obviously, the regular solutions u and v are proportional to confluent hypergeometric functions [6]:

$$u(\gamma, \tau, \xi) = e^{\frac{i}{2}(\gamma-k)\xi} {}_{1}F_{1}\left(\frac{1}{2} + i\tau, 1, -i\gamma\xi\right)$$
(21)

and

$$v(\gamma, \tau, \eta) = e^{-\frac{i}{2}(\gamma-k)\eta} {}_{1}F_{1}(\frac{1}{2} + i(\tau - \tau_{0}), 1, i\gamma\eta)$$

= $e^{\frac{i}{2}(\gamma+k)\eta} {}_{1}F_{1}(\frac{1}{2} + i(\tau_{0} - \tau), 1, -i\gamma\eta),$ (22)

where

$$\mu C = \frac{k^2}{2} - \frac{\gamma^2}{2}, \quad \tau = \frac{k}{\gamma} \left(t + \frac{i}{2} \right), \quad \tau_0 = \frac{k}{\gamma} t_0.$$
(23)

It should be noted that since the representation of the two- and six-dimensional Coulomb Green's functions matrix elements involves an integration over τ from $-\infty$ to ∞ , in the subsequent discussion we assume that τ is real. With this assumption it is readily seen that the solutions (21) and (22) coincide, except for normalization and the phase factors $e^{-\frac{i}{2}k\xi}$ and $e^{\frac{i}{2}k\eta}$, with parabolic Coulomb Sturmians treated in Ref. [7]. In this case γ plays the role of the momentum and $\mathcal{E} = \frac{\gamma^2}{2}$ is the energy. From this we conclude that for $\gamma > 0$ the solutions (21) and (22) correspond to the continuous spectrum of \mathcal{E} .

It is readily verified that the solutions $u(\gamma, \tau, \xi)$ and $v(\gamma, \tau, \eta)$ are expressed by basis set (8) expansions

$$u(\gamma, \tau, \xi) = \frac{2\sqrt{2b}}{2b - i(\gamma - k)} \left(\frac{2b - i(\gamma - k)}{2b + i(\gamma + k)}\right)^{i\tau + \frac{1}{2}} \sum_{n=0}^{\infty} \theta^n p_n(\tau; \zeta) \psi_n(\xi),$$
(24)

$$v(\gamma, \tau, \eta) = \frac{2\sqrt{2b}}{2b - i(\gamma + k)} \left(\frac{2b - i(\gamma + k)}{2b + i(\gamma - k)}\right)^{i(\tau_0 - \tau) + \frac{1}{2}} \sum_{n=0}^{\infty} \lambda^{-n} p_n \left(\tau_0 - \tau; \zeta\right) \psi_n(\eta), \quad (25)$$

where

$$\theta = \frac{2b + i(\gamma - k)}{2b - i(\gamma - k)}, \qquad \lambda = \frac{2b - i(\gamma + k)}{2b + i(\gamma + k)}, \qquad \zeta = \frac{\lambda}{\theta}.$$
(26)

The expansion (24) and (25) coefficients contain the polynomials [4]

$$p_n(\tau; \zeta) = \frac{(-1)^n}{n!} \frac{\Gamma\left(n + \frac{1}{2} - i\tau\right)}{\Gamma\left(\frac{1}{2} - i\tau\right)} {}_2F_1\left(-n, \frac{1}{2} + i\tau; -n + \frac{1}{2} + i\tau; \zeta\right).$$
(27)

The basis set (8) representation of the equation (19) is the three-term recursion relation
[4]

$$a_n y_{n-1} + b_n y_n + d_{n+1} y_{n+1} = 0, \quad n \ge 1$$
(28)

where

$$b_n = (b + \frac{\mu C}{2b} + ik) + 2(b + \frac{\mu C}{2b})n + 2kt,$$

$$a_n = (b - \frac{\mu C}{2b} - ik)n, \quad d_n = (b - \frac{\mu C}{2b} + ik)n.$$
(29)

The functions

$$s_n(t; \ \mu C) = \theta^n \, p_n(\tau; \ \zeta) \tag{30}$$

are the "regular" solutions of (28) with the initial conditions: $s_0 \equiv 1, s_{-1} \equiv 0$. The polynomials p_n (27) of degree n in τ are orthogonal with respect to the weight function [4]

$$\rho(\tau;\zeta) = \frac{\Gamma\left(\frac{1}{2} - i\tau\right)\Gamma\left(\frac{1}{2} + i\tau\right)}{2\pi i} \left(-\zeta\right)^{i\tau + \frac{1}{2}},\tag{31}$$

where it is considered that $|\arg(-\zeta)| < \pi$. The corresponding orthogonality relation reads

$$\frac{i}{\zeta^m} \left(\frac{\zeta - 1}{\zeta}\right) \int_{-\infty}^{\infty} d\tau \,\rho(\tau; \,\zeta) \,p_n(\tau; \,\zeta) \,p_m(\tau; \,\zeta) = \delta_{nm}. \tag{32}$$

B. The discrete spectrum

For $t_0 < 0$ the eigenfunctions of (19) corresponding to the discrete spectrum: $\mathcal{E}^{(\ell)} = \frac{\gamma_{\ell}^2}{2}$, $\gamma_{\ell} = i\kappa_{\ell}, \kappa_{\ell} = -\frac{kt_0}{\ell}, \ell = 1, 2, ..., \infty$, are [8]

$$f_{\ell,m}(\xi,\,\eta) = u_{\ell,m}(\xi)\,v_{\ell,\ell-m-1}(\eta), \qquad m = 0,\,1,\ldots,\,\ell-1,$$
(33)

where

$$u_{\ell,m}(\xi) = e^{-\frac{i}{2}k\xi} e^{-\frac{\kappa_{\ell}\xi}{2}} {}_{1}F_{1}(-m, 1; \kappa_{\ell}\xi) = e^{-\frac{i}{2}k\xi} e^{-\frac{\kappa_{\ell}\xi}{2}} L_{m}(\kappa_{\ell}\xi)$$
(34)

and

$$v_{\ell,m}(\eta) = e^{\frac{i}{2}k\eta} e^{-\frac{\kappa_{\ell}\eta}{2}} {}_{1}F_{1}(-m, 1; \kappa_{\ell}\eta) = e^{\frac{i}{2}k\eta} e^{-\frac{\kappa_{\ell}\eta}{2}} L_{m}(\kappa_{\ell}\eta).$$
(35)

The solutions $f_{\ell,\,m}$ meet the orthogonality relation

$$\frac{\kappa_{\ell}^{3}}{2\ell} \int_{0}^{\infty} \int_{0}^{\infty} (\xi + \eta) d\xi d\eta f_{\ell,m}(\xi, \eta) \left[f_{\ell',m'}(\xi, \eta) \right]^{*} = \delta_{\ell,\ell'} \delta_{m,m'}.$$
(36)

It is readily verified that the expansions of $u_{\ell,m}(\xi)$ and $v_{\ell,m}(\eta)$ in a basis function (8) series are

$$u_{\ell,m}(\xi) = \sum_{n=0}^{\infty} \mathcal{S}_n^{(\ell,m)} \,\psi_n(\xi)$$
(37)

and

$$v_{\ell,m}(\eta) = \sum_{n=0}^{\infty} \left[\mathcal{S}_n^{(\ell,m)} \right]^* \psi_n(\eta), \tag{38}$$

where the coefficients are given by

$$S_{n}^{(\ell,m)} = 2\sqrt{2b} \left(-\right)^{n} \frac{(m+1)_{n}}{n!} \frac{(2b-\kappa_{\ell}-ik)^{n}(2b-\kappa_{\ell}+ik)^{m}}{(2b+\kappa_{\ell}+ik)^{n+m+1}} \times {}_{2}F_{1}\left(-n, -m, -n-m; \frac{(2b+\kappa_{\ell})^{2}+k^{2}}{(2b-\kappa_{\ell})^{2}+k^{2}}\right).$$
(39)

One-dimensional completeness relations

The eigensolution $u(\gamma, \tau, \xi)$ of (19) corresponding to the continuous spectrum $(\gamma > 0)$ for large ξ behaves as

$$u(\gamma, \tau, \xi) \underset{\xi \to \infty}{\sim} \frac{e^{-\frac{i}{2}k\xi} e^{\frac{\pi\tau}{2}}}{\left|\Gamma(\frac{1}{2} + i\tau)\right|} \frac{2i}{\sqrt{\gamma\xi}} \sin\left(\frac{\gamma\xi}{2} - \tau\ln(\gamma\xi) + \frac{\pi}{4} + \sigma\right),\tag{40}$$

where $\sigma = \arg \Gamma \left(\frac{1}{2} + i\tau\right)$. Therefore (see e. g. [9]),

$$\frac{\gamma}{4} \left| \Gamma\left(\frac{1}{2} + i\tau\right) \right|^2 e^{-\pi\tau} \int_0^\infty \xi \, d\xi \, u(\gamma, \, \tau, \, \xi) \, \left[u(\gamma', \, \tau, \, \xi) \right]^* = \pi \delta(\gamma - \gamma'), \tag{41}$$

and for $\tau > 0$

$$\xi \int_{0}^{\infty} \gamma \, d\gamma \, \frac{\left|\Gamma\left(\frac{1}{2} + i\tau\right)\right|^2}{4\pi} \, e^{-\pi\tau} \, u(\gamma, \, \tau, \, \xi) \, \left[u(\gamma, \, \tau, \, \xi')\right]^* = \delta(\xi - \xi'). \tag{42}$$

On the other hand, if the functions $u(\gamma, \tau, \xi)$ are regarded as charge Sturmians [7], i. e. the parameter τ is considered as eigenvalue of the problem, whereas the momentum γ remains constant, the corresponding orthogonality and completeness relations are given by [7]

$$\gamma \left| \Gamma \left(\frac{1}{2} + i\tau \right) \right|^2 e^{-\pi\tau} \int_0^\infty d\xi \, u(\gamma, \, \tau, \, \xi) \left[u(\gamma, \, \tau', \, \xi) \right]^* = 2\pi\delta(\tau - \tau') \tag{43}$$

and

$$\gamma \int_{-\infty}^{\infty} d\tau \, \frac{\left|\Gamma\left(\frac{1}{2} + i\tau\right)\right|^2}{2\pi} e^{-\pi\tau} \, u(\gamma, \, \tau, \, \xi) \left[u(\gamma, \, \tau, \, \xi')\right]^* = \delta(\xi - \xi'). \tag{44}$$

Taking matrix elements of the completeness relation (44) we find

$$\frac{8 b \gamma \theta^{n-m}}{\sqrt{[4b^2 + (\gamma+k)^2] [4b^2 + (\gamma-k)^2]}} \int_{-\infty}^{\infty} d\tau \, \frac{\left|\Gamma\left(\frac{1}{2} + i\tau\right)\right|^2}{2\pi} \left(-\zeta\right)^{i\tau} p_n\left(\tau; \, \zeta\right) \left[p_n\left(\tau; \, \zeta\right)\right]^* = \delta_{n,m}.$$
(45)

It may be noted that (45) is closely related to (32). To see this, let $\gamma > 0$. Then, it follows from the definitions (26) that

$$\frac{(\zeta - 1)}{\zeta} (-\zeta)^{\frac{1}{2}} = \frac{8 \, b \, \gamma}{\sqrt{[4b^2 + (\gamma + k)^2] \, [4b^2 + (\gamma - k)^2]}}.$$
(46)

Further, the regular solution $\theta^n p_n(\tau; \zeta)$ of the three-term recursion relation (28) is an even function of γ , since the coefficients a_n , b_n and d_n depend on γ only through $\mu C = \frac{1}{2}(k^2 - \gamma^2)$. Thus, replacing γ by $-\gamma$, and hence τ by $-\tau$ ($\theta \to \lambda$ and $\lambda \to \theta$) and ζ by ζ^{-1} in Eq. (27) gives

$$\theta^{n} p_{n}(\tau; \zeta) = \lambda^{n} \frac{(-1)^{n}}{n!} \frac{\Gamma\left(n + \frac{1}{2} + i\tau\right)}{\Gamma\left(\frac{1}{2} + i\tau\right)} {}_{2}F_{1}\left(-n, \frac{1}{2} - i\tau; -n + \frac{1}{2} - i\tau; \zeta^{-1}\right).$$
(47)

Comparing Eqs. (27) and (47) then yields the relation

$$\theta^{n} p_{n}(\tau; \zeta) = \lambda^{n} \left[p_{n}(\tau; \zeta) \right]^{*}, \qquad (48)$$

and hence

$$\left[p_n\left(\tau;\,\zeta\right)\right]^* = \zeta^{-n} \, p_n\left(\tau;\,\zeta\right). \tag{49}$$

From the argument above, we conclude that for $\gamma > 0$ the orthogonality relation (32) reduces to (45).

The two-dimensional completeness relation

It follows from the relations (41) and (43) and analogous relations for $v(\gamma, \tau, \xi)$ that the two-dimensional orthogonality relation for $\gamma > 0$ is given by

$$e^{-\pi\tau_0} \frac{\gamma^2}{4} \frac{\left|\Gamma\left(\frac{1}{2}+i\tau\right)\right|^2}{2\pi} \frac{\left|\Gamma\left(\frac{1}{2}+i(\tau_0-\tau)\right)\right|^2}{2\pi} \int_0^\infty \int_0^\infty (\xi+\eta) \, d\xi d\eta \left\{f(\gamma,\,\tau,\,\xi,\,\eta)\right\} \times \left[f(\gamma',\,\tau',\,\xi,\,\eta)\right]^* = \delta(\gamma-\gamma') \,\delta(\tau-\tau').$$
(50)

In turn, in the case $t_0 > 0$ (where there are no bound states) it would appear reasonable that the two-dimensional completeness relation would be given by

$$(\xi + \eta) \left\{ \alpha \int_{0}^{\infty} d\gamma \, \gamma^{2} e^{-\pi\tau_{0}} \int_{-\infty}^{\infty} d\tau \frac{\left| \Gamma\left(\frac{1}{2} + i\tau\right) \right|^{2}}{2\pi} \frac{\left| \Gamma\left(\frac{1}{2} + i(\tau_{0} - \tau)\right) \right|^{2}}{2\pi} \right.$$

$$\left. \times f(\gamma, \, \tau, \, \xi, \, \eta) \left[f(\gamma, \, \tau, \, \xi', \, \eta') \right]^{*} \right\} = \delta(\xi - \xi') \, \delta(\eta - \eta').$$

$$(51)$$

The integration over τ in (51) is performed on the assumption that τ is independent of γ . To test this hypothesis and determine the normalizing factor α , we carried out the following numerical experiments. First with some parameters $t_0 > 0$, k > 0 and b we calculate the matrix elements $A_{n_1, n_2; m_1, m_2}$ for the expression in the figure braces on the right-hand side of (51) in the basis (7):

$$A_{n_1, n_2; m_1, m_2} = \alpha \int_0^\infty d\gamma \, \frac{64 \, b^2 \, \gamma^2 \, (-\zeta)^{i\tau_0}}{[4b^2 + (\gamma + k)^2][4b^2 + (\gamma - k)^2]} \, \theta^{n_1 - m_1} \lambda^{m_2 - n_2} \times \int_{-\infty}^\infty d\tau \frac{\left| \Gamma(\frac{1}{2} + i\tau) \right|^2}{2\pi} \frac{\left| \Gamma(\frac{1}{2} + i(\tau_0 - \tau)) \right|^2}{2\pi} \, p_{n_1}(\tau; \zeta) \, p_{n_2}(\tau_0 - \tau; \zeta) \left[p_{m_1}(\tau; \zeta) \, p_{m_2}(\tau_0 - \tau; \zeta) \right]^*.$$
(52)

It should be noted that the value of $(-\zeta)^{i\tau_0}$ in this formula is determined by the condition $|\arg(-\zeta)| < \pi$. Then the resulting matrix **A** is multiplied by the matrix

$$\mathbf{Q} = \mathbf{Q}_{\xi} \otimes \mathbf{I}_{\eta} + \mathbf{I}_{\xi} \otimes \mathbf{Q}_{\eta} \tag{53}$$

of the operator $(\xi + \eta)$. Finally, using the condition

$$\mathbf{Q} \mathbf{A} = \mathbf{I}_{\xi} \otimes \mathbf{I}_{\eta}. \tag{54}$$

we have obtained that $\alpha = \frac{1}{2}$. Notice that the infinite symmetric matrices \mathbf{Q}_{ξ} and \mathbf{Q}_{η} are tridiagonal [4], therefore the equations on the left hand-side of the linear system (54) each contain no more than six terms.

For for $t_0 < 0$ the completeness relation (51) transforms into

$$(\xi + \eta) \left\{ \frac{1}{2} \int_{0}^{\infty} d\gamma \, \gamma^{2} e^{-\pi\tau_{0}} \int_{-\infty}^{\infty} d\tau \frac{\left| \Gamma\left(\frac{1}{2} + i\tau\right) \right|^{2}}{2\pi} \frac{\left| \Gamma\left(\frac{1}{2} + i(\tau_{0} - \tau)\right) \right|^{2}}{2\pi} f(\gamma, \tau, \xi, \eta) \left[f(\gamma, \tau, \xi', \eta') \right]^{*} + \sum_{\ell=1}^{\infty} \frac{\kappa_{\ell}^{3}}{2\ell} \sum_{m=0}^{\ell-1} f_{\ell,m,\ell-m-1}(\xi, \eta) \left[f_{\ell,m,\ell-m-1}(\xi', \eta') \right]^{*} \right\} = \delta(\xi - \xi') \, \delta(\eta - \eta').$$

$$(55)$$

In this case the matrix **A** with elements

$$A_{n_{1},n_{2};m_{1},m_{2}} = \int_{0}^{\infty} d\gamma \, \frac{32 \, b^{2} \, \gamma^{2} \, (-\zeta)^{i\tau_{0}}}{[4b^{2} + (\gamma + k)^{2}][4b^{2} + (\gamma - k)^{2}]} \, \theta^{n_{1} - m_{1}} \lambda^{m_{2} - n_{2}} \\ \times \int_{-\infty}^{\infty} d\tau \, \frac{\left| \Gamma(\frac{1}{2} + i\tau) \right|^{2}}{2\pi} \frac{\left| \Gamma(\frac{1}{2} + i(\tau_{0} - \tau)) \right|^{2}}{2\pi} \, p_{n_{1}}(\tau; \, \zeta) \, p_{n_{2}}(\tau_{0} - \tau; \, \zeta) \left[p_{m_{1}}(\tau; \, \zeta) \, p_{m_{2}}(\tau_{0} - \tau; \, \zeta) \right]^{*} \quad (56) \\ + \sum_{\ell=1}^{\infty} \frac{\kappa_{\ell}^{3}}{2\ell} \, \sum_{m=0}^{\ell-1} \mathcal{S}_{n_{1}}^{(\ell,m)} \, \mathcal{S}_{m_{2}}^{(\ell,\ell-m-1)} \left[\mathcal{S}_{n_{2}}^{(\ell,\ell-m-1)} \, \mathcal{S}_{m_{1}}^{(\ell,m)} \right]^{*},$$

is also inverse to the matrix \mathbf{Q} (53). The expression (56) can be rewritten, in view of (46) and (49), as

$$A_{n_{1},n_{2};m_{1},m_{2}} = -\frac{1}{2} \int_{0}^{\infty} d\gamma \left\{ \left(\frac{\zeta - 1}{\zeta} \right)^{2} \frac{\theta^{n_{1} + m_{2}}}{\lambda^{n_{2} + m_{1}}} \int_{-\infty}^{\infty} d\tau \rho(\tau; \zeta) \rho(\tau_{0} - \tau; \zeta) \right. \\ \left. \times p_{n_{1}}(\tau; \zeta) p_{n_{2}}(\tau_{0} - \tau; \zeta) p_{m_{1}}(\tau; \zeta) p_{m_{2}}(\tau_{0} - \tau; \zeta) \right\} \\ \left. + \sum_{\ell=1}^{\infty} \frac{\kappa_{\ell}^{3}}{2\ell} \sum_{m=0}^{\ell-1} \mathcal{S}_{n_{1}}^{(\ell,m)} \mathcal{S}_{m_{2}}^{(\ell,\ell-m-1)} \left[\mathcal{S}_{n_{2}}^{(\ell,\ell-m-1)} \mathcal{S}_{m_{1}}^{(\ell,m)} \right]^{*}.$$
(57)

III. CONTOUR INTEGRALS

Notice that expressing the resolvent of the one-dimensional operator $\left[\hat{h}_{\xi} + 2kt + \mu C\xi\right]$ requires two linearly independent solutions of (19). Irregular solutions of (19) are expressed in terms the confluent hypergeometric function [6]

$$w^{(\pm)}(\gamma,\,\tau,\,\xi) = e^{\frac{i}{2}(\pm\gamma-k)\xi} U\left(\frac{1}{2}\pm i\tau,\,1;\,\mp\gamma\xi\right).$$
(58)

The corresponding solutions of the three-term recursion relation (28) are

$$c_n^{(+)}(t; \ \mu C) = \theta^{n+1} \, q_n^{(+)}(\tau; \ \zeta) \,,$$

$$c_n^{(-)}(t; \ \mu C) = \lambda^{n+1} \, q_n^{(-)}(\tau; \ \zeta) \,,$$
(59)

where

$$q_n^{(+)}(\tau; \zeta) = (-)^n \frac{n! \Gamma(\frac{1}{2} + i\tau)}{\Gamma(n + \frac{3}{2} + i\tau)} {}_2F_1\left(\frac{1}{2} + i\tau, n + 1; n + \frac{3}{2} + i\tau; \zeta^{-1}\right),$$

$$q_n^{(-)}(\tau; \zeta) = (-)^n \frac{n! \Gamma(\frac{1}{2} - i\tau)}{\Gamma(n + \frac{3}{2} - i\tau)} {}_2F_1\left(\frac{1}{2} - i\tau, n + 1; n + \frac{3}{2} - i\tau; \zeta\right).$$
(60)

In particular, the functions

$$\widetilde{w}^{(\pm)}(\gamma, \tau, \xi) = \frac{2i\sqrt{2b}}{2b - i(\gamma - k)} \left(-\frac{2b + i(\gamma - k)}{2b - i(\gamma + k)} \right)^{i\tau + \frac{1}{2}} \times \frac{e^{-\pi\tau}}{\theta \,\Gamma\left(\frac{1}{2} \pm i\tau\right)} \sum_{n=0}^{\infty} c_n^{(\pm)}(t; \ \mu C) \,\psi_n(\xi)$$
(61)

tend to $w^{(\pm)}(\gamma, \tau, \xi)$ as $\xi \to \infty$.

The matrix elements of the resolvent of $\left[\hat{h}_{\xi} + 2kt + \mu C\xi\right]$ can be written in the form [4]

$$g_{n,m}^{(+)}(t; \ \mu C) = \frac{i}{2\gamma} \left(\frac{\zeta - 1}{\zeta}\right) \frac{\theta^{n-m}}{\zeta^m} p_{n_{<}}(\tau; \ \zeta) \ q_{n_{>}}^{(+)}(\tau; \ \zeta) \tag{62}$$

and

$$g_{n,m}^{(-)}(t; \ \mu C) = \frac{i}{2\gamma} \left(\frac{\zeta - 1}{\zeta}\right) \frac{\theta^{n-m}}{\zeta^m} p_{n_{\zeta}}(\tau; \ \zeta) \ \zeta^{n_{>}+1} q_{n_{>}}^{(-)}(\tau; \ \zeta), \tag{63}$$

where $n_{>}$ and $n_{<}$ are the greater and lesser of n and m. Notice that $c_n^{(+)}\left(c_n^{(-)}\right)$ are defined in the upper (lower) half of the complex γ -plane where $|\zeta| \geq 1$ ($|\zeta| \leq 1$). To analytically continue $c_n^{(+)}$ onto the lower half of the γ -plane the relation [4]

$$c_n^{(+)}(t; \ \mu C) = c_n^{(-)}(t; \ \mu C) + 2\pi i \rho(\tau; \ \zeta) \ \theta^{n+1} p_n(\tau; \ \zeta)$$
(64)

can be used.

In [4] we obtained the basis set (7) representation of the resolvent for the two-dimensional operator $\left[\hat{h}_{\xi} + 2kt + \mu C\xi\right] + \left[\hat{h}_{\eta} + 2k(t_0 - t) + \mu C\eta\right]$. In particular, the matrix elements of the two-dimensional Green's function can be expressed as the convolution integral

$$G_{n_1,n_2;m_1,m_2}^{(\pm)}(t_0; \ \mu C) = i \left(\frac{\zeta - 1}{\zeta}\right) \frac{\lambda^{m_2 - n_2}}{\zeta^{m_2}} \int_{-\infty}^{\infty} d\tau \ \rho(\tau_0 - \tau; \ \zeta) \times g_{n_1,m_1}^{(\pm)}(t; \ \mu C) \ p_{n_2}(\tau_0 - \tau; \ \zeta) \ p_{m_2}(\tau_0 - \tau; \ \zeta).$$
(65)

Notice that in this case only the regular solutions of (20) discrete analogues $\lambda^{-n}p_n(\tau_0 - \tau; \zeta)$ are used.

Let us consider the integral

$$\mathcal{I}_{1} = \frac{1}{2\pi i} \int_{-\infty}^{\infty} \gamma \, d\gamma \, G_{n_{1}, n_{2}; \, m_{1}, m_{2}}^{(+)}(t_{0}; \, \mu C).$$
(66)

Notice that by replacing $\gamma \to -\gamma$ (and hence $\theta \to \lambda$, $\lambda \to \theta$, $\zeta \to 1/\zeta$, $\tau \to -\tau$ and $\tau_0 \to -\tau_0$) in Eq. (65) $G_{n_1,n_2;m_1,m_2}^{(+)}(t_0; \mu C)$ is transformed to $G_{n_1,n_2;m_1,m_2}^{(-)}(t_0; \mu C)$. Thus, for the integral \mathcal{I}_1 we obtain

$$\mathcal{I}_{1} = \frac{1}{2\pi i} \int_{0}^{\infty} \gamma \, d\gamma \left\{ G_{n_{1}, n_{2}; m_{1}, m_{2}}^{(+)}(t_{0}; \mu C) - G_{n_{1}, n_{2}; m_{1}, m_{2}}^{(-)}(t_{0}; \mu C) \right\}.$$
(67)

Inserting Eqs. (65), (62) and (63) into Eq. (67), we find, in view of (64), that \mathcal{I}_1 coincides with the integral on the right hand-side of (57).

Now, we consider the integral

$$\frac{1}{2\pi i} \int_{\mathcal{C}} d\mathcal{E} \mathbf{G}^{(+)} \left(t_0; \, \frac{k^2}{2} - \mathcal{E} \right), \tag{68}$$

taken along the contour in the complex \mathcal{E} -plane shown in Fig. 1. The contour \mathcal{C} passes in a negative direction (clockwise) round all the points $\mathcal{E}^{(\ell)} = -\frac{\kappa_{\ell}^2}{2}$ (filled circles in Fig. 1 which accumulate at the origin) and the cut along the right half of the real axis and is closed at

infinity (see e. g. [10]). The corresponding matrix element of the integral along the two sides of the cut is equal to \mathcal{I}_1 (66) (this is circumstantial evidence that the normalizing factor α in the completeness relation (51) is equal to $\frac{1}{2}$). On the other hand, the integration along a contour enclosing $\mathcal{E}^{(\ell)}$ reduces to (-1) times the double sum of the residues of the integrand at the points $\tau^{(m)} = i \left(m + \frac{1}{2}\right)$, $m = 0, 1, \ldots$ and $\mathcal{E}^{(\ell)} = -\frac{(kt_0)^2}{2\ell^2}$, $\ell = 1, 2, \ldots$, which are the poles of the gamma functions $\Gamma\left(\frac{1}{2} + i\tau\right)$ and $\Gamma\left(\frac{1}{2} + i(\tau_0 - \tau)\right) = \Gamma\left(m + 1 + \frac{kt_0}{\sqrt{-2\mathcal{E}^{(\ell)}}}\right)$, respectively. It is readily shown that the matrix element of this part of the integral (68) coincides with the double sum in (57). Thus, the integral (68) is equal to the matrix **A**. The contour \mathcal{C} can be deformed, for instance, into a straight line parallel to the real axis. The resulting path \mathcal{C}_1 shown in Fig. 2 runs above the cut and the bound-state poles of $\mathbf{G}^{(+)}\left(t_0; \frac{k^2}{2} - \mathcal{E}\right)$. Then, to make the integral amenable to numerical integration, rotate \mathcal{C}_1 about some point \mathcal{E}_0 on the right half of the real axis through a negative angle φ [10]; see the contour \mathcal{C}_2 in Fig. 2. The part of \mathcal{C}_2 on the unphysical sheet is depicted by the dashed line. Thus, we obtain the following representation of the matrix \mathbf{A} (which is inverse to the matrix \mathbf{Q} (53)):

$$\mathbf{A} = \frac{1}{2\pi i} \int_{\mathcal{C}_2} d\mathcal{E} \, \mathbf{G}^{(+)} \left(t_0; \, \frac{k^2}{2} - \mathcal{E} \right) = \mathbf{Q}^{-1}.$$
(69)

IV. SIX-DIMENSIONAL GREEN'S FUNCTION MATRIX

Using the relation (69) we can rewrite the expression (13) for the six-dimensional Coulomb Green' function matrix as the contour integral

$$\underline{\mathfrak{G}} = \frac{\aleph}{\mu_{23}\mu_{13}} \int_{\mathcal{C}_2} d\mathcal{E}_1 \int_{\mathcal{C}_2} d\mathcal{E}_2 \, \mathbf{G}_1 \left(t_{23}; \, \frac{k_{23}^2}{2} - \mathcal{E}_1 \right) \otimes \mathbf{G}_2 \left(t_{13}; \, \frac{k_{13}^2}{2} - \mathcal{E}_2 \right) \otimes \mathbf{G}_3 \left(t_{12}; \, \frac{k_{12}^2}{2} - \mathcal{E}_3 \right),$$

$$\tag{70}$$

where $\mathcal{E}_j = \frac{k_{ls}^2}{2} - \mu_{ls}C_j$ and $C_1 + C_2 + C_3 = 0$. This also allows us to determine the normalizing factor \aleph . Indeed, it follows from (69) that the third term inside the figure brackets on the right-hand side of (16) is proportional to the unit matrix:

$$\begin{bmatrix} \mathbf{Q}_1 \int\limits_{\mathcal{C}_2} d\mathcal{E}_1 \, \mathbf{G}_1 \left(t_{23}; \, \frac{k_{23}^2}{2} - \mathcal{E}_1 \right) \end{bmatrix} \otimes \begin{bmatrix} \mathbf{Q}_2 \int\limits_{\mathcal{C}_2} d\mathcal{E}_2 \, \mathbf{G}_2 \left(t_{13}; \, \frac{k_{13}^2}{2} - \mathcal{E}_2 \right) \end{bmatrix} \otimes \mathbf{I}_3$$

$$= (2\pi i)^2 \, \mathbf{I}_1 \otimes \mathbf{I}_2 \otimes \mathbf{I}_3.$$
(71)

Consider the first two terms in the figure braces in (16). For the energy $\mathcal{E}_3 = \frac{k_{12}^2}{2} + \mu_{12}(C_1 + C_2)$ we have

$$\mathcal{E}_3 = \frac{k_{12}^2}{2} + \frac{\mu_{12}}{\mu_{23}} \left(\frac{k_{23}^2}{2} - \mathcal{E}_1 \right) + \frac{\mu_{12}}{\mu_{13}} \left(\frac{k_{13}^2}{2} - \mathcal{E}_2 \right).$$
(72)

On the other hand, on the contours C_2 the energy variables \mathcal{E}_j , j = 1, 2 are given by

$$\mathcal{E}_j = \mathcal{E}_{0j} + E_j e^{i\varphi},\tag{73}$$

where $\varphi < 0$, \mathcal{E}_{0j} is an arbitrary positive parameter, E_j is real and runs from $-\infty$ to ∞ . Hence, for the energy \mathcal{E}_3 (72) we obtain

$$\mathcal{E}_{3} = \left[\frac{k_{12}^{2}}{2} + \frac{\mu_{12}}{\mu_{23}}\left(\frac{k_{23}^{2}}{2} - \mathcal{E}_{01}\right) + \frac{\mu_{12}}{\mu_{13}}\left(\frac{k_{13}^{2}}{2} - \mathcal{E}_{02}\right)\right] + \left(-\frac{\mu_{12}}{\mu_{23}}E_{1} - \frac{\mu_{12}}{\mu_{13}}E_{2}\right)e^{i\varphi}.$$
 (74)

Thus, \mathcal{E}_3 can be expressed in the form

$$\mathcal{E}_3 = \mathcal{E}_{03} + E_3 \, e^{i\varphi},\tag{75}$$

where \mathcal{E}_{03} and E_3 denote the term in the square braces and the real factor in front of the exponent in (74), respectively. Since \mathcal{E}_{03} should be positive, the positive parameters \mathcal{E}_{01} and \mathcal{E}_{02} have to satisfy the constraint

$$\frac{\mu_{12}}{\mu_{23}} \mathcal{E}_{01} + \frac{\mu_{12}}{\mu_{13}} \mathcal{E}_{02} < \frac{k_{12}^2}{2} + \frac{\mu_{12}}{\mu_{23}} \frac{k_{23}^2}{2} + \frac{\mu_{12}}{\mu_{13}} \frac{k_{13}^2}{2}.$$
(76)

Now, we consider the integral

$$\mathcal{I}_2 = \int_{\mathcal{C}_2} d\mathcal{E}_1 \mathbf{G}_3 \left(t_{12}; \ \frac{k_{12}^2}{2} - \mathcal{E}_3 \right).$$
(77)

With fixed \mathcal{E}_2 , in view (74), (75) and (69), we see that

$$\mathcal{I}_{2} = e^{i\varphi} \int_{-\infty}^{\infty} dE_{1} \mathbf{G}_{3} \left(t_{12}; \ \frac{k_{12}^{2}}{2} - \mathcal{E}_{3} \right) = -\frac{\mu_{23}}{\mu_{12}} e^{i\varphi} \int_{-\infty}^{\infty} dE_{3} \mathbf{G}_{3} \left(t_{12}; \ \frac{k_{12}^{2}}{2} - \mathcal{E}_{3} \right) \\ = -\frac{\mu_{23}}{\mu_{12}} \int_{\mathcal{C}_{2}} d\mathcal{E}_{3} \mathbf{G}_{3} \left(t_{12}; \ \frac{k_{12}^{2}}{2} - \mathcal{E}_{3} \right) = -2\pi i \frac{\mu_{23}}{\mu_{12}} \mathbf{Q}_{3}^{-1}.$$
(78)

Similarly, we obtain

$$\int_{\mathcal{C}_2} d\mathcal{E}_2 \mathbf{G}_3 \left(t_{12}; \ \frac{k_{12}^2}{2} - \mathcal{E}_3 \right) = -2\pi i \frac{\mu_{13}}{\mu_{12}} \mathbf{Q}_3^{-1}.$$
(79)

Inserting (78), (79) and (71) into (16) then yields

$$\underline{\mathfrak{h}} \ \underline{\mathfrak{G}} = \aleph \, 4\pi^2 \, \mathbf{I}_1 \otimes \mathbf{I}_2 \otimes \mathbf{I}_3. \tag{80}$$

Thus, from (80) we conclude that

$$\aleph = \frac{1}{4\pi^2}.\tag{81}$$

V. CONCLUSION

The Sturmian basis-set representation of the resolvent for the asymptotic three-body Coulomb wave operator is obtained, which can be used in the discrete analog of the Lippmann-Schwinger equation for the three-body continuum wave function. The sixdimensional Green's function matrix is expressed as a convolution integral over separation constants. The integrand of this contour integral involves Green's function matrices corresponding to the two-dimensional operators which are constituents of the full six-dimensional wave operator. The completeness relation of the eigenfunctions of these two-dimensional operators is used to define the appropriate pathes of integration of the convolution integral.

- [1] H. Klar, Z. Phys. D 16, 231 (1990).
- [2] Dz. Belkic, J. Phys. B **11**, 3529 (1978).
- [3] C. R. Garibotti, J. E. Miraglia, Phys. Rev. A 21, 572 (1980).
- [4] S. A. Zaytsev, J. Phys. A, to be published.
- [5] P. C. Ojha, J. Math. Phys. 28, 392 (1987).
- [6] M. Abramowitz and I. A. Stegun, Handbook of mathematical functions (New York: Dover), 1970.
- [7] G. Gasaneo and F. D. Colavecchia, J. Phys. A **36** 8443 (2003).
- [8] L. D. Landau and E. M. Lifshitz, Quantum Mechanics: Non-Relativistic Theory (Oxford: Pergamon), 1965.
- [9] N. Michel, J. Math. Phys. **49** 022109 (2008).
- [10] R. Shakeshaft, Phys. Rev. A, **70**, 042704 (2004).



Figure 1: The path of integration C in (68).



Figure 2: The path of integration C_2 in (69).