Unified $(q; \alpha, \beta, \gamma; \nu)$ -deformation of one-parametric q-deformed oscillator algebras

I. M. Burban

Bogolyubov Institute for Theoretical Physics, Kiev 03680, Ukraine

E-mail: burban@bitp.kiev.ua

Abstract

We define a generalized $(q; \alpha, \beta, \gamma; \nu)$ -deformed oscillator algebra and study the number of its characteristics. We describe the structure function of deformation, analyze the classification of irreducible representations and discuss the asymptotic spectrum behaviour of the Hamiltonian. For a special choice of the deformation parameters we construct the deformed oscillator with discrete spectrum of its "quantized coordinate" operator. We establish its connection with the (generalized) discrete Hermite I polynomials.

PACS numbers: 02.20.Uw, 03.65.G, 03.65.Fd, 05.30.Pr

Short title: Unified deformation

Corresponding author: Ivan Burban

Postal address: Bogolyubov Institute for Theoretical Physics, Metrologichna str. 14b,

Kiev 03680, Ukraine

Phone: 380-44-5213436 Fax: 380-44-5265998

1. Introduction

The investigation of an one-parameter deformed canonical commutation relations in theoretical physics was originated from the study of the dual resonance models of strong interactions. The multi-parameter generalizations of these deformations were actively studied with the appearance quantum groups, quantum algebras, and quantum spaces. These deformations have found applications in the study of exotic, different from the standard Bose-Einstein and Fermi-Dirac, statistics.

On the other hand, many properties of the modified oscillator algebra have found applications in the study of the integrability of the two-particle Calogero models. The modified oscillator algebra has been generalized to C_{λ} -extended oscillator algebra [13] with the hope to exploit these deformations for construction of new integrable models. With the same aim, a "hybrid" model [9, 10] of the modified and q-deformed oscillator algebras has been proposed.

To close this cycle of ideas we consider the generalized $(q; \alpha, \beta, \gamma; \nu)$ -deformed oscillator algebra as "the synthesis" of the $(q; \alpha, \beta, \gamma)$ -deformed [3, 6] and ν -modified oscillator algebras [10].

In Section 2 some of the q-deformed oscillator algebras are placed in the order of their extension. We find the structure function of $(q; \alpha, \beta, \gamma; \nu)$ -deformed oscillator algebra. We show that this algebra is embedded into the deformed C_2 -extended oscillator algebra [13]. In Section 3 we give, following [13, 18], classification of the representations of this algebra. In Section 4 for the special choice of the deformation parameters of the algebra we study the oscillator [24] with the discrete spectrum of its "quantized coordinate" Q. This operator is represented by Jacobi matrix. We find the eigenvalues and eigenfunctions of this operator. The eigenfunctions are expressed by the (generalized) discrete q-Hermite I polynomial [25]. In Section 5 we represent the Hamiltonian of this model as function of the number operator and study its asymptotic energy spectrum behaviour.

2. Oscillator algebra and its generalized deformations

The oscillator algebra of the quantum harmonic oscillator is defined by canonical commutation relations

$$[a, a] = [a^+, a^+] = 0, \quad [a, a^+] = 1, \quad [N, a] = -a, \quad [N, a^+] = a^+.$$
 (1)

It allows the different types of deformations. Some of them have been called *generalized* deformed oscillator algebras [1, 2, 3, 4]. Each of them defines an algebra generated by elements (generators) $\{1, a, a^+, N\}$ and relations

$$[N, a] = -a, \quad [N, a^+] = a^+, \quad a^+ a = f(N), \quad aa^+ = f(N+1),$$
 (2)

where f is called a structure function of the deformation. Among them - the multiparameter generalization of one-parameter deformations [3, 4, 8, 12, 13, 15, 19, 24].

Let us recount some of them.

1. The Arik-Coon q-deformed oscillator algebra [5]

$$aa^{+} - qa^{+}a = 1$$
, $[N, a] = -a$, $[N, a^{+}] = a^{+}$, $q \in \mathbb{R}_{+}$,
$$f(n) = \frac{1 - q^{n}}{1 - q}.$$
 (3)

2. The Biedengarn–Macfarlane q-deformed oscillator algebra [6, 7]

$$aa^{+} - qa^{+}a = q^{-N}, \quad aa^{+} - q^{-1}a^{+}a = q^{N}, \quad [N, a] = -a, \quad [N, a^{+}] = a^{+},$$

$$f(n) = \frac{q^{n} - q^{-n}}{q - q^{-1}} \quad q \in \mathbb{R}_{+}.$$
 (4)

3. The Chung–Nam–Um generalized $(q; \alpha, \beta)$ -deformed oscillator algebra [3]

$$aa^{+} - qa^{+}a = q^{\alpha N + \beta}, \quad [N, a] = -a, \quad [N, a^{+}] = a^{+}, \quad q \in \mathbb{R}_{+}, \quad \alpha, \beta \in \mathbb{R},$$

$$f(n) = \begin{cases} q^{\beta} \frac{q^{\alpha n} - q}{q^{\alpha} - q}, & \text{if } \alpha \neq 1\\ nq^{n-1+\beta}, & \text{if } \alpha = 1. \end{cases}$$

$$(5)$$

4. The generalized $(q; \alpha, \beta, \gamma)$ -deformed oscillator algebra [8]

$$aa^{+} - q^{\gamma}a^{+}a = q^{\alpha N + \beta}, \quad [N, a] = -a, \quad [N, a^{+}] = a^{+}, \quad q \in \mathbb{R}_{+}, \alpha, \beta, \gamma \in \mathbb{R},$$

$$f(n) = \begin{cases} q^{\beta} \frac{q^{\alpha n} - q^{\gamma n}}{q^{\alpha} - q^{\gamma n}}, & \text{if } \alpha \neq \gamma \\ nq^{n-1+\beta}, & \text{if } \alpha = \gamma. \end{cases}$$

$$(6)$$

5. The ν -modified oscillator algebra [9, 10]

$$[a, a^{+}] = 1 + 2\nu K, \quad [N, a] = -a, \quad [N, a^{+}] = a^{+},$$

$$aK = -Ka, \quad a^{+}K = -Ka^{+}, \quad K^{2} = 1, \quad \nu \in \mathbb{R},$$

$$f(n) = \begin{cases} 2k + 1 + 2\nu, & \text{if } n = 2k\\ 2k + 2, & \text{if } n = 2k + 1. \end{cases}$$

$$(7)$$

This oscillator, as it has been shown in [10], is connected with the two-particle Calogero model [11].

6. The deformed C_{λ} -extended oscillator algebra [13] is defined by the relations

$$[a, a^{+}]_{q} \equiv aa^{+} - qa^{+}a = H(N) + K(N) \sum_{k=0}^{\lambda-1} \nu_{k} P_{k}, \quad [N, a] = -a, \quad [N, a^{+}] = a^{+},$$

$$aK = -Ka, \quad a^{+}K = -Ka^{+}, \quad K^{2} = 1, \quad \nu_{k} \in \mathbb{R},$$
(8)

where $\nu_k \in \mathbb{R}$ and H(K), K(N) are real analytic functions. This algebra admits the two Casimir operators $C_1 = e^{2\pi N}$ and $C_2 = \sum_{k=0}^{\lambda-1} e^{-2\pi i(N-k)/\lambda} P_k$.

7. The new $(q; \nu)$ -deformed oscillator [12]

$$aa^{+} - qa^{+}a = (1 + 2\nu K)q^{-N}, \quad [N, a] = -a,$$

$$[N, a^{+}] = a^{+}, \quad Ka = -aK, \quad Ka^{+} = -a^{+}K, \quad K^{2} = 1,$$

$$f(n) = \left(\frac{q^{n} - q^{-n}}{q - q^{-1}} + 2\nu \frac{q^{n} - (-1)^{n}q^{-n}}{q + q^{-1}}\right)$$

$$(9)$$

has been defined by the combination of the idea of Biedenharn– Macfarlane [6, 7] qdeformation with Brink, Hanson and Vasiliev idea [10] of the ν -modification of the
oscillator algebra.

8. In order to close this cycle of ideas we consider a $(q; \alpha, \beta, \gamma; \nu)$ -deformed oscillator algebra – ("hybrid" of the $(q; \alpha, \beta, \gamma)$ -deformed (6) and the ν -modified (7) oscillator algebras), or, more exactly, an oscillator defined by generators $\{I, a, a^+, N, K\}$ and relations

$$aa^{+} - q^{\gamma}a^{+}a = (1 + 2\nu K)q^{\alpha N + \beta}, \quad [N, a] = -a, \quad [N, a^{+}] = a^{+},$$

$$Ka = -a, \quad Ka^{+} = -a^{+}K, \quad [N, K] = 0, \quad N^{+} = N, \quad K^{+} = K,$$
(10)

where $q \in \mathbb{R}_+$, $\alpha, \beta \in \mathbb{R}$, $\nu \in \mathbb{R} - \{0\}$. This model unify all deformations 1. - 7. of the oscillator algebra (1).

3. Generalized $(q; \alpha, \beta, \gamma; \nu)$ -deformed oscillator algebra and its simplest properties

(a) $(q; \alpha, \beta, \gamma; \nu)$ -deformed structure function. Description of an deformed oscillator algebra requires the determination of the deformation structure function f(n).

Equations (2) and (10) imply the recurrence relation

$$f(n+1) - q^{\gamma} f(n) = \left(1 + 2\nu(-1)^n\right) q^{\alpha n + \beta}.$$
 (11)

Its solution is obtained by method of mathematical induction:

0.
$$f(1) = q^{\gamma} f(0) + (1 + 2\nu(-1)^{0}) q^{\alpha 0 + \beta};$$

$$1. \quad f(2) = q^{\gamma} f(1) + \left(1 - 2\nu(-1)^{1}\right) q^{\alpha 1 + \beta} = \left[q^{2\gamma} f(0) + \left(1 + 2\nu(-1)^{0}\right) q^{\gamma} q^{\alpha 0 + \beta} + \left(1 - 2\nu(-1)^{1}\right) q^{\alpha 1 + \beta}\right];$$

$$2. \quad f(3) = q^{\gamma} f(2) + \left(1 - 2\nu(-1)^2\right) q^{\alpha 2 + \beta} = q^{3\gamma} f(0) + \left(1 + 2\nu(-1)^0\right) q^{2\gamma} q^{\alpha 0 + \beta} + \left(1 - 2\nu(-1)^1\right) q^{\gamma} q^{\alpha 1 + \beta} + \left(1 - 2\nu(-1)^2\right) q^{\alpha 2 + \beta};$$

$$3. \quad f(4) = q^{\gamma} f(3) + \left(1 + 2\nu(-1)^{3}\right) q^{\alpha 3 + \beta} = q^{4\gamma} f(0) + \left(1 + 2\nu(-1)^{0}\right) q^{3\gamma} q^{\alpha 0 + \beta} + \left(1 - 2\nu(-1)^{1}\right) q^{2\gamma} q^{\alpha 1 + \beta} + \left(1 - 2\nu(-1)^{2}\right) q^{\gamma} q^{\alpha 2 + \beta} + \left(1 + 2\nu(-1)^{3}\right) q^{\alpha 3 + \beta};$$

•

n.
$$f(n) = q^{n\gamma} f(0) + \sum_{k=0}^{n-1} q^{\gamma(n-k-1)} q^{\alpha k+\beta} + 2\nu \sum_{k=0}^{n-1} q^{\gamma(n-k-1)} (-1)^k q^{\alpha k+\beta}$$
.

The solution of the equation (11) with the initial value f(0) is given by formula

$$f(n) = \begin{cases} f(0)q^{\gamma n} + q^{\beta} \left(\frac{q^{\gamma n} - q^{\alpha n}}{q^{\gamma} - q^{\alpha}} + 2\nu \frac{q^{\gamma n} - (-1)^n q^{\alpha n}}{q^{\gamma} + q^{\alpha}} \right), & \text{if } \alpha \neq \gamma \\ f(0)q^{\gamma n} + nq^{\gamma(n-1) + \beta} + 2\nu q^{\gamma(n-1) + \beta} \left(\frac{1 - (-1)^n}{2} \right), & \text{if } \alpha = \gamma. \end{cases}$$
(12)

(b) Positivity of $(q; \alpha, \beta, \gamma; \nu)$ -deformed structure function. We try to define the values of the parameters $\{\alpha, \beta, \gamma, \nu\}$, where function f(n) is positive.

The inequality

$$\frac{q^{\gamma n} - q^{\alpha n}}{q^{\gamma} - q^{\alpha}} + 2\nu \frac{q^{\gamma n} - (-1)^n q^{\alpha n}}{q^{\gamma} + q^{\alpha}} > 0, \tag{13}$$

can be rewritten as

$$(q^{\gamma n} - q^{\alpha n}) \left(\frac{1}{q^{\gamma} - q^{\alpha}} + 2\nu \frac{1}{q^{\gamma} + q^{\alpha}} \right) > 0 \tag{14}$$

for even n and

$$\frac{q^{\gamma n} - q^{\alpha n}}{q^{\gamma n} + q^{\alpha n}} \frac{q^{\gamma} + q^{\alpha}}{q^{\gamma} - q^{\alpha}} + 2\nu > 0$$
(15)

for odd n.

It follows

$$1 + 2\nu > 0 \tag{16}$$

if q > 1, $\gamma - \alpha > 0$ $(q < 1, \gamma - \alpha < 0)$

$$-1 < 2\nu < -\frac{q^{\gamma} + q^{\alpha}}{q^{\gamma} - q^{\alpha}} \tag{17}$$

if $q < 1, \gamma - \alpha > 0 (q > 1, \gamma - \alpha < 0)$.

The conditions (16) and (17) are sufficient to ensure that the Fock representation of the relation (10) has the Hermitian properties.

(c) Useful formulas. The two formulae for the study of this algebra will be useful. One of them

$$a(a^{+})^{n} - q^{\gamma n} (a^{+})^{n} a = [n; \alpha, \gamma; \nu K](a^{+})^{n-1} q^{\alpha N + \beta}, \tag{18}$$

where $n \geq 1$,

$$[n; \alpha, \gamma; \nu K] = \begin{cases} \left(\frac{q^{\gamma n} - q^{\alpha n}}{q^{\gamma} - q^{\alpha}} + 2\nu K \frac{q^{\gamma n} - (-1)^n q^{\alpha n}}{q^{\gamma} + q^{\alpha}}\right), & \text{if } \alpha \neq \gamma; \\ nq^{\alpha(n-1)} + 2\nu K q^{\alpha(n-1)} \left(\frac{1 - (-1)^n}{2}\right), & \text{if } \alpha = \gamma \end{cases}$$
(19)

is deduced by method of mathematical induction.

Indeed, for n = 1 the relation (18) is true by definition. The assumption that it is true for some n implies

$$\{a(a^{+})^{n+1} = a(a^{+})^{n}a^{+} = \left(q^{\gamma n}(a^{+})^{n}a + [n;\alpha,\gamma;\nu K](a^{+})^{n-1}q^{\alpha N+\beta}\right)a^{+} =$$

$$q^{\gamma n}(a^{+})^{n}aa^{+} + (a^{+})^{n}[n;\alpha,\gamma;-\nu K]q^{\alpha(N+1)+\beta} =$$

$$q^{\gamma n}(a^{+})^{n}\left(q^{\gamma}a^{+}a + (1+2\nu K)q^{\alpha N+\beta}\right) + (a^{+})^{n}[n;\alpha,\gamma;-\nu K]q^{\alpha(N+1)+\beta} =$$

$$q^{\gamma(n+1)}(a^{+})^{(n+1)}a + q^{\gamma n}(a^{+})^{n}\left(1+2\nu K\right)q^{\alpha N+\beta} + q^{\alpha}(a^{+})^{n}[n;\alpha,\gamma;-\nu K]q^{\alpha N+\beta}.$$

The direct calculations leads to (18).

The second formula gives the generated function for $[n; \alpha, \gamma; \nu K]$:

$$\sum_{n=0}^{\infty} [n; \alpha, \gamma; \nu K] z^n = \begin{cases} \frac{z}{1 - q^{\gamma} z} \left(\frac{1}{1 - q^{\alpha} z} + 2\nu K \frac{1}{1 + q^{\alpha} z} \right), & \text{if } \alpha \neq \gamma \\ \frac{z}{(1 - q^{\gamma} z)^2} + 2\nu K \frac{z}{1 - q^{2\gamma} z^2}, & \text{if } \alpha = \gamma. \end{cases}$$
(20)

(d) Deformed C_2 -extended and $(q; \alpha, \beta, \gamma; \nu)$ -deformed oscillator algebras. The defining relations of the deformed C_2 -extended oscillator are given by

$$[N, a^{+}] = a^{+}, \quad [N, P_{k}] = 0, \quad a^{+}P_{k} = P_{k+1}a^{+}, \quad P_{1} + P_{2} = I, \quad P_{k}P_{l} = \delta_{k,l}P_{l},$$

$$aa^{+} - q^{\gamma}a^{+}a = H(N) + \nu \Big(E(N+1) + q^{\gamma}E(N)\Big)(P_{0} - P_{1}), \tag{21}$$

where $q, \nu \in \mathbb{R}, k, l = 1, 2$, and E(N), H(N) are real analytic functions. As we saw above the deformed extended oscillator algebra C_{λ} admits the two Casimir operators C_1, C_2 . In case of the C_2 -extended oscillator algebra they have the form

$$C_1 = e^{2\pi i N}, \quad C_2 = \sum_{k=0}^{1} e^{-2\pi i (N-k)/2} = e^{i\pi N} K.$$
 (22)

Let us define the operator

$$\tilde{C}_3 = q^{-\gamma N} \Big(D(N) + \nu E(N)K - a^+ a \Big), \tag{23}$$

where D(N), E(N) some analytic functions of N. The operator \tilde{C}_3 will be the Casimir operator of the oscillator algebra (21) if the only one condition $[\tilde{C}_3, a] = 0$ holds. It amounts to determination of the solution of the equations

$$D(N+1) - q^{\gamma}D(N) = H(N), \quad E(N+1)\beta_{k+1} - q^{\gamma}E(N)\beta_k = K(N)\nu_k,$$
 (24)

where $\nu_0 = -\nu_1 = \nu$, $\beta_0 = 0$, $\beta_2 = 0$, $\beta_1 = \nu$, k = 0, 1. Putting the solution $E(N) = 2q^{\alpha N + \beta}/(q^{\gamma} + q^{\alpha})$ of the equation of (24) and $H(N) = q^{\alpha N + \beta}$ in (21), we obtain the commutation relations of the $(q; \alpha, \beta, \gamma; \nu)$ -deformed oscillator algebra (10). Moreover, the solution

$$D(N) = \begin{cases} q^{\beta} \left(\frac{q^{\gamma N} - q^{\alpha N}}{q^{\gamma} - q^{\alpha}} + 2\nu \frac{q^{\gamma N}}{q^{\gamma} + q^{\alpha}} \right) & \text{if } \gamma \neq \alpha \\ q^{\beta} \left(q^{\gamma(N-1)} N + \nu q^{-\gamma} \right) & \text{if } \gamma = \alpha \end{cases}$$

of the first equation (24) gives the explicit form of the Casimir operator

$$\tilde{C}_{3} = \begin{cases}
q^{-\gamma N} \left(\left(\frac{q^{\gamma N} - q^{\alpha N}}{q^{\gamma} - q^{\alpha}} + 2\nu \frac{q^{\gamma N} - (-1)^{N} q^{\alpha N}}{q^{\gamma} + q^{\alpha}} \right) q^{\beta} - a^{+} a \right) & \text{if } \alpha \neq \gamma \\
q^{-\gamma N} \left(N + \nu (1 + (-1)^{N}) q^{\gamma N + \beta} - a^{+} a \right) & \text{if } \alpha = \gamma.
\end{cases}$$
(25)

3. Classification of representations of unified $(q;\alpha,\beta,\gamma;\nu)$ -deformed oscillator algebra

As has been shown in the previous Section the $(q; \alpha, \beta, \gamma; \nu)$ -deformed oscillator algebra allows nontrivial center which means that it has irreducible non-equivalent representations [16, 17]. We give a classification of these representations by method similar to the one in the articles [13], [18].

Due to the relations (10) there exists a vector $|0\rangle$ such that

$$a^+a|0\rangle = \lambda_0|0\rangle$$
, $aa^+|0\rangle = \mu_0|0\rangle$, $N|0\rangle = \varkappa_0|0\rangle$, $K|0\rangle = \omega e^{-i\pi\varkappa_0}|0\rangle$,

where ω is the value of the Casimir operator C_2 on the given irreducible representation. By using the formula (18) we find that vectors

$$|n\rangle = \begin{cases} (a^+)^n |0\rangle, & \text{if } n \ge 0\\ (a)^{-n} |0\rangle, & \text{if } n < 0 \end{cases}$$
 (26)

are eigenvectors of the operators a^+a and aa^+ :

$$a^+a|n\rangle = \lambda_n|n\rangle, \quad aa^+|n\rangle = \mu_n|n\rangle.$$

Let us define new system of the orthonormal vectors $\{|n\rangle\}_{n=-\infty}^{n=\infty}$, by

$$|n\rangle = \begin{cases} \left(\prod_{k=1}^{n} \lambda_{k}\right)^{-1/2} (a^{+})^{n} |0\rangle, & \text{if } n \geq 0\\ \left(\prod_{k=1}^{-n} \lambda_{n+k}\right)^{-1/2} (a)^{-n} |0\rangle, & \text{if } n < 0. \end{cases}$$
(27)

Then the relations (10) are represented by operators

$$a^{+}|n\rangle = \sqrt{\lambda_{n+1}}|n+1\rangle, \quad a|n\rangle = \sqrt{\lambda_{n}}|n-1\rangle,$$

$$N|n\rangle = (\varkappa_{0} + n)|n\rangle, \quad K|n\rangle = \frac{(-1)^{n}}{2\nu}B|n\rangle, \tag{28}$$

where $B = 2\nu\omega e^{-i\pi\varkappa_0} \in \mathbb{R}$. Due to non-negativity of the operators a^+a , aa^+ we have $\lambda_n \geq 0$ and $\mu_n \geq 0$. By using (10), we find (28) and $\lambda_n = \mu_{n-1}$ we obtain the recurrence relation

$$\lambda_{n+1} - q^{\gamma} \lambda_n = \left(1 + (-1)^n B \right) q^{\alpha(n+\varkappa_0) + \beta}. \tag{29}$$

Take into account the relation (12) the solution of equation (29) can be represented by

$$\lambda_n = \begin{cases} \lambda_0 q^{\gamma n} + q^{\alpha \varkappa_0 + \beta} \left(\frac{q^{\gamma n} - q^{\alpha n}}{q^{\gamma} - q^{\alpha}} + B \frac{q^{\gamma n} - (-1)^n q^{\alpha n}}{q^{\gamma} + q^{\alpha}} \right), & \text{if } \alpha \neq \gamma; \\ \lambda_0 q^{\gamma n} + n q^{\gamma (n + \varkappa_0 - 1) + \beta} + B q^{\gamma (n + \varkappa_0 - 1) + \beta} \left(\frac{1 - (-1)^n}{2} \right), & \text{if } \alpha = \gamma. \end{cases}$$
(30)

Equivalently, using the expression of the Casimir operator (25) and the representation (28) the solution λ_n of equation (29) can be presented by means of the eigenvalue c_3 of the Casimir operator C_3 in irreducible representation of the relation (10)

$$\lambda_n = \begin{cases} -q^{\gamma n} c_3 + q^{\beta} \left(\frac{q^{\gamma(n+\varkappa_0)} - q^{\alpha(n+\varkappa_0)}}{q^{\gamma} - q^{\alpha}} + 2\nu \frac{q^{\gamma(n+\varkappa_0)} - (-1)^{n+\varkappa_0} q^{\alpha(n+\varkappa_0)}}{q^{\gamma} + q^{\alpha}} \right), & \text{if } \alpha \neq \gamma; \\ \lambda_0 q^{\gamma n} c_3 + n q^{\gamma(n+\varkappa_0-1) + \beta} + B q^{\gamma(n+\varkappa_0-1) + \beta} \left(\frac{1 - (-1)^n}{2} \right), & \text{if } \alpha = \gamma. \end{cases}$$
(31)

It is easy see that λ_n in (30) coincides with the value of (31), where $B = (-1)^{\varkappa_0} 2\nu$.

According to [13] a representations of the generalized oscillator algebra is reduced to the four classes of unireps:

1. Representations bounded from below. They are defined by (28) and $n_1 \in \mathbb{Z}_{\leq 0}$ such that

$$\lambda_n = \begin{cases} \lambda_{n_1} = 0 & \text{if } n_1 \in \{\dots, -2, -1, 0\} \\ \lambda_n > 0 & \text{if } n \in \{n_1 + 1, n_1 + 2, \dots\}. \end{cases}$$
 (32)

(i) Let λ_n be defined by formula (30) for $\gamma - \alpha = 0$, q > 0. The non-negativity of λ_n implies

$$\lambda_n = \lambda_0 + nq^{\gamma(\varkappa_0 - 1) + \beta} + Bq^{\gamma(\varkappa_0 - 1) + \beta} \left(\frac{1 - (-1)^n}{2}\right) \ge 0.$$
 (33)

It follows (32) and due to (28) $a|n_1\rangle = 0$.

After possible renumbering of vectors $|n\rangle$ we obtain $a|0\rangle = 0$ and due to (28) one gets $\lambda_0 = 0$. Therefore the representations are given by formulae (28) with

$$\lambda_n = nq^{\gamma(n+\varkappa_0-1)+\beta} + Bq^{\gamma(n+\varkappa_0-1)+\beta} \left(\frac{1-(-1)^n}{2}\right),\tag{34}$$

where $n \geq 0$. The arbitrary values of the parameter \varkappa_0 , and $B \geq 0$ correspond to nonequivalent representations of (10).

(ii) Let λ_n be defined by (30) for $\gamma - \alpha > 0$, q > 1 ($\gamma - \alpha < 0$, 0 < q < 1). The assumption of the positivity at last one of the numbers $\frac{1}{q^{\gamma} - q^{\alpha}} \pm \frac{B}{q^{\gamma} + q^{\alpha}}$ and the nonnegativity of λ_n

$$\left(\lambda_0 q^{-(\alpha \varkappa + \beta)} + \frac{1}{q^{\gamma} - q^{\alpha}} + \frac{B}{q^{\gamma} + q^{\alpha}}\right) \ge q^{-(\gamma - \alpha)n} \left(\frac{1}{q^{\gamma} - q^{\alpha}} + \frac{(-1)^n B}{q^{\gamma} + q^{\alpha}}\right) \tag{35}$$

for $n \ge 0$ implies (32) and and due to (28) $a|n_1\rangle = 0$.

After possible renumbering of vectors $|n\rangle$ we obtain $a|0\rangle = 0$ and due to (28) one gets $\lambda_0 = 0$. Therefore the representations are given by (28) with

$$\lambda_n = q^{\alpha \varkappa_0 + \beta + n\gamma} \left(\frac{1 - q^{(\alpha - \gamma)n}}{q^{\gamma} - q^{\alpha}} + B \frac{1 - (-1)^n q^{(\alpha - \gamma)n}}{q^{\gamma} + q^{\alpha}} \right), \quad n \ge 0.$$
 (36)

The nonnegativity condition for λ_n implies $B \geq -1$.

The arbitrary values of the parameter \varkappa_0 , and $\lambda_0 = 0$ and B > -1 defined nonequivalent infinite-dimensional representations of (10).

iii) Let λ_n be given by (30) for $\gamma - \alpha > 0$, $0 < q < 1(\gamma - \alpha < 0, q > 1)$.

It can be represented by

$$\lambda_n = q^{\alpha \varkappa_0 + \beta + \gamma n} \times$$

$$\left\{ \left(\lambda_0 q^{-(\alpha \varkappa + \beta)} + \frac{1}{q^{\gamma} - q^{\alpha}} + \frac{B}{q^{\gamma} + q^{\alpha}} \right) - q^{-(\gamma - \alpha)n} \left(\frac{1}{q^{\gamma} - q^{\alpha}} + \frac{(-1)^n B}{q^{\gamma} + q^{\alpha}} \right) \right\}. \tag{37}$$

The nonnegativity of $\lambda_n \geq 0$, negativity of $\left(\lambda_0 q^{-(\alpha \varkappa + \beta)} + \frac{1}{q^{\gamma} - q^{\alpha}} + \frac{B}{q^{\gamma} + q^{\alpha}}\right) < 0$, and the non-positivity of $\left(\frac{1}{q^{\gamma}-q^{\alpha}}+\frac{(-1)^n B}{q^{\gamma}+q^{\alpha}}\right)$ implies (32) and $a|n_1\rangle=0$. The same arguments as in the items i), ii) give $a|0\rangle = 0$ and due to (28) $\lambda_0 = 0$. Therefore the representation is given by (28) with λ_n as (36). The nonnegativity condition for λ_n gives a restriction for possible values of B:

$$-1 \le B < -\frac{q^{\gamma} + q^{\alpha}}{q^{\gamma} + q^{\alpha}} \tag{38}$$

The arbitrary values of the parameter \varkappa_0 , and $-1 \leq B < -\frac{q^{\gamma} + q^{\alpha}}{q^{\gamma} + q^{\alpha}}$ and $\lambda_0 = 0$ distinguished irreducible representation of the relations (10).

2. Representations bounded from above. They are defined by formulae (28) and $n_2 \in \mathbb{N}$ such that

$$\lambda_n = \begin{cases} \lambda_{n_2} = 0 & \text{if } n_2 \in \{1, 2, 3, \dots\} \\ \lambda_n > 0 & \text{if } n \in \{n_2 - 1, n_2 - 2, \dots\}. \end{cases}$$
 (39)

Let λ_n be given by formula (30) $0 < q < 1, (\alpha - \gamma) > 0 \quad (q > 1, (\alpha - \gamma) < 0)$ and both values $\frac{1}{q^{\gamma}-q^{\alpha}} \pm \frac{B}{q^{\gamma}+q^{\alpha}}$ are non-positive (at last one of them must be strictly negative). From the non-negativity condition $\lambda_n \geq 0$ it follows the existence of n_2 such that (39) holds. From formulae (28) we obtain $a^+|n_2\rangle = 0$, and after possible renumbering we may assume that $a^+|0\rangle = 0$, and due to (28) $\lambda_1 = 0$.

The formula (33) implies $\lambda_0 = -q^{\alpha \varkappa_0 + \beta - \gamma} (1+B)$. The condition $\lambda_0 \ge 0$ is equivalent to $B \le -1$. It follows

$$\lambda_n = q^{\alpha \varkappa_0 + \beta + \gamma n} \left(\left(\frac{1 - q^{(\alpha - \gamma)n}}{q^{\gamma} - q^{\alpha}} - q^{-\gamma} \right) + B \left(\frac{1 - q^{(\alpha - \gamma)n}(-1)^n}{q^{\gamma} + q^{\alpha}} - q^{-\gamma} \right) \ge 0, \tag{40}$$

 $n \leq 0$. Therefore the representation is given by formulae (28) with λ_n as (40). If B < -1 the nonnegativity condition for λ_n gives a restriction for possible values of B:

$$B \le -\frac{q^{\gamma} + q^{\alpha}}{q^{\gamma} + q^{\alpha}} \tag{41}$$

The arbitrary values of parameter \varkappa_0 and $\lambda_0 = -q^{\alpha \varkappa_0 + \beta - \gamma} (1+B)$, $B < \frac{q^{\gamma} + q^{\alpha}}{q^{\gamma} + q^{\alpha}}$ distinguish irreducible representation of (10).

3. The finite dimensional representation. They defined by formulae (28) and $n_1 \in \{\ldots, -2, -1, 0\}, n_2 \in \{1, 2, 3, \ldots\}$ such that

$$\lambda_n = \begin{cases} \lambda_{n_1} = \lambda_{n_2} = 0 & \text{if } n_2 \in \{1, 2, 3, \dots\}, n_1 \in \{\dots, -2, -1, 0\} \\ \lambda_n > 0 & \text{if } n \in \{n_1 + 1, n_1 + 2, \dots, n_2 - 1\}. \end{cases}$$
(42)

i) One-dimensional representations. These representations are given by formulae (28) with λ_n calculated from (36), (37) and (40) at B = -1

$$\lambda_n = q^{-\alpha \varkappa_0 + \beta} \left(\frac{q^{\gamma n} - q^{\alpha n}}{q^{\gamma} - q^{\alpha}} - \frac{q^{\gamma n} - (-1)^n q^{\alpha n}}{q^{\gamma} + q^{\alpha}} \right). \tag{43}$$

In this case $n_1 = 0$ $(n_1 = -1)$ and $n_2 = 1$ $(n_2 = 0)$ and the representations (28) are defined by

$$a = a^{+} = 0, \quad N = \varkappa_0, \quad K = -\frac{1}{2\nu}.$$
 (44)

These representations are one-dimensional. Their are parametrized by \varkappa_0 .

ii) Two-dimensional representations. Let λ_n be defined (30) for $\gamma - \alpha > 0$, 0 < q < 1 ($\gamma - \alpha < 0, q > 1$). The values of $\lambda_n, n \ge 0$ in (37) are nonnegative if we assume $\left(\lambda_0 q^{-(\alpha \varkappa + \beta)} + \frac{1}{q^{\gamma} - q^{\alpha}} + \frac{B}{q^{\gamma} + q^{\alpha}}\right) = 0$ and both values $\left(\frac{1}{q^{\gamma} - q^{\alpha}} + \frac{(-1)^n B}{q^{\gamma} + q^{\alpha}}\right)$ are nonpositive. At $B = \pm \frac{q^{\gamma} + q^{\alpha}}{q^{\gamma} - q^{\alpha}}$ we obtain

$$\lambda_n = \frac{q^{\alpha(\varkappa_0 + n) + \beta}}{q^{\alpha} - q^{\gamma}} \Big(1 \pm (-1)^n \Big). \tag{45}$$

The value $\lambda_n = \frac{q^{\alpha(\varkappa_0+n)+\beta}}{q^{\alpha}-q^{\gamma}} \left(1+(-1)^n\right)$ implies (42) for $n_1=-1, n_2=1$ and $\lambda_0 = \frac{2q^{\alpha\varkappa_0+\beta}}{q^{\alpha}-q^{\gamma}}$.

The vector space of this representation spanned by the two-dimensional vectors

$$\left(\begin{array}{c} \psi_{-1} \\ \psi_0 \end{array}\right)$$

and due to (28) the representation are defined by

$$a = \begin{pmatrix} 0 & \sqrt{\frac{2q^{\alpha \varkappa_0 + \beta}}{q^{\alpha} - q^{\gamma}}} \\ 0 & 0 \end{pmatrix}, \qquad a^+ = \begin{pmatrix} 0 & 0 \\ \sqrt{\frac{2q^{\alpha \varkappa_0 + \beta}}{q^{\alpha} - q^{\gamma}}} & 0 \end{pmatrix},$$

$$N = \begin{pmatrix} \chi_0 - 1 & 0 \\ 0 & \chi_0 \end{pmatrix}, \qquad K = \frac{1}{2\nu} \frac{q^{\gamma} + q^{\alpha}}{q^{\alpha} - q^{\gamma}} \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}. \tag{46}$$

These representations are distinguished by the arbitrary values \varkappa_0 , and $B = \pm \frac{q^{\gamma} + q^{\alpha}}{q^{\gamma} - q^{\alpha}}$, $\lambda_0 = \frac{2q^{\alpha(\varkappa_0) + \beta}}{q^{\alpha} - q^{\gamma}}$.

The value $\lambda_n = \frac{q^{\alpha(\varkappa_0+n)+\beta}}{q^{\alpha}-q^{\gamma}} \left(1-(-1)^n\right)$ implies (42) for $n_1=0, n_2=2$. Moreover $\lambda_1 = \frac{2q^{\alpha(\varkappa_0+1)+\beta}}{q^{\alpha}-q^{\gamma}}$.

The vector space of this representation spanned by the two-dimensional vectors

$$\left(\begin{array}{c} \psi_0 \\ \psi_{-1} \end{array}\right)$$

and due to (28) the representations are given by

$$a = \begin{pmatrix} 0 & 0 \\ \sqrt{\frac{q^{\alpha(\varkappa_0+1)+\beta}}{q^{\alpha}-q^{\gamma}}} & 0 \end{pmatrix}, \qquad a^+ = \begin{pmatrix} 0 & \sqrt{\frac{q^{\alpha(\varkappa_0+1)+\beta}}{q^{\alpha}-q^{\gamma}}} \\ 0 & 0 \end{pmatrix},$$

$$N = \begin{pmatrix} \chi_0 & 0 \\ 0 & \chi_0+1 \end{pmatrix}, \qquad K = \frac{1}{2\nu} \frac{q^{\gamma}+q^{\alpha}}{q^{\alpha}-q^{\gamma}} \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix}. \tag{47}$$

These representations are defined by arbitrary values of \varkappa_0 , and $\lambda_0 = 0$, $B = -\frac{q^{\gamma} + q^{\alpha}}{q^{\gamma} - q^{\alpha}}$.

4. Unbounded representations. They defined by (28) with $\lambda_n \geq 0$ for all $n \in \mathbb{Z}$. As follows from (37) they are realized if $0 < q < 1, (\alpha - \gamma) > 0$ $(q > 1, (\alpha - \gamma) < 0)$ and

$$\lambda_0 q^{-\alpha \varkappa_0 - \beta} + \frac{1}{q^{\gamma} - q^{\alpha}} + \frac{B}{q^{\alpha} + q^{\gamma}} \ge 0, \quad |B| < -\frac{q^{\gamma} + q^{\alpha}}{q^{\gamma} - q^{\alpha}},\tag{48}$$

or

$$\lambda_0 q^{-\alpha \varkappa_0 - \beta} + \frac{1}{q^{\gamma} - q^{\alpha}} + \frac{B}{q^{\alpha} + q^{\gamma}} > 0, \quad |B| = -\frac{q^{\gamma} + q^{\alpha}}{q^{\gamma} - q^{\alpha}}. \tag{49}$$

Under the transformations

$$B \to B' = (-1)^n B, \quad \varkappa_0 \to \varkappa' = \varkappa + n,$$

$$\lambda_0 \to \lambda_0' = \lambda_0 q^{\gamma n} + q^{\alpha \varkappa_0 + \beta} \left(\frac{q^{\gamma n} - q^{\alpha n}}{q^{\gamma} - q^{\alpha}} + B \frac{q^{\gamma n} - (-1)^n q^{\alpha n}}{q^{\gamma} + q^{\alpha}} \right)$$
 (50)

the conditions (48) and (49) are preserved

$$\lambda_n' = \lambda_0 q^{2\gamma n} + q^{\alpha \varkappa_0 + \beta} \left(\frac{q^{2\gamma n} - q^{2\alpha n}}{q^{\gamma} - q^{\alpha}} + B \frac{q^{2\gamma n} - q^{2\alpha n}}{q^{\gamma} + q^{\alpha}} \right), \tag{51}$$

$$\lambda_0' q^{-\alpha \varkappa_0' - \beta} + \frac{1}{q^{\gamma} - q^{\alpha}} + \frac{B'}{q^{\alpha} + q^{\gamma}} = q^{(\gamma - \alpha)n} \left(\lambda_0 q^{-\alpha \varkappa_0 - \beta} + \frac{1}{q^{\gamma} - q^{\alpha}} + \frac{B}{q^{\alpha} + q^{\gamma}} \right) \tag{52}$$

and unbounded representations pass to themselves.

4. Spectrum of "quantized coordinate" Q

According to previous section if we put $\omega = 1, \varkappa_0 = 0$, in (36) we obtain the Fock representation of the relations (10)

$$N|n\rangle = n|n\rangle, \quad K|n\rangle = (-1)^{n}|n\rangle,$$

$$a^{+}|2n\rangle = q^{\beta/2} \left(\frac{q^{\gamma(2n+1)} - q^{\alpha(2n+1)}}{q^{\gamma} - q^{\alpha}} + 2\nu \frac{q^{\gamma(2n+1)} + q^{\alpha(2n+1)}}{q^{\gamma} + q^{\alpha}}\right)^{1/2} |2n + 1\rangle,$$

$$a^{+}|2n - 1\rangle = q^{\beta/2} \left(\frac{q^{2\gamma n} - q^{2\alpha n}}{q^{\gamma} - q^{\alpha}} + 2\nu \frac{q^{2\gamma n} - q^{2\alpha n}}{q^{\gamma} + q^{\alpha}}\right)^{1/2} |2n\rangle.$$
(53)

The properties of the quantum harmonic oscillator are closely related to Hermite polynomials. Different q-deformed oscillators generate distinct q-deformed Hermite polynomials and their generalization. Unlike to quantum harmonic oscillator, generalized q-deformed oscillators frequently define the position and momentum operators with the discrete spectrum and their eigenfunctions are represented by means of a q-Hermite polynomials. For some q-deformed oscillators this problem was studied in [20], [22], [21]. In [24] the generalized q-deformed oscillators connected with the discrete (generalize) q-Hermite I and q-Hermite II polynomials have been build. In this Section we show that corresponding oscillator algebras of this models are embedded in the $(q; \alpha, \beta, \gamma : \nu)$ -deformed oscillator algebra.

Let us consider the operator ("quantized coordinate") $Q = a^+ + a$, or

$$Q|n\rangle = r_n|n+1\rangle + r_{n-1}|n-1\rangle, \quad r_n = f^{1/2}(n+1).$$
 (54)

The self-adjointness and spectral properties of this operator is defined by polynomials $P_n(x)$ of first kind for the Jacobi matrix Q.

Defining the generalized eigenfunction $Q|x\rangle = x|x\rangle$, where $|x\rangle = \sum_{n=0}^{\infty} P_n(x)|n\rangle$, we obtain the recurrence relation

$$f^{1/2}(n)P_{n-1}(x) + f^{1/2}(n+1)P_{n+1}(x) = xP_n(x),$$
(55)

where $P_{-1}(x) = 0, P_0 = 1$.

In this Section we consider of the oscillator (10) for special choice of parameters

$$\alpha = 2a, \quad \gamma = 2a + c - 1 \quad \beta = 2a + b, \nu = 0.$$
 (56)

We obtain the relation [24]:

$$aa^{+} - q^{2a+c-1}a^{+}a = q^{2a(N+1)+b},$$
 (57)

or equivalent

$$aa^{+} - q^{2a}a^{+}a = q^{2a(N+1)+b}q^{(c-1)N}$$
(58)

which together with other relations of (10) form the corresponding oscillator algebra.

The self-adjoints of the operator (54) can be established by the Theorem 1.3 Chapter VII in [23]. For this purpose it is sufficient to prove the divergence of the series

$$\sum_{n=0}^{\infty} \frac{1}{r_n} = \infty. \tag{59}$$

The operator Q (54) is symmetric one. If

$$r_n \ge 0, \quad r_{n-1}r_{n+1} \le r_n^2, \quad \sum_{n=0}^{\infty} \frac{1}{r_n} < \infty.$$
 (60)

by the Theorem 1.5 Chapter VII in [23] its closure \bar{Q} is not self-adjoint operator.

In our case the property $r_n \geq 0$, $r_{n-1}r_{n+1} \leq r_n^2$ is satisfied automatically, therefore, the self-adjointness of the operator Q are reduced to the proving of the convergence of the series (59). We obtain the cases

$$q < 1, \quad \begin{cases} a < 0, & c - 1 > 0, & \text{divergent,} \\ a > 0, & c - 1 > 0, & \text{convergent,} \\ a + \frac{c - 1}{2} < 0, & c - 1 < 0, & \text{divergent,} \\ a + \frac{c - 1}{2} > 0, & c - 1 < 0, & \text{convergent.} \end{cases}$$

$$(61)$$

The equation (55) for this case take the form

$$xP_n(x;q) = \left(\frac{1}{1-q'}\right)^{1/2} q^{an+b/2} (1-q'^n)^{1/2} P_{n-1}(x;q)$$

$$+ \left(\frac{1}{1-q'}\right)^{1/2} q^{a(n+1)+b/2} (1-q'^{(n+1)})^{1/2} P_{n+1}(x;q), \tag{62}$$

where $q' = q^{c-1}$. If we change and rescale of the variables $y = (1 - q')^{1/2}x$, $P_n(x;q) = \psi_n((1 - q')^{1/2}x;q)$ then it follows

$$x\psi_n(x;q)$$

$$= q^{a(n+1)+b/2} (1 - q'^n)^{1/2} \psi_{n+1}(x;q) + q^{an+b} (1 - q'^n)^{1/2} \psi_{n-1}(x;q).$$
(63)

Representing $\psi_n(x;q)$ as

$$\psi_n(x;q) = \frac{q^{an^2/2}}{q^{(a+b)n/2}(q';q')_n^{1/2}} h_n(x;q)$$

we get from (63) the recurrence relation for the (generalized) discrete q-Hermite I polynomials $h_n(x;q)$:

$$xh_n(x;q) = h_{n+1}(x;q) + q^{2an+b}(1-q'^n)h_{n-1}(x;q).$$
(64)

The solution of this equation is given by anzatz

$$h_n(x;q) = \sum_{k=0}^{[n/2]} \frac{(q';q')_n}{\left((a_n,c_n);(1,q'^d)\right)_k} (-1)^k q^{(2an+b)k} q'^{k(k-n)} x^{n-2k}$$
(65)

where a_n, c_n, d are unknown quantities. The notation

$$\left((a,c); (p,q) \right)_k = \begin{cases} 1, & \text{if } k = 0; \\ (a-c)(ap-cq)\dots(ap^{k-1}-cq^{k-1}), & \text{otherwise} \end{cases}$$
(66)

we use from [26]. The substitution (65) into (64) leads to the equation

$$1 - q^{n+1} - q^{2an+b}(a_n - c_n q^{d(k-1)})q^{-2(k-1)} = 1 - q^{n-2k+1}$$
(67)

The solutions of this equation

$$a_n = q^{-2an-b}q^{\prime n-1}, c_n = q^{-2an-b}q^{\prime n+1}, d+2$$
 (68)

give the solutions of (64)

$$h_n(x;q) = \sum_{k=0}^{\lfloor n/2 \rfloor} \frac{(q';q')_n}{(q'';q'')_k(q';q')_{n-2k}} (-1)^k q^{(2an+b)k} q'^{k(k-n)} x^{n-2k}.$$
 (69)

They can be represented in terms of the basic hypergeometric function as

$$h_n(x;q) = x^n{}_2\phi_0\left(\begin{array}{cc} q'^{-n}, & q'^{-n+1} \\ - & q'^2; \frac{q^{2an+b}q'^n}{x^2} \end{array}\right).$$
 (70)

As has been shown in [24] these solutions at $a = \frac{1}{2}$, b = -1, c = 2 is reduced to discrete q-Hermite I polynomials and the relations (57) (or (58)) to equation of the corresponding q-deformed oscillator.

The solutions $P_n(x;q)$ of the equations (62) with the initial conditions $P_{-1}(x;q) = 0$, $P_0(x;q) = 1$ can be written as polynomials of degree n in x:

$$P_n(x;q) = \frac{q^{-an^2/2}}{q^{\frac{a+b}{2}n}(q';q')_n^{1/2}} h_n(\sqrt{1-q'}x;q).$$
 (71)

Now we restrict ourselves by the condition a = (c-1)/2 in (71). Then

$$P_n(x;q) = \frac{q^{-an(n-1)/2}}{(q';q')_n^{1/2}} h_n^0(q^{-(2a+b)/2} \sqrt{1-q'}x;q').$$
 (72)

These polynomials are orthogonal with respect to the discrete measure

$$d\omega(x) = \frac{q^{-(2a+b)}\sqrt{1-q'}}{2}(q';q'^2)_{\infty}\delta(x - \frac{q'^0}{q^{-(2a+b)}\sqrt{1-q'}})dx$$

$$+ \sum_{k>0} \frac{q^{-(2a+b)/2}\sqrt{1-q'}|x|}{2} \frac{(q^{-(2a+b)}(q'^2(1-q')x^2,q';q^2)_{\infty}}{(q';q')_{\infty}} \delta(x - \frac{q'^k}{q^{-(2a+b)/2}\sqrt{1-q'}})dx$$

$$+ \sum_{k>0} \frac{q^{-(2a+b)/2}\sqrt{1-q'}|x|}{2} \frac{(q^{-(2a+b)}(q'^2(1-q')x^2,q';q^2)_{\infty}}{(q';q')_{\infty}} \delta(x + \frac{q'^k}{q^{-(2a+b)/2}\sqrt{1-q'}})dx.$$

$$(73)$$

The orthogonality relation has the form

$$\frac{\delta_{mn}}{(q';q')_n} = \frac{1}{2} \frac{(q';q')_{\infty}}{(q'^2;q'^2)_{\infty}} P_m(1;q) P_n(1;q)
+ \sum_{k>0} \{ P_m(q'^k;q) P_n(q'^k;q) + P_m(-q'^k;q) P_n(-q'^k;q) \} \frac{q'^k}{2} \frac{(q'^{2k+2},q';q'^2)_{\infty}(q';q'^2)_{\infty}}{(q'^2;q'^2)_{\infty}}.$$
(74)

It follows that spectrum of the position operator Q is

$$SpQ = \left\{ \frac{\pm q^{(2a+b)/2}}{\sqrt{1-q'}}, \quad \frac{\pm q^{(2a+b)/2}q'}{\sqrt{1-q'}}, \dots, \frac{\pm q^{(2a+b)/2}q'^k}{\sqrt{1-q'}}, \dots; k \ge 0 \right\}.$$
 (75)

4. Hamiltonian and energy spectrum of $(q; \alpha, \beta, \gamma; \nu)$ -deformed oscillator

(g) Energy spectrum of free $(q; \alpha, \beta, \gamma; \nu)$ -deformed oscillator. In general, a Hamiltonian $H(a^+, a, N)$ of an oscillator is a function of the operators a^+, a, N . As an example, we consider the free standard Hamiltonian

$$H = \frac{\hbar\omega_0}{2} \{a^+, a\} = \frac{\hbar\omega_0}{2} (aa^+ + a^+ a) \tag{76}$$

of the $(q; \alpha, \beta, \gamma; \nu)$ -deformed oscillator.

Due to the structure function of deformation the free Hamiltonian H of the $(q; \alpha, \beta, \gamma; \nu)$ deformed oscillator in the Fock representation (53) can be written only as function of
number operator N

$$H = \frac{\hbar\omega_0}{2} q^{\beta} \left\{ \left(\frac{q^{\gamma N} - q^{\alpha N}}{q^{\gamma} - q^{\alpha}} + 2\nu \frac{q^{\gamma N} - (-1)^N q^{\alpha N}}{q^{\gamma} + q^{\alpha}} \right) + \left(\frac{q^{\gamma(N+1)} - q^{\alpha(N+1)}}{q^{\gamma} - q^{\alpha}} + 2\nu \frac{q^{\gamma(N+1)} - (-1)^{(N+1)} q^{\alpha(N+1)}}{q^{\gamma} + q^{\alpha}} \right) \right\}.$$
 (77)

For a study of the spectrum and the nonlinear frequency of this oscillator, it is convenient to use new parametrization

$$q = e^{\tau}, \quad \alpha = \rho - \mu, \quad \gamma = \rho + \mu.$$
 (78)

We have

$$E_{n} = \frac{\hbar\omega_{0}}{2}e^{\tau(\beta+\rho n)} \times \left\{ \frac{\sinh(\tau\mu(n+1))}{\sinh(\tau\mu)} + e^{-\tau\rho} \frac{\sinh(\tau\mu n)}{\sinh(\tau\mu)} + 2\nu \frac{1 - (-1)^{n}}{2} \left(\frac{\sinh(\tau\mu(n+1))}{\cosh(\tau\mu)} + e^{-\tau\rho} \frac{\cosh(\tau\mu n)}{\cosh(\tau\mu)} \right) + 2\nu \frac{1 + (-1)^{n}}{2} \left(\frac{\cosh(\tau\mu(n+1))}{\cosh(\tau\mu)} + e^{-\tau\rho} \frac{\sinh(\tau\mu n)}{\cosh(\tau\mu)} \right) \right\},$$
(79)

or

$$E_{n} = \frac{\hbar\omega_{0}}{2} e^{\tau(\beta-\rho)} \left\{ \frac{1 + e^{\tau(\rho+\mu)}}{2} \frac{e^{\tau(\rho+\mu)n}}{\sinh(\tau\mu)} - \frac{1 + e^{\tau(\rho-\mu)}}{2} \frac{e^{\tau(\rho-\mu)n}}{\sinh(\tau\mu)} + 2\nu \frac{1 - (-1)^{n}}{2} \left(\frac{1 + e^{\tau(\rho+\mu)}}{2} \frac{e^{\tau(\rho+\mu)n}}{\cosh(\tau\mu)} + \frac{1 - e^{\tau(\rho-\mu)}}{2} \frac{e^{\tau(\rho-\mu)n}}{\cosh(\tau\mu)} \right) + 2\nu \frac{1 + (-1)^{n}}{2} \left(\frac{1 + e^{\tau(\rho+\mu)}}{2} \frac{e^{\tau(\rho+\mu)n}}{\cosh(\tau\mu)} - \frac{1 - e^{\tau(\rho-\mu)}}{2} \frac{e^{\tau(\rho-\mu)n}}{\cosh(\tau\mu)} \right) \right\}.$$
(80)

It follows that the number of the energy levels is infinite, and the eigenvalues E_n for $\tau \neq 0, n \to \infty$ depend on the parameters α, γ via the exponential factor $e^{\tau(\rho - |\mu|)}$.

It is convenient to consider the eigenvalue E_n of H for n even and odd separately:

$$E_n = \frac{\hbar\omega_0}{2} e^{\tau(\beta - \rho)} \left\{ \frac{1 + e^{\tau(\rho + \mu)}}{2} \frac{e^{\tau(\rho + \mu)n}}{\sinh(\tau\mu)} - \frac{1 + e^{\tau(\rho - \mu)}}{2} \frac{e^{\tau(\rho - \mu)n}}{\sinh(\tau\mu)} + \frac{1 + e^{\tau(\rho - \mu)}}{2} \frac{e^{\tau(\rho - \mu)n}}{\sinh(\tau\mu)} + \frac{1 + e^{\tau(\rho - \mu)}}{2} \frac{e^{\tau(\rho - \mu)n}}{\sinh(\tau\mu)} + \frac{1 + e^{\tau(\rho - \mu)}}{2} \frac{e^{\tau(\rho - \mu)n}}{\sinh(\tau\mu)} + \frac{1 + e^{\tau(\rho - \mu)}}{2} \frac{e^{\tau(\rho - \mu)n}}{\sinh(\tau\mu)} + \frac{1 + e^{\tau(\rho - \mu)}}{2} \frac{e^{\tau(\rho - \mu)n}}{\sinh(\tau\mu)} + \frac{1 + e^{\tau(\rho - \mu)}}{2} \frac{e^{\tau(\rho - \mu)n}}{\sinh(\tau\mu)} + \frac{1 + e^{\tau(\rho - \mu)}}{2} \frac{e^{\tau(\rho - \mu)n}}{\sinh(\tau\mu)} + \frac{1 + e^{\tau(\rho - \mu)}}{2} \frac{e^{\tau(\rho - \mu)n}}{\sinh(\tau\mu)} + \frac{1 + e^{\tau(\rho - \mu)}}{2} \frac{e^{\tau(\rho - \mu)n}}{\sinh(\tau\mu)} + \frac{1 + e^{\tau(\rho - \mu)}}{2} \frac{e^{\tau(\rho - \mu)n}}{\sinh(\tau\mu)} + \frac{1 + e^{\tau(\rho - \mu)}}{2} \frac{e^{\tau(\rho - \mu)n}}{\sinh(\tau\mu)} + \frac{1 + e^{\tau(\rho - \mu)}}{2} \frac{e^{\tau(\rho - \mu)n}}{\sinh(\tau\mu)} + \frac{1 + e^{\tau(\rho - \mu)}}{2} \frac{e^{\tau(\rho - \mu)n}}{\sinh(\tau\mu)} + \frac{1 + e^{\tau(\rho - \mu)}}{2} \frac{e^{\tau(\rho - \mu)n}}{\sinh(\tau\mu)} + \frac{1 + e^{\tau(\rho - \mu)}}{2} \frac{e^{\tau(\rho - \mu)n}}{\sinh(\tau\mu)} + \frac{1 + e^{\tau(\rho - \mu)}}{2} \frac{e^{\tau(\rho - \mu)n}}{\sinh(\tau\mu)} + \frac{1 + e^{\tau(\rho - \mu)}}{2} \frac{e^{\tau(\rho - \mu)n}}{\sinh(\tau\mu)} + \frac{1 + e^{\tau(\rho - \mu)}}{2} \frac{e^{\tau(\rho - \mu)n}}{\sinh(\tau\mu)} + \frac{1 + e^{\tau(\rho - \mu)}}{2} \frac{e^{\tau(\rho - \mu)n}}{\sinh(\tau\mu)} + \frac{1 + e^{\tau(\rho - \mu)}}{2} \frac{e^{\tau(\rho - \mu)n}}{\sinh(\tau\mu)} + \frac{1 + e^{\tau(\rho - \mu)}}{2} \frac{e^{\tau(\rho - \mu)}}{\sinh(\tau\mu)} + \frac{1 + e^{\tau(\rho - \mu)}}{2} \frac{e^{\tau(\rho - \mu)}}{\sinh(\tau\mu)} + \frac{1 + e^{\tau(\rho - \mu)}}{2} \frac{e^{\tau(\rho - \mu)}}{\sinh(\tau\mu)} + \frac{1 + e^{\tau(\rho - \mu)}}{2} \frac{e^{\tau(\rho - \mu)}}{\sinh(\tau\mu)} + \frac{1 + e^{\tau(\rho - \mu)}}{2} \frac{e^{\tau(\rho - \mu)}}{\sinh(\tau\mu)} + \frac{1 + e^{\tau(\rho - \mu)}}{2} \frac{e^{\tau(\rho - \mu)}}{\sinh(\tau\mu)} + \frac{1 + e^{\tau(\rho - \mu)}}{2} \frac{e^{\tau(\rho - \mu)}}{\sinh(\tau\mu)} + \frac{1 + e^{\tau(\rho - \mu)}}{2} \frac{e^{\tau(\rho - \mu)}}{\sinh(\tau\mu)} + \frac{1 + e^{\tau(\rho - \mu)}}{2} \frac{e^{\tau(\rho - \mu)}}{\sinh(\tau\mu)} + \frac{1 + e^{\tau(\rho - \mu)}}{2} \frac{e^{\tau(\rho - \mu)}}{\sinh(\tau\mu)} + \frac{1 + e^{\tau(\rho - \mu)}}{2} \frac{e^{\tau(\rho - \mu)}}{\sinh(\tau\mu)} + \frac{1 + e^{\tau(\rho - \mu)}}{2} \frac{e^{\tau(\rho - \mu)}}{\sinh(\tau\mu)} + \frac{1 + e^{\tau(\rho - \mu)}}{2} \frac{e^{\tau(\rho - \mu)}}{\sinh(\tau\mu)} + \frac{1 + e^{\tau(\rho - \mu)}}{2} \frac{e^{\tau(\rho - \mu)}}{\sinh(\tau\mu)} + \frac{1 + e^{\tau(\rho - \mu)}}{2} \frac{e^{\tau(\rho - \mu)}}{\sinh(\tau\mu)} + \frac{1 + e^{\tau(\rho - \mu)}}{2} \frac{e^{\tau(\rho - \mu)}}{\sinh(\tau\mu)} + \frac{1 + e^{\tau(\rho - \mu)}}{2} \frac{e^{\tau(\rho - \mu)}}{\sinh(\tau\mu)} + \frac{1 + e^{\tau(\rho - \mu)}}{2} \frac{e^{\tau(\rho - \mu)}}{2}$$

$$2\nu \left(\frac{1 + e^{\tau(\rho + \mu)}}{2} \frac{e^{\tau(\rho + \mu)n}}{\cosh(\tau \mu)} + \frac{1 - e^{\tau(\rho - \mu)}}{2} \frac{e^{\tau(\rho - \mu)n}}{\cosh(\tau \mu)} \right) \right\}$$
(81)

for n odd, and

$$E_{n} = \frac{\hbar\omega_{0}}{2} e^{\tau(\beta-\rho)} \left\{ \frac{1 + e^{\tau(\rho+\mu)}}{2} \frac{e^{\tau(\rho+\mu)n}}{\sinh(\tau\mu)} - \frac{1 + e^{\tau(\rho-\mu)}}{2} \frac{e^{\tau(\rho-\mu)n}}{\sinh(\tau\mu)} + \frac{2\nu \left(\frac{1 + e^{\tau(\rho+\mu)}}{2} \frac{e^{\tau(\rho+\mu)n}}{\cosh(\tau\mu)} - \frac{1 - e^{\tau(\rho-\mu)}}{2} \frac{e^{\tau(\rho-\mu)n}}{\cosh(\tau\mu)} \right) \right\}$$
(82)

for n even. It follows the spectrum of this oscillator is not equidistant and the spacing is equal

$$E_{2n+1} - E_{2n} = \frac{\hbar\omega_0}{2} \frac{\sinh 2\mu\tau}{\sinh(\tau\mu)} e^{\tau(\beta+\rho)} e^{2\gamma\tau n}.$$
 (83)

(h) Asymptotic behaviour of energy spectrum of Hamiltonian H.

According to the analysis given above, we have to give the estimation of the spectrum of Hamiltonian (80) for the parameter $-1 < 2\nu$, if $\tau(\gamma - \alpha) > 0$ and $-1 < 2\nu < \coth(\tau(\gamma - \alpha)/2)$ if $\tau(\gamma - \alpha) < 0$. Three cases must be distinguished: $\tau(\rho + |\mu|) > 0$, $\tau(\rho + |\mu|) < 0$, $\tau(\rho + |\mu|) = 0$.

In special case $\mu = 0$, $(\alpha = \gamma = \rho)$ we have

$$E_n = \frac{\hbar\omega_0}{2} e^{\tau(\beta + \rho n)} \Big[(n+\nu)(1 + e^{-\tau\rho}) + (-1)^n \nu (1 - e^{-\tau\rho}) + 1 \Big]. \tag{84}$$

If $(\tau(\rho + |\mu|) > 0$ and $-1 < 2\nu$, then from (80) it implies that the energy grows to infinity with increasing n.

If $\tau(\rho + |\mu|) = 0$, then in the case $(\rho = \mu)$ we have

$$E_n = \frac{\hbar\omega_0}{2}e^{\tau\beta}e^{\beta-\rho}\left\{\frac{e^{2\tau\rho n} - 1}{e^{\tau\mu} - e^{-\tau\mu}} + \frac{e^{2\tau\rho(n+1)} - 1}{e^{\tau\mu} - e^{-\tau\mu}} + 2\nu\frac{e^{2\rho\tau n} - e^{2\rho\tau(n+1)}}{e^{\tau\mu} - e^{-\tau\mu}}\right\}.$$
 (85)

The same holds for $\rho + \mu = 0$.

It follows that E_n monotonically increase to upper bounds that depends on parameters

$$E_{\max} = \frac{\hbar\omega_0}{2} \frac{e^{\tau(\beta - \rho)}}{\sinh \tau |\mu|}.$$

If $\tau(\rho + |\mu|) < 0$ and $-1 < 2\nu$ then the energies (81) and (82) at first increase and then go to zero as $n \to \infty$. In the particular case $\mu = 0$ in (81) and (82) we have

$$E_n = \frac{\hbar\omega_0}{2} e^{\tau(\beta + \rho n)} \Big\{ n(1 + e^{-\tau \rho}) + 2\nu + 1 \Big\},\,$$

for n odd and

$$E_n = \frac{\hbar\omega_0}{2} e^{\tau(\beta + \rho n)} \left\{ n(1 + e^{-\tau \rho}) + 2\nu e^{-\tau \rho} + 1 \right\}$$

for n even.

Acknowledgements

This research was partially supported by Grant 14.01/016 of the State Foundation of Fundamental Research of Ukraine.

References

- [1] Odaka K., Kishi T., Kamefuchi S., J. Phys. A 24, L591 (1991).
- [2] Daskaloyannis C., J. Phys. A 24, L789 (1991).
- [3] Chung W.-S., K.-S. Chung, S.-T. Nam, Um, C.-I., Phys. Lett. A 183, 363 (1993).
- [4] Meljanac S., Milecovic M., Pallua S., Phys. Lett. B 328, 55 (1994).
- [5] Arik M., Coon D. D., Lett. **75**, 4724 (1995).
- [6] Biedenharn L.C., J. Phys. A: Math. Gen. 22 L873 (1989).
- [7] Macfarlane A. J., J. Phys. A 22, 4581 (1989).
- [8] Borzov, E. V. Damaskinsky, S. V. Yegorov, Some remarks on the representations of the generalized deformed algebra, q-alg/9509022.
- [9] Vasiliev M. A., Int. J. Mod. Phys. A 6 1115 (1991).
- [10] Brink L., Hanson T. H., Vasiliev M. A., Phys. Let. B 276, 109 (1992).
- [11] Calogero F., J. Math. Phys. 10, 2191 (1969).
- [12] Brezinski T., Egusquiza I. L., Macfarlane A., Phys. Lett. B 276, 109 (1992).
- [13] Quesne, C., Vanstineekiste N., C_{λ} -extended oscillator algebras and some of their deformation, math-ph/0003025.
- [14] Quesne, C., Vanstineekiste N., Representation theory of generalized oscillator algebras, q-alg/9701031.
- [15] Burban I. M., Phys. Let. A **366**, 308 (2007).
- [16] Rideau G., Lett. Math. Phys. 24, 147 (1992).
- [17] Chaichian H., Grosse P., J. Phys. Math. Gen. A27, 2045 (1994).
- [18] Kosinski P., Majevski, M., Maslianka, P., Representation of generalized oscillator algebra, q-alg/9501012.
- [19] Mizrahi S. S., Lima J. P., Dodonov V. V., J. Phys. A: Math. Gen. 37, 3707 (2004).
- [20] Burban I. M., Klimyk A. U., Letter in Math. Phys. 29, 13 (1993).
- [21] Borzov V. V., Damaskinsky E. V., Kulish P. P., On position operator spectral measure for deformed oscillator in the case of indetermined Hamburger problem, math.QA/9803099.

- [22] Chung W-S., Klimyk A. U., J. Math. Phys. 37 917 (1996).
- [23] Berezansky Yu. M., Expansion in eigenfunctions of self-adjoint operators. Amer. Math. Soc., Providence, R. I. 1968.
- [24] Burban I. M., Generalized q-deformed oscillator, q-Hermite polynomials, generalized coherent states, math-ph/0607045.
- [25] Koekoek R., Swarttouw R. F., The Askey-Scheme of Hypergeometric Orthogonal Polynomials and Its q-analogue (Delf University of Technology, Report 98-17, 1998).
- [26] Jagannathan, R., Rao K., Two-parameter quantum algebras, twin-basic numbers and associated generalised hypergeometric series, math. NT/06026513.