## Non-linear realization of Poincaré invariance in the graph-representation of extremal hypersurfaces

Jens Hoppe \*
Department of Mathematics,
Royal Institute of Technology,
KTH, 100 44 Stockholm,
Sweden

## Abstract

In the Born-Infeld 'harmonic gauge' description of M-branes moving in  $\mathbb{R}^{M+1}$  the underlying M+2 dimensional Poincaré - invariance gives rise to an interesting system of conservation laws showing signs of integrability.

Varying

$$S[z] = -\int d^{M}x dt \sqrt{1 - \dot{z}^{2} + (\vec{\nabla}z)^{2}}$$
 (1)

yields the quasi-linear PDE

$$(1 - z^{\alpha} z_{\alpha}) \Box z + z^{\alpha} z^{\beta} z_{\alpha\beta} = 0 \tag{2}$$

for 
$$z(t, \vec{x})$$
, with  $z^{\alpha} := \eta^{\alpha\beta} z_{\beta} = \eta^{\alpha\beta} \frac{\partial z}{\partial x^{\beta}}$ ,  $(\alpha, \beta = 0, 1, \dots, M)$ ,  $\eta^{\alpha\beta} = diag(1, -1, \dots, -1)$ ,  $z_{\alpha\beta} = \frac{\partial^2 z}{\partial x^{\alpha} \partial x^{\beta}}$ ,  $\Box z = \eta^{\alpha\beta} z_{\alpha\beta}$ .

At first sight, (2) appears to be a scalar field equation in M+1 - dimensional Minkowski space, but M+2 additional conservation laws follow when

<sup>\*</sup>e-mail: hoppe@math.kth.se

realizing that z is most naturally interpreted as a (for notational convenience, the last) component of a (M + 2) - vector

$$x^{\mu} = \begin{pmatrix} t \\ \vec{x} \\ z(t, \vec{x}) \end{pmatrix} \tag{3}$$

in  $\mathbb{R}^{1,M+1}$  (cp. e.g. [1]). The corresponding world-volume  $\mathcal{M}$ , described by (3)/(2), can easily be shown to have vanishing mean curvature. The M+2-dimensional Poincaré - symmetry (non-linearly realized, as  $x^{M+1}=z(t,\vec{x})$ ) is expressed by

$$\partial_{\alpha} \mathcal{H}^{\alpha \mu} = 0 \quad (\mu = 0, 1, \dots, M + 1) \tag{4}$$

$$\partial_{\alpha}(x^{\mu}\mathcal{H}^{\nu\alpha} - x^{\nu}\mathcal{H}^{\mu\alpha}) = 0 \tag{5}$$

$$\mathcal{H}^{\alpha\beta} := \frac{z^{\alpha}z^{\beta}}{\sqrt{1 - z^{\gamma}z_{\gamma}}} + \eta^{\alpha\beta}\sqrt{1 - z^{\gamma}z_{\gamma}}$$
 (6)

$$\mathcal{H}^{\alpha,M+1} := \frac{z^{\alpha}}{\sqrt{1 - z^{\gamma} z_{\gamma}}} = \mathcal{H}^{M+1,\alpha} \tag{7}$$

(generalising to arbitrary co-dimension is straightforward).

Note in particular that

$$z_{\alpha}\mathcal{H}^{\alpha\beta} = \mathcal{H}^{\alpha,M+1} \tag{8}$$

Let me make a few comments how one may be able to use (4) and (5).

First of all note the monotonicity of

$$\int x^{\mu} \mathcal{H}^{00} = C^{\mu} + tP^{\mu} \tag{9}$$

 $(C^{\mu} \text{ and } P^{\mu} = \int \mathcal{H}^{\mu 0} \text{ being constants, depending only on the initial conditions)};$  and the explicit (polynomial; here linear) appearance of t.

Secondly, one could hope that the conservation laws, in particular the  $\mu \neq M+1$  part of (4) (containing neither  $x^{\alpha}$  nor z!), thought of as a first-order system linear in the derivatives of the M+1-vector

$$z^{\alpha} = \begin{pmatrix} \dot{z} \\ -\vec{\nabla}z \end{pmatrix}, \tag{10}$$

resp.  $u^{\alpha} = \frac{z^{\alpha}}{\sqrt{1-z^{\gamma}z_{\gamma}}}$ , resp.  $\mathcal{H}^{\alpha 0}$  (or other suitably chosen coordinates; e.g. in order to apply the method of characteristics) are *integrable*.

**Acknowledgments** I would like to thank M. Bordemann, B. Gustafsson, E. Langmann, J. Reinhardt, M. Trzetrzelewski and A. Zheltukhin for valuable discussions.

## References

 $[1]\ \, \mathrm{M.\ Bordemann},\ \, \mathrm{J.\ Hoppe},\ \mathrm{PLB\ 325\ (1994)\ 359\text{--}365},$