

On the Initial-Value Problem to the Quantum Dual BBGKY Hierarchy

G. Borgioli[†] and *V. Gerasimenko*[‡]

[†]*Università degli Studi di Firenze, Via S.Marta, 3, 50139 Firenze, Italy*

[‡]*Institute of Mathematics of NAS of Ukraine, Tereshchenkivs'ka Str., 3, 01601 Kyiv-4, Ukraine*

E-mail: [†]*giovanni.borgioli@unifi.it*, [‡]*gerasym@imath.kiev.ua*

We develop a rigorous formalism for the description of the evolution of observables in quantum systems of particles. We construct a solution of the initial-value problem to the quantum dual BBGKY hierarchy of equations as an expansion over particle clusters whose evolution are governed by the corresponding-order dual cumulant (dual semi-invariant) of the evolution operators of finitely many particles. For initial data from the space of sequences of bounded operators the existence and uniqueness theorem is proved.

Key words: quantum dual BBGKY hierarchy; dual cumulant (dual semi-invariant); quantum many-particle system.

2000 Mathematics Subject Classification: 35Q40; 47d06.

Contents

1	Introduction	3
1.1	The initial-value problem to the Heisenberg equation	4
1.2	Quantum many-particle systems	6
2	The quantum dual BBGKY hierarchy	8
2.1	The initial-value problem to the quantum dual BBGKY hierarchy	8
2.2	The formula for a solution and its representations	10
2.3	Dual cumulants of groups of evolution operators	13
3	A solution of the initial-value problem to the quantum dual BBGKY hierarchy	15
3.1	A group of operators for the quantum dual BBGKY hierarchy	16
3.2	The existence and uniqueness theorem	17
4	The existence of the mean value observable functional	19
5	Conclusion	20
	References	20

1 Introduction

Evolution equations of quantum many-particle systems arise in many problems of modern statistical mechanics. [1,10,12,18]. In the theory of such equations, during the last decade, many new results have been obtained, in particular concerning the fundamental problem of the rigorous derivation of quantum kinetic equations [19]- [27].

A description of quantum many-particle systems can be formulated in terms of two sets of objects: observables and states. The mean value defines a duality between observables and states and as a consequence there exist two approaches to the description of the evolution. Usually the evolution of many-particle systems is described, in the framework of the evolution of states, by the BBGKY hierarchy for marginal density operators. In the papers [21]- [27] a solution of the Cauchy problem to the quantum BBGKY hierarchy is constructed in the form of iteration series for initial data in the space of sequences of trace class operators. In [13,14] for the quantum BBGKY hierarchy (for the classical many-particle systems in [15,16]) a solution is represented in the form of series over particle clusters, whose evolution is described by the corresponding order cumulant (semi-invariant) of evolution operators of finitely many particles. Using an analog of Duhamel formulas, such a solution expansion reduces to an iteration series, which is valid for a particular class of initial data and interaction potentials. An equivalent approach for the description of many-particle system evolution is given by the evolution of observables and by the dual BBGKY hierarchy. For classical systems this approach is studied in the paper [2].

In this paper we consider the initial-value problem to the quantum dual BBGKY hierarchy of equations describing the evolution of observables for both finitely and infinitely many quantum particles, obeying Maxwell-Boltzmann statistics. We construct a solution of the Cauchy problem in the form of an expansion over clusters of the decreasing number of particles, whose evolution is governed by the corresponding-order dual cumulant (dual semi-invariant) of the evolution groups of operators of finitely many particles (groups of operators for the Heisenberg equations).

We now outline the structure of the paper and the main results.

In Section 2 we define the dual BBGKY hierarchy for quantum systems with a general type of an interaction potential between particles and construct a solution of the Cauchy problem to this hierarchy. We define the dual cumulants (the dual semi-invariants) of evolution operators for the Heisenberg equations and investigate some of their typical properties. The concept of cumulants of evolution operators forms the basis for the solution expansions of various evolution equations for infinitely many particles [13,16]. In Section 3 we study the properties of a one-parameter group of automorphisms generated by a solution of the initial-value problem to the dual BBGKY hierarchy in the space of sequences of bounded operators and prove the existence and uniqueness theorem of a classical and a generalized solution for the corresponding Cauchy problem. In Section 4 we consider the description of evolution for infinite-particle systems in the Heisenberg picture and prove the existence of mean value functionals for observables. In the final section we give some concluding remarks.

Let us introduce some preliminary definitions and results for the description of quantum many-particle systems of a non-fixed number of particles.

1.1 The initial-value problem to the Heisenberg equation

We consider a quantum system of a non-fixed number of identical (spinless) particles with unit mass $m = 1$ in the space \mathbb{R}^ν , $\nu \geq 1$ (in the terminology of statistical mechanics it is known as non-equilibrium grand canonical ensemble [1]).

Let a sequence $g = (I, g_1, \dots, g_n, \dots)$ be an infinite sequence of self-adjoint bounded operators g_n defined on the Fock space $\mathcal{F}_{\mathcal{H}} = \bigoplus_{n=0}^{\infty} \mathcal{H}^{\otimes n}$ over the Hilbert space \mathcal{H} ($\mathcal{H}^0 = \mathbb{C}$ and I is a unit operator). An operator g_n , defined in the n -particle Hilbert space $\mathcal{H}_n = \mathcal{H}^{\otimes n}$, will be denoted by $g_n(1, \dots, n)$. For a system of identical particles obeying Maxwell-Boltzmann statistics, one has $g_n(1, \dots, n) = g_n(i_1, \dots, i_n)$ for any permutation of indexes $\{i_1, \dots, i_n\} \in \{1, \dots, n\}$.

Let the space $\mathfrak{L}(\mathcal{F}_{\mathcal{H}})$ be the space of sequences $g = (I, g_1, \dots, g_n, \dots)$ of bounded operators g_n (I is a unit operator) defined on the Hilbert space \mathcal{H}_n and satisfying symmetry property: $g_n(1, \dots, n) = g_n(i_1, \dots, i_n)$, if $\{i_1, \dots, i_n\} \in \{1, \dots, n\}$, with an operator norm [5]. We will also consider a more general space $\mathfrak{L}_\gamma(\mathcal{F}_{\mathcal{H}})$ with a norm

$$\|g\|_{\mathfrak{L}_\gamma(\mathcal{F}_{\mathcal{H}})} := \max_{n \geq 0} \frac{\gamma^n}{n!} \|g_n\|_{\mathfrak{L}(\mathcal{H}_n)},$$

where $0 < \gamma < 1$ and $\|\cdot\|_{\mathfrak{L}(\mathcal{H}_n)}$ is an operator norm [5].

An observable of finitely many quantum particles is a sequence of self-adjoint operators from $\mathfrak{L}_\gamma(\mathcal{F}_{\mathcal{H}})$. The case of the unbounded observables can be reduced to the case under consideration [3].

The evolution of observables $A(t) = (I, A_1(t, 1), \dots, A_n(t, 1, \dots, n), \dots)$ is described by the initial-value problem to the Heisenberg equation [3, 12]

$$\frac{d}{dt} A(t) = \mathcal{N} A(t), \quad (1)$$

$$A(t)|_{t=0} = A(0), \quad (2)$$

where, if $g \in \mathcal{D}(\mathcal{N}) \subset \mathfrak{L}(\mathcal{F}_{\mathcal{H}})$, the von Neumann operator $\mathcal{N} = \bigoplus_{n=0}^{\infty} \mathcal{N}_n$ is defined by the formula

$$(\mathcal{N}g)_n := -\frac{i}{\hbar} (g_n H_n - H_n g_n). \quad (3)$$

$h = 2\pi\hbar$ is the Planck constant and the Hamiltonian $H = \bigoplus_{n=0}^{\infty} H_n$ in (3) is a self-adjoint operator with a domain of the definition $\mathcal{D}(H) = \{\psi = \bigoplus_{n=0}^{\infty} \psi_n \in \mathcal{F}_{\mathcal{H}} \mid \psi_n \in \mathcal{D}(H_n) \subset \mathcal{H}_n, \sum_n \|H_n \psi_n\|^2 < \infty\} \subset \mathcal{F}_{\mathcal{H}}$ [5].

Assume $\mathcal{H} = L^2(\mathbb{R}^\nu)$, then an element $\psi \in \mathcal{F}_{\mathcal{H}} = \bigoplus_{n=0}^{\infty} L^2(\mathbb{R}^{\nu n})$ is a sequence of functions $\psi = (\psi_0, \psi_1(q_1), \dots, \psi_n(q_1, \dots, q_n), \dots)$ such that $\|\psi\|^2 = |\psi_0|^2 + \sum_{n=1}^{\infty} \int dq_1 \dots dq_n |\psi_n(q_1, \dots, q_n)|^2 < +\infty$. On the subspace $L_0^2(\mathbb{R}^{\nu n}) \subset L^2(\mathbb{R}^{\nu n})$ of infinitely differentiable functions with compact support the n -particle Hamiltonian H_n acts according to the formula ($H_0 = 0$)

$$H_n \psi_n = -\frac{\hbar^2}{2} \sum_{i=1}^n \Delta_{q_i} \psi_n + \sum_{k=1}^n \sum_{i_1 < \dots < i_k=1}^n \Phi^{(k)}(q_{i_1}, \dots, q_{i_k}) \psi_n, \quad (4)$$

where $\Phi^{(k)}$ is a k -body interaction potential satisfying Kato conditions [5].

On the space $\mathfrak{L}(\mathcal{F}_{\mathcal{H}})$ a solution of the Cauchy problem to the Heisenberg equation (1) is determined by the following one-parameter family of operators

$$\mathbb{R}^1 \ni t \mapsto \mathcal{G}(t)g := \mathcal{U}(t)g\mathcal{U}^{-1}(t), \quad (5)$$

where $\mathcal{U}(t) = \bigoplus_{n=0}^{\infty} \mathcal{U}_n(t)$ and operators $\mathcal{U}_n(t)$, $\mathcal{U}_n^{-1}(t)$ are defined as follows

$$\begin{aligned} \mathcal{U}_n(t) &:= e^{\frac{i}{\hbar}tH_n}, \\ \mathcal{U}_n^{-1}(t) &:= e^{-\frac{i}{\hbar}tH_n} \end{aligned} \quad (6)$$

and $\mathcal{U}_0(t) = I$ is a unit operator.

Proposition 1. *On the space $\mathfrak{L}_{\gamma}(\mathcal{F}_{\mathcal{H}})$ one-parameter mapping (5)*

$$\mathbb{R}^1 \ni t \mapsto \mathcal{G}(t)g := \mathcal{U}(t)g\mathcal{U}^{-1}(t)$$

defines an isometric $$ -weak continuous group of operators, i.e. it is a C_0^* -group. The infinitesimal generator $\mathcal{N} = \bigoplus_{n=0}^{\infty} \mathcal{N}_n$ of this group of operators is a closed operator for the $*$ -weak topology and on its domain of the definition $\mathcal{D}(\mathcal{N}) \subset \mathfrak{L}_{\gamma}(\mathcal{F}_{\mathcal{H}})$ which everywhere dense for the $*$ -weak topology, \mathcal{N} is defined as follows in the sense of the $*$ -weak convergence of the space $\mathfrak{L}_{\gamma}(\mathcal{F}_{\mathcal{H}})$*

$$\text{w}^* - \lim_{t \rightarrow 0} \frac{1}{t} (\mathcal{G}(t)g - g) = -\frac{i}{\hbar} (gH - Hg), \quad (7)$$

where $H = \bigoplus_{n=0}^{\infty} H_n$ is the Hamiltonian (4) and the operator: $\mathcal{N}g = (-i/\hbar)(gH - Hg)$ is defined on the domain $\mathcal{D}(H) \subset \mathcal{F}_{\mathcal{H}}$.

The validity of this statement follows from properties of one-parameter groups (6) [3], [4]. Group of operators (5) preserves the self-adjointness of operators.

On the space $\mathfrak{L}_{\gamma}(\mathcal{F}_{\mathcal{H}})$, for an abstract initial-value problem (1)-(2), the following statement holds

Proposition 2. *A solution of the initial-value problem to the Heisenberg equation (1)-(2) is determined by*

$$A(t) = \mathcal{G}(t)A(0), \quad (8)$$

where the one-parameter family $\{\mathcal{G}(t)\}_{t \in \mathbb{R}}$ of operators is defined by expression (5). For $A(0) \in \mathcal{D}(\mathcal{N}) \subset \mathfrak{L}_{\gamma}(\mathcal{F}_{\mathcal{H}})$, $A(t)$ is a classical solution and for arbitrary initial data $A(0) \in \mathfrak{L}_{\gamma}(\mathcal{F}_{\mathcal{H}})$, $A(t)$ is a generalized solution.

According general theorems about properties of the dual semigroup [4], [6], [7] the validity of these propositions follows also from the fact that the group $\{\mathcal{G}(t)\}_{t \in \mathbb{R}}$ is dual to the strong continuous group of the von Neumann equation defined on the space of sequences of trace class operators [3].

We remark that, in the framework of kernels and symbols of operators, infinitesimal generator (7) and groups (6) are studied in [11].

1.2 Quantum many-particle systems

A positive continuous linear functional on the space of observables $A(t) \in \mathfrak{L}(\mathcal{F}_{\mathcal{H}})$, whose value is interpreted as its mean value (average value of an observable), is defined as follows [4]

$$\langle A(t) | D(0) \rangle := \left(\sum_{n=0}^{\infty} \frac{1}{n!} \text{Tr}_{1, \dots, n} D_n(0) \right)^{-1} \sum_{n=0}^{\infty} \frac{1}{n!} \text{Tr}_{1, \dots, n} A_n(t, 1, \dots, n) D_n(0, 1, \dots, n), \quad (9)$$

where $\text{Tr}_{1, \dots, n}$ is the partial trace over $1, \dots, n$ particles and $\sum_{n=0}^{\infty} \frac{1}{n!} \text{Tr}_{1, \dots, n} D_n(0)$ is a normalizing factor (grand canonical partition function). The sequences $D(0) = (I, D_1(0, 1), \dots, D_n(0, 1, \dots, n), \dots)$ of positive self-adjoint operators D_n , $n \geq 1$, (the density operators whose kernels are known as the density matrices [11]) defined on the n -particle Hilbert space $\mathcal{H}_n = \mathcal{H}^{\otimes n} = L^2(\mathbb{R}^{\nu n})$, describe the states of a quantum system of non-fixed number of particles.

Usually it is assumed that the states belong to the space $\mathfrak{L}_{\alpha}^1(\mathcal{F}_{\mathcal{H}}) = \bigoplus_{n=0}^{\infty} \alpha^n \mathfrak{L}^1(\mathcal{H}_n)$ of sequences $f = (I, f_1, \dots, f_n, \dots)$ of trace class operators $f_n = f_n(1, \dots, n) \in \mathfrak{L}^1(\mathcal{H}_n)$, satisfying the Maxwell-Boltzmann statistics symmetry condition, equipped with the trace norm

$$\|f\|_{\mathfrak{L}_{\alpha}^1(\mathcal{F}_{\mathcal{H}})} := \sum_{n=0}^{\infty} \alpha^n \|f_n\|_{\mathfrak{L}^1(\mathcal{H}_n)} := \sum_{n=0}^{\infty} \alpha^n |\text{Tr}_{1, \dots, n} f_n(1, \dots, n)|,$$

where $\alpha > 1$ is a real number, $\text{Tr}_{1, \dots, n}$ is the partial trace over $1, \dots, n$ particles. We will denote by $\mathfrak{L}_{\alpha, 0}^1$ the everywhere dense set in $\mathfrak{L}_{\alpha}^1(\mathcal{F}_{\mathcal{H}})$ of finite sequences of degenerate operators (finite-rank operators) [5] with infinitely differentiable kernels with compact support. The space $\mathfrak{L}_{\alpha}^1(\mathcal{F}_{\mathcal{H}})$ contains sequences of operators more general than those determining the states of systems. We will also consider the space $\mathfrak{L}^1(\mathcal{F}_{\mathcal{H}}) = \bigoplus_{n=0}^{\infty} \mathfrak{L}^1(\mathcal{H}_n)$.

If $A(t) \in \mathfrak{L}_{\gamma}(\mathcal{F}_{\mathcal{H}})$ and $D(0) \in \mathfrak{L}_{\alpha}^1(\mathcal{F}_{\mathcal{H}})$, then functional (9) exists.

It is well known that there exists an equivalent approach to the description of the evolution of quantum systems, which is given by the evolution of states. In this case the evolution of all possible states $D(t) = (I, D_1(t, 1), \dots, D_n(t, 1, \dots, n), \dots)$ is described by the initial-value problem to the von Neumann equation [3, 12]

$$\frac{d}{dt} D(t) = -\mathcal{N} D(t), \quad (10)$$

$$D(t)|_{t=0} = D(0), \quad (11)$$

where if $f \in \mathfrak{L}_0^1(\mathcal{F}_{\mathcal{H}}) \subset \mathcal{D}(\mathcal{N}) \subset \mathfrak{L}^1(\mathcal{F}_{\mathcal{H}})$ the von Neumann operator $(-\mathcal{N})$ is defined by

$$(-\mathcal{N}f)_n := -\frac{i}{\hbar} (H_n f_n - f_n H_n), \quad (12)$$

The von Neumann equation (10) is dual to the Heisenberg equation (1) with respect to bilinear form (9).

For initial data $D(0)$ from the space $\mathfrak{L}^1(\mathcal{F}_{\mathcal{H}})$ the solution

$$D(t) = \mathcal{G}(-t) D(0)$$

of the Cauchy problem to the von Neumann equation (10)-(11) is determined by the following one-parameter family of operators on $\mathfrak{L}^1(\mathcal{F}_{\mathcal{H}})$

$$\mathbb{R}^1 \ni t \mapsto \mathcal{G}(-t)f := \mathcal{U}(-t)f\mathcal{U}^{-1}(-t), \quad (13)$$

where the operators $\mathcal{U}_n(-t)$, $\mathcal{U}_n^{-1}(-t)$ are defined by (6).

The properties of a one-parameter family $\{\mathcal{G}(-t)\}_{t \in \mathbb{R}}$ of operators (13) follow from the properties of groups (6).

Proposition 3. *On the space $\mathfrak{L}^1(\mathcal{F}_{\mathcal{H}})$ one-parameter mapping (13)*

$$\mathbb{R}^1 \ni t \mapsto \mathcal{G}(-t)f := \mathcal{U}(-t)f\mathcal{U}^{-1}(-t)$$

defines an isometric strongly continuous group of operators, i.e. it is a C_0 -group, which preserves positivity and self-adjointness of operators. If $f \in \mathfrak{L}_0^1(\mathcal{F}_{\mathcal{H}}) \subset \mathcal{D}(-\mathcal{N}) \subset \mathfrak{L}^1(\mathcal{F}_{\mathcal{H}})$ then the infinitesimal generator: $-\mathcal{N} = \bigoplus_{n=0}^{\infty} (-\mathcal{N}_n)$ of this group of operators is determined in the sense of the norm convergence in the space $\mathfrak{L}^1(\mathcal{F}_{\mathcal{H}})$ as follows

$$\lim_{t \rightarrow 0} \frac{1}{t} (\mathcal{G}(-t)f - f) = -\frac{i}{\hbar} (Hf - fH) = -\mathcal{N}f, \quad (14)$$

where $H = \bigoplus_{n=0}^{\infty} H_n$ is the Hamiltonian (4) and the operator: $(-i/\hbar)(Hf - fH)$ is defined on the domain $\mathcal{D}(H) \subset \mathcal{F}_{\mathcal{H}}$.

Thus, if we consider group (5) as the dual to group (13), from the above Proposition it follows the statement about properties of group (5).

For a system of a finite average number of particles there exists an equivalent possibility to describe observables and states, namely, by sequences of marginal observables (the so-called s -particle observables) $G(t) = (G_0, G_1(t, 1), \dots, G_s(t, 1, \dots, s), \dots)$ and marginal states (or s -particle density operators) $F(0) = (I, F_1(0, 1), \dots, F_s(0, 1, \dots, s), \dots)$ [1], [12]. These sequences are correspondingly introduced instead of sequences $A(t)$ and $D(0)$, in such way that mean value (9) does not change, i.e.

$$\langle A(t) | D(0) \rangle = \langle G(t) | F(0) \rangle = \sum_{s=0}^{\infty} \frac{1}{s!} \text{Tr}_{1, \dots, s} G_s(t, 1, \dots, s) F_s(0, 1, \dots, s). \quad (15)$$

Then marginal observables are defined by

$$G_s(t, Y) := \sum_{n=0}^s \frac{(-1)^n}{n!} \sum_{j_1 \neq \dots \neq j_n=1}^s A_{s-n}(t, Y \setminus \{j_1, \dots, j_n\}), \quad s \geq 1, \quad (16)$$

where $Y \equiv (1, \dots, s)$ and the sequence $A(t) = (I, A_1(t, 1), \dots, A_n(t, 1, \dots, n), \dots)$ is solution (8) of the initial-value problem to the Heisenberg equation (1)-(2), i.e. $A(t) = \mathcal{G}(t)A(0)$.

If we introduce the operator \mathfrak{a}^+ (an analog of the creation operator [2])

$$(\mathfrak{a}^+g)_s(Y) := \sum_{j=1}^s g_{s-1}(Y \setminus \{j\}) \quad (17)$$

defined on $\mathfrak{L}_\gamma(\mathcal{F}_\mathcal{H})$, then expression (16) can be rewritten in the following compact form

$$G(t) = e^{-\mathfrak{a}^+} \mathcal{G}(t) A(0).$$

The evolution of marginal observables (16) of both finitely and infinitely many quantum particles is described by the initial-value problem to the dual BBGKY hierarchy. For finitely many particles the quantum dual BBGKY hierarchy is equivalent to the Heisenberg equation (1) (the dual equation to the Heisenberg equation (1) is the von Neumann equation (10)). For systems of classical particles the dual BBGKY hierarchy was examined in [1], [2], [15].

2 The quantum dual BBGKY hierarchy

In order to approach the description of observables of quantum many-particle systems by the marginal observables (s -particle observables) we study the evolution of the system by means of the quantum dual BBGKY hierarchy. We introduce such a hierarchy of evolution equations and construct a solution of the Cauchy problem to this hierarchy.

2.1 The initial-value problem to the quantum dual BBGKY hierarchy

The evolution of marginal observables is described by the initial-value problem for the following hierarchy of evolution equations

$$\begin{aligned} \frac{d}{dt} G_s(t, Y) &= \mathcal{N}_s(Y) G_s(t, Y) + \\ &\sum_{n=1}^s \frac{1}{n!} \sum_{k=n+1}^s \frac{1}{(k-n)!} \sum_{j_1 \neq \dots \neq j_k=1}^s \mathcal{N}_{\text{int}}^{(k)}(j_1, \dots, j_k) G_{s-n}(t, Y \setminus \{j_1, \dots, j_n\}), \end{aligned} \quad (18)$$

$$G_s(t) |_{t=0} = G_s(0), \quad s \geq 1, \quad (19)$$

where, on $\mathcal{D}(\mathcal{N}_k) \subset \mathfrak{L}(\mathcal{H}_k)$, the operator $\mathcal{N}_{\text{int}}^{(n)}$ is defined by

$$\mathcal{N}_{\text{int}}^{(n)} g_n := -\frac{i}{\hbar} (g_n \Phi^{(n)} - \Phi^{(n)} g_n). \quad (20)$$

and the operator $\Phi^{(n)}$ is introduced in Hamiltonian (4). We refer to (18) as the quantum dual BBGKY hierarchy since the canonical BBGKY hierarchy [1] for marginal density operators is the dual hierarchy of evolution equations with respect to bilinear form (15) to evolution equations (18).

In the case of two-body interaction potential (4), hierarchy (18) has the form

$$\frac{d}{dt} G_s(t, Y) = \mathcal{N}_s(Y) G_s(t, Y) + \sum_{j_1 \neq j_2=1}^s \mathcal{N}_{\text{int}}^{(2)}(j_1, j_2) G_{s-1}(t, Y \setminus \{j_1\}), \quad s \geq 1, \quad (21)$$

where the operator $\mathcal{N}_{\text{int}}^{(2)}$ is defined by (20) for $n = 2$. For $\mathcal{H} = L^2(\mathbb{R}^\nu)$, the evolution of kernels of operators $G_s(t)$, $s \geq 1$, for equations (21), is given by

$$\begin{aligned} i\hbar \frac{\partial}{\partial t} G_s(t, q_1, \dots, q_s; q'_1, \dots, q'_s) &= \left(-\frac{\hbar^2}{2} \sum_{i=1}^s (-\Delta_{q_i} + \Delta_{q'_i}) + \right. \\ &\sum_{1=i < j}^s (\Phi^{(2)}(q'_i - q'_j) - \Phi^{(2)}(q_i - q_j)) \Big) G_s(t, q_1, \dots, q_s; q'_1, \dots, q'_s) + \\ &\sum_{1=i \neq j}^s (\Phi^{(2)}(q'_i - q'_j) - \Phi^{(2)}(q_i - q_j)) G_{s-1}(t, q_1, \dots, \overset{j}{\vee}, \dots, q_s; q'_1, \dots, \overset{j}{\vee}, \dots, q'_s), \end{aligned}$$

where $(q_1, \dots, \overset{j}{\vee}, \dots, q_s) \equiv (q_1, \dots, q_{j-1}, q_{j+1}, \dots, q_s)$. The dual BBGKY hierarchy for a system of classical particles stated in [1, 2] is defined by similar recurrence evolution equations.

The quantum dual BBGKY hierarchy (18) can be derived from the sequence of the Heisenberg equations (1) provided that observables of a system are described in terms of marginal operators (s -particle observables) (16).

Remark 1. Another way of looking to the derivation of the quantum dual BBGKY hierarchy consists in the construction of adjoint (dual) equations with respect to the bilinear form (15) to the quantum BBGKY hierarchy. For the sequence $F(t) = (I, F_1(t, 1), \dots, F_s(t, Y), \dots)$ of s -particle density operators (marginal density operators) $F_s(t, Y)$, $s \geq 1$, the quantum BBGKY hierarchy has the form [13]

$$\begin{aligned} \frac{d}{dt} F_s(t, Y) &= -\mathcal{N}_s(Y) F_s(t, Y) + \\ &\sum_{k=1}^s \frac{1}{k!} \sum_{i_1 \neq \dots \neq i_k=1}^s \sum_{n=1}^{\infty} \frac{1}{n!} \text{Tr}_{s+1, \dots, s+n} (-\mathcal{N}_{\text{int}}^{(k+n)})(i_1, \dots, i_k, X \setminus Y) F_{s+n}(t, X), \quad (22) \end{aligned}$$

where $X \equiv \{1, \dots, s+n\}$ and on $\mathfrak{L}_0^1(\mathcal{H}_{s+n}) \subset \mathfrak{L}^1(\mathcal{H}_{s+n})$ the operator $\mathcal{N}_{\text{int}}^{(k+n)}$ is defined by (20). Indeed, hierarchy (18) is dual to hierarchy (22) with respect to bilinear form (15).

Remark 2. In the paper [2], for classical systems of particles with a two-body interaction potential, an equivalent representation for the dual hierarchy generator was used. In the case under consideration, on the subspace $\mathcal{D}(\mathfrak{B}^+) \subset \mathfrak{L}_\gamma(\mathcal{F}_{\mathcal{H}})$, the generator has the following representation

$$\mathfrak{B}^+ = \mathcal{N} + [\mathcal{N}, \mathfrak{a}^+],$$

where $[\cdot, \cdot]$ is a commutator and the operator \mathfrak{a}^+ is defined on the space $\mathfrak{L}_\gamma(\mathcal{F}_{\mathcal{H}})$ by (17). In a general case the generator of the quantum dual BBGKY hierarchy (18) can be represented in the following form

$$\mathfrak{B}^+ = \mathcal{N} + \sum_{n=1}^{\infty} \frac{1}{n!} [\dots [\mathcal{N}, \underbrace{\mathfrak{a}^+}_{n\text{-times}}, \dots, \mathfrak{a}^+]] = e^{-\mathfrak{a}^+} \mathcal{N} e^{\mathfrak{a}^+}, \quad (23)$$

where the operators $e^{\pm \mathbf{a}^+}$ are defined on the space $\mathfrak{L}_\gamma(\mathcal{F}_\mathcal{H})$ by the expansions

$$(e^{\pm \mathbf{a}^+} g)_s(Y) = \sum_{n=0}^s \frac{(\pm 1)^n}{n!} \sum_{j_1 \neq \dots \neq j_n=1}^s g_{s-n}(Y \setminus \{j_1, \dots, j_n\}), \quad s \geq 1.$$

Representation (23) is correct in consequence of definition (17) of the operator \mathbf{a}^+ and the validity of identity

$$\begin{aligned} ([\dots [\mathcal{N}, \underbrace{\mathbf{a}^+}_{\text{n-times}}, \dots, \mathbf{a}^+] g)_s(Y) = \\ \sum_{k=n+1}^s \frac{1}{(k-n)!} \sum_{j_1 \neq \dots \neq j_k=1}^s \mathcal{N}_{\text{int}}^{(k)}(j_1, \dots, j_k) g_{s-n}(Y \setminus \{j_1, \dots, j_n\}), \end{aligned}$$

which for a two-body interaction potential reduces to the following one

$$([\mathcal{N}, \mathbf{a}^+] g)_s(Y) = \sum_{j_1 \neq j_2=1}^s \mathcal{N}_{\text{int}}^{(2)}(j_1, j_2) g_{s-1}(Y \setminus \{j_1\}).$$

2.2 The formula for a solution and its representations

We consider two different approaches to the construction of a solution of the quantum dual BBGKY hierarchy (18). Since hierarchy (18) has the structure of recurrence equations, we deduce that the solution can be constructed by successive integration of the inhomogeneous Heisenberg equations. Indeed, for solutions of the first two equations we obtain

$$\begin{aligned} G_1(t, 1) &= \mathcal{G}_1(t, 1) G_1(0, 1), \\ G_2(t, 1, 2) &= \mathcal{G}_2(t, 1, 2) G_2(0, 1, 2) + \\ &\int_0^t dt_1 \mathcal{G}_2(t - t_1, 1, 2) \mathcal{N}_{\text{int}}^{(2)}(1, 2) \mathcal{G}_1(t_1, 1) \mathcal{G}_1(t_1, 2) (G_1(0, 1) + G_1(0, 2)). \end{aligned} \quad (24)$$

Let us transform the second term on the right hand side of (24) for $G_2(t)$ as follows

$$\begin{aligned} &\int_0^t dt_1 \mathcal{G}_2(t - t_1, 1, 2) \mathcal{N}_{\text{int}}^{(2)}(1, 2) \mathcal{G}_1(t_1, 1) \mathcal{G}_1(t_1, 2) \\ &= - \int_0^t dt_1 \frac{d}{dt_1} (\mathcal{G}_2(t - t_1, 1, 2) \mathcal{G}_1(t_1, 1) \mathcal{G}_1(t_1, 2)) = \mathcal{G}_2(t, 1, 2) - \mathcal{G}_1(t, 1) \mathcal{G}_1(t, 2). \end{aligned} \quad (25)$$

The operator $\mathcal{G}_2(t, 1, 2) - \mathcal{G}_1(t, 1) \mathcal{G}_1(t, 2) := \mathfrak{A}_2^+(t, 1, 2)$ in equality (25) is the $2nd$ -order dual cumulant of evolution operators $\mathcal{G}(t) = \bigoplus_{n=0}^{\infty} \mathcal{G}_n(t)$ (5). Equality (25) is an analog of the Duhamel formula [9], which holds rigorously, for example, for bounded interaction potentials.

Thus, for solutions of the first two equations of hierarchy (18) we finally obtain

$$\begin{aligned} G_1(t, 1) &= \mathfrak{A}_1^+(t, 1) G_1(0, 1), \\ G_2(t, 1, 2) &= \mathfrak{A}_1^+(t, (\{1, 2\})_1) G_2(0, 1, 2) + \mathfrak{A}_2^+(t, 1, 2) (G_1(0, 1) + G_1(0, 2)), \end{aligned}$$

where we introduced the 1st-order dual cumulant $\mathfrak{A}_1^+(t, (\{1, 2\})_1) := \mathcal{G}_2(t, 1, 2)$ and the notation $(\{1, 2\})_1$ denotes a set composed by only one of elements 1 and 2 and it is similarly generalized in next formula (26).

Making use of transformations similar to (25), for $n > 2$, a solution of equations (18), constructed by successive integration of the inhomogeneous Heisenberg equations, is represented by the following expansions

$$G_s(t, Y) = \sum_{n=0}^s \frac{1}{(s-n)!} \sum_{j_1 \neq \dots \neq j_{s-n}=1}^s \mathfrak{A}_{1+n}^+(t, (Y \setminus X)_1, X) G_{s-n}(0, Y \setminus X), \quad s \geq 1. \quad (26)$$

Here we used some abridged notations: $Y \equiv (1, \dots, s)$, $X \equiv Y \setminus \{j_1, \dots, j_{s-n}\}$, the set $(Y \setminus X)_1$ consists of one element from $Y \setminus X = (j_1, \dots, j_{s-n})$, i.e. the set (j_1, \dots, j_{s-n}) is a connected subset of the partition P ($|P| = 1$, $|P|$ denotes the number of considered partitions). The evolution operator $\mathfrak{A}_{1+n}^+(t, (Y \setminus X)_1, X)$ in (26) is the $(1+n)$ th-order dual cumulant $\mathfrak{A}_{1+n}^+(t)$, $n \geq 0$, of groups of operators (5)

$$\mathfrak{A}_{1+n}^+(t, (Y \setminus X)_1, X) := \sum_{P: \{(Y \setminus X)_1, X\} = \cup_i X_i} (-1)^{|P|-1} (|P|-1)! \prod_{X_i \subset P} \mathcal{G}_{|X_i|}(t, X_i), \quad (27)$$

where \sum_P is the sum over all possible partitions P of the set $\{(Y \setminus X)_1, j_1, \dots, j_{s-n}\}$ into $|P|$ nonempty mutually disjoint subsets $X_i \subset \{(Y \setminus X)_1, X\}$. We consider some properties and examples of dual cumulants (27) in the next subsection.

Using the identity

$$\sum_{n=0}^s \frac{1}{(s-n)!} \sum_{j_1 \neq \dots \neq j_{s-n}=1}^s g_{s-n}(j_1, \dots, j_{s-n}) = \sum_{n=0}^s \frac{1}{n!} \sum_{j_1 \neq \dots \neq j_n=1}^s g_{s-n}(Y \setminus \{j_1, \dots, j_n\}) \quad (28)$$

which is valid for the Maxwell-Boltzmann statistics symmetry property, expansion (26) can be rewritten in the following an equivalent form

$$G_s(t, Y) = \sum_{n=0}^s \frac{1}{n!} \sum_{j_1 \neq \dots \neq j_n=1}^s \mathfrak{A}_{1+n}^+(t, (Y \setminus \{j_1, \dots, j_n\})_1, j_1, \dots, j_n) G_{s-n}(0, Y \setminus \{j_1, \dots, j_n\}), \quad s \geq 1.$$

The formula for a solution of the quantum dual BBGKY hierarchy (18) can be also derived from solution (8) of the initial-value problem to the Heisenberg equation (1)-(2) on the basis of expansions (16).

For sake of simplicity we consider the additive-type observables in capacity of initial data, i.e. one-component sequences $G^{(1)}(0) = (0, a_1(1), 0, \dots)$. In this case the expansion for solution (26) attains the form

$$G_s^{(1)}(t, Y) = \mathfrak{A}_s^+(t, 1, \dots, s) \sum_{j=1}^s a_1(j), \quad s \geq 1. \quad (29)$$

To determine the unknown evolution operator $\mathfrak{A}_s^+(t)$ from expansion (29), we equate expansion (29) for the operator $G_s^{(1)}(t, Y)$ with its representation by formula (16) for the

additive-type observables $A(0) = (0, a_1(1), \dots, \sum_{i=1}^n a_1(i), \dots)$, i.e.

$$\begin{aligned} \mathfrak{A}_s^+(t, 1, \dots, s) \sum_{j=1}^s a_1(j) &= \\ \sum_{n=0}^s \frac{(-1)^n}{n!} \sum_{j_1 \neq \dots \neq j_n=1}^s \mathcal{G}_{s-n}(t, Y \setminus \{j_1, \dots, j_n\}) \sum_{i \in Y \setminus \{j_1, \dots, j_n\}} a_1(i), \quad s \geq 1. \end{aligned} \quad (30)$$

Solving recursion relations (30) for arbitrary operators $\sum_{j=1}^s a_1(j)$ we find that the sth -order dual cumulant $\mathfrak{A}_s^+(t, 1, \dots, s)$ is defined by expansion (34), which is particular case of (27). For example, the first two equations from recursion relations (30) have the form

$$\begin{aligned} \mathfrak{A}_1^+(t, 1) a_1(1) &= \mathcal{G}_1(t, 1) a_1(1), \\ \mathfrak{A}_2^+(t, 1, 2) (a_1(1) + a_1(2)) &= \mathcal{G}_2(t, 1, 2) (a_1(1) + a_1(2)) - \mathcal{G}_1(t, 1) a_1(1) - \mathcal{G}_1(t, 2) a_1(2) = \\ &= (\mathcal{G}_2(t, 1, 2) - \mathcal{G}_1(t, 1) \mathcal{G}_1(t, 2)) (a_1(1) + a_1(2)). \end{aligned}$$

Hence a solution of the quantum dual BBGKY hierarchy (18) is defined by expansion (26) where the evolution operators $\mathfrak{A}_{1+n}^+(t)$, $n \geq 0$, are cumulants (27) of groups (5).

Remark 3. In the paper [2], for classical systems of particles, an equivalent representation for the dual hierarchy solution was used. For the case under consideration the solution expansion has the following representation

$$\begin{aligned} G(t) &= \mathcal{G}(t) G(0) + \sum_{n=1}^{\infty} \frac{1}{n!} [\dots [\mathcal{G}(t), \underbrace{\mathfrak{a}^+}_{n\text{-times}}, \dots, \mathfrak{a}^+] G(0) = \\ &= e^{-\mathfrak{a}^+} \mathcal{G}(t) e^{\mathfrak{a}^+} G(0), \end{aligned} \quad (31)$$

where $[\dots, \dots]$ is a commutator, the operator \mathfrak{a}^+ is defined on the space $\mathfrak{L}_\gamma(\mathcal{F}_{\mathcal{H}})$ by (17) and the one-parameter family $\{\mathcal{G}(t)\}_{t \in \mathbb{R}}$ of operators is defined by expression (5).

Representation (31) is true in consequence of definition (17) of the operator \mathfrak{a}^+ and the validity of identity (28) and of the following

$$\begin{aligned} \left(\frac{1}{n!} [\dots [\mathcal{G}(t), \underbrace{\mathfrak{a}^+}_{n\text{-times}}, \dots, \mathfrak{a}^+] g \right)_s(Y) &= \\ \frac{1}{(s-n)!} \sum_{j_1 \neq \dots \neq j_{s-n}=1}^s \mathfrak{A}_{1+n}^+(t, (Y \setminus X)_1, X) g_{s-n}(Y \setminus X), \quad s \geq 1, \end{aligned} \quad (32)$$

where $X \equiv Y \setminus \{j_1, \dots, j_{s-n}\}$ and $Y \setminus X = (j_1, \dots, j_{s-n})$. For example, if $n = 1$ we have

$$\begin{aligned} \left([\mathcal{G}(t), \mathfrak{a}^+] g \right)_s(Y) &= \sum_{j=1}^s (\mathcal{G}_s(t, Y) - \mathcal{G}_{s-1}(t, Y \setminus \{j\})) g_{s-1}(Y \setminus \{j\}) = \\ &= \sum_{j=1}^s \mathfrak{A}_2^+(t, (Y \setminus \{j\})_1, j) g_{s-1}(Y \setminus \{j\}). \end{aligned}$$

Some details of the calculations for identity (32) will be given in a general case in the Remark 4 of next subsection.

2.3 Dual cumulants of groups of evolution operators

We now consider some properties of dual cumulants (27). Let us expand the groups of operators $\mathcal{G}(t) = \bigoplus_{n=0}^{\infty} \mathcal{G}_n(t)$ (5) over new evolution operators as the following cluster expansions

$$\mathcal{G}_n(t, Y) = \sum_{P: Y = \bigcup_i X_i} \prod_{X_i \subset P} \mathfrak{A}_{|X_i|}^+(t, X_i), \quad n = |Y| \geq 0, \quad (33)$$

where \sum_P is the sum over all possible partitions P of the set $Y \equiv (1, \dots, n)$ into $|P|$ nonempty mutually disjoint subsets $X_i \subset Y$. A solution of recurrence relations (33) is determined by the expansions [15]

$$\mathfrak{A}_n^+(t, Y) = \sum_{P: Y = \bigcup_i X_i} (-1)^{|P|-1} (|P| - 1)! \prod_{X_i \subset P} \mathcal{G}_{|X_i|}(t, X_i), \quad n = |Y| \geq 0, \quad (34)$$

where the notations are similar to that in (33). We refer to the evolution operator $\mathfrak{A}_n^+(t)$ as the n th-order dual cumulant (dual semi-invariant) of group of operators (5).

The simplest examples of dual cumulants (34) have the form

$$\begin{aligned} \mathfrak{A}_1^+(t, 1) &= \mathcal{G}_1(t, 1), \\ \mathfrak{A}_2^+(t, 1, 2) &= \mathcal{G}_2(t, 1, 2) - \mathcal{G}_1(t, 1)\mathcal{G}_1(t, 2), \\ \mathfrak{A}_3^+(t, 1, 2, 3) &= \mathcal{G}_3(t, 1, 2, 3) - \mathcal{G}_1(t, 3)\mathcal{G}_2(t, 1, 2) - \mathcal{G}_1(t, 2)\mathcal{G}_2(t, 1, 3) \\ &\quad - \mathcal{G}_1(t, 1)\mathcal{G}_2(t, 2, 3) + 2!\mathcal{G}_1(t, 1)\mathcal{G}_1(t, 2)\mathcal{G}_1(t, 3). \end{aligned}$$

In the case of a quantum system of non-interacting particles ($n \geq 2$) we have: $\mathfrak{A}_n^+(t) = 0$. Indeed, for non-interacting quantum particles obeying Maxwell-Boltzmann statistics it holds

$$\mathcal{G}_n(t, 1, \dots, n) = \prod_{i=1}^n \mathcal{G}_1(t, i)$$

and hence

$$\begin{aligned} \mathfrak{A}_n^+(t, Y) &= \sum_{P: Y = \bigcup_i X_i} (-1)^{|P|-1} (|P| - 1)! \prod_{X_i \subset P} \prod_{l_i=1}^{|X_i|} \mathcal{G}_1(t, l_i) \\ &= \sum_{k=1}^n (-1)^{k-1} s(n, k) (k - 1)! \prod_{i=1}^k \mathcal{G}_1(t, i) = 0, \end{aligned}$$

where $s(n, k)$ are the Stirling numbers of the second kind. Here the following equality is used

$$\sum_{P: Y = \bigcup_i X_i} (-1)^{|P|-1} (|P| - 1)! = \sum_{k=1}^n (-1)^{k-1} s(n, k) (k - 1)! = \delta_{n,1}, \quad (35)$$

where $\delta_{n,1}$ is a Kronecker symbol.

The generator of the 1st-order dual cumulant is given by operator (3), i.e.

$$w^* - \lim_{t \rightarrow 0} \frac{1}{t} (\mathfrak{A}_1^+(t, Y) - I) g_n(Y) = \mathcal{N}_n(Y) g_n(Y),$$

where for $g_n \in \mathcal{D}(\mathcal{N}_n) \subset \mathfrak{L}(\mathcal{H}_n)$ this limit exists in the sense of the $*$ -weak convergence of the space $\mathfrak{L}(\mathcal{H}_n)$ (Proposition 1).

In the general case an infinitesimal generator of the n th-order dual cumulant, $n \geq 2$, is the operator $(\mathcal{N}_{\text{int}}^{(n)})$ defined by n -body interaction potential (4). Indeed, according to equality (35) in the sense of a point-by-point convergence in the space $\mathfrak{L}(\mathcal{H}_n)$ for the n th-order dual cumulant, $n \geq 2$, we have

$$\begin{aligned} \lim_{t \rightarrow 0} \frac{1}{t} \mathfrak{A}_n^+(t, Y) g_n(Y) &= \sum_{P: Y = \bigcup_i X_i} (-1)^{|P|-1} (|P| - 1)! \sum_{X_i \subset P} (\mathcal{N}_{|X_i|}(X_i)) g_n(Y) = \\ &= \sum_{P: Y = \bigcup_i X_i} (-1)^{|P|-1} (|P| - 1)! \sum_{X_i \subset P} \sum_{k=2}^{|X_i|} \sum_{i_1 < \dots < i_k \in \{X_i\}} (\mathcal{N}_{\text{int}}^{(k)}(i_1, \dots, i_k)) g_n(Y), \end{aligned}$$

where $\mathcal{N}_{\text{int}}^{(n)}$ is defined as in (20), i.e.

$$\mathcal{N}_{\text{int}}^{(n)} g_n := -\frac{i}{\hbar} (g_n \Phi^{(n)} - \Phi^{(n)} g_n),$$

where the operator $\Phi^{(n)}$ is defined in Hamiltonian (4).

Summing the coefficients of every operator $\mathcal{N}_{\text{int}}^{(k)}$ we deduce

$$\text{w}^* - \lim_{t \rightarrow 0} \frac{1}{t} \mathfrak{A}_n^+(t, Y) g_n(Y) = \mathcal{N}_{\text{int}}^{(n)}(Y) g_n(Y), \quad (36)$$

Thus, if $g_n \in \mathcal{D}(\mathcal{N}_n) \subset \mathfrak{L}(\mathcal{H}_n)$, the generator of the n th-order dual cumulant is defined by (36) in the sense of the $*$ -weak convergence of the space $\mathfrak{L}(\mathcal{H}_n)$.

We remark that, at the initial time $t = 0$, solution (26) satisfies initial condition (19). Indeed, according to definitions (6) ($\mathcal{U}_n^{\pm 1}(0) = I$ is a unit operator) and of equality (35) for $n \geq 1$, we have

$$\mathfrak{A}_{1+n}^+(0, (Y \setminus X)_1, X) = \sum_{P: \{(Y \setminus X)_1, X\} = \bigcup_i X_i} (-1)^{|P|-1} (|P| - 1)! I = I \delta_{n,1}.$$

Remark 4. Cluster expansions (33) can be put at the basis of all possible solution representations of the quantum dual BBGKY hierarchy (18). In fact we can obtain representation (31) solving recurrence relations (33) with respect to the 1st-order dual cumulants for the separation terms, which are independent from the variable $Y \setminus X \equiv (j_1, \dots, j_{s-n})$

$$\mathfrak{A}_{1+n}^+(t, (Y \setminus X)_1, X) = \sum_{Z \subset X} \mathfrak{A}_1^+(t, Y \setminus X \cup Z) \sum_{P: X \setminus Z = \bigcup_i X_i} (-1)^{|P|} |P|! \prod_{i=1}^{|P|} \mathfrak{A}_1^+(t, X_i),$$

where $\sum_{Z \subset X}$ is a sum over all subsets $Z \subset X$ of the set X . Then, taking into account the identity

$$\sum_{P: X \setminus Z = \bigcup_i X_i} (-1)^{|P|} |P|! \prod_{i=1}^{|P|} \mathfrak{A}_1^+(t, X_i) g_{s-n}(Y \setminus X) = \sum_{P: X \setminus Z = \bigcup_i X_i} (-1)^{|P|} |P|! g_{s-n}(Y \setminus X)$$

and the equality

$$\sum_{P: X \setminus Z = \bigcup_i X_i} (-1)^{|P|} |P|! = (-1)^{|X \setminus Z|}, \quad (37)$$

for expansion (26) we have

$$G_s(t, Y) = \sum_{n=0}^s \frac{1}{(s-n)!} \sum_{j_1 \neq \dots \neq j_{s-n}=1}^s \sum_{\substack{Z \subset X \\ Z \neq \emptyset}} (-1)^{|X \setminus Z|} \mathcal{G}_{s-n+|Z|}(t, Y \setminus X \cup Z) G_{s-n}(0, Y \setminus X).$$

Thus, as a result of the symmetry property for the Maxwell-Boltzmann statistics and of definition (17) of the operator \mathbf{a}^+ , we derive representation (31)

$$G(t) = \sum_{n=0}^{\infty} \frac{1}{n!} \sum_{k=0}^n (-1)^{n-k} \frac{n!}{k!(n-k)!} (\mathbf{a}^+)^{n-k} \mathcal{G}(t) (\mathbf{a}^+)^k G(0) = e^{-\mathbf{a}^+} \mathcal{G}(t) e^{\mathbf{a}^+} G(0).$$

We can obtain one more representation for a solution of the initial-value problem to the quantum dual BBGKY hierarchy if we express the dual cumulants $\mathfrak{A}_{1+n}^+(t)$, $n \geq 1$, of group of operators (5) with respect to the 1st-order and 2nd-order dual cumulants. In fact it holds

$$\mathfrak{A}_{1+n}^+(t, (Y \setminus X)_1, X) = \sum_{\substack{Z \subset X, \\ Z \neq \emptyset}} \mathfrak{A}_2^+(t, Y \setminus X, Z) \sum_{P: X \setminus Z = \cup_i X_i} (-1)^{|P|} |P|! \prod_{i=1}^{|P|} \mathfrak{A}_1^+(t, X_i),$$

where $\sum_{\substack{Z \subset X, \\ Z \neq \emptyset}}$ is a sum over all nonempty subsets $Z \subset X$ of the set X . Then taking into account the identity

$$\sum_{P: X \setminus Z = \cup_i X_i} (-1)^{|P|} |P|! \prod_{i=1}^{|P|} \mathfrak{A}_1^+(t, X_i) g_{s-n}(Y \setminus X) = \sum_{P: X \setminus Z = \cup_i X_i} (-1)^{|P|} |P|! g_{s-n}(Y \setminus X)$$

and equality (37), we get the following representation for expansion (26)

$$G_s(t, Y) = \mathfrak{A}_1^+(t, Y) G_s(0, Y) + \sum_{n=1}^s \frac{1}{(s-n)!} \sum_{j_1 \neq \dots \neq j_{s-n}=1}^s \sum_{\substack{Z \subset X, \\ Z \neq \emptyset}} (-1)^{|X \setminus Z|} \mathfrak{A}_2^+(t, Y \setminus X, Z) G_{s-n}(0, Y \setminus X),$$

where $Y \equiv (1, \dots, s)$, $X \equiv Y \setminus \{j_1, \dots, j_{s-n}\}$, i.e. $Y \setminus X = (j_1, \dots, j_{s-n})$.

3 A solution of the initial-value problem to the quantum dual BBGKY hierarchy

One-parameter mapping generated by solution (26) is not strong continuous on the space $\mathcal{L}_\gamma(\mathcal{F}_\mathcal{H})$. Thus, the constructed solution satisfies the initial-value problem to the dual BBGKY hierarchy (18)-(19) only in the sense of the *-weak convergence of the space $\mathcal{L}_\gamma(\mathcal{F}_\mathcal{H})$.

3.1 A group of operators for the quantum dual BBGKY hierarchy

On the space $\mathfrak{L}_\gamma(\mathcal{F}_\mathcal{H})$ solution (26) of the initial-value problem to the dual BBGKY hierarchy (18)-(19) is determined by a one-parameter mapping with the following properties.

Theorem 1. *If $g \in \mathfrak{L}_\gamma(\mathcal{F}_\mathcal{H})$ and $\gamma < e^{-1}$, then the one-parameter mapping*

$$\begin{aligned} \mathbb{R}^1 \ni t \mapsto (U^+(t)g)_s(Y) := & \quad (38) \\ \sum_{n=0}^s \frac{1}{(s-n)!} \sum_{j_1 \neq \dots \neq j_{s-n}=1}^s \mathfrak{A}_{1+n}^+(t, (Y \setminus X)_1, X) g_{s-n}(Y \setminus X), \quad s \geq 1 \end{aligned}$$

is a C_0^* -group. The infinitesimal generator $\mathfrak{B}^+ = \bigoplus_{n=0}^{\infty} \mathfrak{B}_n^+$ of this group of operators is a closed operator for the $*$ -weak topology and on the domain of the definition $\mathcal{D}(\mathfrak{B}^+) \subset \mathfrak{L}_\gamma(\mathcal{F}_\mathcal{H})$ which is the everywhere dense set for the $*$ -weak topology of the space $\mathfrak{L}_\gamma(\mathcal{F}_\mathcal{H})$ it is defined by the operator

$$\begin{aligned} (\mathfrak{B}^+g)_s(Y) := \mathcal{N}_s(Y)g_s(Y) + & \quad (39) \\ \sum_{n=1}^s \frac{1}{n!} \sum_{k=n+1}^s \frac{1}{(k-n)!} \sum_{j_1 \neq \dots \neq j_k=1}^s \mathcal{N}_{\text{int}}^{(k)}(j_1, \dots, j_k) g_{s-n}(Y \setminus \{j_1, \dots, j_k\}), \end{aligned}$$

where the operator $\mathcal{N}_{\text{int}}^{(k)}$ is given by (20).

Proof. If $g \in \mathfrak{L}_\gamma(\mathcal{F}_\mathcal{H})$, mapping (38) is defined provided that $\gamma < e^{-1}$ and that the following estimate holds

$$\|U^+(t)g\|_{\mathfrak{L}_\gamma(\mathcal{F}_\mathcal{H})} \leq e^2(1 - \gamma e)^{-1} \|g\|_{\mathfrak{L}_\gamma(\mathcal{F}_\mathcal{H})}. \quad (40)$$

This estimate comes out from the inequality

$$\begin{aligned} \|(U^+(t)g)_s\|_{\mathfrak{L}(\mathcal{H}_s)} & \leq \sum_{n=0}^s \frac{1}{(s-n)!} \sum_{j_1 \neq \dots \neq j_{s-n}=1}^s \sum_{P: \{Y \setminus X\}_1, X\} = \bigcup_i X_i} (|P| - 1)! \|g_{s-n}\|_{\mathfrak{L}(\mathcal{H}_{s-n})} \leq \\ \sum_{n=0}^s \|g_{s-n}\|_{\mathfrak{L}(\mathcal{H}_{s-n})} \frac{s!}{n!(s-n)!} \sum_{k=1}^{n+1} s(n+1, k)(k-1)!, \end{aligned}$$

where $s(n+1, k)$ are the Stirling numbers of the second kind for which it holds

$$\sum_{k=1}^{n+1} s(n+1, k)(k-1)! = \sum_{k=1}^{n+1} \frac{1}{k} \sum_{\substack{r_1, \dots, r_k \geq 1 \\ r_1 + \dots + r_k = n+1}} \frac{(n+1)!}{r_1! \dots r_k!} \leq \sum_{k=1}^{n+1} k^n \leq n! e^{n+2}.$$

On the space $\mathfrak{L}_\gamma(\mathcal{F}_\mathcal{H})$ the $*$ -weak continuity property of the group $U^+(t)$ over the parameter $t \in \mathbb{R}^1$ is a consequence of the $*$ -weak continuity of group of operators (5) for the Heisenberg equation (1).

In order to construct an infinitesimal generator of the group $\{U^+(t)\}_{t \in \mathbb{R}}$ we firstly differentiate the n th-term of expansion (38) in the sense of the pointwise convergence of

the space \mathfrak{L}_γ . If $g \in \mathcal{D}(\mathcal{N}) \subset \mathfrak{L}_\gamma(\mathcal{F}_\mathcal{H})$, in a similar way to equality (36) for $(1+n)$ th-order dual cumulant (27), $n \geq 1$, we derive

$$\begin{aligned} \lim_{t \rightarrow 0} \frac{1}{t} \mathfrak{A}_{1+n}^+(t, (Y \setminus X)_1, X) g_{s-n}(Y \setminus X) \psi_s &= \sum_{\substack{Z \subset Y \setminus X, \\ Z \neq \emptyset}} \mathcal{N}_{\text{int}}^{(|Z|+n)}(Z, X) g_{s-n}(Y \setminus X) \psi_s = \\ \sum_{k=1}^{s-n} \frac{1}{k!} \sum_{i_1 \neq \dots \neq i_k \in \{j_1, \dots, j_{s-n}\}} \mathcal{N}_{\text{int}}^{(k+n)}(i_1, \dots, i_k, X) g_{s-n}(Y \setminus X) \psi_s. \end{aligned} \quad (41)$$

Then, according to equalities (7) and (41), for group (38) we obtain

$$\begin{aligned} \lim_{t \rightarrow 0} \frac{1}{t} \left((U^+(t)g)_s - g_s \right) \psi_s &= \lim_{t \rightarrow 0} \frac{1}{t} (\mathfrak{A}_1^+(t)g_s - g_s) \psi_s + \\ \sum_{n=1}^s \frac{1}{(s-n)!} \sum_{j_1 \neq \dots \neq j_{s-n}=1}^s \lim_{t \rightarrow 0} \frac{1}{t} \mathfrak{A}_{1+n}^+(t, (Y \setminus X)_1, X) g_{s-n}(Y \setminus X) \psi_s &= \\ \mathcal{N}_s g_s \psi_s + \sum_{n=1}^s \frac{1}{n!} \sum_{k=n+1}^s \frac{1}{(k-n)!} \sum_{j_1 \neq \dots \neq j_k=1}^s \mathcal{N}_{\text{int}}^{(k)}(j_1, \dots, j_k) g_{s-n}(Y \setminus \{j_1, \dots, j_n\}) \psi_s, \end{aligned}$$

where we used identity (28).

Thus if $g \in \mathcal{D}(\mathfrak{B}^+) \subset \mathfrak{L}_\gamma(\mathcal{F}_\mathcal{H})$ in the sense of the $*$ -weak convergence of the space $\mathfrak{L}_\gamma(\mathcal{F}_\mathcal{H})$ we finally have

$$\text{w}^* - \lim_{t \rightarrow 0} \left(\frac{1}{t} (U^+(t)g - g) - \mathfrak{B}^+ g \right) = 0,$$

where the generator $\mathfrak{B}^+ = \bigoplus_{n=0}^{\infty} \mathfrak{B}_n^+$ of group of operators (38) is given by (39). \square

3.2 The existence and uniqueness theorem

The following theorem holds for abstract initial-value problem (18)-(19) on the space $\mathfrak{L}_\gamma(\mathcal{F}_\mathcal{H})$.

Theorem 2. *A solution of the initial-value problem to the quantum dual BBGKY hierarchy (18)-(19) is determined by the expansion*

$$G_s(t, Y) = \sum_{n=0}^s \frac{1}{(s-n)!} \sum_{j_1 \neq \dots \neq j_{s-n}=1}^s \mathfrak{A}_{1+n}^+(t, (Y \setminus X)_1, X) G_{s-n}(0, Y \setminus X), \quad s \geq 1, \quad (42)$$

where the $(1+n)$ th-order dual cumulant $\mathfrak{A}_{1+n}^+(t, (Y \setminus X)_1, X)$ is defined by (27)

$$\mathfrak{A}_{1+n}^+(t, (Y \setminus X)_1, X) := \sum_{\text{P}: \{(Y \setminus X)_1, X\} = \bigcup_i X_i} (-1)^{|\text{P}|-1} (|\text{P}|-1)! \prod_{X_i \subset \text{P}} \mathcal{G}_{|X_i|}(t, X_i),$$

where \sum_{P} is the sum over all possible partitions P of the set $\{(Y \setminus X)_1, j_1, \dots, j_{s-n}\}$ into $|\text{P}|$ nonempty mutually disjoint subsets $X_i \subset \{(Y \setminus X)_1, X\}$. For $G(0) \in \mathcal{D}(\mathfrak{B}^+) \subset \mathfrak{L}_\gamma(\mathcal{F}_\mathcal{H})$ it is a classical solution and for arbitrary initial data $G(0) \in \mathfrak{L}_\gamma(\mathcal{F}_\mathcal{H})$ it is a generalized solution.

Proof. According to Theorem 1, for the initial data $G(0) \in \mathcal{D}(\mathfrak{B}^+) \subset \mathfrak{L}_\gamma(\mathcal{F}_\mathcal{H})$, sequence (42) is a classical solution of initial-value problem (18)-(19) in the sense of the *-weak convergence of the space $\mathfrak{L}_\gamma(\mathcal{F}_\mathcal{H})$.

Let us now show that in the general case $G(0) \in \mathfrak{L}_\gamma(\mathcal{F}_\mathcal{H})$ expansions (42) give a generalized solution of the initial-value problem to the quantum dual BBGKY hierarchy (18)-(19). To this aim we consider the functional

$$(f, G(t)) := \sum_{s=0}^{\infty} \frac{1}{s!} \text{Tr}_{1, \dots, s} f_s G_s(t), \quad (43)$$

where $f \in \mathfrak{L}_{\alpha,0}^1$ is a finite sequence of the degenerate trace class operators with infinitely times differentiable kernels and with compact support. According to estimate (40) this functional exists provided that $\alpha = \gamma^{-1} > e$ (see Section 4).

Using (42), we can transform functional (43) as follows

$$(f, G(t)) = (f, U^+(t)G(0)) = (U(t)f, G(0)). \quad (44)$$

In this equality the group $U^+(t)$ is defined by expression (38) (Theorem 1) and $U(t)$ is an adjoint mapping to the group $U^+(t)$

$$(U(t)f)_s(Y) = \sum_{n=0}^{\infty} \frac{1}{n!} \text{Tr}_{s+1, \dots, s+n} \mathfrak{A}_{1+n}(t, Y_1, X \setminus Y) f_{s+n}(X), \quad (45)$$

where $X \equiv \{1, \dots, s+n\}$, i.e. $X \setminus Y \equiv \{s+1, \dots, s+n\}$. In expansion (45) the evolution operator $\mathfrak{A}_{1+n}(t, Y_1, X \setminus Y)$ is the $(1+n)$ th-order cumulant of group of operators (13)

$$\mathfrak{A}_{1+n}(t, Y_1, X \setminus Y) = \sum_{P: \{Y_1, X \setminus Y\} = \cup_i X_i} (-1)^{|P|-1} (|P|-1)! \prod_{X_i \subset P} \mathcal{G}_{|X_i|}(-t, X_i),$$

where \sum_P is the sum over all possible partitions P of the set $\{Y_1, X \setminus Y\} = \{Y_1, s+1, \dots, s+n\}$ into $|P|$ nonempty mutually disjoint subsets $X_i \subset \{Y_1, X \setminus Y\}$. If $f \in \mathfrak{L}_{\alpha,0}^1$, series (45) converges, provided that $\alpha > e$ [14] and the functional $(U(t)f, G(0))$ exists.

The one-parameter family of operators $U(t)$ is differentiable with respect to t and for $f \in \mathfrak{L}_{\alpha,0}^1$ an infinitesimal generator of group (45) is defined by the following expression (see also (22))

$$\begin{aligned} (\mathfrak{B}f)_s(Y) &= -\mathcal{N}_s(Y) f_s(Y) + \\ &\sum_{k=1}^s \frac{1}{k!} \sum_{i_1 \neq \dots \neq i_k=1}^s \sum_{n=1}^{\infty} \frac{1}{n!} \text{Tr}_{s+1, \dots, s+n} (-\mathcal{N}_{\text{int}}^{(k+n)})(i_1, \dots, i_k, X \setminus Y) f_{s+n}(X). \end{aligned} \quad (46)$$

Since for bounded interaction potentials (4), if $f \in \mathfrak{L}_{\alpha,0}^1$, the operator $\mathfrak{B}U(t)f$ is a trace class operator, the operator $\mathfrak{B}U(t)fG(0)$ is also a trace class operator then the functional $(\mathfrak{B}U(t)f, G(0))$ exists. Moreover, it holds the equality: $(\mathfrak{B}U(t)f, G(0)) =$

$(U(t)\mathfrak{B}f, G(0))$ and the following result

$$\begin{aligned} & \lim_{t \rightarrow 0} \left| \left(\frac{1}{t} (U(t) - I)f, G(0) \right) - (\mathfrak{B}f, G(0)) \right| = \\ & \lim_{t \rightarrow 0} \left| \sum_{s=0}^{\infty} \frac{1}{s!} \text{Tr}_{1, \dots, s} \left(\frac{1}{t} (U(t)f - f)_s G_s(0) - (\mathfrak{B}f)_s G_s(0) \right) \right| \leq \\ & \|G(0)\|_{\mathfrak{L}_\gamma(\mathcal{F}_\mathcal{H})} \lim_{t \rightarrow 0} \sum_{s=0}^{\infty} \gamma^{-s} \text{Tr}_{1, \dots, s} \left| \frac{1}{t} (U(t)f - f)_s - (\mathfrak{B}f)_s \right| = \\ & \|G(0)\|_{\mathfrak{L}_\gamma(\mathcal{F}_\mathcal{H})} \lim_{t \rightarrow 0} \left\| \frac{1}{t} (U(t)f - f) - \mathfrak{B}f \right\|_{\mathfrak{L}_{\gamma^{-1}}^1(\mathcal{F}_\mathcal{H})} = 0. \end{aligned}$$

Hence equality (44) can be differentiated with respect to time and we get finally

$$\begin{aligned} \frac{d}{dt} (f, G(t)) &= (U(t)\mathfrak{B}f, G(0)) = (\mathfrak{B}f, U^+(t)G(0)) = \\ & (\mathfrak{B}f, G(t)), \end{aligned}$$

where the operator \mathfrak{B} is defined by (46).

These equalities mean that the sequence of operators (42) for arbitrary $G(0) \in \mathfrak{L}_\gamma(\mathcal{F}_\mathcal{H})$ is a generalized solution of the Cauchy problem to the quantum dual BBGKY hierarchy (18)-(19). \square

4 The existence of the mean value observable functional

As it was above mentioned, the functional of the mean value (15) defines a duality between marginal observables and marginal states. If $G(t) \in \mathfrak{L}_\gamma(\mathcal{F}_\mathcal{H})$ and $F(0) \in \mathfrak{L}_\alpha^1$ then, according to estimate (40), the functional

$$\langle G(t) | F(0) \rangle = \sum_{s=0}^{\infty} \frac{1}{s!} \text{Tr}_{1, \dots, s} G_s(t, 1, \dots, s) F_s(0, 1, \dots, s). \quad (47)$$

exists, provided that $\alpha = \gamma^{-1} > e$, and the following estimate holds

$$|\langle G(t) | F(0) \rangle| \leq e^2 (1 - \gamma e)^{-1} \|G(0)\|_{\mathfrak{L}_\gamma(\mathcal{F}_\mathcal{H})} \|F(0)\|_{\mathfrak{L}_{\gamma^{-1}}^1(\mathcal{F}_\mathcal{H})}.$$

Thus, marginal density operators from the space \mathfrak{L}_α^1 describe finitely many quantum particles. Indeed, for such additive-type observable as number of particles, i.e. one-component sequence $N(0) = (0, I, 0, \dots)$, according to definition (34) of dual cumulants, the expansion for solution (26) get the following form (see also (29))

$$(N(t))_s(Y) = \mathfrak{A}_s^+(t, 1, \dots, s) \sum_{j=1}^s I = I \delta_{s,1}, \quad s \geq 1$$

and we have

$$|\langle N(t) | F(0) \rangle| = |\text{Tr}_1 F_1(0, 1)| \leq \|F(0)\|_{\mathfrak{L}_{\gamma^{-1}}^1(\mathcal{F}_\mathcal{H})} < \infty.$$

We have above stated the properties of group (38) defined on $\mathfrak{L}_\gamma(\mathcal{F}_\mathcal{H})$. In order to describe the evolution of infinitely many particles [1] the problem lies in the definition of functional (47) for operators from the suitable Banach spaces. Namely, marginal density operators have to belong to more general spaces than $\mathfrak{L}_\alpha^1(\mathcal{F}_\mathcal{H})$. For example, it can be the space of sequences of bounded operators containing the equilibrium states [17]. In this case every term of expansions for the mean value functional (47) contains the divergent traces [1], [2], [14] and the analysis of such a question for quantum systems remains an open problem.

5 Conclusion

The concept of dual cumulants (34) of groups of operators (5) for the Heisenberg equations (1) or cumulants (45) of groups of operators (13) for the von Neumann equations (10) forms the basis of the groups of operators for quantum evolution equations as well as the quantum dual BBGKY hierarchy and the BBGKY hierarchy for marginal density operators [14].

In the case of quantum systems of particles, obeying Fermi or Bose statistics, group of operators (38) has different structures. The analysis of these cases will be given in a separate paper.

On the space $\mathfrak{L}_\gamma(\mathcal{F}_\mathcal{H})$ one-parameter mapping (38) is not a strong continuous group. The group $\{U^+(t)\}_{t \in \mathbb{R}}$ of operators (38) defined on the space $\mathfrak{L}_\gamma(\mathcal{F}_\mathcal{H})$ is dual to the strong continuous group $\{U(t)\}_{t \in \mathbb{R}}$ of operators (45) for the BBGKY hierarchy defined on the space $\mathfrak{L}_\alpha^1(\mathcal{F}_\mathcal{H})$ and the fact that it is a C_0^* -group follows also from general theorems about properties of the dual semigroups [4, 6].

We have constructed infinitesimal generator (39) on the domain $\mathcal{D}(\mathfrak{B}^+) \subset \mathfrak{L}_\gamma(\mathcal{F}_\mathcal{H})$ which is everywhere dense set for the $*$ -weak topology of the space $\mathfrak{L}_\gamma(\mathcal{F}_\mathcal{H})$. The question of how to define the domain of the definition $\mathcal{D}(\mathfrak{B}^+)$ of generator (39) is an open problem [4, 10].

Acknowledgement

This work was performed under the auspices of the *National Group for Mathematical Physics* of the *Istituto Nazionale di Alta Matematica* and was partially supported by the WTZ grant No M/124 (UA 04/2007), by the *Italian Ministry of University* MIUR National Project "Kinetic and Hydrodynamic Equations of Complex Collisional Systems", PRIN 2006) and by Research Funds of the University of Florence (Italy).

References

- [1] C.Cercignani, V.I.Gerasimenko and D.Ya.Petrina, *Many-Particle Dynamics and Kinetic Equations*. Kluwer Acad. Publ., 1997.
- [2] G.Borgioli and V.Gerasimenko, *The dual BBGKY hierarchy for the evolution of observables*. Riv. Mat. Univ. Parma. **4**, (2001), 251-267.

-
- [3] R.Dautray and J.L.Lions, *Mathematical Analysis and Numerical Methods for Science and Technology*. **5**, Springer-Verlag, 1992.
 - [4] O.Bratelli and D.W.Robinson, *Operator Algebras and Quantum Statistical Mechanics*. **1**, Springer-Verlag, 1979.
 - [5] T.Kato, *Perturbation Theory for Linear Operators*. Springer-Verlag, 1995.
 - [6] A.Pazy, *Semigroups of Linear Operators and Applications to Partial Differential Equations*. Springer-Verlag, 1983.
 - [7] J.A.Goldstein, *Semigroups of Linear Operators and Applications*. Oxford University Press, 1985.
 - [8] A.Belleni-Morante and A.C.McBride, *Applied Nonlinear Semigroups*. John Wiley and Sons, 1998.
 - [9] J.Banasiak and L.Arlotti, *Perturbations of Positive Semigroups with Applications*. Springer, 2006.
 - [10] R.Aliki and K.Lendi, *Quantum dynamical semigroups and applications*. Lect. Notes in Phys. **286**, Springer, 1987.
 - [11] F.A.Berezin and M.A.Shoubin, *Schrödinger Equation*. Kluwer, 1991.
 - [12] D.Ya.Petrina, *Mathematical Foundations of Quantum Statistical Mechanics. Continuous Systems*. Kluwer, 1995.
 - [13] V.I.Gerasimenko and V.O.Shtyk, *Evolution of Correlations of Quantum Many-Particle Systems*. J. Stat. Mech. (3), (2008), P03007, 24p.
 - [14] V.I.Gerasimenko and V.O.Shtyk, *Initial-value problem for the Bogolyubov hierarchy for quantum systems of particles*. Ukrain. Math. J. **58**, (9), (2006), 1175-1191.
 - [15] V.I.Gerasimenko and T.V.Ryabukha, *Cumulant representation of solutions of the BBGKY hierarchy of equations*. Ukrain. Math. J. **54**, (10), (2002), 1583-1601.
 - [16] V.I.Gerasimenko, T.V.Ryabukha and M.O.Stashenko, *On the structure of expansions for the BBGKY hierarchy solutions*. J. Phys. A: Math. Gen. **37**, (2004), 9861-9872.
 - [17] J.Ginibre, *Some applications of functional integrations in statistical mechanics*. (in *Statistical Mechanics and Quantum Field Theory*, eds. S.DeWitt and R.Stora. Gordon and Breach), (1971), 329-427.
 - [18] A.Arnold, *Mathematical properties of quantum evolution equations*. Lect. Notes in Math. **1946**, Springer, 2008.
 - [19] H.Spohn, *Kinetic equations for quantum many-particle systems*. arXiv:0706.0807v1, 2007.
 - [20] D.Benedetto, F.Castella, R.Esposito and M.Pulvirenti, *A short review on the derivation of the nonlinear quantum Boltzmann equations*. Commun. Math. Sci. **5**, (2007), 55-71.

-
- [21] F.Golse, *The mean-field limit for the dynamics of large particle systems*. Journées equations aux dérivées partielles. **9**, (2003), 47p.
- [22] C.Bardos, B.Ducomet, F.Golse, A.Gottlieb and N.Mauser, *The TDHF approximation for Hamiltonians with m -particle interaction potentials*. Commun. Math. Sci. **5**, (2007), 1-9.
- [23] R.Adami, F.Golse and A.Teta, *Rigorous Derivation of the Cubic NLS in Dimension One*. J. Stat. Phys. **127**, (6), (2007), 1193-1220.
- [24] L.Erdős, M.Salmhofer and H.-T.Yau, *On quantum Boltzmann equation*. J. Stat. Phys. **116**, (116), (2004), 367-380.
- [25] L.Erdős, B.Schlein and H.-T.Yau, *Rigorous derivation of the Gross-Pitaevskii equation*. Phys. Rev. Lett. **98**, (4), (2007), 040404.
- [26] L.Erdős, B.Schlein and H.-T.Yau, *Derivation of the cubic non-linear Schrödinger equation from quantum dynamics of many-body systems*. Invent. Math. **167**, (3), (2007), 515-614.
- [27] J.Fröhlich, S.Graffi and S.Schwarz, *Mean-Field- and Classical Limit of Many-Body Schrödinger Dynamics for Bosons*. Commun. Math. Phys., **271**, (2007), 681-697.