

GRÖBNER-SHIRSHOV BASIS FOR THE BRAID SEMIGROUP

L. A. BOKUT*, Y. FONG, W.-F. KE, AND L.-S. SHIAO

ABSTRACT. We found Gröbner-Shirshov basis for the braid semigroup B_{n+1}^+ . It gives a new algorithm for the solution of the word problem for the braid semigroup and so for the braid group.

1. INTRODUCTION

In this paper, we are continuing to use a language of the Gröbner-Shirshov bases for some semigroups presented by defining relations (in papers [31], [32] some groups have been treated as semigroups). Traditionally in this situation, one use some other languages like elementary transformations language ([1],[2]), Newman's Diamond lemma language ([63]), the rewriting systems language ([34]), or some else ([56]). For example, P.M. Cohn used Diamond lemma for semigroups as early as in 1956 ([41], [42]). After him first author used the same language to prove some embedding theorems for semigroups [8],[9]. Also first author used a language of rewriting systems in order to prove that multiplicative semigroups of some rings is embeddable into groups [12], [13] (see also [16], [20]). But here we formulate our main results in terms of Gröbner-Shirshov bases.

During the last few years the Gröbner-Shirshov bases method has been used to study some important examples of associative algebras (quantum enveloping algebras of the type A_n , Hecke algebras of type A_n), Lie algebras (simple Lie algebras of the types $A_n, B_n, C_n, D_n, G_2, F_4, E_6, E_7, E_8$, Kac-Moody algebras of the types $A_n^{(1)}, B_n^{(1)}, C_n^{(1)}, D_n^{(1)}$), Lie superalgebras (simple Lie superalgebras of the types A_n, B_n, C_n, D_n), modules (irreducible modules over simple Lie algebras of the type A_n , Specht modules over Hecke algebras of the type A_n), and groups (Gröbner-Shirshov bases for Coxeter groups of types A_n, B_n, D_n) [25] – [31], [53]–[55], [65] – [68]. Also this method was invented for associative conformal algebras [21], [22].

In recent paper [32], two of us applied the Gröbner-Shirshov bases method to study some groups by P.S. Novikov [64] and W.W. Boone [33] with the algorithmically unsolvable word problem. They revised some papers [10], [11], [17], [18] (see also [19], [28]) that used a rewriting systems language (between lines of HNN-extension theory) using the very language of Gröbner-Shirshov bases. It means that they eliminated any group theory in proving Novikov's and Boone's theorems on unsolvability.

Some other activities on Gröbner-Shirshov (non-commutative Gröbner, standard) bases may be found in papers by T. Mora [62], V.N. Latyshev [58], V. Ufnarovski [73], A.A. Mikhalev and A.A. Zolotyh [61], A.A. Mikhalev and E. Vasilieva [60], V.P. Gerdt and V.V. Kornyak [50].

*Supported in part by the Russia's Fund of Fundamental Research.

A great interest has been done to the word and conjugacy problems for braid groups B_{n+1} . Let us mention a pioneering paper by E. Artin [3], [4], papers by A.A. Markov [59], A.F. Garside [49], P. Deligne [46], W. Thurston [72], K. Tatsuoka [71], E.A. Elrifai and H.R. Morton [48], P. Dehornoy [45], J. Birman, K.H. Ko, and S.J. Lee [6]. In those papers, a fundamental role of a semigroup of positive words of B_{n+1} is proved to be crucial. Actually this semigroup depends of generators of B_{n+1} . Let us denote this semigroup in Artin generators $a_i, 1 \leq i \leq n$ by B_{n+1}^+ and call the braid semigroup. It has the following defining relations as a semigroup:

$$\begin{aligned} a_{i+1}a_i a_{i+1} &= a_i a_{i+1} a_i, \\ a_i a_j &= a_j a_i, i - j \geq 2. \end{aligned}$$

In this paper, we found Gröbner-Shirshov basis of B_{n+1}^+ . It gives a new solution of the word problem for the braid semigroup and as the result for the braid group. We also study a more general concept of braid semigroups. The braid semigroup B_{n+1}^+ is a braid semigroup of type A_n . We formulate a hypothesis on Gröbner-Shirshov bases of any braid semigroup. It is a modification of a hypothesis on Gröbner-Shirshov bases of Coxeter groups [31]. The previous example support our hypothesis.

Let us mentioned that in Birman-Ko-Lee generators $a_{ts}, 1 \leq t < s \leq n + 1$ a semigroup of positive braids of B_{n+1} have the following defining relations [6]:

$$\begin{aligned} a_{ts}a_{rq} &= a_{rq}a_{ts}, (t-r)(t-q)(s-r)(s-q) > 0, \\ a_{ts}a_{sr} &= a_{tr}a_{ts} = a_{sr}a_{tr}. \end{aligned}$$

It would be interesting to find a Gröbner-Shirshov basis of this semigroup too. It may have some connections with a new Public-Key Cryptosystem [57].

2. GRÖBNER-SHIRSHOV BASES

Let \mathbf{X} be a linearly ordered set, k be a field, $k\langle\mathbf{X}\rangle$ be the free associative algebra over \mathbf{X} and k . On the set \mathbf{X}^* of words we impose a monomial well order $>$ (i.e. a well order that agrees with the concatenation of words). Any polynomial $f \in k\langle\mathbf{X}\rangle$ has the leading word \bar{f} . We say that f is monic if \bar{f} occurs in f with coefficient 1. By a composition of intersection $(f, g)_w$ of two monic polynomials relative to some word w , such that $w = \bar{f}b = a\bar{g}$, $\deg(\bar{f}) + \deg(\bar{g}) > \deg(w)$, one means the following polynomial

$$(f, g)_w = fb - ag.$$

By composition of including $(f, g)_w$ two monic polynomials, where $w = \bar{f} = a\bar{g}b$, one means the following polynomial

$$(f, g)_w = f - agb.$$

In the last case the transformation

$$f \mapsto (f, g)_w = f - agb$$

is called the elimination of the leading word (ELW) of g in f .

In both cases the word w is called an *ambiguity* of polynomials (relations) f, g .

A composition $(f, g)_w$ is called trivial relative to some $\mathbf{S} \subset k\langle\mathbf{X}\rangle$ ($(f, g)_w \equiv 0 \pmod{\mathbf{S}, w}$), if

$$(f, g)_w = \sum \alpha_i a_i t_i b_i,$$

where $t_i \in \mathbf{S}$, $a_i, b_i \in \mathbf{X}^*$, and $a_i \bar{t}_i b_i < w$. In particular, if $(f, g)_w$ goes to zero by the ELW's of \mathbf{S} then $(f, g)_w$ is trivial relative to \mathbf{S} .

We assume for some polynomials f_1 and f_2 that

$$f_1 \equiv f_2 \text{ mod}(\mathbf{S}, w),$$

if

$$f_1 - f_2 \equiv 0 \text{ mod}(\mathbf{S}, w).$$

A subset \mathbf{S} of $k\langle \mathbf{X} \rangle$ is called a Gröbner-Shirshov basis if any composition of polynomials from \mathbf{S} is trivial relative to \mathbf{S} .

By $\langle \mathbf{X} | \mathbf{S} \rangle$, the algebra with generators \mathbf{X} and defining relations \mathbf{S} , we will mean the factor-algebra of $k\langle \mathbf{X} \rangle$ by the ideal generated by \mathbf{S} .

The following lemma goes back to the Poincare - Birkhoff-Witt (PBW) theorem, the Diamond Lemma of M.H.A. Newman [63], the Composition Lemma of A.I. Shirshov ([70]) (see also [14], [15], where this Composition Lemma was formulated explicitly and in a current form for Lie algebras and associative algebras respectively), the Buchberger's Theorem ([35]) (published in [36]), the Diamond Lemma of G. Bergman [5] (this lemma was also known to P.M. Cohn, see, for example, [43], [44] and some historical comments to Chapter "Gröbner bases" in [47]):

Lemma 2.1. (*Composition-Diamond Lemma*)

\mathbf{S} is a Gröbner-Shirshov basis if and only if the set

$$\{u \in \mathbf{X}^* \mid u \neq a\bar{f}b, \text{ for any } f \in \mathbf{S}\}$$

of \mathbf{S} -reduced words consists of a linear basis of the algebra $\langle \mathbf{X} | \mathbf{S} \rangle$.

While \mathbf{S} is fixed, we will call \mathbf{S} -reduced words just reduced words. Actually we would like to introduce a new term for this basis - the *PBW - basis* for the algebra relative to the Gröbner-Shirshov basis of relations of this algebra. It would reflect to the fact that as we have mentioned the Composition-Diamond lemma goes back to the Poincare-Birkhoff-Witt (PBW) theorem as well. In fact the PBW theorem is a particular and the most important up to now case of Composition-Diamond lemma when $\mathbf{S} = \{x_i x_j - x_j x_i - \sum x_{ij}^k, i > j\}$ and $[x_i x_j] = \sum x_{ij}^k, i > j$ is the multiplicative table of some Lie algebra. In this case $PBW(\mathbf{S}) = \{x_{i_1} x_{i_2} \dots x_{i_k}, i_1 \leq i_2 \dots \leq i_k\}$.

If a subset \mathbf{S} of $k\langle \mathbf{X} \rangle$ is not a Gröbner-Shirshov basis then one can add to \mathbf{S} all nontrivial compositions of polynomials of \mathbf{S} , and continue this process (infinitely) many times in order to have a Gröbner-Shirshov basis \mathbf{S}^{comp} that contains \mathbf{S} . This procedure is called the Buchberger - Shirshov algorithm [70], [35], [36].

A Gröbner-Shirshov basis \mathbf{S} is called *minimal* if any $s \in \mathbf{S}$ is a linear combination of $\mathbf{S} \setminus \{s\}$ - reduced words. Any ideal of $k\langle \mathbf{X} \rangle$ with a fix monomial order has the unique minimal Gröbner-Shirshov basis.

If \mathbf{S} is a set of "semigroup relations" (that is, polynomials of the form $u - v$, where $u, v \in \mathbf{X}^*$), then any nontrivial composition will have the same form. As a result the set \mathbf{S}^{comp} also consists of semigroup relations.

Let $A = smg\langle \mathbf{X} | \mathbf{S} \rangle$ be a semigroup presentation. Then \mathbf{S} is a subset of $k\langle \mathbf{S} \rangle$ and one can find a Gröbner-Shirshov basis \mathbf{S}^{comp} . The last set does not depend on k , and consists of semigroup relations. We will call \mathbf{S}^{comp} a Gröbner-Shirshov basis of A . It is the same as a Gröbner-Shirshov basis of the semigroup algebra $kA = \langle \mathbf{X} | \mathbf{S} \rangle$.

Let $\{u_i\}$ be a set of words in an alphabet. By $V(u_i)$ we understand any word in $\{u_i\}$. Let $\{w_i\}$ be some other set of words with a surjection $u_i \mapsto w_i$. Then by $V(w_i)$ we understand the same word V , but in the set $\{w_i\}$.

3. A HYPOTHESIS

Here we will formulate a general hypothesis on Gröbner-Shirshov bases for any braid semigroup.

We will present Coxeter group W relative to Coxeter matrix M in the following way. Let S be a linear ordered l -elements set, $M = (m_{ss'})$ be $l \times l$ Coxeter matrix. Then

$$W = smg\langle S \mid s^2 = 1, (ss')^{m_{ss'}} = 1, s \neq s', \forall s, s' \in S \text{ and finite } m_{ss'} \rangle.$$

Let us define for finite $m_{ss'}$,

$$\begin{aligned} m(s, s') &= ss' \dots && \text{(there are } m_{ss'} \text{ alternative letters } s, s'), \\ (m-1)(s, s') &= ss' \dots && \text{(there are } m_{ss'} - 1 \text{ alternative letters } s, s'), \end{aligned}$$

and so on. For example, if $m_{ss'} = 2$, then $m(s, s') = ss'$, $(m-1)(s, s') = s$. If $m_{ss'} = 3$, then $m(s, s') = ss's$, $(m-1)(s, s') = ss'$.

In this notations defining relations of W can be presented in the form

$$(3.1) \quad s^2 = 1, m(s, s') = m(s', s), s > s',$$

for all $s, s' \in S$ and finite $m_{ss'}$.

Definition 3.1. The following semigroup will be called the braid semigroup W^+ relative to the Coxeter matrix M :

$$(3.2) \quad W^+ = smg\langle s_1, \dots, s_l \mid m(s, s') = m(s', s), s > s', s, s' \in S, m_{ss'} < \infty \rangle.$$

Also we will use the following notation:

$$(m, i)(s, s')$$

is a result of removing first i letters in $m(s, s')$, where $1 \leq i \leq m$.

Let us call two words in S equivalent if they are equal module commutativity relations from (3.1). To be more precise, it means that they are equal in the so called free partially commutative semigroup (algebra) generated by S and with commutativity relations from (3.1) [31]. Two relations $a = b$ and $c = d$ of W are called equivalent if a, b are equivalent to c, d respectively.

Hypothesis. A Gröbner-Shirshov basis \mathbf{T} of W^+ include of initial relations (3.2) and relations that are equivalent to the following ones:

$$(3.3) \quad \begin{aligned} &(m - i_0)(s, s')(m - i_1)(s_1, s_2) \dots (m - i_k)(s_{2k-1}, s_{2k})m(s_{2k+1}, s_{2k+2}) \\ &= m(s', s)(m, i_0)(s_2, s_1) \dots (m, i_{k-1})(s_{2k}, s_{2k-1})(m, i_k)(s_{2k+2}, s_{2k+1}), \end{aligned}$$

where $s > s'$, $s_1 < s_2, \dots, s_{2k-1} < s_{2k}, s_{2k+1} < s_{2k+2}$, and

$$\begin{aligned} (m - i_k)(s_{2k-1}, s_{2k})m(s_{2k+2}, s_{2k+1}) &\doteq m(s_{2k-1}, s_{2k})(m, i_k)(s_{2k+2}, s_{2k+1}), \\ (m - i_{k-1})(s_{2k-3}, s_{2k-2})m(s_{2k}, s_{2k-1}) &\doteq m(s_{2k-3}, s_{2k-2})(m, i_{k-1})(s_{2k}, s_{2k-1}), \\ &\cdot \\ &\cdot \\ &\cdot \\ (m - i_1)(s_1, s_2)m(s_4, s_3) &\doteq m(s_1, s_2)(m, i_1)(s_4, s_3), \\ (m - i_0)(s, s')m(s_2, s_1) &\doteq m(s, s')(m, i_0)(s_2, s_1). \end{aligned}$$

Above \doteq means the equality in the free semigroup.

Let us fix left hand and right hand sides of (3.3). If $A = B$ has a form (3.3) then a transformation

$$A \rightarrow B$$

will be called a positive chain, while $B \rightarrow A$ will be called negative.

Any other relation of \mathbf{T} is equivalent to the following one:

$$(3.4) \quad X = Y,$$

where X goes to some X' by a series of negative chains,

$$X \rightarrow X_1 \rightarrow \dots \rightarrow X_k = X',$$

(in the free partially commutative semigroup), $X' = AY_1$, and $Y = BY_1$, where $A \rightarrow B$ is a positive chain.

Remark 3.2. Let us compare this hypothesis with our hypothesis on Gröbner-Shirshov bases of Coxeter groups [31]. Relations (6.2) in [31] is just relations (3.3) with $i_0 = i_1 = \dots = i_k = 1$ and with an extra condition that all neighbor pairs $(s, s'), (s_1, s_2), \dots, (s_{2k+1}, s_{2k+2})$ are different. In particular we don't expect negative chains for Coxeter groups at all.

4. BRAID SEMIGROUP OF THE TYPE A_n

Let B_3^+ is the braid semigroup with 2 generators $a_1, a_2, a_2a_1a_2 = a_1a_2a_1$. We will assume that $a_1 < a_2$, and $u < v$ for words u, v will mean the *deg-lex* order (first we compare words by the degree (length) and then lexicographically).

Lemma 4.1. *The minimal Gröbner-Shirshov basis of B_3^+ consists of the initial relation and relations*

$$(4.1) \quad a_2a_1^l a_2a_1 = a_1a_2a_1^2a_2^{l-1}, l \geq 2$$

of a form (3.3).

Proof. First of all (4.1) is follows from Garside's formulas [49], Lemma 2:

$$a_1\Delta_2 = \Delta_2a_2, a_2\Delta_2 = \Delta_2a_1,$$

where $\Delta_2 = a_1a_2a_1$.

Secondly (4.1) has a form (3.3):

$$\begin{aligned} (m-1)(a_2, a_1)(m-2)(a_1, a_2)(m-2)(a_1, a_2) \dots (m-2)(a_1, a_2)m(a_1, a_2) = \\ m(a_1, a_2)(m, 1)(a_2, a_1)(m, 2)(a_2, a_1) \dots (m, 2)(a_2, a_1)(m, 2)(a_2, a_1). \end{aligned}$$

Now we need to proof that any composition of relations from the lemma is trivial. There are some compositions of intersections relative to the following ambiguities w :

$$\begin{aligned} a_2 a_1 a_2 a_1 a_2, a_2 a_1^l a_2 a_1 a_2, l \geq 2, a_2 a_1 a_2 a_1^l a_2 a_1, l \geq 2, \\ a_2 a_1^l a_2 a_1^k a_2 a_1, l, k \geq 2. \end{aligned}$$

Let us check the last composition for an example:

$$\begin{aligned} (f, g)_w &= (a_2 a_1^l a_2 a_1 - a_1 a_2 a_1^2 a_2^{l-1}) a_1^{k-1} a_2 a_1 - \\ & a_2 a_1^l (a_2 a_1^k a_2 a_1 - a_1 a_2 a_1^2 a_2^{k-1}) = \\ & a_2 a_1^{l+1} a_2 a_1^2 a_2^{k-1} - a_1 a_2 a_1^2 a_2^{l-1} a_1^{k-1} a_2 a_1 \equiv \\ & a_1 a_2 a_1^2 a_1^l a_2^{k-1} - a_1 a_2 a_1^2 a_2^{l-1} a_1^{k-1} a_2 a_1 \equiv 0. \end{aligned}$$

All other compositions can be figure out in the same way. \square

Let us introduce the following notations that are analogous to that of [31]:

$$\begin{aligned} a_{ij} &= a_i a_{i-1} \dots a_j, i \geq j, a_{ii} = a_i, a_{ii+1} = 1, \\ a_{ij}(\mathbf{p}) &= a_i^{p_0} a_{i-1}^{p_1} \dots a_j^{p_{i-j}}, i \geq j, a_{ii}(\mathbf{p}) = a_i^{p_0}, a_{ii+1}(\mathbf{p}) = 1. \end{aligned}$$

where $\mathbf{p} = (p_0, \dots, p_{i-j})$, all p 's are ≥ 1 . We will write $\mathbf{p} > \mathbf{1}$, if at least one of p 's is greater than 1. Otherwise we will write $\mathbf{p} = \mathbf{1}$.

It is easy to see the following equality in B_n^+ :

$$(4.2) \quad \begin{aligned} a_{ij-1} a_{i+1j-1} &= a_{i+1j-1} a_{i+1j}, j \leq i+1. \\ a_{ij-1}(\mathbf{p}) a_{i+1j-1} &= a_{i+1j-1} a_{i+1j}(\mathbf{p}), j \leq i+1. \end{aligned}$$

In fact the first of this equalities has a form:

$$(4.3) \quad \begin{aligned} (m-1)(a_i, a_{i+1})(m-1)(a_{i-1}, a_i) \dots (m-1)(a_j, a_{j+1}) m(a_{j-1}, a_j) = \\ m(a_{i+1}, a_i)(m, 1)(a_{i-1}, a_i) \dots (m, 1)(a_{j+1}, a_j)(m, 1)(a_j, a_{j-1}). \end{aligned}$$

The second of equalities (4.2) has an analogous form:

$$(4.4) \quad \begin{aligned} (m-2)(a_i, a_{i+1})^{p_0-1} (m-1)(a_i, a_{i+1}) \dots (m-2)(a_{j-1}, a_j)^{p_{i-j+1}-1}. \\ m(a_{j-1}, a_j) = m(a_{i+1}, a_i)(m, 2)(a_{i+1}, a_i)^{p_0-1} \dots (m, 1)(a_j, a_{j-1}). \\ (m, 2)(a_j, a_{j-1})^{p_{i-j+1}-1}. \end{aligned}$$

Also we will use the following Garside's formula:

$$(4.5) \quad a_s a_{i+1j} = a_{i+1j} a_{s+1}, j \leq i, j \leq s \leq i.$$

This formula looks like

$$\begin{aligned} m(a_s, a_{i+1}) \dots (m-1)(a_s, a_{s+1}) \dots (m-1)(a_{s+1}, a_j) = \\ (m, 1)(a_{i+1}, a_s) \dots (m, 1)(a_{s+1}, a_s) \dots m(a_j, a_{s+1}). \end{aligned}$$

By $W(a_j, \dots, a_i) = W(j, i)$ we will mean any word in alphabet a_j, a_{j+1}, \dots, a_i if $j \leq i$, and empty word if $j > i$.

Theorem 4.2. *A Gröbner-Shirshov bases of B_{n+1}^+ consists of the following relations :*

$$(4.6) \quad a_{i+1}a_iV(1, i-1)W(j, i)a_{i+1j} = a_i a_{i+1} a_i V(1, i-1) a_{ij} W'(j+1, i+1),$$

$$(4.7) \quad a_s a_k = a_k a_s, s - k \geq 2,$$

where $1 \leq i \leq n-1$, $1 \leq j \leq i+1$, $W' = W(a_{j+1}, \dots, a_{i+1})$, and W begins with a_i if it is not empty.

Remark 4.3. In particular the following relations are in the previous list:

$$a_{i+1}a_i a_{i+1} = a_i a_{i+1} a_i \quad (V = 1, j = i+1),$$

$$a_{i+1}a_i^l a_{i+1} a_i = a_i a_{i+1} a_i^2 a_{i+1}^{l-1}, l \geq 2 \quad (V = 1, W = a_i^{l-1}, j = i),$$

$$a_{i+1}a_i^l a_{i-1j}(\mathbf{p}) a_{i+1j} = a_i a_{i+1} a_i^2 a_{i-1j} a_{i+1}^{l-1} a_{ij+1}(\mathbf{p}), l \geq 2, i-1 \geq j$$

$$(V = 1, W = a_i^{l-1} a_{i-1j}(\mathbf{p})).$$

This relations have a form (3.3) because of Lemma 4.1 and (4.4). In general (if W is not empty) relation (4.6) has a form (3.4). It follows from above and (4.5).

Also the rite hand sides of (4.2) go to the left hand sides using the eliminations of leading words of relations (4.6).

Proof. Let \mathbf{S} is the set of relations (polynomials) from the theorem. We need to proof that any composition of polynomials from \mathbf{S} is trivial *mod* \mathbf{S} .

First of all let us deal with compositions of including. It is easy to see that any composition of including of (4.7) in (4.6) is trivial for the only possible composition of this including should correspond of some including of $a_s a_k$, $s - k \geq 2$ in VW , or else of $a_i a_k$, $i - k \geq 2$ in $a_i V$. So we need to figure out any composition of including relations of the type (4.6). Let f be a relation (polynomial) (4.6) and g be the following relation:

$$a_{i_1+1} a_{i_1} V_1(1, i_1-1) W_1(j_1, i_1) a_{i_1+1j_1} = a_{i_1} a_{i_1+1} a_{i_1} V_1(1, i_1-1) a_{i_1j_1} W_1'(j_1+1, i_1+1),$$

where $j_1 \leq i_1+1$ and W_1 begins with a_{i_1} if not empty. Let u, u_1 be the leading monomials of f, g respectively. Suppose that u_1 is a subword of u . Following possibilities may occur:

$$1) i = i_1. \text{ Then } V = V_1, W = W_1, j_1 \geq j.$$

Then

$$(f, g)_u = a_i a_{i+1} a_i V a_{i j_1} W' a_{j_1-1j} - a_i a_{i+1} a_i V a_{ij} W' \equiv 0,$$

for $W' = W_1' = W(a_{j_1+1}, \dots, a_{i+1})$, hence $W' a_{j_1-1j} \equiv a_{j_1-1j} W'$.

$$2) i = i_1 + 1. \text{ Then}$$

$$u_1 = a_i a_{i-1} V_1(1, i-2) W_1(j_1, i-1) a_{i j_1}.$$

It means that

$$V(1, i-1) = a_{i-1} V_1(1, i-2) W_1(j_1, i-1), W(j, i) = a_{i j_1} W_2(j, i).$$

We have

$$\begin{aligned}
(f, g)_u &= a_{i+1}a_{i-1}a_i a_{i-1} V_1(1, i-2) a_{i-1j_1} W'_1(j_1+1, i) W_2(j, i) a_{i+1j} - \\
& a_i a_{i+1} [a_i a_{i-1} V_1(1, i-2) W_1(j_1, i-1) a_{ij_1}] a_{j_1-1j} a_{i+1j_1+1} W'_2(j+1, i+1) \equiv \\
& a_{i-1} [a_{i+1} a_i a_{i-1} V_1(1, i-2) a_{i-1j_1} W'_1(j_1+1, i) W_2(j, i) a_{i+1j}] - \\
& a_i a_{i+1} a_{i-1} a_i a_{i-1} V_1(1, i-2) a_{i-1j_1} W'_1(j_1+1, i) a_{j_1-1j} a_{i+1j_1+1} W'_2(j+1, i+1) \equiv \\
& a_{i-1} a_i a_{i+1} [a_i a_{i-1} V_1(1, i-2) a_{i-1j_1} a_{ij_1}] a_{j_1-1j} W''_1(j_1+2, i+1) W'_2(j+1, i+1) - \\
& a_i a_{i-1} [a_{i+1} a_i a_{i-1} V_1(1, i-2) a_{i-1j_1} a_{j_1-1j} W'_1(j_1+1, i) a_{i+1j_1+1}] W'_2(j+1, i+1) \equiv \\
& a_{i-1} a_i a_{i+1} a_{i-1} a_i a_{i-1} V_1(1, i-2) a_{i-1j_1} a_{ij_1+1} a_{j_1-1j} W''_1(j_1+2, i+1) W'_2(j+1, i+1) - \\
& a_i a_{i-1} a_i a_{i+1} a_i a_{i-1} V_1(1, i-2) a_{i-1j} a_{ij_1+1} W''_1(j_1+2, i+1) W'_2(j+1, i+1) \equiv 0,
\end{aligned}$$

for $a_{ij_1+1} a_{j_1-1j} \equiv a_{j_1-1j} a_{j_1-1j}$.

3) $i - i_1 \geq 2$. The u_1 is a subword of $V(1, i-1)W(j, i)a_{i+1j}$. Then it should be a subword either V or W . First case is clear. Let us figure out second case:

$$\begin{aligned}
W &= W_2(j, i) a_{i+1} a_{i_1} V_1(1, i_1-1) W_1(j_1, i_1) a_{i_1+1j_1} W_3(j, i), \\
w &= a_{i+1} a_i V(1, i-1) W_2(j, i) a_{i_1+1} a_{i_1} V_1(1, i_1-1) W_1(j_1, i_1) a_{i_1+1j_1} W_3(j, i) a_{i+1j}, \\
(f, g)_w &= a_i a_{i+1} a_i V(1, i-1) a_{ij} W'_2(j+1, i+1) a_{i_1+2} a_{i_1+1} V'_1(2, i_1) W'_1(j_1+1, i_1+1) \cdot \\
& a_{i_1+2j_1+1} W'_3(j+1, i+1) - [a_{i+1} a_i V(1, i-1) W_2(j, i) a_{i_1} a_{i_1+1} a_{i_1} V_1(1, i_1-1) a_{i_1+1j_1} \cdot \\
& W'_1(j_1+1, i_1+1) W_3(j, i) a_{i+1j}] \equiv a_i a_{i+1} a_i V(1, i-1) a_{ij} W'_2(j+1, i+1) [a_{i_1+2} a_{i_1+1} \cdot \\
& V'_1(2, i_1) W'_1(j_1+1, i_1+1) a_{i_1+2j_1+1}] W'_3(j+1, i+1) - a_i a_{i+1} a_i V(1, i-1) a_{ij} \cdot \\
& W'_2(j+1, i+1) a_{i_1+1} a_{i_1+2} a_{i_1+1} V'_1(2, i_1) a_{i_1+1j_1+1} W''_1(j_1+2, i_1+2) W'_3(j+1, i+1) \\
& \equiv 0.
\end{aligned}$$

Now we need to check compositions of intersection.

It is easy to see that all compositions of intersection with (4.7) are trivial. Therefore we only need to consider compositions of intersection of (4.6) relative to the following ambiguities w :

1) It is a case of one letter intersection,

$$\begin{aligned}
w &= a_{i+1} a_i V(1, i-1) W(j, i) a_{i+1j} a_{j-1} V_1(1, j-2) W_1(j_1, j-1) a_{jj_1}, \\
(f, g)_w &= a_i a_{i+1} a_i V(1, i-1) a_{ij} W'(j+1, i+1) a_{j-1} V_1(1, j-2) W_1(j_1, j-1) a_{jj_1} - \\
& a_{i+1} a_i V(1, i-1) W(j, i) a_{i+1j+1} a_{j-1} a_j a_{j-1} V_1(1, j-2) a_{j-1j_1} W'_1(j_1+1, j) \equiv \\
& a_i a_{i+1} a_i V(1, i-1) a_{ij-1} W'(j+1, i+1) V_1(1, j-2) W_1(j_1, j-1) a_{jj_1} - \\
& [a_{i+1} a_i V(1, i-1) (W(j, i) a_{j-1}) a_{i+1j-1}] V_1(1, j-2) a_{j-1j_1} W'_1(j_1+1, j) \equiv \\
& a_i a_{i+1} a_i V(1, i-1) a_{ij-1} W'(j+1, i+1) V_1(1, j-2) W_1(j_1, j-1) a_{jj_1} - \\
& a_i a_{i+1} a_i V(1, i-1) a_{ij-1} (W'(j+1, i+1) a_j) V_1(1, j-2) a_{j-1j_1} W'_1(j_1+1, j) \equiv 0,
\end{aligned}$$

for $a_j V_1(1, j-2) \equiv a_j V_1(1, j-2)$ and $W_1(j_1, j-1) a_{jj_1} \equiv a_{jj_1} W'_1(j_1+1, j)$;

2) It is a case of two letters intersection,

$$\begin{aligned}
w &= a_{i+1}a_iV(1, i-1)W(j, i)a_{i+1j+2}a_{j+1}a_jV_1(1, j-1)W_1(j_1, j)a_{j+1j_1}, \\
(f, g)_w &= a_i a_{i+1} a_i V(1, i-1) a_{ij} W'(j+1, i+1) V_1(1, j-1) W_1(j_1, j) a_{j+1j_1} - \\
& a_{i+1} a_i V(1, i-1) W(j, i) a_{i+1j+2} a_j a_{j+1} a_j V_1(1, j-2) a_{jj_1} W'_1(j_1+1, j+1) \equiv \\
& a_i a_{i+1} a_i V(1, i-1) a_{ij} W'(j+1, i+1) V_1(1, j-1) W_1(j_1, j) a_{j+1j_1} - \\
& [a_{i+1} a_i V(1, i-1) (W(j, i) a_j) a_{i+1j}] V_1(1, j-2) a_{jj_1} W'_1(j_1+1, j+1) \equiv \\
& a_i a_{i+1} a_i V(1, i-1) a_{ij} W'(j+1, i+1) V_1(1, j-1) W_1(j_1, j) a_{j+1j_1} - \\
& a_i a_{i+1} a_i V(1, i-1) a_{ij} (W'(j+1, i+1) a_{j+1}) V_1(1, j-2) a_{jj_1} W'_1(j_1+1, j+1) \equiv 0,
\end{aligned}$$

for $a_{j+1}V_1(1, j-2) \equiv V_1(1, j-2)a_{j+1}$ and $W_1(j_1, j)a_{j+1j_1} \equiv a_{j+1j_1}W'_1(j_1+1, j+1)$.

3) It is a case of more than two letters intersection:

$$\begin{aligned}
w &= a_{i+1}a_iV(1, i-1)W(j, i)a_{i+1}a_i \dots a_{i+1}a_{i_1} \dots a_j V_2(1, i_1-1)W_1(j_1, i_1)a_{i_1+1j_1}, \\
(f, g)_w &= a_i a_{i+1} a_i V(1, i-1) a_{ij} W'(j+1, i+1) V_2(1, i_1-1) W_1(j_1, i_1) = a_{i_1+1j_1} - \\
& a_{i+1} a_i V(1, i-1) W(j, i) a_{i+1i_1+2} a_{i_1} a_{i+1} a_{i_1} a_{i_1-1j} V_2(1, i_1-1) a_{i_1j_1} W'_1(j_1+1, i_1+1) \equiv \\
& a_i a_{i+1} a_i V(1, i-1) a_{ij} W'(j+1, i+1) V_2(1, i_1-1) W_1(j_1, i_1) a_{i_1+1j_1} - \\
& [a_{i+1} a_i V(1, i-1) (W(j, i) a_{i_1}) a_{i+1j}] V_2(1, i_1-1) a_{i_1j_1} W'_1(j_1+1, i_1+1) \equiv \\
& a_i a_{i+1} a_i V(1, i-1) a_{ij} W'(j+1, i+1) V_2(1, i_1-1) W_1(j_1, i_1) a_{i_1+1j_1} - \\
& a_i a_{i+1} a_i V(1, i-1) a_{ij} (W'(j+1, i+1) a_{i_1+1}) V_2(1, i_1-1) a_{i_1j_1} W'_1(j_1+1, i_1+1) \equiv 0,
\end{aligned}$$

for $a_{i_1+1}V_2(1, i_1-1) \equiv V_2(1, i_1-1)a_{i_1+1}$ and $W_1(j_1, i_1)a_{i_1+1j_1} \equiv a_{i_1+1j_1}W'_1(j_1+1, i_1+1)$.

We have checked all compositions and the theorem is proved. \square

Remark 4.4. In view of [49], it is important to have an efficient algorithm to know whether a word v of B_{n+1} begin with the fundamental word Δ in some presentation, where

$$\Delta = \Lambda_1 \Lambda_2 \dots \Lambda_n, \Lambda_i = a_i \dots a_1, i \geq 1.$$

Let us formulate the following statement that easily follows from the previous theorem.

Proposition 4.5. *Let $v \in PBW(B_{n+1}^+)$. Then v begins with Δ in some presentation if and only if*

$$v \doteq \Lambda_1 V_1(1) \Lambda_2 V_2(1, 2) \dots \Lambda_{n-1} V_{n-1}(1, n-1) \Lambda_n u,$$

where $V_i(1, i)$ begins with a_1 or empty. In this case we have

$$v = \Delta V_1^{(n-1)}(n) V_2^{(n-2)}(n-1, n) \dots V_{n-1}'(2, n).$$

Proof. The result follows imediatly from the following formula:

$$PBW(a_{i+1}a_i \dots a_1 w) \doteq V(1, i) a_{i+1} a_i \dots a_1 w', w, w' \in B_{n+1}^+.$$

\square

Remark 4.6. In view of [49], Theorem 4.2 gives rise to a new algorithm of the solution of the word problem for the braid group B_{n+1} . Indeed, any group word $w \in B_{n+1}$ can be presented in a form $\Delta^k v$, where $k \in \mathbb{Z}$, $v \in B_{n+1}^+$. Using Proposition 4.5, we may present v in a form

$$v = \Delta^s u,$$

where $s \in \mathbb{N}$, u does not begin with Δ in any presentation (it means that $u \neq \Delta u_1$ in the braid semigroup). Also we may assume that u is a reduced word, $u \in PBW(B_{n+1}^+)$. Now u is nothing else as \bar{v} in the sense of [49], Theorem 5. As a result, the presentation

$$w = \Delta^{k+s} u$$

is unique due to the same Theorem 5 [49]. Let us call the last presentation the Garsaid normal form of an element of the braid group. Garside normal form is effective due to Theorem 4.2. Now the new algorithm for the solution of the word problem for the braid group can be written down using the ELW's of the Gröbner-Shirshov basis of this group.

REFERENCES

- [1] S.I. Adjan, Defining relations and algorithmical problems for groups and semigroups. Proceedings of the Steklov Math. Institute, **85**, 1958. Moscow:Nauka.
- [2] S.I. Adjan, V.G. Durnev, *Algorithmic problems for groups and semigroups*. Uspehi Math. Nauk. **55** (2000), 3–94.
- [3] E. Artin, *Theory der Zörf.* Abh. math. Sem. Hamburg Univ., **4** (1926), 47–72.
- [4] E. Artin, *Theory of Braids*. Ann. Math. **48** (1947), 101–126.
- [5] G. M. Bergman, *The diamond lemma for ring theory*. Adv. in Math. **29** (1978), 178–218.
- [6] J. Birman, K.H. Ko, S.J. Lee, *A new approach to the word and conjugacy problems in the braid groups*. Adv. Math. **139** (1998), 322–353.
- [7] L. A. Bokut, *Embedding Lie algebras into algebraically closed Lie algebras*. (Russian) Algebra i Logika Sem. **1** (1962), 47–53.
- [8] L. A. Bokut, *Some imbedding theorems for rings and semigroups. I.* (Russian) Sibirsk. Mat. Z. **4** 1963, 500–518.
- [9] L. A. Bokut, *Some embedding theorems for rings and semigroups. II.* (Russian) Sibirsk. Mat. Z. **4** 1963, 729–743.
- [10] L. A. Bokut, *On a property of the Boone groups*. Algebra i Logika Sem. **5** 1966 no. 5, 5–23.
- [11] L. A. Bokut, *On a property of the Boone groups. II.* Algebra i Logika Sem. **6** 1967 no. 1, 15–24.
- [12] L. A. Bokut, *Groups of fractions of multiplicative semigroups of certain rings. I, II, III.* Sibirsk. Mat. Z. **10** (1969), 246–286; *ibid.* **10** (1969), 744–799; *ibid.* **10** 1969 800–819.
- [13] L. A. Bokut, *The problem of Malcev*. Sibirsk. Mat. Z. **10** 1969 965–1005.
- [14] L. A. Bokut, *Unsolvability of the word problem, and subalgebras of finitely presented Lie algebras*. Izv. Akad. Nauk SSSR Ser. Mat. **36** (1972), 1173–1219.
- [15] L. A. Bokut, *Imbeddings into simple associative algebras*. Algebra i Logika **15** (1976), 117–142, 245.
- [16] L. A. Bokut, *Malcev's problem and groups with a normal form*. With the collaboration of D. J. Collins. Stud. Logic Foundations Math., **95**, Word problems, II (Conf. on Decision Problems in Algebra, Oxford, 1976), pp. 29–53, North-Holland, Amsterdam-New York, 1980.
- [17] L. A. Bokut, *New examples of groups with the standard basis*. Dokl. Akad. Nauk SSSR **277** (1984), no. 2, 277–280.
- [18] L. A. Bokut, *The centrally symmetric Novikov group*. Sibirsk. Mat. Zh. **26** (1985), no. 2, 18–28, 221.
- [19] L. A. Bokut, *On the Novikov and Boone—Borisov groups*. Illinois J. Math. **30** (1986), no. 2, 355–359.
- [20] L. A. Bokut, *Abstract semigroups and groups of formal series of dependent variables*. Siberian Advances in Mathematics. Siberian Adv. Math. **6** (1996), no. 3, 1–26.

- [21] L. A. Bokut, Y. Fong and W.-F. Ke, *Gröbner-Shirshov bases and composition lemma for associative conformal algebras: an example*. Contemporary Math. **264** (2000), 63–90.
- [22] L. A. Bokut, Y. Fong and W.-F. Ke, *Composition-Diamond lemma for associative conformal algebras*. J. Algebra, to appear.
- [23] L. A. Bokut, Y. Fong, W.-F. Ke, and P. S. Kolesnikov, *Gröbner and Gröbner-Shirshov bases in algebra and conformal algebras*, Fundamental and applied mathematics, **6** (2000), 669–706.
- [24] L. A. Bokut, S.-J. Kang, K.-H. Lee and P. Malcolmson, *Gröbner-Shirshov bases for Lie superalgebras and their universal enveloping algebras*. J. Algebra. **217** (1999), no. 2, 461–495.
- [25] L. A. Bokut and A. A. Klein, *Serre relations and Gröbner-Shirshov bases for simple Lie algebras. I, II*. Internat. J. Algebra Comput. **6** (1996), 389–400 and 401–412.
- [26] L. A. Bokut and A. A. Klein, *Gröbner-Shirshov bases for exceptional Lie algebras. I.* In “Ring theory. Selected papers from the conference held in Miskolc, July 15–20, 1996,” (L. Rowen., ed). J. Pure Appl. Algebra **133** (1998), 51–57. North-Holland Publishing Co., Amsterdam, 1998.
- [27] L. A. Bokut and A. A. Klein, *Gröbner-Shirshov bases for the exceptional Lie algebras E_6, E_7, E_8* . Algebras and combinatorics (Hong Kong, 1997), 37–46, Springer, Singapore, 1999.
- [28] L. A. Bokut, and G. P. Kukin, *Algorithmic and combinatorial algebra. Mathematics and its Applications*, **255**. Kluwer Academic Publishers Group, Dordrecht, 1994.
- [29] L. A. Bokut, and P. Malcolmson, *Gröbner-Shirshov bases for quantum enveloping algebras*. Israel J. Math. **96** (1996), 97–113.
- [30] L. A. Bokut and P. Malcolmson, *Gröbner-Shirshov bases for relations of a Lie algebra and its enveloping algebra*. Algebras and combinatorics (Hong Kong, 1997), 47–54, Springer, Singapore, 1999.
- [31] L.A. Bokut and L.-S. Shiao, *On Gröbner-Shirshov bases of Coxeter groups*, to appear in Commun. in Algebra.
- [32] L.A. Bokut and L.-S. Shiao, *Gröbner-Shirshov bases for Novikov and Boon groups*,
- [33] W.W. Boone, *The word problem*. Ann. Math. **70** (1959), 207–265.
- [34] R.V. Book and F. Otto, *String-rewriting systems*. Springer-Verlag, 1993.
- [35] B. Buchberger. *An algorithm for finding a basis for the residue class ring of a zero-dimensional polynomial ideal*. (German). Ph. D. thesis, University of Innsbruck, Austria, 1965.
- [36] B. Buchberger, *An algorithmical criteria for the solvability of algebraic systems of equations*. (German). Aequationes Math. **4** (1970), 374–383.
- [37] B. Buchberger, *A theoretical basis for the reduction of polynomials to canonical forms*. ACM SIGSAM Bull. **10** (1976), 19–29.
- [38] B. Buchberger, G. E. Collins, R. Loos, and R. Albrecht (Eds). *Computer algebra. Symbolic and algebraic computation*. Second edition. Springer-Verlag, Vienna, 1983.
- [39] B. Buchberger and F. Winkler (Eds). *Gröbner Bases and Applications*, London Mathematical Lecture Note Series **251**. Cambridge University Press, Cambridge, 1998.
- [40] K.-T. Chen, R. H. Fox and R. C. Lyndon, *Free differential calculus. IV. The quotient groups of the lower central series*. Ann. of Math. **68** (1958), 81–95.
- [41] P. M. Cohn, *Embedding in semigroup with one-sided division*. Journal London Math. Soc. **31** (1956), 169–181.
- [42] P. M. Cohn, *Embedding in sesquilateral division semigroups*. Journal London Math. Soc. **31** (1956), 181–1981.
- [43] P. M. Cohn, *Some remarks on the invariant basis property*. Topology. **5** (1966) 215–228.
- [44] P. M. Cohn, *Review of “The diamond lemma for ring theory” by George M. Bergman*, (Adv. in Math. **29** (1978), 178–218). Amer. Math. Soc. Math. Review 81b:16001.
- [45] Patric Dehornoy, *A fast method for comparing braids*. Adv. in Math. **125** (1997), N2, 200–235.
- [46] P. Deligne, *Les immeubles des groupes de tresses generalises*. Invent. Math. **17** (1972), 273–302.
- [47] David Eisenbud. *Commutative algebra. With a view toward algebraic geometry*. Graduate Texts in Mathematics, **150**. Springer-Verlag, New York, 1995.
- [48] E.A. Elrifai and H.R. Morton. *Algorithms for positive braids*. Quart. J. Math. Oxford **45** (1994), N2, 479–497.
- [49] A.F. Garside, *The braid group and other groups*. Quart. J. Math. Oxford **20** (1969), 235–254.

- [50] V. P. Gerdt and V. V. Korynyak, *Construction of finitely presented Lie algebras and superalgebras*. J. Symbolic Comput. **21** (1996), 337-349.
- [51] H. Hironaka, *Resolution of singularities of an algebraic variety over a field of characteristic zero. I, II*. Ann. of Math. **79** (1964), 109-203; *ibid.* **79** (1964) 205-326.
- [52] K.A. Kalarkoti, *Decision problems in group theory*. Proc. London Math. Soc. **44** (1982), 312-332.
- [53] S.-J. Kang, K.-H. Lee, *Gröbner-Shirshov bases for representation theory*. J. Korean Math. Soc. **37** (2000), 55-72.
- [54] S.-J. Kang, K.-H. Lee, *Gröbner-Shirshov basis theory for irreducible sl_{n+1} -modules*. J. Algebra **232** (2000), 1-20.
- [55] Seok-Jin Kang, In-Sok Lee, Kyu-Hwan Lee and Hyekyung Oh, *Hecke algebras, Specht modules and Gröbner-Shirshov bases*, to appear.
- [56] O.G. Kharlampovich, M.V. Sapir, *Algorithmic problems in varieties*. Inter. J. Algebra Computation, **5** (1995), No.4, 5, 379-602.
- [57] K.H. Ko, *New public-key cryptosystem using braid groups*. Advances in Cryptology-CRYPTO 2000, LNCS **1880** (2000) 166 - 183 (with 5 others)
- [58] V.N. Latyshev, *Canonization and standart bases of filtered structures*. In Lie Algebras, Rings and related Topics, Fong Yuen, A.A. Mikhalev, E. Zelmanov (Eds.), 61-79, Springer-Verlag, Hong Kong, 2000.
- [59] A.A. Markov, *An introduction to the algebraical theory of braids*. (Russian). Proceedings of the Steklov Mast. Ins. RAS, **16** (1945).
- [60] Alexander A. Mikhalev and Ekaterina Vasilieva, *Standard bases of ideals of free supercommutative polynomial algebras (ε -Gröbner bases)*. In Lie Algebras, Rings and related Topics, Fong Yuen, A.A. Mikhalev, E. Zelmanov (Eds.), 61-79, Springer-Verlag, Hong Kong, 2000.
- [61] A. A. Mikhalev and A. A. Zolotykh, *Standard Gröbner-Shirshov bases of free algebras over rings. I. Free associative algebras*. Internat. J. Algebra Comput. **8** (1998), no. 6, 689-726.
- [62] T. Mora, *An introduction to commutative and noncommutative Gröbner bases*. Second International Colloquium on Words, Languages and Combinatorics (Kyoto, 1992). Theoret. Comput. Sci. **134** (1994), 131-173.
- [63] M. H. A. Newman, *On theories with a combinatorial definition of "equivalence."* Ann. of Math. **43** (1942), 223-243.
- [64] P.S. Novikov, *On algorithmic unsolvability of the word problem in the theory of groups*. Moscow:Nauka, 1955 (Trudy MIAN, **44**).
- [65] E. N. Poroshenko, Gröbner-Shirshov bases for the Kac-Moody algebras $A_n^{(1)}$ and $B_n^{(1)}$. Formal power series and algebraic combinatorics, 12th International Conference, FPSAC'00, Moscow, Russia, June 2000. Proceeding, D. Krob, A.A. Mikhalev, A.V. Mikhalev (eds), Springer, Berlin, 2000, 552-563.
- [66] E. N. Poroshenko, Gröbner-Shirshov bases for the Kac-Moody algebras of the type $A_n^{(1)}$. Comm. in Algebra, **43** (2002), 6, 2617-2637.
- [67] E. N. Poroshenko, Gröbner-Shirshov bases for the Kac-Moody algebras of the type $C_n^{(1)}$ and $D_n^{(1)}$. Preprint N15, Novosibirsk State University, Novosibirsk, 2002.
- [68] E. N. Poroshenko, Gröbner-Shirshov bases for the Kac-Moody algebras of the type $B_n^{(1)}$. Preprint N16, Novosibirsk State University, Novosibirsk, 2002.
- [69] A. I. Shirshov, *On free Lie rings*. Mat. Sb. N.S. **45** (87) (1958), 113-122.
- [70] A. I. Shirshov, *Some algorithm problems for Lie algebras*. Sibirsk. Mat. Z. **3** (1962), 292-296. (Translation appears in ACM SIGSAM Bull. **33** (1999).)
- [71] K. Tatsuoka, *An isometric inequality for Artin groups of finite types*. Trans. Amer. Math. Soc. **339** (1993), no. 2, 537-551.
- [72] W. Turstone. Finite state algorithms for the braid group. Preprint, 1988.
- [73] V. Ufnarovski, *Introduction to noncommutative Gröbner bases theory*. In Gröbner bases and applications, B. Buchberger, F. Winkler (Eds). London Math. Soc. Lecture Note Ser. **251**, Cambridge Univ. Press, 1998, 259-280.

SOBOLEV INSTITUTE OF MATHEMATICS, NOVOSIBIRSK 630090, RUSSIA; = AND CHANG JUNG
CHRISTIAN UNIVERSITY, KWAY JEN, TAINAN 711, TAIWAN

E-mail address: bokut@math.nsc.ru

DEPARTMENT OF MATHEMATICS, NATIONAL CHENG-KUNG UNIVERSITY, TAINAN , TAIWAN

E-mail address: yfong@mail.ncku.edu.tw

DEPARTMENT OF MATHEMATICS, NATIONAL CHENG-KUNG UNIVERSITY, TAINAN , TAIWAN

E-mail address: wfke@mail.ncku.edu.tw

CHANG JUNG CHRISTIAN UNIVERSITY, KWAY JEN, TAINAN 711, TAIWAN

E-mail address: lsshiao@mail.cju.edu.tw