

Constructibility in Quantum Mechanics

D.J. BenDaniel

Cornell University, Ithaca, NY 14853, USA

We propose a set theoretical foundation including an axiom of constructibility and derive a generalized quantum mechanics by postulating symmetry of action. The Schrödinger equation proves to be a special case of a nonlinear sigma model. Quantum mechanics is obtained here without the requirement for presupposing the statistical interpretation of the wave function; thus this derivation becomes continuous with prior physics. In this theory we also see space-time as relational and the fields as free of singularities.

We propose the following axioms. The formulae for these axioms are given in the appendix.

TABLE I: Axioms

Extensionality	Two sets with just the same members are equal.
Pairs	For every two sets, there is a set that contains just them.
Union	For every set of sets, there is a set with just all their members.
Infinity	There are infinite ordinals ω^* .
Replacement	Replacing the members of a set one-for-one creates a set (i.e., bijective replacement).
Regularity	Every non-empty set has a minimal member (i.e. “weak” regularity).
Constructibility	The subsets of ω^* are countably constructible.

The first six axioms are the set theory of Zermelo-Frankel (ZF) without the power set axiom and with the axiom schema of subsets (a.k.a., separation) deleted from the axioms of regularity and replacement. Because of the deletion of the axiom schema of subsets, a minimal ω^* , usually denoted by ω and called the set of all finite natural numbers, cannot be shown to exist in this theory; instead this set theory is uniformly dependent on ω^* and then all the finite as well as infinitely many infinite natural numbers are included in ω^* . These infinite numbers are one-to-one with ω^* ; a finite natural number is any member of ω^* that is not infinite.

We can now adjoin to this sub-theory of ZF an axiom asserting that the subsets of ω^* are constructible. By constructible sets we mean sets that are generated sequentially by some process, one after the other, so that the process well-orders the sets. Gödel has shown that an axiom asserting that all sets are constructible can be consistently added to ZF [1], giving a theory usually called ZFC⁺. It is well known that no more than countably many subsets of ω^* can be shown to exist in ZFC⁺. This result will, of course, hold for the sub-theory ZFC⁺ minus the axiom schema of subsets and the power set axiom. Thus we can now add a new

axiom which says that the subsets of ω^* are countably constructible. This axiom, combined with the axiom schema of bijective replacement, creates a set of constructible subsets of ω^* and deletion of the power set axiom then assures that no other subsets of ω^* exist in this theory.

We shall refer to these seven axioms as T . All the sets of finite natural numbers in T are finite. The general continuum hypothesis holds in this theory because all sets are countable. We now will also show that this theory is rich enough to contain functions of a real variable effectively governing physical fields.

We first show T has a countable real line. Recall the definition of “rational numbers” as the set of ratios, in ZF called Q , of any two members of the set ω . In T , we can likewise, using the axiom of unions, establish for ω^* the set of ratios of any two of its natural numbers, finite or infinite. This will become an “enlargement” of the rational numbers and we shall call this enlargement Q^* . Two members of Q^* are called “identical” if their ratio is 1. We now employ the symbol “ \equiv ” for “is identical to.” Furthermore, an “infinitesimal” is a member of Q^* “equal” to 0, i.e., letting y signify the member and employing the symbol “ $=$ ” to signify equality, $y = 0 \leftrightarrow \forall k[y < 1/k]$, where k is a finite natural number. The reciprocal of an infinitesimal is “infinite”. A member of Q^* that is not an infinitesimal and not infinite is “finite”. The constructibility axiom well-orders the set of constructible subsets of ω^* and gives a metric space. These subsets then represent the binimals forming a countable real line R^* . In this theory R^* is a subset of Q^* .

An **equality-preserving** bijective map $\phi(x, u)$ between intervals X and U of R^* in which $x \in X$ and $u \in U$ such that $\forall x_1, x_2, u_1, u_2[\phi(x_1, u_1) \wedge \phi(x_2, u_2) \rightarrow (x_1 - x_2 = 0 \leftrightarrow u_1 - u_2 = 0)]$ creates pieces which are biunique and homeomorphic. Note that $U = 0$ if and only if $X = 0$, i.e., the piece is inherently relational.

We can now define “functions of a real variable in T ”. $u(x)$ is a function of a real variable in T only if it is a constant or a sequence in x of continuously connected biunique pieces such that the derivative of u with respect to x is also a function of a real variable in T . These functions are thus of bounded variation and locally homeomorphic. If some derivative is a constant, they are polynomials. If no derivative

is a constant, these functions do not *per se* exist in T but can, however, always be represented as closely as required for physics by a sum of polynomials of sufficiently high degree obtained by an iteration of:

$$\int_a^b \left[p \left(\frac{du}{dx} \right)^2 - qu^2 \right] dx \equiv \lambda \int_a^b ru^2 dx \quad (1)$$

where λ is minimized subject to:

$$\int_a^b ru^2 dx \equiv \text{const} \quad (2)$$

where:

$$a \neq b, \quad u \left(\frac{du}{dx} \right) \equiv 0 \quad (3)$$

at a and b ; p , q , and r are functions of the real variable x .

Letting n denote the n^{th} iteration, $\forall k \exists n [\lambda_{n-1} - \lambda_n < 1/k]$ where k is a finite natural number. So, a polynomial such that, say, $1/k < 10^{-50}$ should be sufficient for physics as it is effectively a Sturm-Liouville “eigenfunction”. These can be decomposed, since they are polynomials, into biunique “irreducible eigenfunction pieces” obeying the boundary conditions.

As a bridge to physics, let x_1 be space and x_2 be time. We now note that in the theory T we can obtain Hamilton’s Principle for a one-dimensional string $\Psi = u_1(x_1)u_2(x_2)$ just from an identity:

$$\begin{aligned} \int \left[\left(\frac{\partial \Psi}{\partial x_1} \right)^2 - \left(\frac{\partial \Psi}{\partial x_2} \right)^2 \right] dx_1 dx_2 &\equiv 0 \\ \rightarrow \delta \int \left[\left(\frac{\partial \Psi}{\partial x_1} \right)^2 - \left(\frac{\partial \Psi}{\partial x_2} \right)^2 \right] dx_1 dx_2 &\equiv 0 \\ \rightarrow \delta \int \left[\left(\frac{\partial \Psi}{\partial x_1} \right)^2 - \left(\frac{\partial \Psi}{\partial x_2} \right)^2 \right] dx_1 dx_2 &= 0 \end{aligned} \quad (4)$$

The eigenvalues λ_{1m} are determined by the spatial boundary conditions. For each eigenstate m , we can use this identity to iterate the eigenfunctions u_{1m} and u_{2m} constrained by the indicial expression $\lambda_{1m} \equiv \lambda_{2m}$.

A more general string in finitely many space-like and time-like dimensions can likewise be produced. Let $u_{\ell mi}(x_i)$ and $u_{\ell mj}(x_j)$ be eigenfunctions with non-negative eigenvalues $\lambda_{\ell mi}$ and $\lambda_{\ell mj}$ respectively.

We define a “field” as a sum of eigenstates:

$$\underline{\Psi}_m = \sum_{\ell} \Psi_{\ell m} i_{\ell}, \quad \Psi_{\ell m} = C \prod_i u_{\ell mi} \prod_j u_{\ell mj} \quad (5)$$

with the postulate: **for every eigenstate m in a compactified space the integral of the Lagrange density is identically null.**

The physics behind this will become clear. Let ds represent $\prod_i r_i dx_i$ and $d\tau$ represent $\prod_j r_j dx_j$. Then for all m ,

$$\begin{aligned} \int \sum_{\ell i} \frac{1}{r_i} \left[P_{\ell mi} \left(\frac{\partial \Psi_{\ell m}}{\partial x_i} \right)^2 - Q_{\ell mi} \Psi_{\ell m}^2 \right] ds d\tau \\ - \int \sum_{\ell j} \frac{1}{r_j} \left[P_{\ell mj} \left(\frac{\partial \Psi_{\ell m}}{\partial x_j} \right)^2 - Q_{\ell mj} \Psi_{\ell m}^2 \right] ds d\tau \equiv 0 \end{aligned}$$

In this integral expression the P , Q , and R can be functions of any of the x_i and x_j , thus of any $\Psi_{\ell m}$ as well.

This is a **nonlinear sigma model**. As we have seen in the case of a one-dimensional string, these Ψ_m can in principle be obtained by iterations constrained by an indicial expression, $\sum_{\ell i} \lambda_{\ell mi} \equiv \sum_{\ell j} \lambda_{\ell mj}$ for all m .

A proof in T that the integrals in the nonlinear sigma model have only discrete values will now be shown. Let expressions (6) and (7) both be represented by α , since they are identical:

$$\sum_{\ell mi} \int \frac{1}{r_i} \left[P_{\ell mi} \left(\frac{\partial \Psi_{\ell m}}{\partial x_i} \right)^2 - Q_{\ell mi} \Psi_{\ell m}^2 \right] ds d\tau \quad (6)$$

$$\sum_{\ell mj} \int \frac{1}{r_j} \left[P_{\ell mj} \left(\frac{\partial \Psi_{\ell m}}{\partial x_j} \right)^2 - Q_{\ell mj} \Psi_{\ell m}^2 \right] ds d\tau \quad (7)$$

- I. α is positive and must be closed to addition and to the absolute value of subtraction; in the theory T , α is a natural number times a unit that is either infinitesimal or finite.
- II. If the field is not present, $\alpha \equiv 0$; otherwise, if the field is present, then in T , α must be finite so that $\alpha \neq 0$; thus $\alpha = 0 \leftrightarrow \alpha \equiv 0$.
- III. $\therefore \alpha \equiv n\kappa$, where n is a natural number and κ is some finite positive constant such that $\alpha = 0 \leftrightarrow n \equiv 0$.

If $\pm\kappa$ is named “action,” the physical meaning of the postulate is an assertion of **symmetry of action**.

With this result and without any additional physical postulates, we can now obtain quantum mechanics from this nonlinear sigma model in one time-like dimension and finitely many space-like dimensions.

Let $\ell = 1, 2$, $r_i = P_{1mt} = P_{2mt} = 1$, $Q_{1mt} = Q_{2mt} = 0$, $\tau = \omega_m t$ and we normalize Ψ as follows:

$$\Psi_m = \sqrt{(C/2\pi)} \prod_i u_{im}(x_i) [u_{1m}(\tau) + i \cdot u_{2m}(\tau)] \quad (8)$$

where $i = \sqrt{-1}$ with

$$\int \sum_m \prod_i u_{im}^2 ds (u_{1m}^2 + u_{2m}^2) \equiv 1 \quad (9)$$

then:

$$\frac{du_{1m}}{d\tau} = -u_{2m} \quad \text{and} \quad \frac{du_{2m}}{d\tau} = u_{1m} \quad (10)$$

or

$$\frac{du_{1m}}{d\tau} = u_{2m} \quad \text{and} \quad \frac{du_{2m}}{d\tau} = -u_{1m} \quad (11)$$

For the minimal non-vanishing field, the action integral α over each irreducible eigenfunction piece of $u_{1m}(\tau)$ and $u_{2m}(\tau)$ is κ ; over a cycle it is 4κ . Thus,

$$(C/2\pi) \sum_m \oint \int \left[\left(\frac{du_{1m}}{d\tau} \right)^2 + \left(\frac{du_{2m}}{d\tau} \right)^2 \right] \prod_i u_{im}^2(x_i) ds d\tau \equiv C \equiv 4\kappa \quad (12)$$

Substituting the Planck h for 4κ , the action integral α can now be put into the conventional Lagrangian form for the Schrödinger equation,

$$\frac{1}{2i} \sum_m \oint \int \left[\Psi_m^* \left(\frac{\partial \Psi_m}{\partial t} \right) - \left(\frac{\partial \Psi_m^*}{\partial t} \right) \Psi_m \right] ds dt \quad (13)$$

Therefore, the Schrödinger equation is a special case of a nonlinear sigma model in one time dimension and finitely-many spatial dimensions.

In the process of this discussion, we have also shown that:

- Quantum mechanics is obtained without requiring an additional assumption of the statistical interpretation of the wave function, thereby resolving a long-standing controversy [3].
- Quantum mechanics is instead derived in a constructible theory using an action symmetry postulate.
- There are inherently no singularities in the physical fields obtained in this theory.

In addition, though we do not have the opportunity to discuss these points, we note that:

- Space-time is here emergent, relational and its differential properties fulfill the strict requirements of Einstein-Weyl causality [4], suggesting a possible foundation for quantum gravity.

- The solution to the QED divergence problem posed by Dyson [5] is provided, since the actual convergence or divergence of the essential perturbation series is undecidable in this theory.

- Wigner's metaphysical question regarding the apparent unreasonable effectiveness of mathematics in physics [6] is directly answered, since the foundations of mathematics and physics are now linked.

Appendix: ZF - Axiom Schema of Subsets - Power Set + Constructibility

Extensionality. Two sets with just the same members are equal. $\forall x \forall y [\forall z [z \in x \leftrightarrow z \in y] \rightarrow x = y]$

Pairs. For every two sets, there is a set that contains just them. $\forall x \forall y \exists z [\forall w [w \in z \leftrightarrow w = x \vee w = y]]$

Union. For every set of sets, there is a set with just all their members. $\forall x \exists y \forall z [z \in y \leftrightarrow \exists u [z \in u \wedge u \in x]]$

Infinity. There are infinite ordinals ω^* (i.e., sets are transitive and well-ordered by \in -relation). $\exists \omega^* [O \in \omega^* \wedge \forall x [x \in \omega^* \rightarrow x \cup \{x\} \in \omega^*]]$

Replacement. Replacing members of a set one-for-one creates a set (i.e., “bijective” replacement). Let $\phi(x, y)$ a formula in which x and y are free, $\forall z \forall x \in z \forall y [\phi(x, y) \wedge \forall u \in z \forall v [\phi(u, v) \rightarrow u = x \leftrightarrow y = v]] \rightarrow \exists r \forall t [t \in r \leftrightarrow \exists s \in z \phi(s, t)]$

Regularity. Every non-empty set has a minimal member (i.e. “weak” regularity). $\forall x [\exists y y \in x \rightarrow \exists y [y \in x \wedge \forall z \neg [z \in x \wedge z \in y]]]$

Constructibility. The subsets of ω^* are countably constructible. $\forall \omega^* \exists S [(\omega^*, O) \in S \wedge \forall y \forall z [(y, z) \in S \rightarrow [[\exists m_y [m_y \in y \wedge \forall v \neg [v \in y \wedge v \in m_y]] \wedge \exists t_y \forall u [u \in t_y \leftrightarrow u \in y \wedge u \neq m_y]] \leftrightarrow (t_y \cup m_y, z \cup \{z\}) \in S]]]$

-
- [1] Gödel, K., The consistency of the axiom of choice and of the generalized continuum hypothesis. *Annals of Math. Studies*, 1940.
- [2] Tarski, A., Mostowski, A., Robinson, R., *Undecidable Theories*. North Holland, Amsterdam, 1953.
- [3] Gottfried, K., Inferring the statistical interpretation of quantum mechanics from the classical limit. *Nature*, 2000, 405.
- [4] Sen, R.N., Causality, Measurement Theory and the

Differentiable Structure of Space-Time, Cambridge, 2010.

- [5] Dyson, F. J., Divergence of perturbation theory in quantum electrodynamics. *Phys.Rev.*, 1952, 85.
- [6] Wigner, E. P., The unreasonable effectiveness of mathematics in the natural sciences. *Comm. Pure and Appl. Math.* 1960, 13.