

Scaling of the Atmosphere of Self-Avoiding Walks

A L Owczarek¹ and T Prellberg²

¹ Department of Mathematics and Statistics,
The University of Melbourne, Victoria 3010, Australia.
a.owczarek@ms.unimelb.edu.au

² School of Mathematical Sciences
Queen Mary, University of London
Mile End Road, London E1 4NS, UK
t.prellberg@qmul.ac.uk

November 11, 2018

Abstract

The number of free sites next to the end of a self-avoiding walk is known as the *atmosphere*. The average atmosphere can be related to the number of configurations. Here we study the distribution of atmospheres as a function of length and how the number of walks of fixed atmosphere scale. Certain bounds on these numbers can be proved. We use Monte Carlo estimates to verify our conjectures. Of particular interest are walks that have zero atmosphere, which are known as *trapped*. We demonstrate that these walks scale in the *same* way as the full set of self-avoiding walks, barring an overall constant factor.

Subsequently they have generalised this elegant idea to self-avoiding walks near walls [2] and recently to self-avoiding polygons [3].

Motivating this paper is recent interest in various subsets of walks that do not *trap* [4, 5] including prudent walks [6, 7]. The question that naturally arises for each class of walks is “Does the numbers of walks in the class scale in the same way as full set of self-avoiding walks”; that is, as in equation (2) with the same value of μ and the same value of γ .

It is then natural to consider the limits

$$p^{(a)} = \lim_{n \rightarrow \infty} \frac{c_n^{(a)}}{c_n} . \quad (4)$$

In this paper we demonstrate results in the literature [8, 9] can be used to prove

$$\liminf_{n \rightarrow \infty} \frac{c_n^{(a)}}{c_n} > 0 , \quad (5)$$

and so that if one assumes the limit exists then one has immediately that

$$p^{(a)} > 0 . \quad (6)$$

Going further and assuming the scaling in equation (2) we can therefore predict

$$c_n^{(a)} \sim A_1^{(a)} \mu^n n^{\gamma-1} , \quad (7)$$

where

$$A_1^{(a)} = p^{(a)} A_1 . \quad (8)$$

Hence walks that trap, that is, those with atmosphere zero will have the same connective constant and same value of exponent γ as all self-avoiding walks. Conversely the same is true for walks that do not trap.

Furthermore we use the Monte Carlo Algorithm flatPERM[10] to provide estimates for $p_n^{(a)}$ up to length $n = 512$ and use the corrections-to-scaling to extrapolate high-precision estimates of $p^{(a)}$.

Theory Intuitively one would expect that the probability that the end of a long self-avoiding walk ω ends at the corner of a cube Q of fixed size, but otherwise does not intersect with the cube, is non-zero. Therefore one can append to this walk a short walk contained within the cube with fixed atmosphere. Using a related argument, we shall prove below that the *limes inferior* of $p_n^{(a)}$ is bounded away from zero.

Our main result is the following theorem.

Theorem 1. *Let $0 \leq a < 2d$. Then*

$$\liminf_{n \rightarrow \infty} p_n^{(a)} > 0 ,$$

and

$$\limsup_{n \rightarrow \infty} p_n^{(a)} < 1 .$$

This theorem is very similar to a result concerning tail patterns of self-avoiding walks.

Definition 2. *A self-avoiding walk $P = (p_0, p_1, \dots, p_k)$ is called a tail pattern if there is a self-avoiding walk $\omega = (\omega_0, \omega_1, \dots, \omega_n)$ such that $P = (\omega_{n-k}, \omega_{n-k+1}, \dots, \omega_n)$. P is called a proper tail pattern if for all sufficiently large n there is a self-avoiding walk having P as a tail pattern.*

Let $c_n[P]$ be the number of n -step self-avoiding walks with tail pattern P . We have the following result, which is taken from [8].

Proposition 3. *If P is a proper tail pattern then*

$$\liminf_{n \rightarrow \infty} \frac{c_n[P]}{c_n} > 0 .$$

When applying this proposition for the proof of Theorem 1, we find that the main difficulty is that the atmosphere of a proper tail pattern P and the atmosphere of a self-avoiding walk having P as a tail pattern can be different.

Proof of Theorem 1. It is sufficient to show that for any atmosphere a with $0 \leq a < 2d$ there exist a proper tail pattern P_a with atmosphere a and the additional property that any sufficiently long self-avoiding walk having P_a as a tail pattern has also atmosphere a . The first inequality then follows as an immediate consequence of Proposition 3, as $c_n^{(a)} \geq c_n[P_a]$, and therefore

$$\liminf_{n \rightarrow \infty} p_n^{(a)} = \liminf_{n \rightarrow \infty} \frac{c_n^{(a)}}{c_n} \geq \liminf_{n \rightarrow \infty} \frac{c_n[P_a]}{c_n} > 0 .$$

The upper bound follows now from the observation that $\sum_{a=0}^{2d-1} p_n^{(a)} = 1$, as

$$\begin{aligned} \limsup_{n \rightarrow \infty} p_n^{(a)} &= \limsup_{n \rightarrow \infty} \left(1 - \sum_{a' \neq a} p_n^{(a')} \right) \\ &\leq 1 + \sum_{a' \neq a} \limsup_{n \rightarrow \infty} \left(-p_n^{(a')} \right) \\ &= 1 - \sum_{a' \neq a} \liminf_{n \rightarrow \infty} p_n^{(a')} < 1 . \end{aligned}$$

The pattern P_a is constructed as follows. Consider a self-avoiding walk ω which starts at the origin, visits precisely $2d - a$ neighbouring vertices, and ends on the corner of a cube Q containing it. Continue ω by a self-avoiding walk ω' on the set of vertices ∂Q which have distance 1 from Q , and which is Hamiltonian on ∂Q . The pattern P_a which is obtained by reversal of steps is a proper tail pattern. It has atmosphere a , and by construction the atmosphere of any sufficiently long self-avoiding walk having P as a tail pattern must also be a . \square

Note that the idea of this proof generalises easily to different notions of atmosphere, such as a k -step atmosphere given by the number of $(n + k)$ -step self-avoiding walks which can be grown from an n -step self-avoiding walk.

Simulation results Backed up by the rigorous results, we now consider results of simulations for self-avoiding walks on the square lattice \mathbb{Z}^2 . Using flatPERM, a flat-histogram kinetic growth algorithm, we have grown 10^9 walks at length 512. As a consequence of using a flat-histogram method with respect to the atmosphere, we could boost the occurrence of zero-atmosphere walks roughly by a factor of three, with minimal computational overhead.

From Figure 2 it is clear that the quantities $p_n^{(a)}$ approach an asymptotic value quickly as $n \rightarrow \infty$ with only small corrections to scaling.

Figure 3 shows that these corrections are asymptotically linear in $1/n$. We therefore conjecture on the basis of our simulations that the limit

$$p^{(a)} = \lim_{n \rightarrow \infty} \frac{c_n^{(a)}}{c_n}$$

indeed exists. A linear fit can be used to obtain estimates for $p^{(a)}$, and we find that

$$p^{(0)} = 0.009096(4) \tag{9}$$

$$p^{(1)} = 0.05476(1) \tag{10}$$

$$p^{(2)} = 0.22500(2) \tag{11}$$

$$p^{(3)} = 0.71114(3) \tag{12}$$

for self-avoiding walks on the square lattice \mathbb{Z}^2 .

Acknowledgements

Financial support from the Australian Research Council via its support for the Centre of Excellence for Mathematics and Statistics of Complex Systems is gratefully acknowledged




Figure 2: A plot of the probability $p_n^{(a)}$ of an n -step self-avoiding walk on the square lattice having atmosphere a . Shown are $p_n^{(a)}$ for $a = 0, 1, 2, 3$ (from bottom to top) on a logarithmic scale versus $1/n$. We note that there are only small corrections to scaling.

by the authors.

References

- [1] A. Rechnitzer and E. J. Janse van Rensburg, *J. Phys. A: Math. Gen.* **35** L60512, 2002
- [2] A. Rechnitzer and E. J. Janse van Rensburg, *J. Phys. A: Math. Gen.* **37** 68756898, 2004
- [3] E. J. Janse van Rensburg and A. Rechnitzer, *J. Phys. A: Math. Theor.* **41** 105002, 2008
- [4] A. J. Guttmann, J. Dethridge and I. Jensen, Prudent self-avoiding walks and polygons. In *Random Polymers*, EURANDOM (June, 2007)




Figure 3: A plot of the probability $p_n^{(a)}$ of an n -step self-avoiding walk on the square lattice having atmosphere a . Shown are $p_n^{(a)}$ for $a = 0$ (bottom right), 1 (bottom left), 2 (top right), 3 (top left) on a linear scale versus $1/n$, together with linear fits of $p_n^{(a)}$ to $1/n$ computed from all walks with at least 100 steps.

- [5] M. Bousquet-Mélou, Families of Prudent Self-avoiding walks *arXiv:0804.4843v1 [math.CO]*, 2008
- [6] L. Turban and J.-M. Debievre. *J. Phys. A*, 20(3):679–686, 1987.
- [7] S. B. Santra, W. A. Seitz, and D. J. Klein. *Phys. Rev. E*, 63(6):067101–1–4, 2001.
- [8] N. Madras and G. Slade. *The self-avoiding walk*. Probability and its Applications. Birkhäuser Boston Inc., Boston, MA, 1993.
- [9] N. Madras, *J. Stat. Phys.* **53** 689–701, 1988.
- [10] Prellberg T and Krawczyk J, 2004 *Phys. Rev. Lett.* **92** 120602