

A TAMED 3D NAVIER-STOKES EQUATION IN DOMAINS

XICHENG ZHANG

*School of Mathematics and Statistics
The University of New South Wales, Sydney, 2052, Australia
Department of Mathematics, Huazhong University of Science and Technology,
Wuhan, Hubei 430074, P.R.China
Email: XichengZhang@gmail.com*

ABSTRACT. In this paper, we analyze a tamed 3D Navier-Stokes equation in uniform C^2 -domains (not necessarily bounded), which obeys the scaling invariance principle, and prove the existence and uniqueness of strong solutions to this tamed equation. In particular, if there exists a bounded solution to the classical 3D Navier-Stokes equation, then this solution satisfies our tamed equation. Moreover, the existence of a global attractor for the tamed equation in bounded domains is also proved. As simple applications, some well known results for the classical Navier-Stokes equations in unbounded domains are covered.

1. INTRODUCTION

The motion of a viscous incompressible fluid in a domain $\Omega \subset \mathbb{R}^3$ is described by the Navier-Stokes equation (NSE) as follows (with homogeneous boundary):

$$\begin{cases} \partial_t \mathbf{u} = \nu \Delta \mathbf{u} - (\mathbf{u} \cdot \nabla) \mathbf{u} + \nabla P + \mathbf{f}, \\ \operatorname{div}(\mathbf{u}) = 0, \quad (t, x) \in [0, \infty) \times \Omega, \\ \mathbf{u}(t, x) = 0, \quad t \geq 0, \quad x \in \partial\Omega, \quad \mathbf{u}(0) = \mathbf{u}_0, \end{cases} \quad (1)$$

where $\nu > 0$ is the kinematic viscosity constant, $\mathbf{u}(t, x) = (u_1(t, x), u_2(t, x), u_3(t, x))$ represents the velocity field, $P = P(t, x)$ is the pressure (an unknown scalar function), \mathbf{f} is a known external force.

The study of 3D NSEs has a long history. In their pioneering works, Leray [11] and Hopf [9] proved the existence of a weak solution to equation (1). Since then, there are many papers devoted to the study of regularities of Leray-Hopf weak solutions (cf. [10, 19, 17, etc.]). Up to now, one knows that the singular set of the Leray-Hopf weak solutions has Lebesgue measure zero (cf. [11, 8, 7]). Moreover, a deep result obtained by Scheffer [16] and Caffarelli, Kohn and Nirenberg [3] says that the singular set for a class of weak solutions (satisfying a generalized energy inequality) has one dimensional Hausdorff measure zero (see also [12]). However, the uniqueness and regularity of Leray-Hopf weak solutions are still big open problems.

Most of the source of difficulties to solve equation (1) comes from the nonlinear term $(\mathbf{u} \cdot \nabla) \mathbf{u}$ (cf. [7]). In order to counteract this term, the authors in [15] analyzed the following modified (called tamed therein) 3D NSE in $\Omega = \mathbb{R}^3$:

$$\begin{cases} \partial_t \mathbf{u} = \nu \Delta \mathbf{u} - (\mathbf{u} \cdot \nabla) \mathbf{u} + \nabla P - g'_N(|\mathbf{u}|^2) \mathbf{u} + \mathbf{f}, \\ \operatorname{div}(\mathbf{u}) = 0, \quad t \geq 0, \quad x \in \mathbb{R}^3, \quad \mathbf{u}(0) = \mathbf{u}_0, \end{cases} \quad (2)$$

where $|\mathbf{u}|^2 := \sum_{j=1}^3 |u_j|^2$ and for $N > 0$

$$g_N^\nu(r) := (r - N) \cdot 1_{\{r \geq N\}} / \nu. \quad (3)$$

The existence of a unique smooth solution to equation (2) was proved in [15] when the initial velocity is smooth (in Sobolev spaces). The main feature of equation (2) is that there exists a bounded solution (say bounded by \sqrt{N} for some large N) to the classical NSE, then this solution must satisfy equation (2). Therein, the property that the Leray projection operator onto the space of divergence free vector fields commutes with the derivatives plays a key role. But, when we consider NSE (1) in a domain, this property does not hold in general (cf. [13, p.83-85]).

In order to deal with the Dirichlet boundary problem and keep the same feature as equation (2), in the present paper, we consider the following globally tamed scheme (assuming $\mathbf{f} = 0$ for simplicity):

$$\partial_t \mathbf{u} = \nu \Delta \mathbf{u} - (\mathbf{u} \cdot \nabla) \mathbf{u} + \nabla P - g_N^{\nu, \kappa} (\|\mathbf{u} - \mathbf{U}\|_\infty^2) (\mathbf{u} - \mathbf{U}), \quad (4)$$

where $\|\mathbf{u}\|_\infty := \sup_{x \in \Omega} |\mathbf{u}(x)|$, \mathbf{U} is a reference velocity field and for $\kappa, N \geq 1$

$$g_N^{\nu, \kappa}(r) := \kappa \cdot (r - N) 1_{\{r \geq N\}} / \nu.$$

Here, $\kappa \geq 1$ is a dimensionless constant and \sqrt{N} has the velocity dimension.

Let $(\mathbf{u}_{N, \mathbf{U}}, P_{N, \mathbf{U}})$ be a solution pair of equation (4). Simple calculations show that $(\mathbf{u}_{N, \mathbf{U}}, P_{N, \mathbf{U}})$ has the following properties:

(A) (Galilean invariance): for any constant velocity vector $\mathbf{v} \in \mathbb{R}^3$

$$\begin{aligned} \mathbf{u}_{N, \mathbf{U}}^{\mathbf{v}}(t, x) &:= \mathbf{u}_{N, \mathbf{U} + \mathbf{v}}(t, x - \mathbf{v}t) + \mathbf{v}, \\ P_{N, \mathbf{U}}^{\mathbf{v}}(t, x) &:= P_{N, \mathbf{U} + \mathbf{v}}(t, x - \mathbf{v}t) \end{aligned}$$

is also a solution pair of equation (4).

(B) (Rotation symmetry): for any orthogonal matrix \mathcal{Q} (i.e. $\mathcal{Q}\mathcal{Q}^t = I$)

$$\begin{aligned} \mathbf{u}_{N, \mathbf{U}}^{\mathcal{Q}}(t, x) &:= \mathcal{Q}^t \mathbf{u}_{N, \mathcal{Q}^t \mathbf{U}}(t, \mathcal{Q}x), \\ P_{N, \mathbf{U}}^{\mathcal{Q}}(t, x) &:= P_{N, \mathcal{Q}^t \mathbf{U}}(t, \mathcal{Q}x) \end{aligned}$$

is also a solution pair of equation (4).

(C) (Scale invariance): for any $\lambda > 0$

$$\begin{aligned} \mathbf{u}_{N, \mathbf{U}}^\lambda(t, x) &:= \lambda \mathbf{u}_{\lambda^{-2}N, \lambda^{-1}\mathbf{U}}(\lambda^2 t, \lambda x), \\ P_{N, \mathbf{U}}^\lambda(t, x) &:= \lambda^2 P_{\lambda^{-2}N, \lambda^{-1}\mathbf{U}}(\lambda^2 t, \lambda x) \end{aligned}$$

is also a solution pair of equation (4).

These three properties are exhibited by the classical Navier-Stokes equations (cf. [2]).

Intuitively, when the maximum of the fluid velocity is larger than \sqrt{N} , the dissipative term $g_N^{\nu, \kappa} (\|\mathbf{u}\|_\infty^2) \mathbf{u}$ (regarded as some extra force) will enter into the equation and restrain the flux of the liquid. In this sense, the value of N plays the role of a valve. On the other hand, when we realize equation (4) on a computer, the value of N can be reset as an arbitrarily large number along with the process of calculations as long as there is no explosion. So, the term involving $g_N^{\nu, \kappa}$ plays the role of some kind of adjustment. The parameter κ can be understood as the extent of the extra dissipative force, and will be used to give a better estimate for $\|\mathbf{u}\|_\infty$ in terms of N (see part (III) of Theorem 2.4).

In contrast with equation (2), the tamed equation (4) in domain Ω is global since $g_N^{\nu, \kappa} (\|\mathbf{u}\|_\infty^2)$ depends on all values of \mathbf{u} in Ω . But, better than (2), it is easy to write down

the vorticity equation: Let $\omega = \operatorname{curl} \mathbf{u} = \nabla \wedge \mathbf{u}$. Then

$$\partial_t \omega = \nu \Delta \omega + (\omega \cdot \nabla) \mathbf{u} - (\mathbf{u} \cdot \nabla) \omega - g_N^{\nu, \kappa} (\|\mathbf{u} - \mathbf{U}\|_\infty^2) \omega.$$

We remark that in [4], Caraballo, Real and Kloeden studied the following globally modified NSE in a bounded regular domain Ω :

$$\begin{cases} \partial_t \mathbf{u}(t) = \Delta \mathbf{u} - \min\{1, N/\|\nabla \mathbf{u}\|_{\mathbf{L}^2}\} (\mathbf{u} \cdot \nabla) \mathbf{u} + \nabla P, \\ \operatorname{div}(\mathbf{u}) = 0, \quad (t, x) \in [0, \infty) \times \Omega, \\ \mathbf{u}(t, x) = 0, \quad t \geq 0, \quad x \in \partial\Omega, \quad \mathbf{u}(0) = \mathbf{u}_0, \end{cases} \quad (5)$$

and they proved the existence of a unique strong solution to this modified equation as well as the existence of a global attractor. Nevertheless, equation (5) does not enjoy the above properties **(A)**-**(C)**.

This paper is organized as follows: In Section 2, all main results are announced. In Section 3, we prepare some necessary lemmas for later use. In the remaining sections, we shall give the proofs of main results. We want to emphasize that for the proof of existence of strong solutions (see Section 4), not using the usual Galerkin approximation, we only use the linearized equations and simple Picard's iteration. Moreover, the semigroup method used in Fujita-Kato [6] (cf. [17]) will be used to improve the regularity of strong solutions (see Section 5). The existence of a global attractor for the evolution semigroup determined by equation (4) will follow by proving some asymptotic compactness (cf. [4, 20, etc.]).

2. ANNOUNCEMENT OF MAIN RESULTS

Throughout this paper, all \mathbb{R}^3 -valued functions and spaces of such functions will be denoted by boldfaced letters, and we use the following convention: the letter C with or without subscripts will denote a positive constant whose value may change in different occasions.

Let Ω be a uniform C^3 -regular domain of \mathbb{R}^3 (see [1, p.84] for the definition of regular domains). Let $\mathbf{C}_0^\infty(\Omega)$ denote the set of all smooth functions from Ω to \mathbb{R}^3 with compact supports in Ω , and $\mathbf{C}_{0,\sigma}^\infty(\Omega) \subset \mathbf{C}_0^\infty(\Omega)$ the set of all smooth vector fields of divergence free. For $p > 1$, let $\mathbf{L}^p(\Omega)$ be the usual \mathbb{R}^3 -valued L^p -space with the norm denoted by $\|\cdot\|_{\mathbf{L}^p(\Omega)} = \|\cdot\|_{\mathbf{L}^p}$, and $\mathbf{L}_\sigma^2(\Omega)$ the closure of $\mathbf{C}_{0,\sigma}^\infty(\Omega)$ in $\mathbf{L}^2(\Omega)$. For $k \in \mathbb{N}$ and $p > 1$, let $\mathbf{W}^{k,p}(\Omega)$ be the space of \mathbb{R}^3 -valued functions with finite norm:

$$\|\mathbf{u}\|_{\mathbf{W}^{k,p}(\Omega)} := \|\mathbf{u}\|_{\mathbf{W}^{k,p}} := \left(\sum_{j=0}^k \int_{\Omega} |\nabla^j \mathbf{u}(x)|^p dx \right)^{\frac{1}{p}} < +\infty,$$

where ∇^j denotes the j -th order generalized derivative operator. The space $\mathbf{W}_0^{1,2}(\Omega)$ (resp. $\mathbf{W}_{0,\sigma}^{1,2}(\Omega)$) denotes the completion of $\mathbf{C}_0^\infty(\Omega)$ (resp. $\mathbf{C}_{0,\sigma}^\infty(\Omega)$) with respect to the above norm with $k = 1$ and $q = 2$.

Let \mathcal{P} be the orthogonal projection from $\mathbf{L}^2(\Omega)$ to $\mathbf{L}_\sigma^2(\Omega)$. By A (called the Stokes operator) we denote the self-adjoint operator in $\mathbf{L}_\sigma^2(\Omega)$ formally given by

$$A := -\mathcal{P}\Delta.$$

More precisely, $\mathbf{u} \in \mathcal{D}(A)$ if and only if for some $\mathbf{w} \in \mathbf{L}_\sigma^2(\Omega)$ (written as $A\mathbf{u} = \mathbf{w}$), it holds that

$$\langle \nabla \mathbf{u}, \nabla \mathbf{v} \rangle_{\mathbf{L}^2} = \langle \mathbf{w}, \mathbf{v} \rangle_{\mathbf{L}^2}, \quad \forall \mathbf{v} \in \mathbf{W}_{0,\sigma}^{1,2}(\Omega).$$

In particular, $\mathcal{D}(A^{\frac{1}{2}}) = \mathbf{W}_{0,\sigma}^{1,2}(\Omega)$ and

$$\|A^{\frac{1}{2}} \mathbf{u}\|_{\mathbf{L}^2} = \|\nabla \mathbf{u}\|_{\mathbf{L}^2}, \quad \mathbf{u} \in \mathbf{W}_{0,\sigma}^{1,2}(\Omega). \quad (6)$$

Moreover, it is well known that (cf. [17, p.129])

$$\mathcal{D}(A) = \mathbf{W}^{2,2}(\Omega) \cap \mathbf{W}_{0,\sigma}^{1,2}(\Omega).$$

Since A is a positive self-adjoint operator in $\mathbf{L}_\sigma^2(\Omega)$, for $\alpha \in (-1, 1)$, the fractional power A^α is well defined via the spectral representation. For $\beta \in [0, 2]$, define the Hilbert space

$$\mathbf{H}^\beta(\Omega) := \mathbf{H}^\beta := \mathcal{D}(A^{\beta/2})$$

with the norm $\|\cdot\|_{\mathbf{H}^\beta}$ generated by inner product

$$\langle \mathbf{u}, \mathbf{v} \rangle_{\mathbf{H}^\beta} := \langle \mathbf{u}, \mathbf{v} \rangle_{\mathbf{L}^2} + \langle A^{\beta/2} \mathbf{u}, A^{\beta/2} \mathbf{v} \rangle_{\mathbf{L}^2}.$$

We introduce the following bilinear form B on $\mathbf{W}_{0,\sigma}^{1,2}(\Omega) = \mathbf{H}^1$:

$$B(\mathbf{v}, \mathbf{u}) := -\mathcal{P}((\mathbf{v} \cdot \nabla) \mathbf{u}). \quad (7)$$

Using \mathcal{P} to act on both sides of equation (4), we can and shall consider the following equivalent abstract equation

$$\partial_t \mathbf{u} = -\nu A \mathbf{u} + B(\mathbf{u}, \mathbf{u}) - g_N^{\nu,\kappa} (\|\mathbf{u}\|_\infty^2) \mathbf{u}, \quad \mathbf{u}(0) = \mathbf{u}_0. \quad (8)$$

We give the following definition of strong solutions to the above equation.

Definition 2.1. Let $\mathbf{u}_0 \in \mathbf{H}^1$. A continuous function

$$\mathbb{R}_+ \ni t \mapsto \mathbf{u}(t) \in \mathbf{H}^1$$

is called a strong solution of equation (8) if $\mathbf{u} \in L_{loc}^2(\mathbb{R}_+; \mathbf{H}^2)$ and for all $t \geq 0$

$$\mathbf{u}(t) = \mathbf{u}_0 - \nu \int_0^t A \mathbf{u} ds + \int_0^t B(\mathbf{u}, \mathbf{u}) ds - \int_0^t g_N^{\nu,\kappa} (\|\mathbf{u}\|_\infty^2) \mathbf{u} ds \quad \text{in } \mathbf{L}_\sigma^2(\Omega). \quad (9)$$

Our first main result is stated as follows:

Theorem 2.2. Let Ω be a uniform C^2 -domain of \mathbb{R}^3 . For any $\mathbf{u}_0 \in \mathbf{H}^1$, there exists a unique strong solution $\mathbf{u}(t) = \mathbf{u}_N(t)$ to equation (8) in the sense of Definition 2.1, which satisfies that for any $t \geq 0$

$$\|\mathbf{u}(t)\|_{\mathbf{L}^2}^2 + 2\nu \int_0^t \|\nabla \mathbf{u}\|_{\mathbf{L}^2}^2 dt \leq \|\mathbf{u}_0\|_{\mathbf{L}^2}^2, \quad (10)$$

$$\|\nabla \mathbf{u}(t)\|_{\mathbf{L}^2}^2 + \nu \int_0^t \|A \mathbf{u}\|_{\mathbf{L}^2}^2 dt \leq \frac{\kappa N}{\nu^2} \|\mathbf{u}_0\|_{\mathbf{L}^2}^2 + \|\nabla \mathbf{u}_0\|_{\mathbf{L}^2}^2 \quad (11)$$

and for some $T^* = T^*(\nu, \Omega, \|\mathbf{u}_0\|_{\mathbf{L}^2})$ and $C = C(\nu, \Omega, \|\mathbf{u}_0\|_{\mathbf{L}^2})$

$$\|\nabla \mathbf{u}(t)\|_{\mathbf{L}^2} \leq C/\sqrt{t}, \quad \forall t \geq T^*. \quad (12)$$

Moreover, letting $\mathbf{u}_N(t)$ (resp. $\mathbf{v}_M(t)$) be the solution of equation (8) with initial value $\mathbf{u}_0 \in \mathbf{H}^1$ (resp. $\mathbf{v}_0 \in \mathbf{H}^1$) and taming function $g_N^{\nu,\kappa}$ (resp. $g_M^{\nu,\kappa}$), we have for any $T > 0$

$$\begin{aligned} & \sup_{t \in [0, T]} \|\mathbf{u}_N(t) - \mathbf{v}_M(t)\|_{\mathbf{H}^1}^2 + \int_0^T \|\mathbf{u}_N - \mathbf{v}_M\|_{\mathbf{H}^1}^2 ds \\ & \leq C(\nu, N, M, \|\mathbf{u}_0\|_{\mathbf{H}^1}, \|\mathbf{v}_0\|_{\mathbf{H}^1}, T) \cdot (|N - M|^2 + \|\mathbf{u}_0 - \mathbf{v}_0\|_{\mathbf{H}^1}^2), \end{aligned} \quad (13)$$

where the constant $C(\nu, N, M, \|\mathbf{u}_0\|_{\mathbf{H}^1}, \|\mathbf{v}_0\|_{\mathbf{H}^1}, T)$ continuously depends on its parameters.

Remark 2.3. For $T > 0$ and $N \geq 1$, define

$$\mathbb{T}_N^T := \{t \in [0, T] : \|\mathbf{u}(t)\|_\infty \geq \sqrt{N}\}.$$

By (10), (11) and (21) below, we have

$$\begin{aligned} \lambda(\mathbb{T}_N^T) &\leq \frac{1}{N} \int_0^T \|\mathbf{u}\|_\infty^2 ds \leq \frac{C_\Omega}{N} \int_0^T \|\mathbf{u}\|_{\mathbf{H}^2} \cdot \|\nabla \mathbf{u}\|_{\mathbf{L}^2} ds \leq \\ &\leq \frac{C_\Omega}{N} \left(\int_0^T \|\mathbf{u}\|_{\mathbf{H}^2}^2 ds \right)^{1/2} \cdot \left(\int_0^T \|\nabla \mathbf{u}\|_{\mathbf{L}^2}^2 ds \right)^{1/2} \\ &\leq \frac{C_\Omega \cdot \|\mathbf{u}_0\|_{\mathbf{L}^2} \cdot \left(\frac{\kappa N}{\nu^2} \|\mathbf{u}_0\|_{\mathbf{L}^2}^2 + \|\nabla \mathbf{u}_0\|_{\mathbf{L}^2}^2 + \nu T \|\mathbf{u}_0\|_{\mathbf{L}^2}^2 \right)^{1/2}}{\sqrt{2\nu}N}, \end{aligned}$$

where $\lambda(\mathbb{T}_N^T)$ denotes the Lebesgue measure of \mathbb{T}_N^T . This gives an estimate of the length of the time for which \mathbf{u}_N does not satisfy equation (1). In particular,

$$\lim_{N \rightarrow \infty} \lambda(\mathbb{T}_N^T) = 0,$$

which shows that as N goes to infinity, \mathbf{u}_N satisfies equation (1) at ‘‘almost all’’ times.

We are now interested in the estimation of $\|\mathbf{u}\|_\infty$ in terms of N and prove the following result.

Theorem 2.4. *Let Ω be a uniform C^2 -domain and $\mathbf{u}_0 \in \mathbf{H}^2$. Let \mathbf{u}_N^κ be the unique strong solution in Theorem 2.2. We have the following conclusions:*

- (I) *There exist two continuous function $K_1 : \mathbb{R}_+^2 \rightarrow \mathbb{R}_+$ and $K_2 : \mathbb{R}_+^3 \rightarrow \mathbb{R}_+$ such that for all $t \geq 0$ and $N \geq 1$*

$$\|\mathbf{u}_N^\kappa(t)\|_\infty \leq K_1(t, \|\mathbf{u}_0\|_{\mathbf{H}^2}) + K_2(t, \nu, \|\mathbf{u}_0\|_{\mathbf{L}^2}) \cdot N^3, \quad (14)$$

where $K_1(t, r), K_2(t, \nu, r) \rightarrow 0$ as $t \rightarrow 0$ or $\nu \rightarrow \infty$ or $r \rightarrow 0$. In particular, for $T > 0$, if one of the following conditions is satisfied, then there is a unique strong solution in $[0, T]$ for equation (1):

- (i) T is small; (ii) $\|\mathbf{u}_0\|_{\mathbf{H}^2}$ is small; (iii) ν is large.

- (II) *Let $\Omega = \mathbb{R}^3$ or be a bounded uniform C^4 -domain and $\mathbf{u} = \mathbf{u}_N^\kappa$. Then*

$$\mathbf{u} \in C([0, \infty) \times \bar{\Omega}; \mathbb{R}^3)$$

and for $i, j = 1, 2, 3$

$$\partial_t \mathbf{u}, \partial_i \mathbf{u}, \partial_i \partial_j \mathbf{u} \in C((0, \infty) \times \bar{\Omega}; \mathbb{R}^3).$$

Moreover, for some $P \in C((0, \infty) \times \bar{\Omega}; \mathbb{R})$ (with $\int_\Omega P(x) dx = 0$), it holds that

$$\partial_t \mathbf{u} = \Delta \mathbf{u} - (\mathbf{u} \cdot \nabla) \mathbf{u} + \nabla P - g_N(\|\mathbf{u}\|_\infty^2) \mathbf{u}, \quad \forall (t, x) \in (0, \infty) \times \Omega. \quad (15)$$

- (III) *Let $\Omega = \mathbb{R}^3$ and $\nu > 0$. For any $\alpha > \frac{1}{2}$, there exist $\kappa > 0$ and two functions $K_{1,\alpha,\kappa}, K_{2,\alpha,\kappa}$ as in (I) such that for all $t \geq 0$ and $N \geq 1$*

$$\|\mathbf{u}_N^\kappa(t)\|_\infty \leq K_{1,\alpha,\kappa}(t, \|\mathbf{u}_0\|_{\mathbf{H}^2}) + K_{2,\alpha,\kappa}(t, \nu, \|\mathbf{u}_0\|_{\mathbf{L}^2}) \cdot N^\alpha. \quad (16)$$

Remark 2.5. We do not know whether the α in (III) can be smaller than $1/2$. If this can be proven, then (1) will have a classical solution. In fact, even for $\alpha = 1/2$, it seems also hard to prove (16).

Remark 2.6. Fix $T > 0$ and $N_1 \geq 1$. Define a sequence of real numbers recursively as follows:

$$N_{k+1} := \sup_{t \in [0, T]} \|\mathbf{u}_{N_k}(t)\|_\infty^2, \quad k \in \mathbb{N}.$$

It is easy to see that equation (1) has a explosion solution in $[0, T]$ if and only if

$$N_1 < N_2 < N_3 < \cdots < N_k \rightarrow \infty.$$

The strict monotonicity is clear. Assume that $\lim_{k \rightarrow \infty} N_k = N_\infty < \infty$. By the continuous dependence of \mathbf{u}_N with respect to N (see (13)), we have

$$\lim_{k \rightarrow \infty} \sup_{t \in [0, T]} \|\mathbf{u}_{N_k}(t) - \mathbf{u}_{N_\infty}(t)\|_\infty^2 = 0.$$

Therefore,

$$N_\infty = \sup_{t \in [0, T]} \|\mathbf{u}_{N_\infty}(t)\|_\infty^2 < \infty,$$

which implies that $\mathbf{u}_{N_\infty}(t)$ satisfies (1), no explosion.

For $\mathbf{u}_0 \in \mathbf{H}^1$, let $\{\mathbf{u}(t; \mathbf{u}_0); t \geq 0\}$ be the unique strong solution of equation (8), which defines a nonlinear evolution semigroup:

$$S(t)\mathbf{u}_0 := \mathbf{u}(t; \mathbf{u}_0) : \mathbf{H}^1 \rightarrow \mathbf{H}^1. \quad (17)$$

By Theorem 2.2, $\{S(t); t \geq 0\}$ has the following properties:

- (i) $S(0) = I$ identity map on \mathbf{H}^1 ;
- (ii) $S(t+s) = S(t)S(s)$ for any $t, s \geq 0$;
- (iii) $[0, \infty) \times \mathbf{H}^1 \ni (t, \mathbf{u}_0) \mapsto S(t)\mathbf{u}_0 \in \mathbf{H}^1$ is continuous.

Definition 2.7. A compact subset $\mathcal{A} \subset \mathbf{H}^1$ is called a global attractor of the evolution semigroup $\{S(t); t \geq 0\}$ if

- (i) \mathcal{A} is invariant under $S(t)$, i.e., for any $t > 0$, $S(t)\mathcal{A} = \mathcal{A}$;
- (ii) \mathcal{A} attracts all bounded set $\mathcal{U} \subset \mathbf{H}^1$, i.e.,

$$\lim_{t \rightarrow \infty} \rho(S(t)\mathcal{U}, \mathcal{A}) = 0,$$

$$\text{where } \rho(\mathcal{A}_1, \mathcal{A}_2) := \sup_{\mathbf{u} \in \mathcal{A}_1} \inf_{\mathbf{v} \in \mathcal{A}_2} \|\mathbf{u} - \mathbf{v}\|_{\mathbf{H}^1}.$$

We have the following existence of global attractors of $\{S(t), t \geq 0\}$.

Theorem 2.8. Let Ω be a bounded uniform C^2 -domain of \mathbb{R}^3 . Then there exists a global attractor $\mathcal{A} \subset \mathbf{H}^1$ to $\{S(t); t \geq 0\}$ defined by (17).

3. PRELIMINARIES

In this section, we collect some necessary materials for later use. The following lemma is from [8, Lemma 6].

Lemma 3.1. Let $\phi : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ be an absolute continuous function and $g : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ a locally Lipschitz continuous function. Suppose that $\Lambda := \int_0^\infty \phi(t) dt < +\infty$ and $g(\phi) \leq \alpha \phi^2$ for $\phi \leq \beta$, where $\alpha, \beta > 0$. If

$$\phi'(t) \leq g(\phi(t)), \quad \forall t \geq 0,$$

then for $t \geq (\Lambda/\beta) \exp(\alpha\Lambda)$

$$\phi(t) \leq (e^{\alpha\Lambda} - 1)/(\alpha t).$$

Let $\{E_\lambda, \lambda > 0\}$ be the spectrum decomposition of A in $\mathbf{L}_\sigma^2(\Omega)$. The Stokes semigroup is then defined by

$$e^{-tA} := \int_0^\infty e^{-t\lambda} dE_\lambda,$$

and for $\alpha \in [-1, 1]$, A^α is given by

$$A^\alpha := \int_0^\infty \lambda^\alpha dE_\lambda.$$

The following lemma is easily derived from the above representations (cf. [17]).

Lemma 3.2. (i) For any $\alpha \in [0, 1]$ and $\mathbf{u} \in \mathbf{L}_\sigma^2(\Omega)$, we have $e^{-tA}\mathbf{u} \in \mathcal{D}(A^\alpha)$ and

$$\|A^\alpha e^{-tA}\mathbf{u}\|_{\mathbf{L}^2} \leq t^{-\alpha} \|\mathbf{u}\|_{\mathbf{L}^2}, \quad \forall t > 0.$$

(ii) For all $\mathbf{u} \in \mathcal{D}(A^\alpha)$ and $t \geq 0$

$$A^\alpha e^{-tA}\mathbf{u} = e^{-tA}A^\alpha\mathbf{u}, \quad \|e^{-tA}\mathbf{u} - \mathbf{u}\|_{\mathbf{L}^2} \leq C_\alpha t^\alpha \|A^\alpha\mathbf{u}\|_{\mathbf{L}^2}.$$

(iii) For any $0 \leq \alpha < \gamma < \beta \leq 1$ and $\mathbf{u} \in \mathcal{D}(A^\beta)$

$$\|A^\gamma\mathbf{u}\|_{\mathbf{L}^2} \leq \|A^\beta\mathbf{u}\|_{\mathbf{L}^2}^{\frac{\gamma-\alpha}{\beta-\alpha}} \cdot \|A^\alpha\mathbf{u}\|_{\mathbf{L}^2}^{\frac{\beta-\gamma}{\beta-\alpha}} \leq \frac{\gamma-\alpha}{\beta-\alpha} \|A^\beta\mathbf{u}\|_{\mathbf{L}^2} + \frac{\beta-\gamma}{\beta-\alpha} \|A^\alpha\mathbf{u}\|_{\mathbf{L}^2}.$$

We recall the following well known results (cf. [17, Lemma 2.4.2 (p.142), Lemma 2.5.2 (p.152) and Lemma 2.4.3 (p.143)]).

Lemma 3.3. (i) For $\alpha \in [0, 1/2]$ and $q = \frac{6}{3-4\alpha}$, there exists a constant $C = C(\alpha, q) > 0$ such that for any $\mathbf{u} \in \mathbf{H}^{2\alpha}$

$$\|\mathbf{u}\|_{\mathbf{L}^q} \leq C \|A^\alpha\mathbf{u}\|_{\mathbf{L}^2}. \quad (18)$$

(ii) For $\alpha \in [0, 1/2]$ and $q = \frac{6}{3+4\alpha}$, there exists a constant $C = C(\alpha, q) > 0$ such that for any $\mathbf{u} \in \mathbf{L}^q(\Omega)$

$$\|A^{-\alpha}\mathcal{P}\mathbf{u}\|_{\mathbf{L}^2} \leq C \|\mathbf{u}\|_{\mathbf{L}^q}. \quad (19)$$

(iii) For $\alpha \in [1/2, 1]$ and $q = \frac{6}{5-4\alpha}$, there exists a constant $C = C(\Omega, \alpha, q) > 0$ such that for any $\mathbf{u} \in \mathbf{H}^{2\alpha}$

$$\|\mathbf{u}\|_{\mathbf{W}^{1,q}} \leq C (\|A^\alpha\mathbf{u}\|_{\mathbf{L}^2} + \|\mathbf{u}\|_{\mathbf{L}^2}). \quad (20)$$

This lemma has the following conclusions.

Lemma 3.4. Let Ω be a uniform C^2 -domain. For some $C_\Omega > 0$ and any $\mathbf{u} \in \mathbf{H}^2(\Omega)$

$$\|\mathbf{u}\|_{\mathbf{L}^\infty(\Omega)}^2 \leq C_\Omega \cdot \|\mathbf{u}\|_{\mathbf{H}^2(\Omega)} \cdot \|\nabla\mathbf{u}\|_{\mathbf{L}^2(\Omega)}, \quad (21)$$

and for $\frac{3}{4} < \alpha \leq 1$, some $C_{\alpha,\Omega} > 0$ and any $\mathbf{u} \in \mathbf{H}^{2\alpha}(\Omega)$

$$\|\mathbf{u}\|_{\mathbf{L}^\infty(\Omega)} \leq C_{\alpha,\Omega} \cdot (\|A^\alpha\mathbf{u}\|_{\mathbf{L}^2(\Omega)} + \|\mathbf{u}\|_{\mathbf{L}^2(\Omega)}). \quad (22)$$

Proof. Since Ω is a uniform C^2 -domain, by [1, p. 154, Theorem 5.24] there exists a bounded linear operator $E : \mathbf{W}^{k,p}(\Omega) \mapsto \mathbf{W}^{k,p}(\mathbb{R}^3)$ such that $E\mathbf{u} = \mathbf{u}$ a.e. on Ω . Recall the following Gagliardo-Nirenberg inequality (cf. [5, p.24, Theorem 9.3]): Let $1 \leq p, q \leq \infty$ and $\alpha \in [0, 1]$ with $p \neq 3$ and

$$\frac{1}{r} = \alpha \left(\frac{1}{p} - \frac{1}{3} \right) + (1 - \alpha) \frac{1}{q}.$$

Then, for some $C = C(r, p, q)$ and all $\mathbf{u} \in \mathbf{W}^{1,p}(\mathbb{R}^3) \cap \mathbf{L}^q(\mathbb{R}^3)$

$$\|\mathbf{u}\|_{\mathbf{L}^r(\mathbb{R}^3)} \leq C \|\mathbf{u}\|_{\mathbf{W}^{1,p}(\mathbb{R}^3)}^\alpha \cdot \|\mathbf{u}\|_{\mathbf{L}^q(\mathbb{R}^3)}^{1-\alpha}. \quad (23)$$

Thus, by (18) we have

$$\begin{aligned}\|\mathbf{u}\|_{\mathbf{L}^\infty(\Omega)}^2 &\leq \|E\mathbf{u}\|_{\mathbf{L}^\infty(\mathbb{R}^3)}^2 \leq C\|E\mathbf{u}\|_{\mathbf{W}^{2,2}(\mathbb{R}^3)} \cdot \|E\mathbf{u}\|_{\mathbf{L}^6(\mathbb{R}^3)} \\ &\leq C_\Omega\|\mathbf{u}\|_{\mathbf{W}^{2,2}(\Omega)} \cdot \|\mathbf{u}\|_{\mathbf{L}^6(\Omega)} \\ &\leq C_\Omega\|\mathbf{u}\|_{\mathbf{H}^2(\Omega)} \cdot \|A^{\frac{1}{2}}\mathbf{u}\|_{\mathbf{L}^2(\Omega)}\end{aligned}$$

and for $q = \frac{6}{5-4\alpha} > 3$

$$\begin{aligned}\|\mathbf{u}\|_{\mathbf{L}^\infty(\Omega)} &\leq \|E\mathbf{u}\|_{\mathbf{L}^\infty(\mathbb{R}^3)} \leq C_q\|E\mathbf{u}\|_{\mathbf{W}^{1,q}(\mathbb{R}^3)} \leq C_{q,\Omega}\|\mathbf{u}\|_{\mathbf{W}^{1,q}(\Omega)} \\ (20) \quad &\leq C_{\alpha,\Omega} \cdot (\|A^\alpha\mathbf{u}\|_{\mathbf{L}^2(\Omega)} + \|\mathbf{u}\|_{\mathbf{L}^2(\Omega)}).\end{aligned}$$

The proof is complete. \square

The contents below in this section are only used in Section 5.

Lemma 3.5. *For some $C, C_\Omega > 0$ and all $\mathbf{u} \in \mathbf{H}^2 = \mathcal{D}(A)$, we have*

$$\|A^{-\frac{1}{4}}B(\mathbf{u}, \mathbf{u})\|_{\mathbf{L}^2} \leq C\|A^{\frac{1}{2}}\mathbf{u}\|_{\mathbf{L}^2}^2, \quad (24)$$

$$\|B(\mathbf{u}, \mathbf{u})\|_{\mathbf{L}^2} \leq C_\Omega(\|A^{\frac{5}{8}}\mathbf{u}\|_{\mathbf{L}^2}^2 + \|\mathbf{u}\|_{\mathbf{L}^2}^2). \quad (25)$$

Proof. By Hölder's inequality, we have

$$\|A^{-\frac{1}{4}}B(\mathbf{u}, \mathbf{u})\|_{\mathbf{L}^2} \stackrel{(19)}{\leq} C\|(\mathbf{u} \cdot \nabla)\mathbf{u}\|_{\mathbf{L}^{3/2}} \leq C\|\mathbf{u}\|_{\mathbf{L}^6} \cdot \|\nabla\mathbf{u}\|_{\mathbf{L}^2} \stackrel{(18)}{\leq} C\|A^{\frac{1}{2}}\mathbf{u}\|_{\mathbf{L}^2}^2$$

and

$$\begin{aligned}\|B(\mathbf{u}, \mathbf{u})\|_{\mathbf{L}^2} &\leq \|(\mathbf{u} \cdot \nabla)\mathbf{u}\|_{\mathbf{L}^2} \leq \|\mathbf{u}\|_{\mathbf{L}^{12}} \cdot \|\mathbf{u}\|_{\mathbf{W}^{1,12/5}} \\ &\leq C_\Omega\|\mathbf{u}\|_{\mathbf{W}^{1,12/5}}^2 \stackrel{(20)}{\leq} C_\Omega(\|A^{\frac{5}{8}}\mathbf{u}\|_{\mathbf{L}^2}^2 + \|\mathbf{u}\|_{\mathbf{L}^2}^2),\end{aligned}$$

where the third inequality is due to $\mathbf{W}^{1,12/5}(\Omega) \subset \mathbf{L}^{12}(\Omega)$. \square

Lemma 3.6. *For any $\frac{3}{4} < \gamma < \beta \leq 1$, there are three positive continuous functions $F_1, F_3 : \mathbb{R}_+^2 \rightarrow \mathbb{R}_+$ and $F_2 : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ such that for all $\mathbf{u}, \mathbf{v} \in \mathbf{H}^{2\beta}$*

$$\begin{aligned}\|B(\mathbf{u}, \mathbf{u}) - B(\mathbf{v}, \mathbf{v})\|_{\mathbf{L}^2} + \|g_N^{\nu,\kappa}(\|\mathbf{u}\|_\infty^2)\mathbf{u} - g_N^{\nu,\kappa}(\|\mathbf{v}\|_\infty^2)\mathbf{v}\|_{\mathbf{L}^2} \\ \leq F_1(\|\mathbf{u}\|_{\mathbf{H}^{2\beta}}, \|\mathbf{v}\|_{\mathbf{H}^{2\beta}}) \cdot \|A^\beta(\mathbf{u} - \mathbf{v})\|_{\mathbf{L}^2}^{\frac{\gamma}{\beta}} \cdot \|\mathbf{u} - \mathbf{v}\|_{\mathbf{L}^2}^{\frac{\beta-\gamma}{\beta}} \\ + F_2(\|\mathbf{u}\|_{\mathbf{H}^{2\beta}}) \cdot \|A^\beta(\mathbf{u} - \mathbf{v})\|_{\mathbf{L}^2}^{\frac{1}{2\beta}} \cdot \|\mathbf{u} - \mathbf{v}\|_{\mathbf{L}^2}^{1-\frac{1}{2\beta}} \\ + F_3(\|\mathbf{u}\|_{\mathbf{H}^{2\beta}}, \|\mathbf{v}\|_{\mathbf{H}^{2\beta}}) \cdot \|\mathbf{u} - \mathbf{v}\|_{\mathbf{L}^2}.\end{aligned}$$

Proof. Note that by (iii) of Lemma 3.2

$$\|\nabla(\mathbf{u} - \mathbf{v})\|_{\mathbf{L}^2} = \|A^{\frac{1}{2}}(\mathbf{u} - \mathbf{v})\|_{\mathbf{L}^2} \leq \|A^\beta(\mathbf{u} - \mathbf{v})\|_{\mathbf{L}^2}^{\frac{1}{2\beta}} \cdot \|\mathbf{u} - \mathbf{v}\|_{\mathbf{L}^2}^{1-\frac{1}{2\beta}}$$

and

$$\begin{aligned}\|\mathbf{u} - \mathbf{v}\|_\infty &\stackrel{(22)}{\leq} C_\Omega(\|A^\gamma(\mathbf{u} - \mathbf{v})\|_{\mathbf{L}^2} + \|\mathbf{u} - \mathbf{v}\|_{\mathbf{L}^2}) \\ &\leq C_\Omega(\|A^\beta(\mathbf{u} - \mathbf{v})\|_{\mathbf{L}^2}^{\frac{\gamma}{\beta}} \cdot \|\mathbf{u} - \mathbf{v}\|_{\mathbf{L}^2}^{\frac{\beta-\gamma}{\beta}} + \|\mathbf{u} - \mathbf{v}\|_{\mathbf{L}^2}).\end{aligned}$$

The result now follows from

$$\|B(\mathbf{u}, \mathbf{u}) - B(\mathbf{v}, \mathbf{v})\|_{\mathbf{L}^2} \leq \|\mathbf{u}\|_\infty \cdot \|\nabla(\mathbf{u} - \mathbf{v})\|_{\mathbf{L}^2} + \|\mathbf{u} - \mathbf{v}\|_\infty \cdot \|\nabla\mathbf{v}\|_{\mathbf{L}^2}$$

and

$$\|g_N^{\nu,\kappa}(\|\mathbf{u}\|_\infty^2)\mathbf{u} - g_N^{\nu,\kappa}(\|\mathbf{v}\|_\infty^2)\mathbf{v}\|_{\mathbf{L}^2} \leq \frac{\kappa}{\nu}\|\mathbf{u} - \mathbf{v}\|_\infty \cdot (\|\mathbf{u}\|_\infty + \|\mathbf{v}\|_\infty) \cdot \|\mathbf{v}\|_{\mathbf{L}^2}$$

$$+\frac{\kappa}{\nu}\|\mathbf{u}\|_{\infty}^2 \cdot \|\mathbf{u} - \mathbf{v}\|_{\mathbf{L}^2}.$$

□

We introduce some notations. Let I be a closed interval of t , and let \mathbb{X} be a Banach space. By $C(I; \mathbb{X})$ we denote the set of all continuous \mathbb{X} -valued functions defined on I . For $0 < \theta < 1$, $C^\theta(I; \mathbb{X})$ means the set of all functions which are strongly Hölder continuous with the exponent θ . If I is not closed, $v \in C^\theta(I; \mathbb{X})$ means that $v \in C^\theta(I_1; \mathbb{X})$ for any closed interval I_1 contained in I .

The following lemma is easily deduced from Lemma 3.2 (cf. [6, 14]).

Lemma 3.7. *For $T > 0$, let $\mathbf{f} : [0, T] \mapsto \mathbf{H}^0 = \mathbf{L}_\sigma^2(\Omega)$ be continuous and consider*

$$\mathbf{w}(t) := \int_0^t e^{-(t-s)A} \mathbf{f}(s) ds.$$

(i) *For any $0 \leq \alpha < \theta < 1$*

$$A^\alpha \mathbf{w} \in C^{1-\theta}([0, T], \mathbf{H}^0), \quad \|A^\alpha \mathbf{w}(t)\|_{\mathbf{L}^2} \leq C_\alpha \cdot t^{1-\alpha} \cdot \sup_{s \in [0, T]} \|\mathbf{f}(s)\|_{\mathbf{L}^2}.$$

(ii) *If $\mathbf{f} \in C([0, T], \mathbf{H}^0) \cap C^\alpha((0, T], \mathbf{H}^0)$ for some $\alpha \in (0, 1)$, then for any $0 < \theta < \alpha$*

$$A \mathbf{w} \in C^\theta((0, T], \mathbf{H}^0), \quad \partial_t \mathbf{w} \in C((0, T], \mathbf{H}^{2\theta}).$$

Moreover, $(0, T]$ can be replaced by $[0, T]$ in the above condition and conclusions.

Proof. The first conclusion is direct from Lemma 3.2. For the second, fixing $\delta \in (0, T)$, we write

$$\mathbf{w}(t) = e^{-(t-\delta)A} \mathbf{w}(\delta) + \int_\delta^t e^{-(t-s)A} \mathbf{f}(s) ds =: \Psi_\delta(t) + \Phi_\delta(t).$$

It is easy to see that

$$\Psi_\delta(\cdot) \in C^\infty((\delta, T], \mathbf{H}^0)$$

and

$$\begin{aligned} \partial_t \Phi_\delta(t) &= e^{-(t-\delta)A} \mathbf{f}(t) - \int_\delta^t A e^{-(t-s)A} (\mathbf{f}(s) - \mathbf{f}(t)) ds \\ &= -A \Phi_\delta(t) + \mathbf{f}(t), \quad \delta \leq t \leq T. \end{aligned}$$

(ii) now follows from Lemma 3.2. □

For $\alpha \in [0, 1]$, let $\mathbf{W}^{k+\alpha, 2}(\Omega)$ be the complex interpolation space between $\mathbf{W}^{k, 2}(\Omega)$ and $\mathbf{W}^{k+1, 2}(\Omega)$. The following lemma is easily derived by [19, p.23, Proposition 2.2] and the interpolation theorem (cf. [21]).

Lemma 3.8. *Let $k \in \mathbb{N} \cup \{0\}$ and $\Omega \subset \mathbb{R}^3$ be a bounded domain of class C^{k+2} . For any $\mathbf{f} \in \mathbf{W}^{\alpha, 2}(\Omega)$, $0 \leq \alpha \leq k$, there exist unique functions $\mathbf{u} \in \mathbf{W}^{2+\alpha, 2}(\Omega)$ and $P \in W^{1+\alpha, 2}(\Omega)$ (with $\int_\Omega P dx = 0$), which solve the following Stokes problem in the distribution sense:*

$$\nu \Delta \mathbf{u} = \nabla P + \mathbf{f}, \quad \operatorname{div}(\mathbf{u}) = 0, \quad \mathbf{u}|_{\partial\Omega} = 0.$$

Moreover, there exists a constant $C_{\alpha, \nu} > 0$ such that

$$\|\mathbf{u}\|_{\mathbf{W}^{2+\alpha, 2}(\Omega)} + \|P\|_{W^{1+\alpha, 2}(\Omega)} \leq C_{\alpha, \nu} \cdot \|\mathbf{f}\|_{\mathbf{W}^{\alpha, 2}(\Omega)}.$$

4. PROOF OF THEOREM 2.2

In this section, we use the following equivalent norm in \mathbf{H}^β ($\beta \in [0, 2]$)

$$\|\mathbf{u}\|_{\mathbf{H}^\beta} = \|(I + A)^{\beta/2}\mathbf{u}\|_{\mathbf{L}^2}, \quad \mathbf{u} \in \mathbf{H}^\beta.$$

We first prove:

Lemma 4.1. *For any $\mathbf{u}, \mathbf{v}, \mathbf{u}', \mathbf{v}' \in \mathbf{H}^2$ we have*

$$\begin{aligned} & \langle B(\mathbf{u}', \mathbf{u}) - g_N^{\nu, \kappa}(\|\mathbf{u}'\|_\infty^2)\mathbf{u} - B(\mathbf{v}', \mathbf{v}) + g_N^{\nu, \kappa}(\|\mathbf{v}'\|_\infty^2)\mathbf{v}, (I + A)(\mathbf{u} - \mathbf{v}) \rangle_{\mathbf{L}^2} \quad (26) \\ & \leq \frac{3\nu}{4}\|\mathbf{u} - \mathbf{v}\|_{\mathbf{H}^2}^2 + \frac{\nu}{8}\|\mathbf{u}' - \mathbf{v}'\|_{\mathbf{H}^2}^2 + \frac{\kappa N}{\nu} \cdot \|\mathbf{w}\|_{\mathbf{H}^1}^2 \\ & \quad + \frac{C_\Omega}{\nu^3}\|\mathbf{u}' - \mathbf{v}'\|_{\mathbf{H}^1}^2 \cdot ((1 + 4\kappa)\|\mathbf{v}\|_{\mathbf{H}^1}^2 + \kappa\|\mathbf{u}\|_{\mathbf{H}^1}^2)^2. \end{aligned}$$

Proof. Set

$$\mathbf{w} = \mathbf{u} - \mathbf{v}, \quad \mathbf{w}' = \mathbf{u}' - \mathbf{v}',$$

and write (26) as the following four terms' sum

$$\begin{aligned} I_1 & := \langle B(\mathbf{u}', \mathbf{w}), (I + A)\mathbf{w} \rangle_{\mathbf{L}^2}, \\ I_2 & := \langle B(\mathbf{w}', \mathbf{v}), (I + A)\mathbf{w} \rangle_{\mathbf{L}^2}, \\ I_3 & := -\langle g_N^{\nu, \kappa}(\|\mathbf{u}'\|_\infty^2)\mathbf{w}, (I + A)\mathbf{w} \rangle_{\mathbf{L}^2}, \\ I_4 & := -\langle [g_N^{\nu, \kappa}(\|\mathbf{u}'\|_\infty^2) - g_N^{\nu, \kappa}(\|\mathbf{v}'\|_\infty^2)]\mathbf{v}, (I + A)\mathbf{w} \rangle_{\mathbf{L}^2}. \end{aligned}$$

By $ab \leq \frac{\nu}{4\epsilon}a^2 + \frac{\epsilon}{\nu}b^2$, we have

$$\begin{aligned} I_1 & \leq \|B(\mathbf{u}', \mathbf{w})\|_{\mathbf{L}^2} \cdot \|(I + A)\mathbf{w}\|_{\mathbf{L}^2} \\ & \leq \frac{1}{2\nu}\|(\mathbf{u}' \cdot \nabla)\mathbf{w}\|_{\mathbf{L}^2}^2 + \frac{\nu}{2}\|\mathbf{w}\|_{\mathbf{H}^2}^2 \\ & \leq \frac{1}{2\nu}\|\mathbf{u}'\|_\infty^2\|\nabla\mathbf{w}\|_{\mathbf{L}^2}^2 + \frac{\nu}{2}\|\mathbf{w}\|_{\mathbf{H}^2}^2 \end{aligned}$$

and similarly,

$$I_2 \leq \frac{1}{\nu}\|\mathbf{w}'\|_\infty^2\|\nabla\mathbf{v}\|_{\mathbf{L}^2}^2 + \frac{\nu}{4}\|\mathbf{w}\|_{\mathbf{H}^2}^2.$$

For I_3 , by $g_N^{\nu, \kappa}(r) \geq \frac{\kappa}{\nu}(r - N)$ we have

$$\begin{aligned} I_3 & = -g_N^{\nu, \kappa}(\|\mathbf{u}'\|_\infty^2) \cdot \|(I + A)^{1/2}\mathbf{w}\|_{\mathbf{L}^2}^2 \\ & \leq -\frac{\kappa}{\nu}\|\mathbf{u}'\|_\infty^2 \cdot \|\mathbf{w}\|_{\mathbf{H}^1}^2 + \frac{\kappa N}{\nu} \cdot \|\mathbf{w}\|_{\mathbf{H}^1}^2. \end{aligned}$$

For I_4 , by $|g_N^{\nu, \kappa}(r) - g_N^{\nu, \kappa}(r')| \leq \frac{\kappa}{\nu}|r - r'|$ we have

$$\begin{aligned} I_4 & \leq \frac{\kappa}{\nu}\|\mathbf{w}'\|_\infty(\|\mathbf{u}'\|_\infty + \|\mathbf{v}'\|_\infty) \cdot \|\mathbf{v}\|_{\mathbf{H}^1} \cdot \|\mathbf{w}\|_{\mathbf{H}^1} \\ & \leq \frac{\kappa}{\nu}\|\mathbf{w}'\|_\infty \cdot (2\|\mathbf{u}'\|_\infty + \|\mathbf{w}'\|_\infty) \cdot \|\mathbf{v}\|_{\mathbf{H}^1} \cdot \|\mathbf{w}\|_{\mathbf{H}^1} \\ & = \frac{2\kappa}{\nu}(\|\mathbf{u}'\|_\infty\|\mathbf{w}\|_{\mathbf{H}^1}) \cdot (\|\mathbf{w}'\|_\infty\|\mathbf{v}\|_{\mathbf{H}^1}) \\ & \quad + \frac{\kappa}{\nu}\|\mathbf{w}'\|_\infty^2 \cdot \|\mathbf{v}\|_{\mathbf{H}^1} \cdot \|\mathbf{w}\|_{\mathbf{H}^1} \\ & \leq \frac{\kappa}{2\nu}\|\mathbf{u}'\|_\infty^2\|\mathbf{w}\|_{\mathbf{H}^1}^2 + \frac{2\kappa}{\nu}\|\mathbf{w}'\|_\infty^2\|\mathbf{v}\|_{\mathbf{H}^1}^2 \\ & \quad + \frac{\kappa}{\nu}\|\mathbf{w}'\|_\infty^2 \cdot \|\mathbf{v}\|_{\mathbf{H}^1} \cdot (\|\mathbf{u}\|_{\mathbf{H}^1} + \|\mathbf{v}\|_{\mathbf{H}^1}). \end{aligned}$$

Combining the above calculations, we obtain

$$\begin{aligned}
I_1 + I_2 + I_3 + I_4 &\leq \frac{3\nu}{4} \|\mathbf{w}\|_{\mathbf{H}^2}^2 + \frac{\kappa N}{\nu} \cdot \|\mathbf{w}\|_{\mathbf{H}^1}^2 + \frac{(1-\kappa)}{2\nu} \|\mathbf{u}'\|_{\infty}^2 \cdot \|\mathbf{w}\|_{\mathbf{H}^1}^2 \\
&\quad + \frac{1}{\nu} \|\mathbf{w}'\|_{\infty}^2 \cdot ((1+3\kappa)\|\mathbf{v}\|_{\mathbf{H}^1}^2 + \kappa\|\mathbf{v}\|_{\mathbf{H}^1} \cdot \|\mathbf{u}\|_{\mathbf{H}^1}) \\
(21) \quad &\leq \frac{3\nu}{4} \|\mathbf{w}\|_{\mathbf{H}^2}^2 + \frac{\kappa N}{\nu} \cdot \|\mathbf{w}\|_{\mathbf{H}^1}^2 + \frac{(1-\kappa)}{2\nu} \|\mathbf{u}'\|_{\infty}^2 \cdot \|\mathbf{w}\|_{\mathbf{H}^1}^2 \\
&\quad + \frac{C_{\Omega}}{\nu} \|\mathbf{w}'\|_{\mathbf{H}^2} \|\nabla \mathbf{w}'\|_{\mathbf{L}^2} \cdot ((1+4\kappa)\|\mathbf{v}\|_{\mathbf{H}^1}^2 + \kappa\|\mathbf{u}\|_{\mathbf{H}^1}^2) \\
&\stackrel{(\kappa \geq 1)}{\leq} \frac{3\nu}{4} \|\mathbf{w}\|_{\mathbf{H}^2}^2 + \frac{\nu}{8} \|\mathbf{w}'\|_{\mathbf{H}^2}^2 + \frac{\kappa N}{\nu} \cdot \|\mathbf{w}\|_{\mathbf{H}^1}^2 \\
&\quad + \frac{C_{\Omega}^2}{\nu^3} \|\mathbf{w}'\|_{\mathbf{H}^1}^2 \cdot ((1+4\kappa)\|\mathbf{v}\|_{\mathbf{H}^1}^2 + \kappa\|\mathbf{u}\|_{\mathbf{H}^1}^2)^2,
\end{aligned}$$

which produces the desired estimate. \square

4.1. Proof of Existence. Let $\mathbf{v} \in C([0, \infty); \mathbf{H}^1) \cap L_{loc}^2(\mathbb{R}_+; \mathbf{H}^2)$. We first consider the following linearized equation:

$$\partial_t \mathbf{u} = -\nu A \mathbf{u} + B(\mathbf{v}, \mathbf{u}) - g_N^{\nu, \kappa}(\|\mathbf{v}\|_{\infty}^2) \mathbf{u}, \quad \mathbf{u}(0) = \mathbf{u}_0 \in \mathbf{H}^1.$$

By the standard theory of PDE, there is a unique strong solution \mathbf{u} to above equation with

$$\mathbf{u} \in C([0, \infty); \mathbf{H}^1) \cap L_{loc}^2(\mathbb{R}_+; \mathbf{H}^2).$$

Let us construct the approximation sequence of equation (8) as follows: Set $\mathbf{u}_1(t) \equiv 0$. For $k = 2, 3, \dots$, let

$$\mathbf{u}_k(t) \in C([0, \infty); \mathbf{H}^1) \cap L_{loc}^2(\mathbb{R}_+; \mathbf{H}^2) \quad (27)$$

solve the following equation

$$\partial_t \mathbf{u}_k = -\nu A \mathbf{u}_k + B(\mathbf{u}_{k-1}, \mathbf{u}_k) - g_N^{\nu, \kappa}(\|\mathbf{u}_{k-1}\|_{\infty}^2) \mathbf{u}_k, \quad \mathbf{u}_k(0) = \mathbf{u}_0. \quad (28)$$

Firstly, note that

$$\langle A \mathbf{u}_k, \mathbf{u}_k \rangle_{\mathbf{L}^2} = \|\nabla \mathbf{u}_k\|_{\mathbf{L}^2}^2$$

and

$$\langle B(\mathbf{u}_{k-1}, \mathbf{u}_k), \mathbf{u}_k \rangle_{\mathbf{L}^2} = -\langle (\mathbf{u}_{k-1} \cdot \nabla) \mathbf{u}_k, \mathbf{u}_k \rangle_{\mathbf{L}^2} = -\frac{1}{2} \langle \operatorname{div} \mathbf{u}_{k-1}, |\mathbf{u}_k|^2 \rangle_{\mathbf{L}^2} = 0.$$

By the chain rule, we have from (28) that

$$d\|\mathbf{u}_k\|_{\mathbf{L}^2}^2/dt = -2\nu \|\nabla \mathbf{u}_k\|_{\mathbf{L}^2}^2 - 2g_N^{\nu, \kappa}(\|\mathbf{u}_{k-1}\|_{\infty}^2) \|\mathbf{u}_k\|_{\mathbf{L}^2}^2 \leq -2\nu \|\nabla \mathbf{u}_k\|_{\mathbf{L}^2}^2. \quad (29)$$

Integrating both sides of (29) yields that

$$\|\mathbf{u}_k(t)\|_{\mathbf{L}^2}^2 + 2\nu \int_0^t \|\nabla \mathbf{u}_k\|_{\mathbf{L}^2}^2 dt \leq \|\mathbf{u}_0\|_{\mathbf{L}^2}^2, \quad \forall t \geq 0. \quad (30)$$

Secondly, for any $T > 0$ we have

$$\begin{aligned}
\int_0^T |g_N^{\nu, \kappa}(\|\mathbf{u}_{k-1}\|_{\infty}^2)|^2 \cdot \|\mathbf{u}_k\|_{\mathbf{L}^2}^2 dt &\leq \frac{\kappa}{\nu} \sup_{t \in [0, T]} \|\mathbf{u}_k(t)\|_{\mathbf{L}^2}^2 \int_0^T \|\mathbf{u}_{k-1}\|_{\infty}^4 dt \\
(21) \quad &\leq C \cdot \sup_{t \in [0, T]} \|\mathbf{u}_k(t)\|_{\mathbf{L}^2}^2 \cdot \sup_{t \in [0, T]} \|\mathbf{u}_{k-1}(t)\|_{\mathbf{H}^1}^2 \int_0^T \|\mathbf{u}_{k-1}\|_{\mathbf{H}^2}^2 dt \stackrel{(27)}{<} +\infty
\end{aligned}$$

and

$$\begin{aligned} \int_0^T \|B(\mathbf{u}_{k-1}, \mathbf{u}_k)\|_{\mathbf{L}^2}^2 dt &\leq C \int_0^T \|\mathbf{u}_{k-1}\|_\infty^2 \|\nabla \mathbf{u}_k\|_{\mathbf{L}^2}^2 dt \\ &\stackrel{(22)}{\leq} C \int_0^T \|\mathbf{u}_{k-1}\|_{\mathbf{H}^2}^2 \cdot \|\nabla \mathbf{u}_k\|^2 dt \stackrel{(27)}{<} +\infty. \end{aligned} \quad (31)$$

Thus, recalling $\mathbf{H}^0 = \mathbf{L}_\sigma^2(\Omega)$, from (28) one has

$$\partial_t \mathbf{u}_k \in L_{loc}^2(\mathbb{R}_+; \mathbf{H}^0).$$

Consider the evolution triple

$$\mathbf{H}^2 \subset \mathbf{H}^1 \subset \mathbf{H}^0.$$

By the chain rule (cf. [19, p.176, Lemma 1.2]) and Young's inequality, we have

$$\begin{aligned} \frac{1}{2} \frac{d\|\nabla \mathbf{u}_k\|_{\mathbf{L}^2}^2}{dt} &= -\nu \|A\mathbf{u}_k\|_{\mathbf{L}^2}^2 + \langle B(\mathbf{u}_{k-1}, \mathbf{u}_k), A\mathbf{u}_k \rangle_{\mathbf{L}^2} - g_N^{\nu, \kappa}(\|\mathbf{u}_{k-1}\|_\infty^2) \|\nabla \mathbf{u}_k\|_{\mathbf{L}^2}^2 \\ &\leq -\frac{\nu}{2} \|A\mathbf{u}_k\|_{\mathbf{L}^2}^2 + \frac{1}{2\nu} \|\mathbf{u}_{k-1}\|_\infty^2 \|\nabla \mathbf{u}_k\|_{\mathbf{L}^2}^2 - g_N^{\nu, \kappa}(\|\mathbf{u}_{k-1}\|_\infty^2) \|\nabla \mathbf{u}_k\|_{\mathbf{L}^2}^2 \\ &\leq -\frac{\nu}{2} \|A\mathbf{u}_k\|_{\mathbf{L}^2}^2 + \frac{1-2\kappa}{2\nu} \|\mathbf{u}_{k-1}\|_\infty^2 \|\nabla \mathbf{u}_k\|_{\mathbf{L}^2}^2 + \frac{\kappa N}{\nu} \|\nabla \mathbf{u}_k\|_{\mathbf{L}^2}^2, \end{aligned} \quad (32)$$

where the last step is due to

$$g_N^{\nu, \kappa}(r) \geq \frac{\kappa}{\nu}(r - N).$$

Integrating both sides of (32) and using (30) and $\kappa \geq 1$, we obtain

$$\|\nabla \mathbf{u}_k(t)\|_{\mathbf{L}^2}^2 + \nu \int_0^t \|A\mathbf{u}_k\|_{\mathbf{L}^2}^2 ds \leq \frac{\kappa N}{\nu^2} \|\mathbf{u}_0\|_{\mathbf{L}^2}^2 + \|\nabla \mathbf{u}_0\|_{\mathbf{L}^2}^2, \quad \forall t \geq 0. \quad (33)$$

Now set

$$\mathbf{w}_{k,m}(t) := \mathbf{u}_k(t) - \mathbf{u}_m(t).$$

Then

$$\begin{aligned} \partial_t \mathbf{w}_{k,m} &= -A\mathbf{w}_{k,m} + B(\mathbf{u}_{k-1}, \mathbf{w}_{k,m}) + B(\mathbf{w}_{k-1,m-1}, \mathbf{v}_m) \\ &\quad - g_N^{\nu, \kappa}(\|\mathbf{u}_{k-1}\|_\infty^2) \mathbf{w}_{k,m} - [g_N^{\nu, \kappa}(\|\mathbf{u}_{k-1}\|_\infty^2) - g_N^{\nu, \kappa}(\|\mathbf{v}_{m-1}\|_\infty^2)] \mathbf{v}_m. \end{aligned}$$

Again, by [19, p.176, Lemma 1.2] and Lemma 4.1 we have

$$\begin{aligned} \frac{1}{2} \frac{d\|\mathbf{w}_{k,m}\|_{\mathbf{H}^1}^2}{dt} &= -\nu \|\mathbf{w}_{k,m}\|_{\mathbf{H}^2}^2 + \nu \|\mathbf{w}_{k,m}\|_{\mathbf{H}^1}^2 \\ &\quad + \langle B(\mathbf{u}_{k-1}, \mathbf{w}_{k,m}), (I+A)\mathbf{w}_{k,m} \rangle_{\mathbf{L}^2} \\ &\quad + \langle B(\mathbf{w}_{k-1,m-1}, \mathbf{v}_m), (I+A)\mathbf{w}_{k,m} \rangle_{\mathbf{L}^2} \\ &\quad - g_N^{\nu, \kappa}(\|\mathbf{u}_{k-1}\|_\infty^2) \langle \mathbf{w}_{k,m}, (I+A)\mathbf{w}_{k,m} \rangle_{\mathbf{L}^2} \\ &\quad - [g_N^{\nu, \kappa}(\|\mathbf{u}_{k-1}\|_\infty^2) - g_N^{\nu, \kappa}(\|\mathbf{v}_{m-1}\|_\infty^2)] \langle \mathbf{v}_m, \mathbf{w}_{k,m} \rangle_{\mathbf{H}^1} \\ &\leq -\frac{\nu}{4} \|\mathbf{w}_{k,m}\|_{\mathbf{H}^2}^2 + \frac{\nu}{8} \|\mathbf{w}_{k-1,m-1}\|_{\mathbf{H}^2}^2 + (\nu + \frac{\kappa N}{\nu}) \|\mathbf{w}_{k,m}\|_{\mathbf{H}^1}^2 \\ &\quad + \frac{C_\Omega}{\nu^3} \|\mathbf{w}_{k-1,m-1}\|_{\mathbf{H}^1}^2 \cdot ((1+4\kappa)\|\mathbf{u}_k\|_{\mathbf{H}^1}^2 + \kappa\|\mathbf{u}_m\|_{\mathbf{H}^1}^2)^2. \end{aligned} \quad (34)$$

Integrating this inequality and using (30) and (33), we get

$$\|\mathbf{w}_{k,m}(t)\|_{\mathbf{H}^1}^2 + \frac{\nu}{2} \int_0^t \|\mathbf{w}_{k,m}\|_{\mathbf{H}^2}^2 ds$$

$$\begin{aligned} &\leq \frac{\nu}{4} \int_0^t \|\mathbf{w}_{k-1,m-1}\|_{\mathbf{H}^2}^2 ds + C_{\nu,N} \int_0^t \|\mathbf{w}_{k,m}\|_{\mathbf{H}^1}^2 ds \\ &\quad + C_{\nu,N,\|\mathbf{u}_0\|_{\mathbf{H}^1}} \cdot \int_0^t \|\mathbf{w}_{k-1,m-1}\|_{\mathbf{H}^1}^2 ds. \end{aligned}$$

Set

$$h(t) := \overline{\lim}_{k,m \rightarrow \infty} \sup_{s \in [0,t]} \|\mathbf{w}_{k,m}(s)\|_{\mathbf{H}^1}^2$$

and

$$f(t) := \overline{\lim}_{k,m \rightarrow \infty} \int_0^t \|\mathbf{w}_{k,m}\|_{\mathbf{H}^2}^2 ds.$$

Then by (30), (33) and Fatou's lemma, we have

$$\frac{\nu}{2} f(t) \leq \frac{\nu}{4} f(t) + C_{\nu,N,\|\mathbf{u}_0\|_{\mathbf{H}^1}} \int_0^t h(s) ds$$

and

$$h(t) \leq \frac{\nu}{4} f(t) + C_{\nu,N,\|\mathbf{u}_0\|_{\mathbf{H}^1}} \int_0^t h(s) ds \leq 2C_{\nu,N,\|\mathbf{u}_0\|_{\mathbf{H}^1}} \int_0^t g(s) ds.$$

Hence, by Gronwall's inequality we have

$$h(t) = f(t) = 0, \quad \forall t \geq 0.$$

Thus, there exists a function $\mathbf{u} \in C([0, \infty); \mathbf{H}^1) \cap L_{loc}^2(\mathbb{R}_+; \mathbf{H}^2)$ such that for any $T > 0$

$$\overline{\lim}_{k \rightarrow \infty} \sup_{s \in [0,T]} \|\mathbf{u}_k - \mathbf{u}\|_{\mathbf{H}^1}^2 + \overline{\lim}_{k \rightarrow \infty} \int_0^T \|\mathbf{u}_k(s) - \mathbf{u}(s)\|_{\mathbf{H}^2}^2 ds = 0.$$

Lastly, taking limits $k \rightarrow \infty$ for

$$\mathbf{u}_k(t) = \mathbf{u}_0 - \int_0^t A\mathbf{u}_k ds + \int_0^t B(\mathbf{u}_{k-1}, \mathbf{u}_k) ds - \int_0^t g(\|\mathbf{u}_{k-1}\|_{\infty}^2) \mathbf{u}_k ds$$

and inequalities (30) and (33), we can see that $\mathbf{u}(t)$ satisfies (9), (10) and (11).

4.2. Proof of Decay Estimate (12). Following the method of Heywood [8], by the chain rule and [8, p.649 (14)], we have

$$\begin{aligned} \frac{d\|\nabla \mathbf{u}\|_{\mathbf{L}^2}^2}{dt} &= -2\nu \|A\mathbf{u}\|_{\mathbf{L}^2}^2 + 2\langle B(\mathbf{u}, \mathbf{u}), A\mathbf{u} \rangle_{\mathbf{L}^2} - 2g_N^{\nu,\kappa}(\|\mathbf{u}\|_{\infty}^2) \|\nabla \mathbf{u}\|_{\mathbf{L}^2}^2 \\ &= -\nu \|A\mathbf{u}\|_{\mathbf{L}^2}^2 + C_{\nu,\Omega} \|\nabla \mathbf{u}\|_{\mathbf{L}^2}^4 + C'_{\nu,\Omega} \|\nabla \mathbf{u}\|_{\mathbf{L}^2}^6. \end{aligned}$$

Note that

$$\int_0^{\infty} \|\nabla \mathbf{u}_k\|_{\mathbf{L}^2}^2 dt \leq \frac{\|\mathbf{u}_0\|_{\mathbf{L}^2}^2}{2\nu} =: \Lambda.$$

In Lemma 3.1, if we take $\phi(t) = \|\nabla \mathbf{u}(t)\|_{\mathbf{L}^2}^2$, $\beta = 1/(C'_{\nu,\Omega}\Lambda)$ and $\alpha = C_{\nu,\Omega} + 1/\Lambda$, then

$$\|\nabla \mathbf{u}(t)\|_{\mathbf{L}^2}^2 \leq \frac{e^{C_{\nu,\Omega}\Lambda+1} - 1}{(C_{\nu,\Omega} + 1/\Lambda)t} \leq C_3 t^{-1}$$

for $t \geq (C'_{\nu,\Omega}\Lambda)e^{C_{\nu,\Omega}\Lambda+1} = T^*$. Thus, (12) follows.

4.3. **Proof of Continuous Dependence (13).** Set

$$\mathbf{w}_{N,M}(t) := \mathbf{u}_N(t) - \mathbf{v}_M(t).$$

Once again, by the chain rule (cf. [19, p.176, Lemma 1.2]) we have

$$\begin{aligned} \frac{1}{2} \frac{d\|\mathbf{w}_{N,M}\|_{\mathbf{H}^1}^2}{dt} &= -\|\mathbf{w}_{N,M}\|_{\mathbf{H}^2}^2 + \|\mathbf{w}_{N,M}\|_{\mathbf{H}^1}^2 + \langle B(\mathbf{u}_N, \mathbf{w}_{N,M}), \mathbf{w}_{N,M} \rangle_{\mathbf{H}^1} \\ &\quad + \langle B(\mathbf{w}_{N,M}, \mathbf{v}_M), \mathbf{w}_{N,M} \rangle_{\mathbf{H}^1} - g_N^{\nu,\kappa}(\|\mathbf{u}_N\|_{\infty}^2) \|\mathbf{w}_{N,M}\|_{\mathbf{H}^1}^2 \\ &\quad - [g_N^{\nu,\kappa}(\|\mathbf{u}_N\|_{\infty}^2) - g_N^{\nu,\kappa}(\|\mathbf{v}_M\|_{\infty}^2)] \langle \mathbf{v}_M, \mathbf{w}_{N,M} \rangle_{\mathbf{H}^1} \\ &\quad - [g_N^{\nu,\kappa}(\|\mathbf{v}_M\|_{\infty}^2) - g_M^{\nu,\kappa}(\|\mathbf{v}_M\|_{\infty}^2)] \langle \mathbf{v}_M, \mathbf{w}_{N,M} \rangle_{\mathbf{H}^1}. \end{aligned}$$

Noting that

$$|g_N^{\nu,\kappa}(r) - g_M(r)| \leq \frac{\kappa}{\nu} |N - M|, \quad \forall r \geq 0, \quad N, M \geq 1,$$

as in the proof of existence, by Lemma 4.1 and Young's inequality we find that

$$\begin{aligned} \frac{1}{2} \frac{d\|\mathbf{w}_{N,M}\|_{\mathbf{H}^1}^2}{dt} &\leq -\frac{\nu}{8} \|\mathbf{w}_{N,M}\|_{\mathbf{H}^2}^2 + \left(\nu + \frac{\kappa N}{\nu}\right) \|\mathbf{w}_{N,M}\|_{\mathbf{H}^1}^2 \\ &\quad + \frac{C_{\Omega}}{\nu^3} \|\mathbf{w}_{N,M}\|_{\mathbf{H}^1}^2 \cdot ((1 + 4\kappa) \|\mathbf{u}_N\|_{\mathbf{H}^1}^2 + \kappa \|\mathbf{v}_M\|_{\mathbf{H}^1}^2)^2 \\ &\quad + \frac{\kappa}{\nu} |N - M| \cdot \|\mathbf{v}_M\|_{\mathbf{L}^2} \cdot \|\mathbf{w}_{N,M}\|_{\mathbf{H}^2} \\ (10)(11) \quad &\leq -\frac{\nu}{16} \|\mathbf{w}_{N,M}\|_{\mathbf{H}^2}^2 + C_{\nu,N,M,\mathbf{u}_0,\mathbf{v}_0} \|\mathbf{w}_{N,M}\|_{\mathbf{H}^1}^2 + C_{\nu,\mathbf{v}_0} |N - M|^2, \end{aligned}$$

where $C_{\nu,\mathbf{v}_0} = 4\kappa^2 \|\mathbf{v}_0\|_{\mathbf{L}^2}^2 / \nu^3$,

$$C_{\nu,N,M,\mathbf{u}_0,\mathbf{v}_0} = \left(\nu + \frac{\kappa N}{\nu}\right) + \frac{C_{\Omega}}{\nu^3} ((1 + 4\kappa) K_{\nu,N,\mathbf{u}_0} + \kappa K_{\nu,M,\mathbf{v}_0})^2$$

and

$$K_{\nu,N,\mathbf{u}_0} := \frac{\kappa N}{\nu^2} \|\mathbf{u}_0\|_{\mathbf{L}^2}^2 + \|\mathbf{u}_0\|_{\mathbf{H}^1}^2.$$

The estimate (13) now follows by Gronwall's inequality.

5. PROOF OF THEOREM 2.4

5.1. **Proof of Part (I).** Let $\mathbf{u}(t)$ be the unique strong solution of equation (8). By Duhamel's formula, we may write

$$\begin{aligned} \mathbf{u}(t) &= e^{-At} \mathbf{u}_0 + \int_0^t e^{-(t-s)A} B(\mathbf{u}, \mathbf{u}) ds - \int_0^t e^{-(t-s)A} g_N^{\nu,\kappa}(\|\mathbf{u}\|_{\infty}^2) \mathbf{u} ds \\ &=: \mathbf{w}_1(t) + \mathbf{w}_2(t) + \mathbf{w}_3(t). \end{aligned} \quad (36)$$

First of all, it is clear that $\mathbf{w}_1 \in C^{\infty}((0, T]; \mathbf{H}^2)$ and

$$\|A\mathbf{w}_1(t)\|_{\mathbf{L}^2} \leq \|A\mathbf{u}_0\|_{\mathbf{L}^2}. \quad (37)$$

For $\mathbf{w}_2(t)$, by (i) of Lemma 3.2 we have

$$\begin{aligned} \|A^{\frac{5}{8}} \mathbf{w}_2(t)\|_{\mathbf{L}^2} &\leq \int_0^t \|A^{\frac{7}{8}} e^{-(t-s)A} A^{-\frac{1}{4}} B(\mathbf{u}, \mathbf{u})\|_{\mathbf{L}^2} ds \\ &\leq \int_0^t \frac{\|A^{-\frac{1}{4}} B(\mathbf{u}, \mathbf{u})\|_{\mathbf{L}^2}^2}{(t-s)^{\frac{7}{8}}} ds \stackrel{(21)}{\leq} C_{\Omega} \cdot \int_0^t \frac{\|A^{\frac{1}{2}} \mathbf{u}\|_{\mathbf{L}^2}^2}{(t-s)^{\frac{7}{8}}} ds \\ (11) \quad &\leq C_{\Omega} K_{\nu,N,\mathbf{u}_0} \cdot t^{\frac{1}{8}}, \end{aligned} \quad (38)$$

where

$$K_{\nu, N, \mathbf{u}_0} := \frac{\kappa N}{\nu^2} \|\mathbf{u}_0\|^2 + \|\nabla \mathbf{u}_0\|^2.$$

For $\mathbf{w}_3(t)$, recalling (6) and by Lemma 3.2, we have for $\alpha \in [1/2, 1)$

$$\begin{aligned} \|A^\alpha \mathbf{w}_3(t)\|_{\mathbf{L}^2} &\leq \int_0^t g_N^{\nu, \kappa} (\|\mathbf{u}\|_\infty^2) \cdot \|A^\alpha e^{-(t-s)A} \mathbf{u}\|_{\mathbf{L}^2} ds \\ &\leq \int_0^t \|\mathbf{u}\|_\infty^2 \cdot \frac{1}{(t-s)^{\alpha-\frac{1}{2}}} \cdot \|A^{\frac{1}{2}} \mathbf{u}\|_{\mathbf{L}^2} ds \\ &\stackrel{(21)}{\leq} C_\Omega \int_0^t \|\mathbf{u}\|_{\mathbf{H}^2} \cdot \|\nabla \mathbf{u}\|_{\mathbf{L}^2}^2 \cdot \frac{1}{(t-s)^{\alpha-\frac{1}{2}}} ds \\ &\leq C_\Omega \int_0^t \|A\mathbf{u}\|_{\mathbf{L}^2} \cdot \|\nabla \mathbf{u}\|_{\mathbf{L}^2}^2 \cdot \frac{1}{(t-s)^{\alpha-\frac{1}{2}}} ds \\ &\quad + C_\Omega \int_0^t \|\mathbf{u}\|_{\mathbf{L}^2} \cdot \|\nabla \mathbf{u}\|_{\mathbf{L}^2}^2 \cdot \frac{1}{(t-s)^{\alpha-\frac{1}{2}}} ds \\ &=: I_1 + I_2. \end{aligned}$$

By (10), (11) and Hölder's inequality we have

$$I_1 \leq C_\Omega K_{\nu, N, \mathbf{u}_0} t^{1-\alpha} \left(\int_0^t \|A\mathbf{u}(s)\|_{\mathbf{L}^2}^2 ds \right)^{1/2} \leq C_\Omega K_{\nu, N, \mathbf{u}_0}^{3/2} \cdot t^{1-\alpha}$$

and

$$I_2 \leq C_\Omega \|\mathbf{u}_0\|_{\mathbf{L}^2} K_{\nu, N, \mathbf{u}_0}^{\frac{1}{2}} t^{1-\alpha} \left(\int_0^t \|\nabla \mathbf{u}(s)\|_{\mathbf{L}^2}^2 ds \right)^{1/2} \leq C_\Omega \|\mathbf{u}_0\|_{\mathbf{L}^2}^2 K_{\nu, N, \mathbf{u}_0}^{1/2} \cdot t^{1-\alpha}.$$

Hence

$$\|A^\alpha \mathbf{w}_3(t)\|_{\mathbf{L}^2} \leq C_\Omega \left(\|\mathbf{u}_0\|_{\mathbf{L}^2}^2 K_{\nu, N, \mathbf{u}_0}^{1/2} + K_{\nu, N, \mathbf{u}_0}^{3/2} \right) \cdot t^{1-\alpha}. \quad (39)$$

Combining (36), (37), (38) and (39), we find that

$$\begin{aligned} \|A^{\frac{5}{8}} \mathbf{u}(t)\|_{\mathbf{L}^2} &\leq \|A^{\frac{5}{8}} \mathbf{u}_0\|_{\mathbf{L}^2} + C_\Omega K_{\nu, N, \mathbf{u}_0} \cdot t^{\frac{1}{8}} \\ &\quad + C_\Omega \left(\|\mathbf{u}_0\|_{\mathbf{L}^2}^2 K_{\nu, N, \mathbf{u}_0}^{1/2} + K_{\nu, N, \mathbf{u}_0}^{3/2} \right) \cdot t^{3/8} \\ &=: M_0(t, \nu, N, \mathbf{u}_0) \end{aligned}$$

and by (25) and (10)

$$\begin{aligned} \|B(\mathbf{u}(t), \mathbf{u}(t))\|_{\mathbf{L}^2} &\leq C_\Omega (\|A^{\frac{5}{8}} \mathbf{u}(t)\|_{\mathbf{L}^2}^2 + \|\mathbf{u}(t)\|_{\mathbf{L}^2}^2) \\ &\leq C_\Omega \cdot (M_0(t, N, \mathbf{u}_0))^2 + \|\mathbf{u}_0\|_{\mathbf{L}^2}^2. \end{aligned}$$

By (1⁰) of Lemma 3.7, we have for any $\frac{3}{4} < \gamma < 1$

$$\begin{aligned} \|A^\gamma \mathbf{u}(t)\|_{\mathbf{L}^2} &\leq \|A^\gamma \mathbf{u}_0\|_{\mathbf{L}^2} + C_\Omega \cdot (M_0(t, \nu, N, \mathbf{u}_0))^2 + \|\mathbf{u}_0\|_{\mathbf{L}^2}^2 \cdot t^{1-\gamma} \\ &\quad + C_\Omega \left(\|\mathbf{u}_0\|_{\mathbf{L}^2}^2 K_{\nu, N, \mathbf{u}_0}^{1/2} + K_{\nu, N, \mathbf{u}_0}^{3/2} \right) \cdot t^{1-\gamma}, \end{aligned} \quad (40)$$

which then yields the estimate (14) by (22).

5.2. **Proof of Part (II).** In this subsection, we assume $\Omega = \mathbb{R}^3$ or Ω is a bounded uniform C^4 -domain. Our proof is concentrated on the case of bounded domain. Clearly, it also works for $\Omega = \mathbb{R}^3$.

Below, fix $T > 0$ and set

$$\mathbf{f}(s) := B(\mathbf{u}(s), \mathbf{u}(s)) - g_N^{\nu, \kappa}(\|\mathbf{u}(s)\|_\infty^2) \mathbf{u}(s).$$

Then by Lemma 3.6 and (22), (40)

$$[0, T] \ni s \mapsto \mathbf{f}(s) \in \mathbf{H}^0 = \mathbf{L}_\sigma^2(\Omega) \text{ is continuous.}$$

By (i) of Lemma 3.7, we have for any $\beta \in (0, 1)$ and $0 < \theta < 1$

$$\mathbf{u} \in C^\theta([0, T], \mathbf{H}^0) \cap C([0, T]; \mathbf{H}^{2\beta}). \quad (41)$$

Thus, by Lemma 3.6 and (22), for any $\frac{3}{4} < \gamma < \beta \leq 1$, there are constants $C_1, C_2, C_3 > 0$ such that for all $t, s \in [0, T]$

$$\begin{aligned} \|\mathbf{f}(t) - \mathbf{f}(s)\|_{\mathbf{L}^2} &\leq C_1 \cdot \|A^\beta(\mathbf{u}(t) - \mathbf{u}(s))\|_{\mathbf{L}^2}^{\frac{\gamma}{\beta}} \cdot \|\mathbf{u}(t) - \mathbf{u}(s)\|_{\mathbf{L}^2}^{1-\frac{\gamma}{\beta}} \\ &\quad + C_2 \cdot \|A^\beta(\mathbf{u}(t) - \mathbf{u}(s))\|_{\mathbf{L}^2}^{\frac{1}{2\beta}} \cdot \|\mathbf{u}(t) - \mathbf{u}(s)\|_{\mathbf{L}^2}^{1-\frac{1}{2\beta}} \\ &\quad + C_3 \cdot \|\mathbf{u}(t) - \mathbf{u}(s)\|_{\mathbf{L}^2}. \end{aligned} \quad (42)$$

Choosing β close to 1 and γ close to $\frac{3}{4}$ and using (40) and (41), we find that for any $0 < \alpha < \frac{1}{4}$

$$\mathbf{f} \in C^\alpha([0, T], \mathbf{H}^0).$$

Thus, by (ii) of Lemma 3.7 and (36) we have, for any $0 < \alpha < \frac{1}{4}$

$$A\mathbf{u} \in C^\alpha((0, T], \mathbf{H}^0), \quad \partial_t \mathbf{u} \in C((0, T], \mathbf{H}^{2\alpha})$$

Using induction and (42) with $\beta = 1$ as well as (41), one finds that for any $n \in \mathbb{N}$ and $0 < \alpha < 1 - \left(\frac{3}{4}\right)^{n+1}$

$$\mathbf{f} \in C^\alpha((0, T], \mathbf{H}^0)$$

and

$$A\mathbf{u} \in C^\alpha((0, T], \mathbf{H}^0), \quad \partial_t \mathbf{u} \in C((0, T], \mathbf{H}^{2\alpha}).$$

In particular, by $\mathbf{H}^\alpha \subset \mathbf{W}^{\alpha, 2}(\Omega)$ for $\alpha \in [0, 2]$ we have

$$\mathbf{u} \in C^\alpha((0, T], \mathbf{W}^{2, 2}(\Omega)), \quad \partial_t \mathbf{u} \in C((0, T], \mathbf{W}^{9/5, 2}(\Omega)). \quad (43)$$

Set

$$\mathbf{b}(t) := (\mathbf{u}(t) \cdot \nabla) \mathbf{u}(t) + g_N^{\nu, \kappa}(\|\mathbf{u}(t)\|_\infty^2) \mathbf{u}(t).$$

As in the proof of Lemma 3.6, it is not hard to verify by (43) that

$$\mathbf{b}(t) \in C((0, T]; \mathbf{W}^{1, 2}(\Omega)). \quad (44)$$

Consider the Stokes equation:

$$\begin{cases} \nu \Delta \mathbf{u} + \nabla P = \partial_t \mathbf{u} + \mathbf{b} & \text{in } \Omega, \\ \mathbf{d}\mathbf{u} = 0 & \text{in } \Omega, \quad \mathbf{u}|_{\partial\Omega} = 0. \end{cases}$$

By (43), (44) and Lemma 3.8 with $\alpha = 1$, we have

$$\mathbf{u}(t) \in C((0, T], \mathbf{W}^{3, 2}(\Omega)).$$

As above, a simple calculation shows that

$$\mathbf{b}(t) \in C((0, T]; \mathbf{W}^{2, 2}(\Omega)). \quad (45)$$

By (43), (45) and Lemma 3.8 with $\alpha = \frac{9}{5}$ again, we further have

$$\mathbf{u}(t) \in C((0, T], \mathbf{W}^{\frac{19}{5}, 2}(\Omega)) \quad (46)$$

and

$$P \in C((0, T], W^{\frac{14}{5}, 2}(\Omega)). \quad (47)$$

By (43), (46), (47) and the Sobolev embedding theorem (cf. [21, Theorem 4.6.1]), we finally obtain that

$$\partial_t \mathbf{u}, \partial_i \mathbf{u}, \partial_i \partial_j \mathbf{u}, P \in C((0, T] \times \bar{\Omega}; \mathbb{R}^3)$$

and (15) holds.

5.3. Proof of Part (III). In this subsection, we assume $\Omega = \mathbb{R}^3$.

Lemma 5.1. *For fixed $q \geq 2$ and $r \geq 1$, there exists $\kappa := \kappa(q) := Cq^4$, where C is a universal constant, such that for any $N \geq 1$ and $t \geq 0$*

$$\|\mathbf{u}_N^\kappa(t)\|_{\mathbf{L}^q}^r \leq \|\mathbf{u}_0\|_{\mathbf{L}^q}^r + \frac{r\kappa}{\nu} \cdot N \int_0^t \|\mathbf{u}_N^\kappa\|_{\mathbf{L}^q}^r ds \quad (48)$$

and

$$\int_0^t \|\mathbf{u}_N^\kappa\|_{\infty}^2 \cdot \|\mathbf{u}_N^\kappa\|_{\mathbf{L}^q}^r ds \leq \frac{2\nu}{r\kappa} \|\mathbf{u}_0\|_{\mathbf{L}^q}^r + 2N \int_0^t \|\mathbf{u}_N^\kappa\|_{\mathbf{L}^q}^r ds. \quad (49)$$

Proof. Let $\mathbf{u} := \mathbf{u}_N^\kappa$. Taking the scalar product for both sides of equation (15) with $q|\mathbf{u}|^{q-2}\mathbf{u}$, and then integrating over \mathbb{R}^3 , we find by the integration by parts formula

$$\begin{aligned} \frac{d\|\mathbf{u}\|_{\mathbf{L}^q}^q}{dt} &= -q\nu \|\nabla \mathbf{u}\|_{\mathbf{L}^2} \|\mathbf{u}\|_{\mathbf{L}^2}^{(q-2)/2} \|\mathbf{u}\|_{\mathbf{L}^2}^2 - \frac{4(q-2)\nu}{q} \|\nabla |\mathbf{u}|^{q/2}\|_{\mathbf{L}^2}^2 \\ &\quad + q \langle \nabla P, |\mathbf{u}|^{q-2}\mathbf{u} \rangle_{\mathbf{L}^2} - q \cdot g_N^{\nu, \kappa} (\|\mathbf{u}\|_{\infty}^2) \|\mathbf{u}\|_{\mathbf{L}^q}^q, \end{aligned}$$

where we have used that

$$q \langle (\mathbf{u} \cdot \nabla) \mathbf{u}, |\mathbf{u}|^{q-2}\mathbf{u} \rangle_{\mathbf{L}^2} = \langle \mathbf{u}, \nabla |\mathbf{u}|^{q/2} \rangle_{\mathbf{L}^2} = 0.$$

Let f be an increasing smooth function on $[0, \infty)$. We further have

$$\begin{aligned} \frac{df(\|\mathbf{u}\|_{\mathbf{L}^q}^q)}{dt} &= f'(\|\mathbf{u}\|_{\mathbf{L}^q}^q) \left[-q\nu \|\nabla \mathbf{u}\|_{\mathbf{L}^2} \|\mathbf{u}\|_{\mathbf{L}^2}^{(q-2)/2} \|\mathbf{u}\|_{\mathbf{L}^2}^2 - \frac{4(q-2)\nu}{q} \|\nabla |\mathbf{u}|^{q/2}\|_{\mathbf{L}^2}^2 \right. \\ &\quad \left. + q \langle \nabla P, |\mathbf{u}|^{q-2}\mathbf{u} \rangle_{\mathbf{L}^2} - q \cdot g_N^{\nu, \kappa} (\|\mathbf{u}\|_{\infty}^2) \|\mathbf{u}\|_{\mathbf{L}^q}^q \right]. \quad (50) \end{aligned}$$

On the other hand, taking the divergence for equation (15) we have

$$\Delta P = \operatorname{div}[(\mathbf{u} \cdot \nabla) \mathbf{u}],$$

which gives

$$P = -(-\Delta)^{-1} \operatorname{div}[(\mathbf{u} \cdot \nabla) \mathbf{u}] = -(-\Delta)^{-1} \partial_j \partial_i (u^i \cdot u^j).$$

So, by the Calderón-Zygmund inequality we get for any $\gamma \geq 2$ (cf. [18])

$$\|P\|_{L^\gamma} \leq C_1 \cdot \gamma \cdot \|\mathbf{u}\|_{\mathbf{L}^{2\gamma}}^2, \quad (51)$$

Here and below, $C_i, i = 1, 2, 3$ are universal constants. Thus, by Young's inequality and Hölder's inequality we have

$$\begin{aligned} q \langle \nabla P, |\mathbf{u}|^{q-2}\mathbf{u} \rangle_{\mathbf{L}^2} &= q \langle P, |\mathbf{u}|^{q-2} \operatorname{div} \mathbf{u} \rangle_{\mathbf{L}^2} + q \langle P, \nabla |\mathbf{u}|^{q-2} \cdot \mathbf{u} \rangle_{\mathbf{L}^2} \\ &\leq q\nu \|\nabla \mathbf{u}\|_{\mathbf{L}^2} \cdot \|\mathbf{u}\|_{\mathbf{L}^2}^{(q-2)/2} \|\mathbf{u}\|_{\mathbf{L}^2}^2 + \frac{C_2 q^3}{\nu} \|\nabla P\|_{\mathbf{L}^2} \cdot \|\mathbf{u}\|_{\mathbf{L}^2}^{(q-2)/2} \|\mathbf{u}\|_{\mathbf{L}^2}^2 \end{aligned}$$

$$\begin{aligned}
&\leq q\nu\|\nabla\mathbf{u}\cdot|\mathbf{u}|^{(q-2)/2}\|_{\mathbf{L}^2}^2 + \frac{C_2q^3}{\nu}\|\mathbf{u}\|_{\mathbf{L}^{q+2}}^{q-2}\cdot\|P\|_{\mathbf{L}^{(q+2)/2}}^2 \\
(51) \quad &\leq q\nu\|\nabla\mathbf{u}\cdot|\mathbf{u}|^{(q-2)/2}\|_{\mathbf{L}^2}^2 + \frac{C_3q^5}{\nu}\|\mathbf{u}\|_{\mathbf{L}^{q+2}}^{q+2} \\
&\leq q\nu\|\nabla\mathbf{u}\cdot|\mathbf{u}|^{(q-2)/2}\|_{\mathbf{L}^2}^2 + \frac{C_3q^5}{\nu}\|\mathbf{u}\|_{\infty}^2\cdot\|\mathbf{u}\|_{\mathbf{L}^q}^q.
\end{aligned}$$

Substituting this estimate into (50), we get

$$\begin{aligned}
&\frac{df(\|\mathbf{u}\|_{\mathbf{L}^q}^q)}{dt} + \frac{4(q-2)\nu}{q}f'(\|\mathbf{u}\|_{\mathbf{L}^q}^q)\cdot\|\nabla|\mathbf{u}|^{q/2}\|_{\mathbf{L}^2}^2 \\
&\leq f'(\|\mathbf{u}\|_{\mathbf{L}^q}^q)\cdot\left[\frac{C_3q^5}{\nu}\cdot\|\mathbf{u}\|_{\infty}^2\cdot\|\mathbf{u}\|_{\mathbf{L}^q}^q - q\cdot g_N^{\nu,\kappa}(\|\mathbf{u}\|_{\infty}^2)\|\mathbf{u}\|_{\mathbf{L}^q}^q\right].
\end{aligned}$$

Now noticing that

$$g_N^{\nu,\kappa}(r) \geq \frac{\kappa(r-N)}{\nu}, \quad r \geq 0,$$

we find that if

$$\kappa = 2C_3q^4, \quad (52)$$

then

$$\begin{aligned}
&\frac{df(\|\mathbf{u}\|_{\mathbf{L}^q}^q)}{dt} + \frac{4(q-2)\nu}{q}f'(\|\mathbf{u}\|_{\mathbf{L}^q}^q)\cdot\|\nabla|\mathbf{u}|^{q/2}\|_{\mathbf{L}^2}^2 \\
&\leq f'(\|\mathbf{u}\|_{\mathbf{L}^q}^q)\cdot\left[\frac{q\kappa}{\nu}\cdot N\cdot\|\mathbf{u}\|_{\mathbf{L}^q}^q - \frac{q\kappa}{2\nu}\cdot\|\mathbf{u}\|_{\infty}^2\cdot\|\mathbf{u}\|_{\mathbf{L}^q}^q\right].
\end{aligned} \quad (53)$$

Lastly, taking $f_{\epsilon}(x) := (\epsilon + x)^{r/q}$ in (53), then integrating with respect to t and letting $\epsilon \downarrow 0$ yield (48) and (49). \square

Lemma 5.2. *Fix $r_0 \geq 1$ and $q_0 \geq 2$. Let $\mathbf{u}_0 \in \mathbf{H}_2^2$ and set $N_0 := C\|\mathbf{u}_0\|_{\mathbf{H}_2^2}^2$ for some universal constant C . There exists $n_0 := n_0(\nu, N_0, q_0, r_0)$ large enough such that for all $n \geq n_0$, $N \geq N_0 \vee 1$ and $t \geq 0$*

$$\|\mathbf{u}_N^{\kappa_n}(t)\|_{\mathbf{L}^{q_0+2q_0n/r_0}} \leq 2N^{\frac{n}{2n+r_0}} + 2N^{\frac{n+1}{2n+r_0}} \left[\int_0^t \|\mathbf{u}_N^{\kappa_n}\|_{\mathbf{L}^{q_0}}^{r_0} ds \right]^{\frac{1}{2n+r_0}}, \quad (54)$$

where $\kappa_n = 2C_3 \cdot [(2n+r_0)q_0/r_0]^4$ (see (52)). In particular, there is an $n_0 := n_0(\nu, N_0)$ large enough such that for all $n \geq n_0$, $N \geq N_0 \vee 1$ and $t \geq 0$

$$\|\mathbf{u}_N^{\kappa_n}(t)\|_{\mathbf{L}^{6(n+1)}} \leq 3N^{\frac{1}{2}}. \quad (55)$$

Proof. Let $\mathbf{u} := \mathbf{u}_N^{\kappa_n}$. First of all, by the Gagliardo-Nirenberg inequality (23), there is a universal constant $C_0 \geq 1$ such that for any $q \geq 2$

$$\|\mathbf{u}_0\|_{\mathbf{L}^q} \leq \|\mathbf{u}_0\|_{\infty}^{1-2/q} \cdot \|\mathbf{u}_0\|_{\mathbf{L}^2}^{2/q} \leq C_0\|\mathbf{u}_0\|_{\mathbf{H}_2^2} =: N_0^{1/2}. \quad (56)$$

Define

$$q_n := q_{n-1} + 2q_0/r_0 = (2n+r_0)q_0/r_0$$

and

$$r_n := r_0q_n/q_0 = 2n+r_0.$$

Then we have

$$\int_0^t \|\mathbf{u}\|_{\mathbf{L}^{q_{n+1}}}^{r_{n+1}} ds \leq \int_0^t \|\mathbf{u}\|_{\infty}^2 \cdot \|\mathbf{u}\|_{\mathbf{L}^{q_n}}^{r_n} ds$$

$$\begin{aligned}
& \text{(by (49))} \leq \frac{2\nu}{r_n \cdot \kappa(q_n)} \|\mathbf{u}_0\|_{\mathbf{L}^{q_n}}^{r_n} + 2N \int_0^t \|\mathbf{u}\|_{\mathbf{L}^{q_n}}^{r_n} ds \\
& \text{(by } r_n \kappa(q_n) \geq 2) \leq \nu \|\mathbf{u}_0\|_{\mathbf{L}^{q_n}}^{r_n} + 2N \int_0^t \|\mathbf{u}\|_{\mathbf{L}^{q_n}}^{r_n} ds \\
& \text{(by iterating)} \leq \nu \sum_{k=0}^n (2N)^k \|\mathbf{u}_0\|_{\mathbf{L}^{q_{n-k}}}^{r_{n-k}} + (2N)^{n+1} \int_0^t \|\mathbf{u}\|_{\mathbf{L}^{q_0}}^{r_0} ds \\
& \text{(by (56))} \leq \nu \sum_{k=0}^n (2N)^k \cdot N_0^{r_{n-k}/2} + (2N)^{n+1} \int_0^t \|\mathbf{u}\|_{\mathbf{L}^{q_0}}^{r_0} ds \\
& \text{(by } r_n = 2n + r_0) = \frac{\nu((2N)^{n+1} N_0^{r_0/2} - N_0^{n+1+r_0/2})}{2N - N_0} \\
& \quad + (2N)^{n+1} \int_0^t \|\mathbf{u}\|_{\mathbf{L}^{q_0}}^{r_0} ds \\
& \text{(by } N \geq N_0 \vee 1) \leq \frac{\nu N_0^{r_0/2} (2N)^{n+1}}{N} + (2N)^{n+1} \int_0^t \|\mathbf{u}\|_{\mathbf{L}^{q_0}}^{r_0} ds.
\end{aligned}$$

Hence, by (48)

$$\begin{aligned}
\|\mathbf{u}(t)\|_{\mathbf{L}^{q_n}}^{r_n} & \leq \|\mathbf{u}_0\|_{\mathbf{L}^{q_n}}^{r_n} + r_n \kappa(q_n) \cdot N_0^{r_0/2} (2N)^n \\
& \quad + \frac{r_n \kappa(q_n) N}{\nu} \cdot (2N)^n \int_0^t \|\mathbf{u}\|_{\mathbf{L}^{q_0}}^{r_0} ds.
\end{aligned}$$

Now taking the root $1/r_n$ and noting that

$$\lim_{n \rightarrow \infty} (r_n \kappa(q_n))^{1/r_n} \stackrel{(52)}{=} \lim_{n \rightarrow \infty} (2C_3 r_n q_n^4)^{1/r_n} = 1,$$

we obtain the desired estimate (54).

As for (55), it follows by taking $r_0 = 2$ and $q_0 = 6$ in (54) and noting that

$$\int_0^t \|\mathbf{u}\|_{\mathbf{L}^6}^2 ds \stackrel{(18)}{\leq} C \int_0^t \|\nabla \mathbf{u}\|_{\mathbf{L}^2}^2 ds \stackrel{(10)}{\leq} \frac{C}{2\nu} \|\mathbf{u}_0\|_{\mathbf{L}^2}^2.$$

The proof is complete. \square

We are now in a position to give

Proof of (16): By (20), (40) and (10) we have

$$\|\mathbf{u}_N^\kappa(t)\|_{\mathbf{W}^{1,4}} \leq C(\|A^{7/8} \mathbf{u}_N^\kappa(t)\|_{\mathbf{L}^2} + \|\mathbf{u}_N^\kappa(t)\|_{\mathbf{L}^2}) \tag{57}$$

$$\begin{aligned}
& \leq C\|A^{7/8} \mathbf{u}_0\|_{\mathbf{L}^2} + C \cdot (M_0(t, \nu, N, \mathbf{u}_0)^2 + \|\mathbf{u}_0\|_{\mathbf{L}^2}^2) \cdot t^{1/8} \\
& \quad + C_\Omega \left(\|\mathbf{u}_0\|_{\mathbf{L}^2}^2 K_{\nu, N, \mathbf{u}_0}^{1/2} + K_{\nu, N, \mathbf{u}_0}^{3/2} \right) \cdot t^{1/8} + C\|\mathbf{u}_0\|_{\mathbf{L}^2}. \tag{58}
\end{aligned}$$

By the Gagliardo-Nirenberg inequality (23) and (55), we have

$$\|\mathbf{u}_N^\kappa(t)\|_\infty \leq C \|\mathbf{u}_N^\kappa(t)\|_{\mathbf{W}^{1,4}}^{\frac{2}{n+3}} \cdot \|\mathbf{u}_N^\kappa(t)\|_{\mathbf{L}^{6(n+1)}}^{\frac{n+1}{n+3}}.$$

Letting n be large enough, the estimate (16) follows from (55) and (58).

6. PROOF OF THEOREM 2.8

We need the following simple lemma. For the reader's convenience, a short proof is provided here.

Lemma 6.1. *Let $(\mathbb{X}, \|\cdot\|_{\mathbb{X}})$ be a uniformly convex Banach space and $K \subset \mathbb{X}$. Then K is relatively compact in \mathbb{X} if and only if there exists a family of finite dimensional subspaces $\{\mathbb{X}_n, n \in \mathbb{N}\}$ of \mathbb{X} such that*

$$\sup_{n \in \mathbb{N}} \sup_{x \in K} \|\Pi_n x\|_{\mathbb{X}} < +\infty \quad (59)$$

and

$$\lim_{n \rightarrow \infty} \sup_{x \in K} \|(I - \Pi_n)x\|_{\mathbb{X}} = 0, \quad (60)$$

where Π_n is the projection operator from \mathbb{X} to \mathbb{X}_n , i.e., $\Pi_n x \in \mathbb{X}_n$ is the unique element such that

$$\|x - \Pi_n x\|_{\mathbb{X}} = \inf_{y \in \mathbb{X}_n} \|x - y\|_{\mathbb{X}}.$$

Proof. (“Only if”:) Let K be relatively compact in \mathbb{X} . For any $n \in \mathbb{N}$, there are finite points $\{x_1, \dots, x_m\} \subset K$ such that

$$K \subset \cup_{i=1}^m B_{1/n}(x_i),$$

where $B_{1/n}(x_k)$ denotes the ball in \mathbb{X} with center x_k and radius $1/n$. Now put

$$\mathbb{X}_n := \text{span}\{x_1, \dots, x_m\}.$$

It is easy to see that the corresponding Π_n satisfy (59) and (60).

(“If”:) Fix any sequence $\{x_k, k \in \mathbb{N}\} \subset K$. It suffices to prove that there is a subsequence x_{k_l} such that x_{k_l} converges to some point $x \in \mathbb{X}$. For any $n \in \mathbb{N}$, since \mathbb{X}_n is finite dimensional, by (59) there is a subsequence $x_{k_l^{(n)}}$ and $y_n \in \mathbb{X}_n$ such that $\Pi_n x_{k_l^{(n)}}$ converges to y_n as $l \rightarrow \infty$. By the diagonalization method, one can find a common subsequence x_{k_l} such that for any $n \in \mathbb{N}$

$$\lim_{l \rightarrow \infty} \|\Pi_n x_{k_l} - y_n\|_{\mathbb{X}} = 0.$$

Noting that

$$\|y_n - y_m\|_{\mathbb{X}} \leq \|\Pi_n x_{k_l} - y_n\|_{\mathbb{X}} + \|\Pi_n x_{k_l} - y_n\|_{\mathbb{X}} + \|\Pi_n x_{k_l} - P_m y_n\|_{\mathbb{X}},$$

we have by (60) that $\{y_n, n \in \mathbb{N}\}$ is a Cauchy sequence in \mathbb{X} . So, there is an $x \in \mathbb{X}$ such that y_n converges to x in \mathbb{X} . By (60) again, it is easy to find that x_{k_l} converges to x in \mathbb{X} . The proof is complete. \square

Since we have assumed that Ω is a bounded domain in Theorem 2.8, $\mathbf{H}^1 = \mathbf{W}_{0,\sigma}^{1,2}(\Omega)$ is compactly embedded in $\mathbf{H}^0 = \mathbf{L}_{\sigma}^2(\Omega)$. Let $0 < \lambda_1 \leq \lambda_2 \leq \dots \leq \lambda_k \rightarrow \infty$ be the eigenvalues of A , and $\mathcal{E} := \{\mathbf{e}_k; k \in \mathbb{N}\}$ the corresponding orthonormal eigenvectors, i.e.,

$$A\mathbf{e}_k = \lambda_k \mathbf{e}_k, \quad \langle \mathbf{e}_k, \mathbf{e}_j \rangle_{\mathbf{L}^2} = \delta_{kj}. \quad (61)$$

From this, one knows that the following Poincaré inequality holds:

$$\sqrt{\lambda_1} \cdot \|\mathbf{u}\|_{\mathbf{L}^2} \leq \|A^{\frac{1}{2}} \mathbf{u}\|_{\mathbf{L}^2}, \quad \forall \mathbf{u} \in \mathbf{H}^1. \quad (62)$$

Moreover, by (21) and (62) we have

$$\|\mathbf{u}\|_{\infty}^2 \leq C_0 \cdot \|A\mathbf{u}\|_{\mathbf{L}^2} \cdot \|\nabla \mathbf{u}\|_{\mathbf{L}^2}. \quad (63)$$

We have:

Lemma 6.2. For $\epsilon > 0$, let $\mathcal{B}_\epsilon := \{\mathbf{v} \in \mathbf{H}^1 : \|\mathbf{v}\|_{\mathbf{H}^1} \leq \epsilon\}$. Then \mathcal{B}_ϵ is an absorbing set of $\{S(t); t \geq 0\}$, i.e., for any bounded set $\mathcal{U} \subset \mathbf{H}^1$, there exists $t_{\mathcal{U}} > 0$ such that for any $t > t_{\mathcal{U}}$

$$S(t)\mathcal{U} \subset \mathcal{B}_\epsilon.$$

Proof. By the chain rule and (62), we have

$$d\|\mathbf{u}\|_{\mathbf{L}^2}^2/dt = -2\nu\|\nabla\mathbf{u}\|_{\mathbf{L}^2}^2 - 2g_N^{\nu,\kappa}(\|\mathbf{u}\|_{\infty}^2)\|\mathbf{u}\|_{\mathbf{L}^2}^2 \leq -2\nu\lambda_1\|\mathbf{u}\|_{\mathbf{L}^2}^2,$$

which implies

$$\|\mathbf{u}(t)\|_{\mathbf{L}^2}^2 \leq \|\mathbf{u}_0\|_{\mathbf{L}^2}^2 e^{-2\nu\lambda_1 t}. \quad (64)$$

As the calculation of (32), by Young's inequality we have

$$\begin{aligned} \frac{d\|A^{\frac{1}{2}}\mathbf{u}\|_{\mathbf{L}^2}^2}{dt} &\leq -\nu\|A\mathbf{u}\|_{\mathbf{L}^2}^2 + \frac{2\kappa N}{\nu}\|A^{\frac{1}{2}}\mathbf{u}\|_{\mathbf{L}^2}^2 \\ (63) \quad &\leq -\nu\|A\mathbf{u}\|_{\mathbf{L}^2}^2 + C_{\nu,\kappa} \cdot N \cdot \|A\mathbf{u}\|_{\mathbf{L}^2} \cdot \|\mathbf{u}\|_{\mathbf{L}^2} \\ &\leq -\frac{\nu}{2}\|A\mathbf{u}\|_{\mathbf{L}^2}^2 + C_{\nu,\kappa} \cdot N^2 \cdot \|\mathbf{u}\|_{\mathbf{L}^2}^2 \\ (62) \quad &\leq -\frac{\nu\lambda_1}{2} \cdot \|A^{\frac{1}{2}}\mathbf{u}\|_{\mathbf{L}^2}^2 + C_{\nu,\kappa} \cdot N^2 \cdot \|\mathbf{u}_0\|_{\mathbf{L}^2}^2 \cdot e^{-2\nu\lambda_1 t}. \end{aligned}$$

Integrating this differential inequality yields that

$$\|A^{\frac{1}{2}}\mathbf{u}(t)\|_{\mathbf{L}^2}^2 \leq e^{-\nu\lambda_1 t/2} \left[\|A^{\frac{1}{2}}\mathbf{u}_0\|_{\mathbf{L}^2}^2 + C_{\nu,\kappa} \cdot N^2 \cdot \|\mathbf{u}_0\|_{\mathbf{L}^2}^2 \cdot (1 - e^{-3\nu\lambda_1 t/2})/(\nu\lambda_1) \right]. \quad (65)$$

Hence, for any $\mathbf{u}_0 \in \mathbf{H}^1$

$$\lim_{t \rightarrow \infty} \|S(t)\mathbf{u}_0\|_{\mathbf{H}^1}^2 = \lim_{t \rightarrow \infty} \|\mathbf{u}(t)\|_{\mathbf{H}^1}^2 = 0.$$

The result follows. \square

We now use Lemma 6.1 to prove the following compactness result.

Lemma 6.3. For any $t > 0$, $S(t)$ is a compact operator from \mathbf{H}^1 to \mathbf{H}^1 , i.e., maps a bounded set in \mathbf{H}^1 into a relatively compact in \mathbf{H}^1 .

Proof. Let $\mathcal{U} \subset \mathbf{H}^1$ be a bounded set. Let Π_n be the projection operator from \mathbf{H}^1 to $\text{span}\{\mathbf{e}_k : k = 1, \dots, n\}$, i.e.,

$$\Pi_n \mathbf{v} := \sum_{k=1}^n \langle \mathbf{v}, \mathbf{e}_k \rangle_{\mathbf{L}^2} \mathbf{e}_k. \quad (66)$$

First of all, by (65) we have

$$\sup_{n \in \mathbb{N}} \sup_{\mathbf{u}_0 \in \mathcal{U}} \|\Pi_n S(t)\mathbf{u}_0\|_{\mathbf{H}^1} \leq \sup_{\mathbf{u}_0 \in \mathcal{U}} \|S(t)\mathbf{u}_0\|_{\mathbf{H}^1} < +\infty. \quad (67)$$

Write

$$\Pi_n^c := I - \Pi_n$$

By (61) and (66) we have

$$\Pi_n^c A = A \Pi_n^c.$$

Thus, from (9) we get

$$\Pi_n^c \mathbf{u}(t) = \Pi_n^c \mathbf{u}_0 - \nu \int_0^t A \Pi_n^c \mathbf{u} ds + \int_0^t \Pi_n^c B(\mathbf{u}, \mathbf{u}) ds - \int_0^t g_N^{\nu,\kappa}(\|\mathbf{u}\|_{\infty}^2) \Pi_n^c \mathbf{u} ds.$$

By the chain rule (cf. [19, p.176, Lemma 1.2]) we have

$$\begin{aligned} \frac{d}{dt} \|\Pi_n^c \mathbf{u}(t)\|_{\mathbf{H}^1}^2 &= -2\nu \|A\Pi_n^c \mathbf{u}\|_{\mathbf{L}^2}^2 + 2\langle \Pi_n^c B(\mathbf{u}, \mathbf{u}), \Pi_n^c \mathbf{u} \rangle_{\mathbf{H}^1} \\ &\quad - 2g_N^{\nu, \kappa}(\|\mathbf{u}\|_{\infty}^2) \cdot \|\Pi_n^c \mathbf{u}\|_{\mathbf{H}^1}^2 \\ &\leq -\nu \|A\Pi_n^c \mathbf{u}\|_{\mathbf{L}^2}^2 + \frac{1}{\nu} \|\Pi_n^c B(\mathbf{u}, \mathbf{u})\|_{\mathbf{L}^2}^2. \end{aligned}$$

Noting that

$$\|\Pi_n^c \mathbf{u}\|_{\mathbf{H}^1}^2 = \|A^{\frac{1}{2}} \Pi_n^c \mathbf{u}\|_{\mathbf{L}^2}^2 \leq \frac{1}{\lambda_n} \|A\Pi_n^c \mathbf{u}\|_{\mathbf{L}^2}^2$$

and

$$\|\Pi_n^c B(\mathbf{u}, \mathbf{u})\|_{\mathbf{L}^2}^2 \leq \|\mathbf{u}\|_{\infty}^2 \cdot \|\nabla \mathbf{u}\|_{\mathbf{L}^2}^2 \leq C \cdot \|A\mathbf{u}\|_{\mathbf{L}^2} \cdot \|\nabla \mathbf{u}\|_{\mathbf{L}^2}^3 =: h(t),$$

we have

$$\frac{d}{dt} \|\Pi_n^c \mathbf{u}(t)\|_{\mathbf{H}^1}^2 + \nu \lambda_n \|\Pi_n^c \mathbf{u}\|_{\mathbf{H}^1}^2 \leq \frac{h(t)}{\nu}.$$

Solving this differential inequality yields that

$$\begin{aligned} \|\Pi_n^c \mathbf{u}(t)\|_{\mathbf{H}^1}^2 &\leq e^{-\nu \lambda_n t} \left(\|\Pi_n^c \mathbf{u}_0\|_{\mathbf{H}^1}^2 + \int_0^t \frac{1}{\nu} e^{\nu \lambda_n s} h(s) ds \right) \\ &\leq e^{-\nu \lambda_n t} \left(\|\mathbf{u}_0\|_{\mathbf{H}^1}^2 + \frac{1}{\nu} \left(\int_0^t e^{2\nu \lambda_n s} ds \right)^{1/2} \left(\int_0^t h(s)^2 ds \right)^{1/2} \right) \\ &\leq e^{-\nu \lambda_n t} \|\mathbf{u}_0\|_{\mathbf{H}^1}^2 + \frac{1}{\nu \sqrt{2\nu \lambda_n}} \left(\int_0^t h(s)^2 ds \right)^{1/2}. \end{aligned}$$

On the other hand, by (11) we have

$$\int_0^t h(s)^2 ds \leq \sup_{s \in [0, t]} \|\nabla \mathbf{u}(s)\|_{\mathbf{L}^2}^6 \int_0^t \|A\mathbf{u}\|_{\mathbf{L}^2}^2 ds \leq \frac{1}{\nu} \left(\frac{\kappa N}{\nu^2} \|\mathbf{u}_0\|_{\mathbf{L}^2}^2 + \|\nabla \mathbf{u}_0\|_{\mathbf{L}^2}^2 \right)^4.$$

Hence, by $\lambda_n \uparrow \infty$ we obtain

$$\limsup_{n \rightarrow \infty} \sup_{\mathbf{u}_0 \in \mathcal{U}} \|\Pi_n^c S(t) \mathbf{u}_0\|_{\mathbf{H}^1}^2 = \limsup_{n \rightarrow \infty} \sup_{\mathbf{u}_0 \in \mathcal{U}} \|\Pi_n^c \mathbf{u}(t)\|_{\mathbf{H}^1}^2 = 0,$$

which combined with (67) yields by Lemma 6.1 that $S(t)\mathcal{U}$ is relatively compact in \mathbf{H}^1 . \square

Proof of Theorem 2.8: It follows from [20, p. 23 Theorem 1.1 and (1.12')] and Lemmas 6.2 and 6.3.

Acknowledgements:

The author would like to thank Professor Benjamin Goldys for providing him an excellent environment to work in the University of New South Wales. His work is supported by ARC Discovery grant DP0663153 of Australia.

REFERENCES

- [1] R.A. Adams and J.F. Fournier: *Sobolev Spaces*. Second Edition, Academic Press, 2003.
- [2] A.L. Bertozzi, A.J. Majda: *Vorticity and Incompressible Flow*. Cambridge Texts in Applied Mathematics, Cambridge University Press, 2002.
- [3] L. Caffarelli, R. Kohn, L. Nirenberg: Partial regularity of suitable weak solutions of the Navier-Stokes equations. *Comm. Pure Appl. Math.* 35 (1982), no. 6, 771–831.

- [4] T. Caraballo, J. Real and P.E. Kloeden: Unique Strong Solutions and V -Attractors of a Three Dimensional System of Globally Modified Navier-Stokes Equations. *Advanced Nonlinear Studies*, 6(2006), 411-436.
- [5] A. Friedman: Partial Differential Equations, Holt, Rinehart and Winston, INC., New York, 1969.
- [6] H. Fujita, T. Kato: On the Navier-Stokes Initial Value Problem. I. *Arch. Rational Mech. Anal.*, 16 (1964), 269–315.
- [7] G.P. Galdi: An introduction to the Navier-Stokes initial-boundary value problem. Fundamental directions in mathematical fluid mechanics, 1–70, Adv. Math. Fluid Mech., Birkhäuser, Basel, 2000.
- [8] J.G. Heywood: The Navier-Stokes Equations: On the Existence, Regularity and Decay of Solutions, *Indiana University Mathematics Journal*, Vol.29, No.5 (1980), 639-681.
- [9] E. Hopf: Über die Anfangswertaufgabe für die hydrodynamischen Grundgleichungen, *Math. Nachr.*, 4, 1950-1951, 213-231.
- [10] O.A. Ladyzhenskaya: *The mathematical theory of viscous incompressible flow*. Second English edition, revised and enlarged. Translated from the Russian by Richard A. Silverman and John Chu. Mathematics and its Applications, Vol. 2 Gordon and Breach, Science Publishers, New York-London-Paris 1969 xviii+224.
- [11] J. Leray: Sur le mouvement d'un liquide visqueux emplissant l'espace, *Acta Math.*, 63(1934), 193-248.
- [12] F. Lin: A new proof of the Caffarelli-Kohn-Nirenberg theorem. *Comm. Pure Appl. Math.* 51 (1998), no. 3, 241–257.
- [13] P.L. Lions: Mathematical Topics in Fluid Mechanics, Volume 1, Incompressible Models. Oxford Lect. Series in Math. and its App. 3, 1996.
- [14] A. Pazy: *Semi-groups of linear operators and applications*. Berlin: Springer-Verlag, 1985.
- [15] M. Röckner, X. Zhang: Tamed 3D Navier-Stokes Equation: Existence, Uniqueness and Regularity. Preprint.
- [16] V. Scheffer: Hausdorff measure and the Navier-Stokes equations. *Comm. Math. Phys.* 55 (1977), no. 2, 97–112.
- [17] H. Sohr: The Navier-Stokes Equations: An Elementary Functional Analytic Approach. Advanced Texts, Birkhäuser Verlag, 2001.
- [18] E.M. Stein: Singular Integrals and Differentiability Properties of Functions. Princeton University Press, Princeton, New Jersey, 1970.
- [19] R. Temam: *Navier-Stokes equations: Theory and numerical analysis*. Originally published: Amsterdam, North-Holland, 1977. AMS, Providence, Rhode Island, 2001.
- [20] R. Temam: *Infinite-Dimensional Dynamical Systems in Mechanics and Physics*. Applied Math. Sci., 68. Springer-Verlag, Berlin, 1997
- [21] H. Triebel: *Interpolation Theory, Function Spaces, Differential Operators*. North-Holland Publishing Company, Amsterdam-New York-Oxford, 1978.