

# Activity-dependent stochastic resonance in recurrent neuronal networks

Vladislav Volman<sup>1,2,\*</sup> and Herbert Levine<sup>1</sup>

<sup>1</sup>Center for Theoretical Biological Physics, University of California at San Diego, La Jolla, CA 92093-0319, USA

<sup>2</sup>Computational Neurobiology Laboratory, The Salk Institute for Biological Studies, La Jolla, CA 92037, USA

We use a biophysical model of a local neuronal circuit to study the implications of synaptic plasticity for the detection of weak sensory stimuli. Networks with fast plastic coupling show behavior consistent with stochastic resonance. Addition of an additional slow coupling that accounts for the asynchronous release of neurotransmitter results in qualitatively different properties of signal detection, and also leads to the appearance of transient post-stimulus bistability. Our results suggest testable hypothesis with regard to the self-organization and dynamics of local neuronal circuits.

PACS numbers: Valid PACS appear here

Stochastic resonance (SR) refers to the condition in which noise and nonlinearity combine together to amplify otherwise undetectable stimuli [1]. This simple, yet important, phenomenon, has received much attention due to its apparent ubiquity in many nonlinear abiotic [1] and biological [2] systems. In particular, a number of studies have raised the possibility that neurons [3, 4] and neuronal cell assemblies [5] might utilize SR in order to detect weak sensory stimuli [2].

For these studies, the noise felt by individual neurons has been assumed to arise from the random summation of a large number of synaptic stimuli [4, 6]. There is however another important source of noise, that of the stochastic nature of synaptic transmission. In particular, there can occur spontaneous asynchronous release (AR) of neurotransmitter at a rate that is strongly dependent on the pre-synaptic  $Ca^{2+}$  concentration and hence strongly dependent on the rate of spike-induced  $Ca^{2+}$  intake [7]. Since a high probability of release can last for  $> 0.1$  sec, AR constitutes a challenging example of slow time-scale, activity-dependent noise.

The purpose of this work is to show that SR for *local* circuits consisting of roughly 100 neurons (a "micro-column" [8]) coupled via noisy plastic synapses takes a dramatically different form from that seen in investigations to date. As we will see, the coherence of the response continues to depend non-trivially on the coupling strength and the assembly size. Furthermore, the circuit can exhibit short-term memory, by which we mean that spiking will continue to occur for a transient period following removal of the stimulus. These results can be directly tested in experiments on cultured networks [7, 9] and offer some new insights into the way neuronal systems can be organized for optimal information processing. From the dynamical systems point of view, this work represents a new example of how SR phenomenology can depend on the specific type of noise; this has been considered in only a few examples to date [10]

To proceed, we use a network model that has recently been developed to account for the occurrence of rhythmic reverberatory responses in hippocampal cultures [7, 11]. The neurons in the network obey Morris-Lecar like dy-

namics [12] with the membrane voltage given by

$$C\dot{V} = -I_{ion} + I_{bg} + I_{syn} + I_{stim} \quad (1)$$

In eqn.1, the ionic current  $I_{ion}$  describes the contribution from membrane channels [13]. The term  $I_{bg}$  is a background current that represents summation of a large number of synaptic stimuli from neurons that are not part of the specific local circuit. Rather than explicitly modeling a very large network and imposing a connectivity pattern which embodies the local circuit notion, we instead include these neurons *implicitly* by assuming (as in [6]) that their contribution is described by a Langevin equation  $\dot{I}_{bg} = -I_{bg}/\tau_n + \sqrt{D/\tau_n}\mathcal{N}(0, 1)$ , with  $\tau_n = 10$  msec and  $\mathcal{N}(0, 1)$  being uncorrelated Gaussian noise with zero mean and unitary variance. The synaptic current due to the local circuit is modeled as  $I_i^{syn} = -\sum \bar{g}Y_{ij}(t)V_i$ , with  $\bar{g} \in [0.5, 0.8]$  mS/cm<sup>2</sup> being the maximal value of synaptic conductance, the sum running over the set of input channels, and the term  $Y_{ij}$  as described below. With the parameters as given in [13], the transition from quiescence to regular spiking occurs through a Hopf bifurcation.

To capture the dynamical aspects of synaptic coupling, we assume that at any time, presynaptic resources can be in a recovered state (described by the state variable  $X$  in equations below), in an active state (described by the state variable  $Y$ ), or in an inactive state (described as  $Z = 1 - X - Y$ ) [14]. The dynamics for the  $j \rightarrow i$  presynaptic terminal are

$$\dot{X}_{ij} = \frac{Z_{ij}}{\tau_r} - X_{ij}(U\delta(t - t_s^j) + \xi\delta(t - t_a^j)) \quad (2a)$$

$$\dot{Y}_{ij} = \frac{-Y_{ij}}{\tau_d} + X_{ij}(U\delta(t - t_s^j) + \xi\delta(t - t_a^j)) \quad (2b)$$

$$\dot{Z}_{ij} = \frac{Y_{ij}}{\tau_d} - \frac{Z_{ij}}{\tau_r} \quad (2c)$$

$$\eta(c) = \eta_{max} \frac{c^4}{c^4 + K_a^4} \quad (2d)$$

$$\dot{c} = \frac{-\beta c^2}{c^2 + K_{Ca}^2} + \gamma \log\left(\frac{c_o}{c}\right)\delta(t - t_s^j) + I_p \quad (2e)$$

At each presynaptic terminal of the  $i$ -th neuron, the

fraction of active resources ( $Y_{ij}$ ) experiences a brief increase of magnitude  $UX_{ij}$  when, at time  $t_s^j$ , an action potential from  $j$ -th neuron invades the presynaptic terminal. Alternatively, a relatively small amount of resource can be maintained in an active state by the asynchronous release of synaptic resource that occurs at times  $t_a^j$  with  $Ca^{2+}$ -dependent rate  $\eta(c)$  and amplitude  $\xi$ . The rate of asynchronous release (probability to observe an event during the interval  $[t_a, t_a + dt]$ , modeled as Poisson process) is taken to be a Hill function of the presynaptic residual  $Ca^{2+}$  concentration,  $c$  [11, 15]. This residual  $Ca^{2+}$  accumulates at presynaptic terminals in an activity-dependent way that is proportional to electrochemical gradient across the membrane, and is extruded into the extra-cellular space by a non-linear pump. The term  $I_p$  ensures that the minimal  $Ca^{2+}$  concentration is  $\approx 60$  nM. Parameters are given in [16]. Note that the phasic ( $UX_{ij}$ ) and asynchronous ( $\xi X_{ij}$ ) terms are both proportional to the amount of available resource,  $X_{ij}$ , underscoring the activity-dependent competition between these two different coupling modes [17].

To assess the extent to which an individual neuron and/or a network can exhibit coherent activity, we use the coherence of spiking (COS) measure [4, 18]. Given a weak external sub-threshold stimulation of period  $T$ ,  $I_{stim}(t) = 1 \frac{nA}{cm^2} \sin(2\pi \frac{t}{T})$ , the COS measure is defined here as  $C_S \equiv \frac{N(0.9T <= ISI <= 1.1T)}{N(ISI)}$ , that is, the fraction of inter-spike-intervals (ISIs) that are within 20% of stimulus period,  $T = 0.1$  sec. All results, unless otherwise indicated, are for a network of  $N = 100$  neurons that have probability  $p = 0.1$  to establish connections with their peers.

We first analyze the response of an uncoupled neuronal network to weak sub-threshold periodic stimulation and different (controlled) intensities of synaptic background noise,  $I_{bg}$ . In agreement with previous studies [3], we find that there exists an optimal level of noise for which a model neuron exhibits a maximal coherence of spiking (Fig.1A, dashed line). Coupling the model neurons by activity-dependent synapses (as in eqs. 2) while setting  $\eta_{max} = 0$  (no asynchronous release) moves the resonant peak towards lower noise intensities. As Fig.1A (insets) shows, the location and the height of the new peak is largely independent of the coupling parameter,  $U$ . This observation is consistent with the notion of efficient signal propagation on random graphs - once  $U$  is above critical coupling threshold, an SR-like activation of one neuron will quickly spread the word to other neurons, regardless of the exact value of  $U$ .

Introduction of activity-dependent asynchronous release of neurotransmitter results in a qualitatively different picture. The coherence measure as a function of evoked and asynchronous release is shown in Fig.2A. It is clear that the spiking coherence increases significantly for higher values of the resource utilization parameter




FIG. 1: **Stochastic resonance in dynamically coupled neuronal networks.** A) An uncoupled cell ensemble exhibits a broad-peak stochastic resonance with relatively weak coherence of spiking (dashed line). Introduction of dynamic coupling enables the efficient exchange of stimulus-related information, and moves the resonance peak to lower noise intensities. Once above a minimal coupling threshold, different coupling strengths induce nearly the same coherence-noise curves (superimposed lines). Both optimal noise intensity (top inset) and peak coherence (bottom inset) are nearly independent of synaptic coupling. B) Sample neuronal membrane voltage for  $D = 1.25 \cdot 10^{-2}$ ,  $U = 0.4$ ,  $\eta_{max} = 0$ .

$U$ . The optimal level of AR needed to produce maximal coherence (peaks in Fig.2A) also depends on the value of  $U$ . Stronger evoked transmission will quickly deplete the available resources; therefore, since asynchronous and evoked releases draw from the same pool of vesicles, higher rate of spontaneous events is needed to achieve significant spiking coherence (top inset of Fig.2A). For higher values of  $\eta_{max}$ , when the combined action of AR and  $I_{bg}$  masks the stimulus by making the cell spike more frequently, the coherence measure converges to low values. On the other hand, strong coupling and fast depletion of resources provide a constraint for spiking activity, resulting in higher overall coherence for higher resource usage (bottom inset of Fig.2A).

The distinctive effect of AR (as compared with  $I_{bg}$ ) is further assessed by analyzing the collective dynamics for high  $\eta_{max}$  (vs. high  $D$ ). Subjecting the network to high-intensity dynamics-independent noise (Fig.3A) results in high-rate, weakly correlated, activity. On the other hand, as Fig.3B shows, the combined action of strong AR and synaptic depression significantly sharpens the network's

response to the stimulus. Further, the prolonged time-scale of AR enables the network to "remember" the stimulus seconds after its cessation (Fig.3C).

The observation that coherence of spiking depends on the strength of dynamic coupling prompted us to explore how networks' parameters affect its ability to detect weak stimuli. To this end, we considered the performance of different size networks, for a range of AR rates. For easier interpretation of results, we assume here that, for all cases,  $U = 0.3$ . Figure 4A shows that the profiles of COS curves are different for different network sizes. Due to the  $p = const$  constraint, neurons in larger networks are subject to higher levels of asynchronous release in their inputs; as a result, the resonant peak moves toward lower values of  $\eta_{max}$ . Conversely, fixing the value of  $\eta_{max}$  and plotting the COS measure as a function of network size (as is in Fig.4B) reveals that the optimal network size (giving maximal coherence) depends on the level of AR at individual model synapses. Thus, in a network with plastic coupling, synaptic parameters might provide constraints for the sizes of cell assembly.

Stochastic resonance relies on the cooperativity between noise, nonlinearity and a weak sub-threshold stimulus [1]. In most examples, the noise is taken to be independent of the characteristics of the weak sub-threshold stimulus (but see [10]). Here, we have investigated the properties of signal processing in local recurrent neuronal networks with plastic processing and asynchronous release of neurotransmitter, where the noise is inherently coupled to the signal. We found that in plastic networks without AR, the characteristics of stochastic resonance (location and height of peak coherence) only weakly depend on the strength of synaptic coupling. On the contrary, introduction of AR leads to a strong dependence of SR properties on network parameters.

These observations suggest that asynchronous release of neurotransmitter might play an important role in neuronal dynamics [19]. Information that is contained in weak signals should not only be detected and amplified by brain circuitry; a network has to have the ability to transiently "hold" knowledge about preceding events. As shown in [7, 11], a brief stimulus delivered to the network evokes reverberatory activity that is sustained by the asynchronous release of neurotransmitter and lasts for several seconds. Our results (Fig.3C) together with experimental observations [7] and prior modeling [11], suggest that AR can be instrumental in detection, amplification, and transient holding of weak sensory stimuli.

This study leads to several potentially interesting conclusions. First, we showed here that the ability of a neuron (that is embedded in a neuronal network) to detect and amplify weak stimuli might crucially depend on the form of feedback from the network, and in particular on the plasticity features of the effective connectivity. Second, our results suggest that the plasticity of synaptic connections might provide an important con-




FIG. 2: **Stochastic resonance in a network with asynchronous release of transmitter.** A) When a slow asynchronous mode of synaptic transmission is introduced in addition to the fast phasic coupling, the extent of output spiking coherence depends on the strength of the phasic coupling ( $U$ ). For clarity of presentation, only the cases  $U = 0.1, 0.3, 0.5, 0.7$  are shown. Both the location (top inset) and magnitude (bottom inset) of the coherence peak are positively correlated with the strength of evoked synaptic transmission, underscoring the fact that both kinds of synaptic transmission share the same pool of synaptic resources. B) Sample neuronal membrane voltage for  $\eta_{max} = 0.28, U = 0.4, D = 0.64 \cdot 10^{-2}$ .

straint for the optimal number of neurons in a local circuit. With this perspective, the local network with strong inter-connectivity is optimized for signal detection, with distant neurons providing contextual information in the form of an overall background noise signal. Cultured networks can provide an adequate framework to test the validity of our conclusions. State of the art techniques allow one to grow small networks of controlled size, geometry and connectivity [9]. Future experiments will determine how these parameters affect the ability of a network to process weak stimuli.

We thank W.J. Rappel and T.J. Sejnowski for stimulating discussions. This research has been supported by the NSF-sponsored Center for Theoretical Biological Physics (grant nos. PHY-0216576 and PHY-0225630).

\* Communicating author: volman@salk.edu

[1] L. Gammitoni *et al.*, Rev. Mod. Phys. **70**, 223 (1998)  
 [2] R.J. Douglas *et al.*, Nature **365**, 337 (1993); J.J. Collins




FIG. 3: **AR induces correlated collective dynamics.** Raster plots show the firing activity of a neuronal network with  $U = 0.4$ . *A*) A network with  $D = 7.84 \cdot 10^{-2}$ ,  $\eta_{max} = 0$  exhibits high-rate, uncorrelated activity. On the contrary, a network with  $D = 0$ ,  $\eta_{max} = 0.8$  (*B*) exhibits burst-like, correlated collective activity. *C*) A network driven by AR shows transient bi-stability in its activity (top panel) that persists for seconds after stimulus removal (bottom panel).

*et al.*, J. Neurophysiol. **76**(1), 642 (1996); J.E. Levin and J.P. Miller, Nature **380**, 165 (1996); B. Gluckman *et al.*, Phys. Rev. Lett. **77**(19), 4098 (1996); F. Jaramillo and K. Wiesenfeld, Nature Neurosci. **1**, 384 (1998); W.C. Stacey and D.M. Durand, J. Neurophysiol. **83**, 1394 (2000); F. Moss *et al.*, Clin. Neurophysiol. **115**, 267 (2004)

- [3] A. Longtin, J. Stat. Phys. **70**(1,2), 309 (1993)
- [4] M. Rudolph and A. Destexhe, Phys. Rev. Lett. **86**(16), 3662 (2001)
- [5] A.R. Bulsara and G. Schnera, Phys. Rev. E **47**, 3734 (1993); W.J. Rappel and A. Karma, Phys. Rev. Lett. **77**(15), 3256 (1996); D. Chialvo *et al.*, Phys. Rev. E **55**(2), 1798 (1997); G. Mato, Phys. Rev. E **58**, 876 (1998); Y. Yu *et al.*, Phys. Rev. E **63**, 021907 (2001); C. Zhou *et al.*, Phys. Rev. Lett. **87**(9), 098101 (2001)
- [6] A. Destexhe *et al.*, Neurosci. **107**(1) (2001)
- [7] P. Lau and G. Bi, Proc. Natl. Acad. Sci. U.S.A. **102**, 10333 (2005)
- [8] E.G. Jones, Proc. Natl. Acad. Sci. U.S.A. **97**(10), 5019 (2000)
- [9] R. Sorkin *et al.*, J. Neural Eng., **3**, 95 (2007)
- [10] S. Bezikov and I. Vodyanoy, Nature **385**, 319 (1997); B. Lindner and L. Schimansky-Geier, Phys. Rev. Lett. **86**(14), 2934 (2001)
- [11] V. Volman *et al.*, Phys. Biol., **4**, 91(2007)
- [12] C. Morris and H. Lécarré, Biophys. J. **35**, 193 (1981); S.A. Prescott *et al.*, J. Neurosci. **25**(36), 9084 (2006)
- [13] For each model neuron, the ionic current was  $I_{ion} = G_{Na}m_{\infty}(V - E_{Na}) + G_Kw(V)(V - E_K) + G_L(V - E_L)$ .

The fraction of open  $K^+$  channels evolved as  $dw/dt =$




FIG. 4: **System size and connectivity affect coherence of spiking.** *A*) For a network with a uniform connection probability ( $p = 0.1$ ), the profile of coherence as a function of maximal AR rate depends on the number of neurons. With a  $p = const$  scheme, larger networks lead to higher per cell number of afferents that affects the effective level of asynchronous release. *B*) Optimal network size (peaks) that gives rise to maximal coherence depends on the maximal rate of AR at model synapses. In all cases, the intensity of background synaptic noise is  $D = 0.64 \cdot 10^{-2}$ .

$0.15(w_{\infty}(V) - w(V))\cosh((V - V_3)/2V_4)$ . The steady-state fraction of  $Na^+$  and  $K^+$  channels were, correspondingly,  $m_{\infty} = 0.5(1 + \tanh((V - V_1)/V_2))$  and  $w_{\infty} = 0.5(1 + \tanh((V - V_3)/V_4))$ . Values of parameters were:  $E_{Na} = 50$  mV,  $E_K = -100$  mV,  $E_L = -55.8$  mV,  $V_1 = -1.2$  mV,  $V_2 = 23$  mV,  $V_3 = -2$  mV,  $V_4 = 21$  mV,  $G_{Na} = 11$  mS/cm<sup>2</sup>,  $G_K = 10$  mS/cm<sup>2</sup>,  $G_L = 1.5$  mS/cm<sup>2</sup>,  $C = 1$   $\mu$ F/cm<sup>2</sup>. Equations were solved using second-order Runge-Kutta method with  $dt = 0.1$  msec.

- [14] M. Tsodyks *et al.*, J. Neurosci. **20** (2000)
- [15] R. Ravin *et al.*, J. Physiol. **501**(2), 251 (1997) S. Kirischuk and R. Grantyn, J. Physiol. **548**(3), 754 (2003).
- [16] The parameters used to model synaptic transmission were  $\tau_d = 5$  msec,  $\tau_r = 0.6$  sec,  $K_a = 0.1$   $\mu$ M,  $K_{Ca} = 0.4$   $\mu$ M,  $\beta = 2$   $\mu$ M/sec,  $\gamma = 80$  nM,  $C_0 = 2$  mM,  $I_p = 0.11$   $\mu$ M/sec,  $\xi = 10^{-3}$
- [17] D. Hagler and Y. Goda, J. Neurophysiol. **85**, 2324 (2001); Y. Otsu *et al.*, J. Neurosci. **24**(2), 420 (2004)
- [18] D.R. Chialvo and A.V. Apkarian, J. Stat. Phys. **70**, 375 (1993)
- [19] J. Jones *et al.*, J. Neurophysiol. **97**, 3812 (2007); K.J. Iremonger and J.S. Bains, J. Neurosci. **27**(25), 6684 (2007)