ON THE EXCEDANCE SETS OF COLORED PERMUTATIONS

ELI BAGNO, DAVID GARBER, AND ROBERT SHWARTZ

ABSTRACT. We define the excedence set and the excedance word on $G_{r,n}$, generalizing a work of Ehrenborg and Steingrimsson and use the inclusion-exclusion principle to calculate the number of colored permutations having a prescribed excedance word. We show some symmetric properties as Log concavity and unimodality of a specific sequence of excedance words.

1. INTRODUCTION

Let S_n be the symmetric group on n letters. The parameter *excedance*, which is defined on a permutation $\pi \in S_n$ by

$$\exp(\pi) = |\{i \in [n] \mid \pi(i) > i\}|,\$$

is well-known. $([n] = \{1, \ldots, n\}, as usual).$

One can also define the *excedance set* of a permutation π :

$$\operatorname{Exc}(\pi) = \{ i \in [n] \mid \pi(i) > i \}.$$

One can encode the excedance set of a permutation π as a word w_{π} of length n-1 in the letters a, b, where a in the *i*th place means that $i \notin \text{Exc}(\pi)$, while b in the *i*th place means that $i \in \text{Exc}(\pi)$. For example, the excedance set of the permutation:

$$\pi = \left(\begin{array}{rrrrr} 1 & 2 & 3 & 4 & 5 \\ 3 & 5 & 1 & 4 & 2 \end{array}\right) \in S_5$$

is encoded by the word: *bbaa*.

Let w be a word in a, b. Ehrenborg and Steingrimsson [2] defined:

$$[w] = \#\{\pi \in S_n \mid w_\pi = w\}$$

They give some recursive and explicit formulas for [w]. Moreover, they consider the unimodality of the series $\{[b^k a^{n-1-k}] \mid 0 \le k \le n-1\}$ as well as some log-concavity results.

Date: October 30, 2018.

In this paper, we generalize this idea to the colored permutation group $G_{r,n} = \mathbb{Z}_r \wr S_n$ (defined in Sections 2. We consider the series

$$\{ [b^k a^{rn-1-k}] \mid 0 \le k \le rn-1 \},\$$

show its log-concavity and conclude its unimodality (since all elements are positive) (See Section 4). The proof is based on finding an explicit formula for the number $[b^k a^{rn-1-k}]$. Moreover, in Section 5, we supply an algorithm to compute the number of colored permutations having a fixed excedance, based on the inclusion-exclusion approach presented in [2].

2. The group of colored permutations

Definition 2.1. Let r and n be positive integers. The group of colored permutations of n digits with r colors is the wreath product:

$$G_{r,n} = \mathbb{Z}_r \wr S_n = \mathbb{Z}_r^n \rtimes S_n,$$

consisting of all the pairs (\vec{z}, τ) where \vec{z} is an *n*-tuple of integers between 0 and r-1 and $\tau \in S_n$. The multiplication is defined by the following rule: for $z = (z_1, \ldots, z_n)$ and $\vec{z}' = (z'_1, \ldots, z'_n)$

$$(\vec{z},\tau) \cdot (\vec{z}',\tau') = ((z_1 + z'_{\tau^{-1}(1)}, \dots, z_n + z'_{\tau^{-1}(n)}), \tau \circ \tau')$$

(here + is taken modulo r).

Note that the symmetric group $S_n = G_{1,n}$ and the group of signed permutations $B_n = G_{2,n} = C_2 \wr S_n$ are special cases of $G_{r,n}$.

We use some conventions along this paper. For an element $\pi = (\vec{z}, \tau) \in G_{r,n}$ with $\vec{z} = (z_1, \ldots, z_n)$ we write $z_i(\pi) = z_i$. For $\pi = (\vec{z}, \tau)$, we denote $|\pi| = (\vec{0}, \tau), (\vec{0} \in \mathbb{Z}_r^n)$. We also define $c_i(\pi) = z_i(\pi^{-1})$ and $\vec{c} = (c_1, \ldots, c_n)$. Using this notation, the element $(\vec{c}, \tau) = \left((0, 1, 2, 3), \begin{pmatrix} 1 & 2 & 3 & 4 \\ 2 & 1 & 4 & 3 \end{pmatrix}\right) \in G_{3,4}$ will be written as $(2\overline{1}\overline{4}\overline{3})$.

Here is another way to present $G_{r,n}$: Consider the alphabet $\Sigma = \{1, \ldots, n, \overline{1}, \ldots, \overline{n}, \ldots, 1^{[r-1]}, \ldots, n^{[r-1]}\}$ as the set [n] colored by the colors $0, \ldots, r-1$. Then, an element of $G_{r,n}$ is a colored permutation, i.e., a bijection $\pi : \Sigma \to \Sigma$ satisfying the following condition: if $\pi(i^{[\alpha]}) = j^{[\beta]}$ then $\pi(i^{[\alpha+1]}) = j^{[\beta+1]}$.

Define the *color order* on Σ :

$$1^{[r-1]} < \dots < n^{[r-1]} < 1^{[r-2]} < 2^{[r-2]} < \dots < n^{[r-2]} < \dots < 1 < \dots < n.$$

The complete notation of π will be presented by an example: Let $\pi = (\vec{z}, \tau) = \left((1, 2, 0), \begin{pmatrix} 1 & 2 & 3 \\ 3 & 1 & 2 \end{pmatrix} \right) \in G_{3,3}$. The complete notation of

 π is:

 $\begin{pmatrix} \bar{\bar{1}} & \bar{\bar{2}} & \bar{\bar{3}} & \bar{1} & \bar{2} & \bar{3} & 1 & 2 & 3 \\ \bar{\bar{3}} & 1 & \bar{2} & \bar{3} & \bar{\bar{1}} & 2 & 3 & \bar{1} & \bar{\bar{2}} \end{pmatrix}.$

We will use the complete notation throughout this paper.

3. Statistics on $G_{r,n}$

We start by defining the excedance set for colored permutations. Let $\pi \in G_{r,n}$. Write π in its complete notation and define:

$$\operatorname{Exc}(\pi) = \{ x \in \Sigma \mid \pi(x) > x \}.$$

Other definitions can be found in [3, 4].

We associate to π the matrix $M(\pi) = (t_i^j)$ where $1 \le i \le n, j$ varies from r-1 down to 0 and $t_i^j \in \{a, b\}$ in the following way:

$$t_i^j = \begin{cases} a & \pi(i^{[j]}) \le i^{[j]} \\ b & \pi(i^{[j]}) > i^{[j]} \end{cases}$$

Example 3.1. Let

$$\pi = \begin{pmatrix} \bar{1} & \bar{2} & \bar{3} & \bar{1} & \bar{2} & \bar{3} & 1 & 2 & 3 \\ \bar{3} & 1 & \bar{2} & \bar{3} & \bar{1} & 2 & 3 & \bar{1} & \bar{2} \end{pmatrix}.$$

The associated matrix is:

$$M(\pi) = \begin{pmatrix} b & b & b \\ b & a & b \\ b & a & a \end{pmatrix}.$$

Note that $t_n^0 = a$ always.

Let $\pi \in G_{r,n}$. The excedance word of π , w_{π} is the word obtained by reading the entries of $M(\pi) = (t_i^j)$ row by row from left to right, ignoring the last place. Thus for π defined above one has $w_{\pi} = bbbbabba$. We usually insert separators between the rows of $M(\pi)$. In our example, we get $w_{\pi} = (bbb|bab|ba)$.

For a word $w \in \{a, b\}^{rn-1}$, define [w] to be the number of elements $\pi \in G_{r,n}$ such that $w_{\pi} = w$.

4. Unimodality and Log-concavity of the sequence $\{[b^k a^{rn-1-k}] \mid 0 \le k \le rn-1\}$

We recall the following definitions:

Definition 4.1. A sequence of positive real numbers a_0, a_1, \ldots is *unimodal* if for some integers $0 \le k \le n, t \ge 0$, we have

$$a_0 \leq a_1 \leq \cdots \leq a_k \geq \cdots \geq a_n.$$

Definition 4.2. A sequence of real numbers a_0, a_1, \ldots is Log-concave if for any k > 0 one has: $a_{k-1} \cdot a_{k+1} \leq a_k^2$.

It is known that if the elements of the sequence are positive then Log-concavity implies unimodality.

The following two observations are essential for the sequal.

Observation 4.3. Let $\pi \in G_{r,n}$ and let $i \in \{1, \ldots, n\}$. Then $c_i(\pi) = 0$ if and only if all the entries of the *i*-th column of $M(\pi)$ are equal. In this case, $t_i^{r-1} = a$ if and only if $\pi(i) \leq i$.

Observation 4.4. Let $\pi \in G_{r,n}$ and let $i \in \{1, \ldots, n\}$ be such that $c_i(\pi) \neq 0$. Then the *i*-th column of $M(\pi)$ is of the form: $t_i^{r-1} \cdots t_i^0 = b^u a^{r-u}$ if and only if $u = c_i(\pi)$.

Theorem 4.5. For each $0 \le k \le rn - 1$, one has:

$$[b^{k}a^{rn-1-k}] = \begin{cases} (k+1)^{n-k}k! & 0 \le k \le n\\ n! & n+1 \le k \le n(r-1)\\ (nr-k)^{k-nr+n}(nr-k)! & n(r-1)+1 \le k \le rn-1 \end{cases}$$

Proof. Let $w = (b^k a^{rn-1-k})$.

We start with the case k = 0. Then $w = a^{rn-1}$. The only element $\pi \in G_{r,n}$ having no excedances at all is the identity element and thus $[a^{rn-1}] = 1$.

For $1 \leq k \leq n$, we have $w = (b^k a^{n-k} |a^n| a^n | \cdots |a^n| a^{n-1})$. Let π be such that $w_{\pi} = w$. This means that for each $1 \leq i \leq k$ we have $c_i(\pi) = 1$ by Observation 4.4, while for $k < i \leq n$, $c_i(\pi) = 0$ and $\pi(i) \leq i$ by Observation 4.3. Thus $[w] = (k+1)^{n-k}k!$. (Indeed, first fix $\pi(i)$ for $k+1 \leq i \leq n$ and then fix the rest k places). Similar computations can be found at [1] and [2].

For $n+1 \leq k \leq n(r-1)$, by Observation 4.4, $c_i(\pi) \neq 0$ for each $i \in \{1, \ldots, n\}$, and thus [w] = n!.

If
$$n(r-1) + 1 \le k \le rn-1$$
, then $w = (\underbrace{b^n | \cdots | b^n}_{r-1} | b^{k-n(r-1)} a^{rn-1-k})$.

Denote k' = k - n(r - 1). Then for $1 \le i \le k'$ one has $c_i(\pi) = 0$ and $\pi(i) > i$ and for $k' + 1 \le i \le n$, $c_i(\pi) \ne 0$. Hence $[w] = (n - k')^{k'}(n - k')!$.

Corollary 4.6. The sequence $\{[b^k a^{rn-1-k}] \mid k \in \{0, \ldots, rn-1\}\}$ is log-concave and unimodal.

Proof. By the symmetry property of the excedance, appearing in [1] one has for each $k \in \{0, \ldots, rn-1\}$, $[a^k b^{rn-1-k}] = [b^k a^{rn-1-k}]$, hence it is sufficient to prove that the sequence $[b^k a^{rn-1-k}]$ is increasing for $k \in \{0, \ldots, n\}$, but this is trivial.

Since the sequence is positive, the sequence is unimodal too. \Box

5. A direct computation for [w] using inclusion-exclusion

Recall that for each $\pi \in G_{r,n}$, the word w_{π} is actually a monomial in a, b. By a + b in the location i of w_{π} we mean that we do not care if i is an excedance of π or not. In this way, we can speak about expressions $w \in \mathbb{Z}[a, b]$.

For each word $u = u_1 \cdots u_t$, define $E(u) = \{i \mid u_i = b\}$. Let $(n_1, \ldots, n_{k+1}) \in \mathbb{N}^{k+1}$ and define $w = a^{n_1} \cdot (a+b) \cdot a^{n_2} \cdot (a+b) \cdots (a+b) \cdot a^{n_{k+1}}$. Then one has

$$\{\pi \in G_{r,n} \mid w_{\pi} = a^{n_1} \cdot (a+b) \cdot a^{n_2} \cdot (a+b) \cdots (a+b) \cdot a^{n_k+1}\} = \{\pi \in G_{r,n} \mid E(w_{\pi}) \subseteq E(u)\}$$

where $u = a^{n_1} b a^{n_2} b a^{n_3} b \cdots a^{n_k} b a^{n_k+1}$.

Ehrenborg and Steingrimsson [2] computed a formula for the number of permutations of S_n having a given excedance set using the inclusionexclusion principle. Their main results are the following:

Lemma 5.1 (Ehrenborg-Steingrimsson). For any vector $(n_1, \ldots, n_{k+1}) \in \mathbb{N}^{k+1}$ one has:

$$[a^{n_1} \cdot (a+b) \cdot a^{n_2} \cdot (a+b) \cdots (a+b) \cdot a^{n_{k+1}}] = 1^{n_1+1} \cdot 2^{n_2+1} \cdots (k+1)^{n_{k+1}+1}.$$

Theorem 5.2 (Ehrenborg-Steingrimsson). Let $w = a^{n_1}ba^{n_2}b\cdots a^{n_k}ba^{n_{k+1}}$ be an ab-word with exactly k b's. Then

$$[w] = \sum_{\mathbf{r} \in R_k} (-1)^{h(\mathbf{r})} \cdot \mathbf{r}^{\mathbf{n}(w)+1}$$

where

$$R_{k} = \{ \mathbf{r} = (r_{1}, \dots, r_{k+1}) \mid r_{1} = 1, r_{i+1} - r_{i} \in \{0, 1\} \},\$$
$$h(\mathbf{r}) = |\{i \in \{1, \dots, k\} \mid r_{i} = r_{i+1}\}|,\$$

 $\mathbf{n}(w) = (n_1, \dots, n_{k+1}) \text{ and } \mathbf{1} = (1, 1, \dots, 1).$

In this section we present a similar formula for excedance words in $G_{r,n}$, based on Theorem 5.2.

Define

$$\Psi: M_{r,n}(\{a,b\}) \to \{a,b,a+b\}^{n-1}$$

by $M = (t_i^j) \mapsto w' = (w_1' \cdots w_{n-1}')$ where

$$w'_{i} = \begin{cases} a & t^{j}_{i} = a, \forall j \in \{0, \dots, r-1\} \\ b & t^{j}_{i} = b, \forall j \in \{0, \dots, r-1\} \\ a+b & \text{Otherwise} \end{cases}$$

Example 5.3. Let

$$M = \begin{pmatrix} a & b & b & b \\ a & b & a & b \\ a & b & a & a \end{pmatrix}$$

which can be also written as a word: w = (abbb|abab|aba). Then $\Psi(w) = (a \ b \ a+b)$.

Now, for each $M \in M_{r,n}(\{a, b\})$ which comes from some $\pi \in G_{r,n}$ (or equivalently, for the associated word w) define another map:

$$\varphi: \{\pi \in G_{r,n} \mid w_{\pi} = w\} \to \{\pi \in S_n \mid w_{\pi} = \Psi(w)\}\$$

by $\varphi(\pi) = |\pi|$. It can be easily shown that φ is a bijection. Thus, instead of computing [w], we compute $[\Psi(w)]$.

In order to compute $[\Psi(w)]$, following [2], we write each b in $w' = \Psi(w)$ as b = (a+b) - a and use the distributivity of the ring $\mathbb{Z}[a, b]$ to expand w' into a sum of monomials in a and a + b. Then we can use Lemma 5.1 to calculate [w'].

We first illustrate this idea by the following example:

Example 5.4. Let w be such that

$$w' = \Psi(w) = (a \quad b \quad a+b \quad a \quad a \quad a+b \quad b \quad a).$$

Then:

$$\begin{bmatrix} w' \end{bmatrix} = \begin{bmatrix} a & b & a+b & a & a & a+b & b & a \end{bmatrix}$$

$$= \begin{bmatrix} a & (a+b) - a & a+b & a & a & a+b & (a+b) - a & a \end{bmatrix} =$$

$$= \begin{bmatrix} a & a+b & a+b & a & a & a+b & a+b & a \end{bmatrix} -$$

$$- \begin{bmatrix} a & a+b & a & b & a & a+b & a & a \end{bmatrix} -$$

$$- \begin{bmatrix} a & a+b & a & a & a+b & a & a \end{bmatrix} +$$

$$+ \begin{bmatrix} a & a & a+b & a & a & a+b & a & a \end{bmatrix} =$$

$$= \begin{bmatrix} a^1 & a+b & a^0 & a+b & a^2 & a+b & a^0 & a+b & a^1 \end{bmatrix} -$$

$$- \begin{bmatrix} a^1 & a+b & a^0 & a+b & a^2 & a+b & a^2 \end{bmatrix} -$$

$$- \begin{bmatrix} a^2 & a+b & a^2 & a+b & a^0 & a+b & a^1 \end{bmatrix} +$$

$$+ \begin{bmatrix} a^2 & a+b & a^2 & a+b & a^2 \end{bmatrix} =$$

$$= 1^2 \cdot 2^1 \cdot 3^3 \cdot 4^1 \cdot 5^2 - 1^2 \cdot 2^1 \cdot 3^3 \cdot 4^3 - 1^3 \cdot 2^3 \cdot 3^1 \cdot 4^2 + 1^3 \cdot 2^3 \cdot 3^3 =$$

$$= 1^2 \cdot 2^1 \cdot 3^3 \cdot 4^1 \cdot 5^2 - 1^2 \cdot 2^1 \cdot 3^3 \cdot 4^1 \cdot 4^2 -$$

$$-1^2 \cdot 1^1 \cdot 2^3 \cdot 3^1 \cdot 4^2 + 1^2 \cdot 1^1 \cdot 2^3 \cdot 3^1 \cdot 3^2 = 1776$$

In general, if $\Psi(w) = (w'_1 \cdots w'_{n-1})$ then define

$$I(w) = \{i \in \{1, \dots, n-1\} \mid w'_i = a+b\}$$

and let k be the number of letters in $\Psi(w)$ which are not a. In order to separate between the digit a and the other digits appearing in $\Psi(w)$, we write $\Psi(w) = (a^{n_1}x_1a^{n_2}x_2\cdots x_ka^{n_{k+1}})$ where $x_i \in \{a+b,b\}$.

Define:

$$R_{k} = \left\{ \mathbf{r} = (r_{1}, \dots, r_{k+1}) \middle| \begin{array}{c} r_{1} = 1, r_{i+1} - r_{i} \in \{0, 1\}, \\ r_{i+1} - r_{i} = 1 \text{ if } i \in I(w) \end{array} \right\},$$

and $h(\mathbf{r}) = |\{i \in \{1, \dots, k\} | r_i = r_{i+1}\}|, \mathbf{n}(w) = (n_1, \dots, n_{k+1})$ and $\mathbf{1} = (1, 1, \dots, 1).$

In the previous example, we have k = 4 and n(w) = (1, 0, 2, 0, 1). $I(w) = \{2, 3\}$, so :

$$R_k = \{(1, 2, 3, 4, 5), (1, 2, 3, 4, 4), (1, 1, 2, 3, 4), (1, 1, 2, 3, 3)\}$$

Theorem 5.5. Let $M \in M_{r,n}(\{a, b\})$ and let w be the associated word. Then

$$[w] = [\Psi(w)] = \sum_{\mathbf{r} \in R_k} (-1)^{h(\mathbf{r})} \cdot \mathbf{r}^{\mathbf{n}(w)+1}.$$

Proof. When we substitute b = (a + b) - a in $\Psi(w)$ and expand it into a sum of monomials in $\{a, a + b\}$, we get 2^p monomials where p is the number of appearances of b in $\Psi(w)$. Each monomial is obtained by substituting a or a + b in each instance of b in $\Psi(w)$.

We can index these monomials in the following way: Let $m = (a^{n_1}y_1a^{n_2}\cdots y_ka^{n_{k+1}})$ be a monomial appearing in the expansion of $\Psi(w)$, where $y_i \in \{a, a+b\}, y_i = x_i$ if $x_i = a + b$. Define $\mathbf{r} = (r_1, \ldots, r_{k+1}) \in R_k$ by: $r_1 = 1$ and:

$$r_{i+1} = \begin{cases} r_i & y_i = a \\ r_i + 1 & y_i = a + b \end{cases}$$

This gives us the 2^p vectors of R_k . Each vector can be seen as a lattice walk of length k where $r_{i+1} - r_i = 1$ corresponds to a vertical step while $r_{i+1} = r_i$ corresponds to a horizontal step. In this view, $h(\mathbf{r})$ is just the number of horizontal steps which is also the number of appearances of a in m. The sign of each monomial depends on the parity of $h(\mathbf{r})$ and by Lemma 5.1 each monomial contributes $\mathbf{r}^{\mathbf{n}(w)+1}$ to [w].

References

- [1] E. Bagno, D. Garber and R. Shwartz, *Recursions for excedance number in some permutations groups*, preprint (arXiv: math.CO/0702452).
- [2] R. Ehrenborg and E. Steingrimsson, The excedance set of a permutation, Adv. Appl. Math. 24 (2000), 284–299.
- [3] M. Fire, Statistics on wreath products, preprint (arXiv: math.CO/0409421).
- [4] E. Steingrimsson, Permutation statistics of indexed permutations, Europ. J. Combin. 15 (1994), 187–205.

THE JERUSALEM COLLEGE OF TECHNOLOGY, JERUSALEM, ISRAEL *E-mail address:* bagnoe@jct.ac.il

DEPARTMENT OF APPLIED MATHEMATICS, FACULTY OF SCIENCES, HOLON INSTITUTE OF TECHNOLOGY, PO BOX 305, 58102 HOLON, ISRAEL *E-mail address*: garber@hit.ac.il

Department of Mathematics, Bar-Ilan University, 52900 Ramat-Gan, Israel

E-mail address: shwartr1@macs.biu.ac.il