Entanglement spectrum in one-dimensional systems

Pasquale Calabrese¹ and Alexandre Lefevre²

¹Dipartimento di Fisica dell'Università di Pisa and INFN, Pisa, Italy.

²IPhT, Orme des Merisiers, CEA Saclay, 91191 Gif sur Yvette Cedex, France.

(Dated: October 22, 2018)

We derive the distribution of eigenvalues of the reduced density matrix of a block of length ℓ in a one-dimensional system in the scaling regime. The resulting "entanglement spectrum" is described by a universal scaling function depending only on the central charge of the underlying conformal field theory. This prediction is checked against exact results for the XX chain. We also show how the entanglement gap closes when ℓ is large.

PACS numbers:

The interest in quantifying the entanglement in extended quantum systems has been growing in recent times at an impressive rate, mainly because of its ability to detect the scaling behavior in proximity of quantum critical points (see e.g. Refs. [1] as reviews). Among the various measures, the so-called entanglement entropy has by far been the most studied. By partitioning an extended quantum system into two blocks, the entanglement entropy is defined as the von Neumann entropy of the reduced density matrix ρ_A of one of the two blocks. The success of this quantity can be understood because it is a single number able to capture the main features of the scaling behavior. In fact, in one-dimensional (1D) critical ground-states, when the block A is a segment of length ℓ in an infinite system, the entanglement entropy diverges with the logarithm of the block size as [2-4]

$$S_A \equiv -\operatorname{Tr} \rho_A \ln \rho_A = \frac{c}{3} \ln \ell + c_1', \qquad (1)$$

where c is the central charge of the associated conformal field theory (CFT) and c'_1 a non-universal constant. Away from the critical point, S_A saturates to a constant value [3] proportional to the logarithm of the correlation length [4].

However, the reduced density matrix (at least in principle) contains more information than the entanglement entropy. This information should be encoded in the full spectrum of the reduced density matrix, which we shortly call "entanglement spectrum", following Ref. [5]. In this letter we calculate the entanglement spectrum for 1D systems in the scaling regime, i.e. at or close to a quantum critical point. The study of the distribution of eigenvalues provides clearly a deeper theoretical understanding of entanglement and correlations in extended systems, but not only. Indeed, all numerical algorithms based on the so-called matrix product states (among which density matrix renormalization group [6] is the best known) give as first output a truncated spectrum of the reduced density matrix and from this all the other quantities are derived. Consequently, the knowledge of some scaling properties of this distribution provides an optimal check for the convergence and the accuracy of the numerics and could be used for putting accurate bounds on the efficiency of these methods as already done from other quantities in Refs. [7].

The scaling behavior of the entanglement spectrum can be related to the properties of the moments of the reduced density matrix: $R_{\alpha} \equiv \text{Tr} \rho_A^{\alpha} = \sum_i \lambda_i^{\alpha}$, where λ_i are the eigenvalues of ρ_A . In the scaling regime, R_{α} can be written as

$$R_{\alpha} = c_{\alpha} L_{\text{eff}}^{-c(\alpha - 1/\alpha)/6} \,, \tag{2}$$

where c_{α} is a non-universal constant and L_{eff} is the relevant length in the considered regime. For example, L_{eff} equals the length block ℓ/a if A is part of an infinite gapless system [2, 4], $L_{\text{eff}} = \frac{L}{a} \sin \frac{\pi \ell}{L}$ if A is in a finite gapless system of length L, and $L_{\text{eff}} = \xi/a$ [4] if the system is gapped [valid when the correlation length ξ (the inverse mass gap) is large, but smaller than all the other lengths like ℓ, L]. Everywhere a stands for the scale setting the microscopic length, e.g. the lattice spacing. The same α dependence is found in the case of open systems (the exponent of L_{eff} in Eq. (2) halves and so does S_A [4]) and when A consists of more disjoint intervals [4] and also in some non-equilibrium situations [8]. For practical reasons, it is convenient to write

$$R_{\alpha} = c_{\alpha} e^{-b(\alpha - 1/\alpha)}, \text{ with } b = \frac{c}{6} \ln L_{\text{eff}} > 0, \quad (3)$$

where by simple inspection b is related to the maximum eigenvalue: $b = -\ln \lambda_{\max}$, as well known [9].

In order to characterize the entanglement spectrum, we define the distribution of eigenvalues $P(\lambda) = \sum_{i} \delta(\lambda - \lambda)$ λ_i), which is normalized to m^{ℓ} , where m is the dimension of the local Hilbert space, e.g. m = 2 for spin 1/2systems. We determine this distribution for a 1D system in the scaling regime where Eq. (2) applies. The only assumption in what follows is that the α -dependence of the non-universal part of c_{α} can be ignored, i.e. that we can write $c_{\alpha} = a^{c(\alpha-1/\alpha)/6} f_{\alpha}$, with f_{α} a constant function (i.e. the effect of c_{α} is to replace the lengths appearing in R_{α} with the dimensionless quantity L_{eff}). A priori this can appear as a very crude approximation, but it is not the case. For the gapless XX chain, c_{α} is known analytically [10], and it is easy to show that f_{α} varies less then 1% as soon as $\alpha \geq 2$. Physically, this is equivalent to stating that the main contribution of c_{α} is to set the microscopic length a to be used in the continuum description in terms of the lattice one. We will check a

posteriori for the XX chain that the result obtained ignoring c_{α} describes accurately the actual entanglement spectrum when L_{eff} is large. A very precise numerical determination of c_{α} for the XXZ chain with $-1 < \Delta \leq 1$ [11] shows that the same scenario is true on the full critical line of the model. In few other cases c_{α} is also known [12–14] and similar properties can be found.

Thus, ignoring temporarily c_{α} , we will compute the entanglement spectrum, using the simple observation that $\lambda P(\lambda) = \lim_{\alpha} \operatorname{Im} f(\lambda - i\epsilon)$, with

$$f(z) = \frac{1}{\pi} \sum_{n=1}^{\infty} R_n z^{-n} = \frac{1}{\pi} \int d\lambda \frac{\lambda P(\lambda)}{z - \lambda}.$$
 (4)

Here, $f(\lambda - i\epsilon)$ has an imaginary part when $\epsilon \to 0$ only on the support of $P(\lambda)$, due to the pole in the r.h.s. of (4). The calculation of f(z) is straightforward:

$$f(z) = \frac{1}{\pi} \sum_{k=0}^{\infty} \frac{b^k}{k!} \sum_{n=1}^{\infty} \frac{(e^{-b}/z)^n}{n^k} = \frac{1}{\pi} \sum_{k=0}^{\infty} \frac{b^k}{k!} \mathrm{Li}_k(e^{-b}/z),$$
(5)

where $\operatorname{Li}_k(y)$ is the polylogarithm function, which is analytic on the complex plane, with a cut on the real axis for $y \geq 1$ (that once again is just $b = -\ln \lambda_{\max}$). The discontinuity along the cut is given by $\lim_{\epsilon \to 0} \operatorname{Im} \operatorname{Li}_k(y + i\epsilon) = \pi(\ln y)^{k-1}/\Gamma(k)$ for $k \geq 1$. The sum can be explicitly done and we end up with

$$P(\lambda) = \delta(\lambda_{\max} - \lambda) + \frac{b \theta(\lambda_{\max} - \lambda)}{\lambda \sqrt{b \ln(\lambda_{\max} / \lambda)}} I_1(2\sqrt{b \ln(\lambda_{\max} / \lambda)}), \quad (6)$$

where the delta peak comes from the contribution of k = 0 in (5) and $I_k(x)$ stands for the modified Bessel function of the first kind. In this derivation, only R_{α} with positive integers α enter. For these values (as we already discussed) c_{α} in general does not vary significantly. This gives an argument that explains why Eq. (6) works well for large enough L_{eff} in the following calculations for the XX model and why the same is expected in general.

 $P(e^{-t})$ can also be obtained by considering the inverse Laplace transform of R_{α} in the variable α . Using standard results Eq. (6) is easily recovered. We preferred to give the previous derivation because it highlights the role played by positive integer α , where c_{α} can be ignored. Conversely the inverse Laplace transform requires an integral on the complex plane, over a contour where there is less control on the values of α that are contributing relevantly. We noticed this because numerical inverse Laplace transform could be done in principle in more complicated cases (like those in Refs. [10–14]) where the previous analytic reasoning does not work.

Let us discuss now the main properties of $P(\lambda)$:

(i) The mean number of eigenvalues larger than a given λ is

$$n(\lambda) = \int_{\lambda}^{\lambda_{\max}} d\lambda P(\lambda) = I_0(2\sqrt{b\ln(\lambda_{\max}/\lambda)}).$$
 (7)

Note that for $\lambda \to 0$, $n(\lambda)$ diverges, as it should, because in the continuum the number of eigenvalues is infinite. In the lattice models, this can be regularized by the finite number of degrees of freedom (as e.g. in spin chains), but not always (e.g. bosons have always infinitely many eigenvalues).

(ii) The normalization $\sum \lambda_i = 1$ corresponds to $\int \lambda P(\lambda) = 1$, and follows directly from Eq. (6).

(iii) The entanglement entropy is given by

$$S = -\int_{0}^{\lambda_{\max}} \lambda \ln \lambda P(\lambda) d\lambda = -2 \ln \lambda_{\max} , \qquad (8)$$

reproducing the result that the single copy entanglement equals one-half of the entanglement entropy [9].

(iv) Majorization is a relation between two probability distributions $\lambda \equiv \{\lambda_i\}$ and $\mu \equiv \{\mu_i\}$ whose elements are ordered $\lambda_1 > \lambda_2 \cdots > \lambda_N$ (and similarly for μ): it is said that λ majorizes μ if $\sum_{i=1}^{M} \lambda_i \ge \sum_{i=1}^{M} \mu_i$ for any $M = 1, \ldots, N$ and $\sum_{i=1}^{N} \lambda_i = \sum_{i=1}^{N} \mu_i = 1$. It has been argued, observed numerically and in some instances proven analytically, that with increasing L_{eff} the resulting distribution of eigenvalues is majorized by the ones at smaller scaling lengths [15, 16] (sometimes this is referred to as majorization along renormalization group flow). From the previous result it is straightforward that majorization holds in the scaling regime when Eq. (6) for $P(\lambda)$ applies. In fact, we have

$$s(M) \equiv \sum_{i=1}^{M} \lambda_i \to \lambda_{\max} \left[1 + \int_0^{I_0^{-1}(M)} dy e^{-y^2/4b} I_1(y) \right],$$
(9)

which, at fixed M, is a monotonous function of λ_{\max} (that is a monotonous function of L_{eff}). This proves majorization in a very easy way. It is also simple to check majorization directly by making the numerical integral at fixed M and varying λ_{\max} .

We now compare the scaling function for the entanglement spectrum with the eigenvalues of lattice models, in order to show its predictivity and to highlight its limits. As a prototype of lattice models we consider the gapless XX chain defined by the Hamiltonian

$$H_{XX} = \sum_{i} [\sigma_{i}^{x} \sigma_{i+1}^{x} + \sigma_{i}^{y} \sigma_{i+1}^{y}], \qquad (10)$$

where $\sigma_i^{x,y}$ are the Pauli matrices at site *i*. The reduced density matrix of a block of ℓ contiguous spins in an infinite chain (so, among the cases of before, this corresponds to $L_{\text{eff}} = \ell/a$) can be obtained by exploiting the mapping to free fermions [3, 17]:

$$\rho_{\ell} \propto \exp\left[-\sum_{1 \le i, j \le \ell} h_{ij} c_i^{\dagger} c_j\right], \qquad (11)$$

where c_i are spinless fermion annihilation operators, and

$$h = \ln[(1-C)/C], \qquad C_{ij} = \frac{\sin \pi (i-j)/2}{\pi (i-j)}.$$
 (12)



FIG. 1: Sum of the first M eigenvalues of the XX model up to M = 100: 1 - s(M) as function of M for $\ell =$ 10, 100, 1000, 10000 (black dots). The red line is the conformal field theory prediction Eq. (9), in which λ_{\max} has been fixed to the maximum eigenvalue obtained numerically.

Calculating the eigenvalues ν_i of the matrix C_{ij} requires only the diagonalization of an $\ell \times \ell$ matrix. In terms of the ν_i , the 2^{ℓ} eigenvalues of the reduced density matrix are the products

$$\lambda_A = \prod_{i \in A, j \in B} \nu_i (1 - \nu_j), \qquad (13)$$

where A is a subset of $\{1, 2, \ldots, \ell\}$ and B the complement. While it is possible to obtain all the ν_i up to very large ℓ , making all of the 2^{ℓ} products to obtain the full spectrum requires far too much memory on a personal computer. Thus we calculate the full eigenvalue spec-



FIG. 2: Inverse function of λ_i (value of the *i*-th eigenvalue of the reduced density matrix) for $\ell = 10,100,1000$ for the XX chain. The plot is shown in terms of the scaling variable $2\sqrt{b \ln(\lambda_{\max}/\lambda)}$. The full line is the CFT prediction.



FIG. 3: The same as in Fig. 2 for all $10 \le \ell \le 28$. Even (odd) ℓ are shown on the left (right) panel.

trum only up to $\ell = 28$ (which corresponds to almost 300 millions of eigenvalues), and for larger ℓ we truncated the fermionic spectrum to 24 modes (i.e. we diagonalize the C matrix finding all the ν_i , but in the products we do not vary the $\ell - 24$ eigenvalues that are closer to 0 or 1 and generate the smaller λ_i). This reproduces in the exact order the first few thousands eigenvalues, to which we limit our analysis in the truncated cases.

We start this analysis from the check of the "sumrule" given by Eq. (9). In Fig. 1 we report 1 minus the sum of the first M (up to 100) eigenvalues for $\ell = 10, 100, 1000, 10000$. It is evident that increasing ℓ the sum is well described by Eq. (9) for larger and larger values of M. The (negative) deviations from the CFT take place at a value of M that roughly scales like $\ln \ell$ and they are obviously due to lattice effects, because the sum rule must be saturated by a finite sum up to 2^{ℓ} that cutoffs the integral of the continuum limit.

A stronger test of our predictions is provided by Eq. (7) that gives the total number of eigenvalues between λ and λ_{max} . This formula provides the natural scaling variable $x = 2\sqrt{b \ln(\lambda_{\max}/\lambda)}$. Once λ_{\max} is determined from the numerics, $n(\lambda)$ is just given by $I_0(x)$, independently from any other detail of the model under consideration and, more surprisingly, it is also independent from the model itself. Instead of plotting the number of eigenvalues larger than a given λ , we prefer to show the inverse function of λ_i that is the value of the *i*-th eigenvalue. In addition to represent $n(\lambda)$, this also has the advantage of giving information about the degeneracy of all eigenvalues, as the number of points at the same position on the horizontal axis. For $\ell = 10, 100, 1000$ this function is shown in Fig. 2 and compared with $n(\lambda)$ in Eq. (7). It is evident that when the spectrum becomes almost continuous, it is well described by the CFT prediction up to a given $\lambda_{\min}(\ell)$ (i.e. a given $x_{\max}(\ell)$). Increasing ℓ the number of eigenvalues described by Eq. (7) increases and at $\ell = 1000$, approximately thousands eigenvalues fall on the CFT scaling curve. The negative deviations for eigenvalues smaller than $\lambda_{\min}(\ell)$ are due to lattice effects as in the previous case of the partial sumrule. The degeneracies of the eigenvalues are not reproduced by our approach. In Fig. 3 we plot the entanglement spectrum

for all ℓ from 10 to 28 in order to show both finite-size and parity effects. When ℓ is odd (a case rarely considered in the literature) the approach to the asymptotic scaling is much slower. This can be understood because of the double degeneracy of *all* the eigenvalues (included the largest), which moves a large weight toward x = 0that, because of the sumrule, must be compensated by a smaller weight at large x (i.e. small eigenvalues).

There is a final interesting feature that is clearly visible in Figs. 2 and 3. When plotting in terms of $x = 2\sqrt{b \ln(\lambda_{\text{max}}/\lambda)}$, the first few eigenvalues do not change their positions with changing ℓ . The logarithms of these discrete eigenvalues result to be equispaced, i.e.

$$b\ln\frac{\lambda_{\mu}}{\lambda_{\nu}} \simeq k \Rightarrow \quad \frac{\lambda_{\nu}}{\lambda_{\mu}} \simeq e^{-\frac{6k}{\ln\ell/a}},$$
 (14)

where μ and ν are indices referring to two non-degenerate consecutive eigenvalues and k is a constant that is different for even and odd ℓ (and we used $b = -\ln \lambda_{\max} =$ $c/6 \ln \ell$ and c = 1). When $\lambda_{\mu} = \lambda_{\max}$, this simple formula tells us how the gap between the first and the second eigenvalues closes when increasing ℓ , which, following Ref. [5], we call "entanglement gap". This scaling of the discrete part of the spectrum can be related to an old result about the scaling of the eigenvalues of the corner transfer matrix [18]. In fact, because $b \ln(\lambda_{\mu}/\lambda_{\nu})$ is almost independent of ℓ , for this part of the spectrum the result should be the same as that of a segment ℓ in a system of length 2ℓ . In this case the reduced transfer matrix can be seen as the fourth power of a corner transfer matrix [4] (of angle $\pi/2$), for which the analogous of Eq. (14) has been derived from CFT [18]. Using this correspondence the asymptotic behavior of $P(\lambda \ll 1)$ for *qapped* systems has been already derived in Ref. [19]. However our result goes far beyond, explaining also the reason of such a simple result.

In conclusions, the main result of this letter is the an-

alytic derivation of the *universal* entanglement spectrum for 1D models in the scaling regime, given by Eq. (6). This turns out to depend only on the central charge of the underlying conformal field theory. We found that Eq. (6) describes accurately the continuum part of the spectrum of the XX chain for large enough L_{eff} , and is expected to work for any model at or close to a quantum critical point. This can be seen as a surprise because it means that the continuum part of the entanglement spectrum does not contain more information than the entanglement entropy that is just one possible average over the spectrum. The reason for this (maybe unexpected) result can be traced back to the fact that conformal invariance in 1D is so strong that it fixes completely the shape of the full spectrum, leaving only one parameter (the central charge) free. Such parameter can be fixed by of a single "measure" like the entanglement entropy, the largest eigenvalue, etc.

Oppositely, the discrete part of the spectrum is only reproduced in average by Eq. (6). As a consequence, the location and the degeneracies of the low-lying eigenvalues of the reduced density matrix can still be a tool for extracting further universal information on the model under investigation that could not be captured by the entanglement entropy. For example, it is known that the degeneracies of eigenvalues of the isotropic Heisenberg antiferromagnetic chain are larger than the XXZ ones (because of the larger symmetry), but they all have c = 1 and so the same continuum part of the spectrum. A careful study of these issues is needed, but it goes far beyond the goals of this letter.

Acknowledgments. This work started from a discussion with J. Cardy. We thank L. Tagliacozzo for pointing out the important Ref. [18]. We thank J. Cardy, A. Celi, S. Iblisdir, J. I. Latorre, B. Nienhuis, and L. Tagliacozzo for stimulating discussions. PC benefited of a travel grant from ESF (INSTANS activity).

- L. Amico et al., Rev. Mod. Phys. 80, 517 (2008); J. Cardy, Eur. Phys. J. B, to appear.
- [2] C. Holzhey, F. Larsen, and F. Wilczek, Nucl. Phys. B 424, 443 (1994).
- [3] G. Vidal et al., Phys. Rev. Lett. 90, 227902 (2003); J. I.
 Latorre et al., Quant. Inf. and Comp. 4, 048 (2004).
- [4] P. Calabrese and J. Cardy, J. Stat. Mech. P06002 (2004); Int. J. Quant. Inf. 4, 429 (2006).
- [5] H. Li and F. D. M. Haldane, Phys. Rev. Lett. 101, 010504 (2008).
- [6] S. White, Phys. Rev. Lett. **69**, 2863 (1992); U. Schollwoeck, Rev. Mod. Phys. **77**, 259 (2005).
- [7] N. Schuch et al., Phys. Rev. Lett. 100, 030504 (2008); D.
 Perez-Garcia et al., Quantum Inf. Comput. 7, 401 (2007);
 L. Tagliacozzo et al., Phys. Rev. B 78, 024410 (2008).
- [8] P. Calabrese and J. Cardy, J. Stat. Mech. P04010 (2005);
 J. Stat. Mech. P10004 (2007).
- [9] J. Eisert and M. Cramer, Phys. Rev. A **72**, 042112 (2005); I. Peschel and J. Zhao, J. Stat. Mech. P11002

(2005); R. Orus et al., Phys. Rev. A 73, 060303 (2006).

- [10] B.-Q. Jin, V. E. Korepin, J. Stat. Phys. 116, 79 (2004).
- [11] P. Calabrese, M. Campostrini, B. Nienhuis, to appear.
- [12] H. Casini and M. Huerta, J. Stat. Mech. P05012 (2005);
 H. Casini et al., J. Stat. Mech. P05007 (2005).
- [13] J. L. Cardy et al., J. Stat. Phys. 130, 129 (2008).
- [14] F. Franchini et al., J. Phys. A **41**, 025302 (2008).
- [15] J. I. Latorre et al., Phys. Rev. A **71**, 034301 (2005); R. Orus, Phys. Rev. A **71**, 052327 (2005); **73**, 019904(E) (2006).
- [16] H.-Q. Zhou et al., Phys. Rev. A 74, 050305(R) (2006).
- [17] I. Peschel, J. Stat. Mech. P06004 (2004); I. Peschel, J. Phys. A **36**, L205 (2003); M.-C. Chung and I. Peschel, Phys. Rev. B **64**, 064412 (2001); I. Peschel et al., Ann. Physik (Leipzig) **8**, 153 (1999).
- [18] I. Peschel and T. T. Truong, Z. Phys. B 69, 385 (1987).
- [19] K. Okunishi et al., Phys. Rev. E 59, R6227 (1999).