A concept of Kinetic Energy in Quantum Mechanics should be corrected

Yuri Kornyushin

Maître Jean Brunschvig Research Unit, Chalet Shalva, Randogne, CH-3975

A concept of Kinetic Energy in Quantum Mechanics is analyzed. It is shown that what is called Kinetic Energy in Non-Relativistic Quantum Mechanics is a more general concept, which includes Kinetic Energy. It is really Inhomogeneity Quantum Energy. It is not zero in many cases where there are no motion and flux.

1. Introduction

Let us write down Schrödinger equation for a free particle [1]:

$$-(\hbar^2/2m)\Delta\psi = E\psi,\tag{1}$$

where \hbar is Planck constant, divided by 2π , *m* is the mass of a particle, Δ is Laplace operator, ψ is a wave function, and *E* is the energy of the stationary state of a particle. A well-known solution of Eq. (1) is a plane wave [1], $\psi(x) = \text{const}e^{ikx}$, where *i* is the imaginary unit, and *k* is the wave vector. This solution corresponds to a non-localized free particle. The energy of this particle is well known to be $E(k) = \hbar^2 k^2/2m$ [1]. This is a Kinetic Energy of a non-localized particle.

2. Kinetic Energy in general case

Kinetic Energy is the energy of a motion [1]. A matter flux and electric current accompany the motion of a charged matter. No motion without a flux of a matter exists. In the ground state of any particle the wave function is real, it contains no imaginary part [1]. Real wave function yields no flux and no electric current for a charged particle [1]. So the question arises: *is not the Kinetic Energy of a particle in a ground state zero, or is it?* As an example let us take an electron in a ground state of a hydrogen atom. Corresponding wave function is [1] $\psi(r) = (g^3/\pi)^{1/2} \exp{-gr}$, where $g = me^2/\hbar^2$ and *r* is the distance from proton [1].

The operator of the momentum is $-i\hbar\nabla$, where ∇ is gradient operator. The Kinetic Energy *T* operator is considered to be the operator of the momentum in square, divided by 2m [1]. This operator $-(\hbar^2/2m)\Delta$, averaged over the whole space, is regarded to be an averaged Kinetic Energy [1]. From this follows that for the electron in a hydrogen atom $T = me^4/2\hbar^2$. So we have here a very essential Kinetic Energy without any motion. This casts some doubt upon that the operator $-(\hbar^2/2m)\Delta$ is really and always the Kinetic Energy operator [2].

There is another question: does the averaged over the whole space operator $-(\hbar^2/2m)\Delta$ has to be positive? It is assumed tacitly that this is the case. It may be not always so [2].

The operator $-(\hbar^2/2m)\Delta$, averaged over the whole space gives some energy. This energy is not zero because the wave function in inhomogeneous; it depends on coordinates. So the regarded energy can be better called the *Inhomogeneity Quantum Energy*. This energy is a real part of a total energy, not the *Kinetic Energy*. As is well known, the *Inhomogeneity Quantum Energy* really exists and it is a subject of the Energy Conservation Law. Just it should not be called a Kinetic Energy, because it is not. *Kinetic Energy* is some integral of something, proportional to the square of a flux. It is hence zero for any ground state. And it is really of no interest.

4. Discussion

The concept of *Kinetic Energy* in Non-Relativistic Quantum Mechanics should be reconsidered, as presently used concept gives non-zero values for many cases with no motion and flux. Instead a concept of *Inhomogeneity Quantum Energy* should be used.

References

L. D. Landau and E. M. Lifshitz, *Quantum Mechanics*, Fizmatgiz, Moscow, 1963.
Yuri Kornyushin, <u>http://arxiv.org/abs/0806.3232v1</u>.