Yuichi Itto · Sumiyoshi Abe

On the Geodesic Nature of Wegner's Flow

Received: date / Accepted: date

Abstract Wegner's method of flow equations offers a useful tool for diagonalizing a given Hamiltonian and is widely used in various branches of quantum physics. Here, generalizing this method, a condition is derived, under which the corresponding flow of a quantum state becomes geodesic in a submanifold of the projective Hilbert space, independently of specific initial conditions. This implies the geometric optimality of the present method as an algorithm of generating stationary states. The result is illustrated by analyzing some physical examples.

Keywords Wegner's method of flow equations · Diagonalization of Hamiltonian · Geodesic curve

1 Introduction

In recent years, much attention has been focused on Wegner's method of flow equations [1] (for reviews, see [2]). This method offers a powerful tool for diagonalizing a given quantum-mechanical Hamiltonian, which has been devised in analogy with the theory of renormalization groups and has widely been applied to a variety of problems in condensed matter physics, nuclear and particle physics, and quantum information. Examples include the effective Hamiltonian of the Anderson impurity model [3], a Dirac particle in an external electromagnetic field [4], an effective spin-spin coupling arising from spin-phonon chains [5], the t-t' Hubbard model for high-temperature superconductivity [6], the Tomonaga-Luttinger model for interacting spinless electrons in one dimension [7], localized superfluidity [8], the Lipkin-Meshkov-Glick model in nuclear physics [9], quantum phase transition in the interacting boson model [10], light-cone quantum chromodynamics [11], electron-electron and electron-phonon interactions in the Hubbard-Holstein model [12], and stationary photon-atom entanglement [13].

Wegner's method employs a continuous unitary transformation represented by the operator, U(l), where $l \in [0,\infty)$ is referred to as the flow parameter. U(l) transforms the original Hamiltonian H = H(0) to $H(l) = U(l)HU^{\dagger}(l)$, which satisfies $dH(l)/dl = [\eta(l), H(l)]$, where $\eta(l)$ is the anti-Hermitian generator given by $\eta(l) = [dU(l)/dl]U^{\dagger}(l)$. Wegner's choice for $\eta(l)$ to diagonalize (or, block-diagonalize) the Hamiltonian reads

$$\eta(l) = \eta^{W}(l) \equiv [H_{d}(l), H_{o-d}(l)],$$
(1)

where $H_{\rm d}(l)$ and $H_{\rm o-d}(l)$ stand for the diagonal and off-diagonal parts of H(l), respectively. It can be shown [1,2] that $H_{\rm o-d}(l)$ tends to vanish in the limit $l \to \infty$. Mathematically, this procedure corresponds to the Jacobi

S.A. was supported in part by a Grant-in-Aid for Scientific Research from the Japan Society for the Promotion of Science.

Y. Itto

Science Division, Center for General Education, Aichi Institute of Technology, Aichi 470-0392, Japan E-mail: itto@aitech.ac.jp

S. Abe

Department of Physical Engineering, Mie University, Mie 514-8507, Japan

E-mail: suabe@sf6.so-net.ne.jp

algorithm for eigenvalue problems, which defines a steepest-descent flow in the space of matrices with the Frobenius norm [14].

Although Wegner's method has been applied to many problems as mentioned above, its inherent properties do not seem to be explored well. It is our opinion that this method is more than just a mathematical tool for diagonalizing a Hamiltonian. The purpose of this paper is to reveal an interesting geometric property hidden behind Wegner's flow equations. For it, we first generalize Wegner's method itself. The original method of Wegner is a special case within this new framework. Next, we consider the flow of a given quantum state through a submanifold of the projective Hilbert space composed of rays explained below. Then, we find a condition, under which the flow becomes geodesic in the submanifold, independently of specific "initial" conditions at l=0. This remarkable property is illustrated by analyzing some physical examples. Our discussion may also have significance for laboratory experiments, since one often need realize a stationary state from an initially prepared state via quantum operations [13], and in this context the unitary transformation in Wegner's method is now shown to offer the optimal strategy for it in a certain class of systems (satisfying the condition discovered here).

2 Generalization of Wegner's Method of Flow Equations

Let us start our discussion with considering a quantum system in a d-dimensional Hilbert space, where d is either finite or infinite. Its unitary-transformed Hamiltonian, H(l), is decomposed into two parts: H(l) $H_{\rm d}(l)+H_{\rm o-d}(l)$, where $H_{\rm d}(l)$ and $H_{\rm o-d}(l)=\sum_{a=1}^AH_{\rm o-d}^{(a)}(l)$ $(1\leq A\leq d-1)$ are the diagonal and off-diagonal parts, respectively. Using the complete set of the normalized eigenstates of $H_{\rm d}(l)$, $\{\mid u_n\rangle\}_{n=1,2,\ldots,d}$, which is a natural basis such as the Fock basis of the harmonic oscillator, we write them in the following forms:

$$H_{\rm d}(l) = \sum_{n=1}^{d} \varepsilon_n(l) \mid u_n \rangle \langle u_n \mid, \tag{2}$$

$$H_{\text{o-d}}^{(a)}(l) = \sum_{n=1}^{d} \left[C_{n+i_a}^{(a)}(l) \mid u_{n+i_a} \rangle \langle u_n \mid + C_{n+i_a}^{(a)*}(l) \mid u_n \rangle \langle u_{n+i_a} \mid \right], \tag{3}$$

where $\varepsilon_n(l)$ is the nth eigenvalue of $H_{\rm d}(l)$ and $C_{n+i_a}^{(a)}(l)$'s are the complex expansion coefficients. The nonzero index i_a describes the off-diagonality and is ordered without loss of generality as follows: $0 < i_1 < i_2 < \cdots < i_A$. It is understood that $C_{n+i_a}^{(a)}(l) = 0$ ($C_{n-i_a}^{(a)}(l) = 0$), if $n+i_a > d$ ($n-i_a \le 1$). Here, we generalize Wegner's method as follows. Instead of taking the whole of $H_{\text{o-d}}(l)$, we employ only

a single term, say $H_{o-1}^{(a)}(l)$:

$$\eta^{(a)}(l) = \left[H_{d}(l), H_{o-d}^{(a)}(l) \right].$$
(4)

Clearly, Wegner's choice is $\eta^W(l) = \sum_{a=1}^A \eta^{(a)}(l)$. It can be found after a straightforward calculation using Eqs. (2)-(4) that the corresponding generalized flow equation

$$\frac{dH(l)}{dl} = \left[\eta^{(a)}(l), H(l)\right] \tag{5}$$

gives rise to

$$\frac{d}{dl} \sum_{m,n=1}^{d} |\langle u_m | H(l) | u_n \rangle|^2 = -\frac{d}{dl} \sum_{n=1}^{d} \varepsilon_n^2(l)$$

$$= -4 \sum_{n=1}^{d} \left[\varepsilon_{n+i_a}(l) - \varepsilon_n(l) \right]^2 \left| C_{n+i_a}^{(a)}(l) \right|^2, \tag{6}$$

implying that the off-diagonal elements of H(l) tend to decay as l increases. (The first equality in Eq. (6) comes from the fact that $(d/dl) \text{Tr}[H^2(l)] = 0$.) η^W is obtained from the set $\{\eta^{(a)}\}_{a=1,2,\dots,A}$, but the reverse is obviously not possible if $A \ge 2$. In this sense, the present method is regarded as a generalization of Wegner's. Closing this section, we emphasize the following point. Like in Wegner's method, reduction of off-diagonal elements of a Hamiltonian does not necessarily mean exact vanishing of the elements. In Wegner's case, it tends to be block-diagonalization of a Hamiltonian, in general, whereas Eq. (6) implies reduction of *targeted* off-diagonal elements, not all off-diagonal elements. The present generalized method enables one to look into the details of the structure of a Hamiltonian.

3 Condition for Geodesic Flow

Next, let us translate the flow of the Hamiltonian into the flow of a state. Given a normalized state $|\psi\rangle$, the stationary Schrödinger equation reads $H |\psi(\infty)\rangle = E |\psi(\infty)\rangle$, where $|\psi(l)\rangle = U^{\dagger}(l) |\psi\rangle$. As well known, two states different from each other only by total phases are equivalent in quantum mechanics, and therefore a physical state is represented by a ray. Accordingly, the quantum-state space is the projective Hilbert space that is generically a curved space. Eq. (5) determines the flows of the physical coefficients contained in the Hamiltonian, which depend on the parameters appearing in the unitary operator. Let us explicitly write as follows: $U = U(\alpha)$, $|\psi(\alpha)\rangle = U^{\dagger}(\alpha) |\psi\rangle$, where $\alpha \equiv (\alpha^1, \alpha^2, \dots, \alpha^k)$. The set of parameters, α , defines a local coordinate on the submanifold of the projective Hilbert space. This submanifold is referred to as the quantum evolution submanifold [15], on which a quantum state flows. Then, the Fubini-Study metric [15–17] induced on this submanifold is, up to the second-order infinitesimals, given by

$$ds^{2} = 1 - |\langle \psi(\alpha) | \psi(\alpha + d\alpha) \rangle|^{2} \equiv g_{ij}(\alpha) d\alpha^{i} d\alpha^{j}, \tag{7}$$

where the metric tensor is expressed in terms of the anti-Hermitian operator $G_i(\alpha) = [\partial_i U(\alpha)]U^{\dagger}(\alpha)$ ($\partial_i = \partial/\partial \alpha^i; i = 1, 2, ..., k$) as follows:

$$g_{ij}(\alpha) = -\frac{1}{2} \langle \psi \mid G_i(\alpha) G_j(\alpha) + G_j(\alpha) G_i(\alpha) \mid \psi \rangle + \langle \psi \mid G_i(\alpha) \mid \psi \rangle \langle \psi \mid G_j(\alpha) \mid \psi \rangle. \tag{8}$$

Here and hereafter, Einstein's convention is understood for the repeated indices. The equation for a geodesic curve parametrized by the arc length, $\alpha(s)$, is given by $d^2\alpha^h/ds^2 + \Gamma^h_{ij}(d\alpha^i/ds)(d\alpha^j/ds) = 0$, where Γ^h_{ij} is Christoffel's symbol defined by $\Gamma_{hij} = g_{hl}\Gamma^l_{ij} = (1/2)\left(\partial_i g_{hj} + \partial_j g_{ih} - \partial_h g_{ij}\right)$.

Let us parametrize the curve by the flow parameter, instead of the arc length, i.e., $\alpha(l)$, and consider the functional

$$S[\alpha] = \int_{l_1}^{l_2} dl \ L(\alpha, \dot{\alpha}), \tag{9}$$

where

$$L(\alpha, \dot{\alpha}) = \frac{ds}{dl} = \sqrt{g_{ij}(\alpha)\dot{\alpha}^i\dot{\alpha}^j}$$
 (10)

and $\dot{\alpha}^i \equiv d\alpha^i(l)/dl$. The quantity in Eq. (9) is the arc length in the interval $[l_1, l_2]$. The variation of S with respect to $\alpha(l)$ is calculated to be

$$\frac{\delta S[\alpha]}{\delta \alpha^i} = -\frac{1}{4L^3} X_i,\tag{11}$$

where

$$X_{i} = \left(2\langle\psi\mid\eta\mid\psi\rangle\langle\psi\mid\frac{\partial\eta}{\partial\dot{\alpha}^{i}}\mid\psi\rangle - \langle\psi\mid\frac{\partial\eta^{2}}{\partial\dot{\alpha}^{i}}\mid\psi\rangle\right)\frac{d}{dl}\left(\langle\psi\mid\eta^{2}\mid\psi\rangle - \langle\psi\mid\eta\mid\psi\rangle^{2}\right)$$

$$+2\left(\langle\psi\mid\left\{\frac{d\eta}{dl},\frac{\partial\eta}{\partial\dot{\alpha}^{i}}\right\} + \left[\eta^{2},\frac{\partial\eta}{\partial\dot{\alpha}^{i}}\right]\mid\psi\rangle - 2\langle\psi\mid\frac{d\eta}{dl}\mid\psi\rangle\langle\psi\mid\frac{\partial\eta}{\partial\dot{\alpha}^{i}}\mid\psi\rangle\right)$$

$$-2\langle\psi\mid\eta\mid\psi\rangle\langle\psi\mid\left[\eta,\frac{\partial\eta}{\partial\dot{\alpha}^{i}}\right]\mid\psi\rangle\right)\left(\langle\psi\mid\eta^{2}\mid\psi\rangle - \langle\psi\mid\eta\mid\psi\rangle^{2}\right)$$
(12)

with

$$\eta = \eta(\alpha(l)) = \frac{dU(\alpha(l))}{dl} U^{\dagger}(\alpha(l)) = G_i(\alpha(l)) \dot{\alpha}^i, \tag{13}$$

provided that the following relation has been used:

$$\frac{d}{dl}\left(\frac{\partial \eta}{\partial \dot{\alpha}^{i}}\right) = \frac{\partial \eta}{\partial \alpha^{i}} + \left[\eta, \frac{\partial \eta}{\partial \dot{\alpha}^{i}}\right]. \tag{14}$$

Up to this stage, we have purely discussed the variational problem for the functional in Eq. (9), which is naturally assumed to be analytic, and have not used the flow equation yet.

We are now going to prove that under a certain condition the generalized flow equation (5) makes Eq. (12) vanish and defines a geodesic curve associated with the "initial state" $|\psi\rangle = |u_n\rangle$, which is an eigenstate of $H_{\rm d}(l)$. (This is the state of relevance, because we are considering the flow to an exact stationary state.) That is, we are going to show that, under the condition found later, the above X_i vanishes for $\eta = \eta^{(a)}$ in Eq. (4). Clearly, $\langle u_n | \eta^{(a)} | u_n \rangle$ and its derivatives with respect to l and $\dot{\alpha}^i$ vanish. Furthermore, it can be shown by

Clearly, $\langle u_n \mid \eta^{(a)} \mid u_n \rangle$ and its derivatives with respect to l and $\dot{\alpha}^i$ vanish. Furthermore, it can be shown by using Eqs. (2)-(4) that $\langle u_n \mid \left[\eta^{(a)^2}, \, \partial \eta^{(a)} / \partial \dot{\alpha}^i \right] \mid u_n \rangle$ also vanishes. Therefore, X_i for $\mid \psi \rangle = \mid u_n \rangle$ and $\eta = \eta^{(a)}$ is reduced to

$$X_{i} = 2\langle u_{n} \mid \eta^{(a)^{2}} \mid u_{n} \rangle \langle u_{n} \mid \left\{ \frac{d\eta^{(a)}}{dl}, \frac{\partial \eta^{(a)}}{\partial \dot{\alpha}^{i}} \right\} \mid u_{n} \rangle - \langle u_{n} \mid \frac{d\eta^{(a)^{2}}}{dl} \mid u_{n} \rangle \langle u_{n} \mid \frac{\partial \eta^{(a)^{2}}}{\partial \dot{\alpha}^{i}} \mid u_{n} \rangle. \tag{15}$$

This equation can be rewritten by using Eqs. (2)-(4) again as follows:

$$X_{i} = 4\left(\left|D_{n+i_{a}}^{(a)}\right|^{2} + \left|D_{n}^{(a)}\right|^{2} \sum_{m=1}^{d} \delta_{n,m+i_{a}}\right) \operatorname{Re}\left[\frac{dD_{n+i_{a}}^{(a)}}{dl} \frac{\partial D_{n+i_{a}}^{(a)^{*}}}{\partial \dot{\alpha}^{i}} + \frac{dD_{n}^{(a)}}{dl} \frac{\partial D_{n}^{(a)^{*}}}{\partial \dot{\alpha}^{i}} \sum_{m=1}^{d} \delta_{n,m+i_{a}}\right] - \left(\frac{d\left|D_{n+i_{a}}^{(a)}\right|^{2}}{dl} + \frac{d\left|D_{n}^{(a)}\right|^{2}}{dl} \sum_{m=1}^{d} \delta_{n,m+i_{a}}\right) \left(\frac{\partial\left|D_{n+i_{a}}^{(a)}\right|^{2}}{\partial \dot{\alpha}^{i}} + \frac{\partial\left|D_{n}^{(a)}\right|^{2}}{\partial \dot{\alpha}^{i}} \sum_{m=1}^{d} \delta_{n,m+i_{a}}\right),$$
(16)

where $D_{n+i_a}^{(a)} \equiv \left(\varepsilon_{n+i_a}(l) - \varepsilon_n(l)\right) C_{n+i_a}^{(a)}(l)$. The *l*-derivatives appearing in this expression should be calculated from the generalized flow equation in Eq. (5). It is also noted that α^i and $\dot{\alpha}^i$ are not independent in Eqs. (15) and (16) any more, since Eq. (4) has already been used.

Now, we present our discovery. If the labels a and a' satisfying

$$i_{a'} \neq 2i_a, 3i_a \quad (1 \le a' \le A) \tag{17}$$

can be taken for $\eta^{(a)}(l)$, then sandwiching the flow equation in Eq. (5) by three pairs, $\langle u_n \mid$ and $\mid u_{n-i_a} \rangle$, $\langle u_{n+i_a} \mid$ and $\mid u_n \rangle$, $\langle u_{n+i_a} \mid$ and $\mid u_{n-i_a} \rangle$, we have

$$\langle u_n \mid \left[\eta^{(a)}(l), \sum_{a'=1}^A H_{\text{o-d}}^{(a')}(l) \right] \mid u_{n-i_a} \rangle = 0,$$
 (18)

$$\langle u_{n+i_a} \mid \left[\eta^{(a)}(l), \sum_{a'=1}^{A} H_{\text{o-d}}^{(a')}(l) \right] \mid u_n \rangle = 0,$$
 (19)

$$\langle u_{n+i_a} | \left[\eta^{(a)}(l), \sum_{a'=1, a' \neq a}^{A} H_{\text{o-d}}^{(a')}(l) \right] | u_{n-i_a} \rangle = 0,$$
 (20)

respectively. And, correspondingly, we obtain the following sandwiched flow equations:

$$\frac{dC_n^{(a)}(l)}{dl} = -\left(\varepsilon_n(l) - \varepsilon_{n-i_a}(l)\right)^2 C_n^{(a)}(l),\tag{21}$$

$$\frac{dC_{n+i_a}^{(a)}(l)}{dl} = -\left(\varepsilon_{n+i_a}(l) - \varepsilon_n(l)\right)^2 C_{n+i_a}^{(a)}(l),\tag{22}$$

$$\left(\varepsilon_{n+i_a}(l) + \varepsilon_{n-i_a}(l) - 2\varepsilon_n(l)\right) C_n^{(a)}(l) C_{n+i_a}^{(a)}(l) = 0.$$
(23)

It is mentioned that, from Eqs. (21) and (22), the phases of $C_n^{(a)}(l)$ and $C_{n+l}^{(a)}(l)$ are found to be independent of l.

Case-A: If $C_n^{(a)}(l) = C_{n+i_a}^{(a)}(l) = 0$, then X_i in Eq. (16) obviously vanishes.

Case-B: If $C_n^{(a)}(l) = 0$ and $C_{n+i_a}^{(a)}(l) \neq 0$ (or, $C_n^{(a)}(l) \neq 0$ and $C_{n+i_a}^{(a)}(l) = 0$), then X_i vanishes, due to the fact that the phases of $D_n^{(a)}(l)$ and $D_{n+i_a}^{(a)}(l)$ are independent of l.

Case-C: If both $C_n^{(a)}(l)$ and $C_{n+i_a}^{(a)}(l)$ are nonzero, then Eq. (23) yields $\varepsilon_{n+i_a}(l) + \varepsilon_{n-i_a}(l) - 2\varepsilon_n(l) = 0$.

Combining the relation in Case-C with Eqs. (21) and (22), we obtain a crucial result that $\left|C_{n+i_a}^{(a)}(l)\right|$ is proportional to $|C_n^{(a)}(l)|$. And, this makes X_i vanish again.

Therefore, we conclude that the flow equation with the generator $\eta^{(a)}(l)$ with the label a satisfying the condition in Eq. (17) gives rise to the geodesic flow of $|u_n\rangle$, independently of a specific initial condition. This is the main result of the present work.

We emphasize that Eq. (17) does not necessarily lead to the geodesic nature of Wegner's flow generated by $\eta^W \equiv \sum_{a=1}^A \eta^{(a)}$: it guarantees the geodesic nature of the flow generated by $\eta^{(a)}$. However, according to our experience, there exist physically important examples, in which A=1, that is, the present method becomes reduced to Wegner's. So, in such a case, Eq. (17) works for establishing the geodesic nature of Wegner's flow. In the next section, some of such examples are discussed.

4 Physical Examples

In what follows, we illustrate our result by analyzing some physical examples. The first example we consider is the generalized harmonic oscillator. The Hamiltonian reads

$$H = \omega a^{\dagger} a + \lambda a^{\dagger^{2}} + \lambda^{*} a^{2} + \mu a^{\dagger} + \mu^{*} a + \nu.$$
 (24)

Here, a^{\dagger} and a are the creation and annihilation operators obeying the algebra: $[a, a^{\dagger}] = 1$, $[a^{\dagger}, a^{\dagger}] = [a, a] = 0$. ω (>0) and ν are real constants, whereas λ and μ are complex. In particular, the condition, $\omega > 2|\lambda|$, is to be satisfied. Here and hereafter, \hbar is set equal to unity. We analyze the following two cases: (i) $\mu \neq 0$, $\lambda = 0$ and (ii) $\mu = 0$, $\lambda \neq 0$. [In the case when both μ and λ are nonzero, one can first eliminate the terms linear in a^{\dagger} and a by the unitary transformation with the displacement operator, $\exp(\alpha a^{\dagger} - \alpha^* a)$, where α is a complex c-number variable satisfying $\omega \alpha + 2\lambda \alpha^* - \mu = 0$. Therefore, this case is reduced to (ii).] In case (i), the Hamiltonian is transformed by the displacement operator

$$U(l) = \exp\left[z(l)a^{\dagger} - z^{*}(l)a\right]$$
(25)

with the complex z(l) satisfying z(0) = 0. For the normalized ground state, $|0\rangle$, satisfying $a |0\rangle = 0$, as the "initial state" at l = 0, the flow of the state is along the coherent state. The corresponding Fubini-Study metric is Euclidean [15]:

$$ds^2 = \frac{1}{2}(dx^2 + dp^2) \tag{26}$$

with the parametrization, $z = (x+ip)/\sqrt{2}$. Then, the condition in Eq. (17) is satisfied, and the flow is, in fact, geodesic: a straight line in the space with a global coordinate (x, p). In case (ii), the squeezing operator

$$U(l) = \exp\left[\frac{1}{2}\left(\xi(l)a^{\dagger^{2}} - \xi^{*}(l)a^{2}\right)\right]$$
 (27)

is to be considered, where the complex coefficient $\xi(l)$ is parametrized as $\xi(l)=r(l)e^{-2i\phi(l)}$ $(0\leq r(l),0\leq \phi(l)<2\pi)$. Accordingly, $(\alpha^1,\alpha^2)\equiv (r,\phi)$. The condition U(0)=I (with the identity operator I) leads to r(0) = 0. The initial state is taken to be the number state, $|n\rangle = (n!)^{-1/2} a^{\dagger^n} |0\rangle$. The corresponding metric is [15]:

$$ds^{2} = \frac{1}{2}(n^{2} + n + 1)\left[dr^{2} + \left(\sinh^{2}2r\right)d\phi^{2}\right],$$
(28)

which shows that the manifold is the Lobachevsky space. The transformed Hamiltonian is written as follows:

$$H(l) = \omega(l)a^{\dagger}a + \lambda(l)a^{\dagger^{2}} + \lambda^{*}(l)a^{2} + v(l).$$
(29)

The coefficients appearing here depend not only on their original values at l=0 but also on $\xi(l)$. In this case,

$$H_{\text{o-d}}(l) = \lambda(l)a^{\dagger^2} + \lambda^*(l)a^2$$
 (30)

is the one and only off-diagonal part. Therefore, the generator in Eq. (4) is identical to Wegner's choice

$$\eta^{W}(l) = 2\omega(l) \left[\lambda(l)a^{\dagger^{2}} - \lambda^{*}(l)a^{2} \right], \tag{31}$$

and the condition in Eq. (17) is automatically fulfilled. The flow equation for $\lambda(l)$ is given by

$$\frac{d\lambda(l)}{dl} = -4\omega^2(l)\lambda(l),\tag{32}$$

showing that the phase of $\lambda(l)$ does not depend on l. Then, one can explicitly find that Eq. (15) indeed vanishes. Comparing $[dU(l)/dl]U^{\dagger}(l)$ with $\eta^W(l)$, we obtain $\phi(l)=$ const, which in fact turns out to make both $\delta S/\delta r(l)$ and $\delta S/\delta \phi(l)$ vanish, where S is the arc-length functional defined in terms of the above metric. We also mention that the spectrum of $H_{\rm d}(l)=\omega(l)a^{\dagger}a+v(l)$ is equally spaced, and accordingly Case-C in the preceding section is realized. Thus, Wegner's flow is geodesic.

The second example is a spin-s, $\mathbf{S} = (S_x, S_y, S_z)$, in a constant external magnetic field \mathbf{B} . The Hamiltonian reads

$$H = \mathbf{S} \cdot \mathbf{B} \tag{33}$$

in an appropriate unit. The unitary operator to be considered is

$$U(l) = \exp[\sigma(l)S_{+} - \sigma^{*}S_{-}], \tag{34}$$

where $S_{\pm} \equiv S_x \pm iS_y$ and the complex coefficient $\sigma(l)$ is parametrized as $\sigma(l) = [\theta(l)/2]e^{-i\varphi(l)}$ ($0 \le \theta(l) < \pi, 0 \le \varphi(l) < 2\pi$) with $\theta(0) = 0$. The basic commutation relations satisfied by the spin operators are as follows: $[S_z, S_{\pm}] = \pm S_{\pm}, [S_+, S_-] = 2S_z$. The local coordinate is given by $(\alpha^1, \alpha^2) \equiv (\theta, \varphi)$. The initial state is taken to be $|m\rangle_s$ ($m = -s, -s + 1, \ldots, 0, \ldots, s - 1, s$), which satisfies $S_z |m\rangle_s = m |m\rangle_s$. The metric is found to be given by [15]:

$$ds^{2} = \frac{1}{2}(s^{2} + s - m^{2}) \left[d\theta^{2} + (\sin^{2}\theta) d\varphi^{2} \right], \tag{35}$$

which is of a sphere. The transformed Hamiltonian is written as

$$H(l) = \beta_{z}(l)S_{z} + \beta(l)S_{+} + \beta^{*}(l)S_{-}.$$
(36)

 $\beta_z(l)$ is real, whereas $\beta(l)$ is complex. They depend not only on **B** but also on $\sigma(l)$. The one and only off-diagonal part is

$$H_{\text{o-d}}(l) = \beta(l)S_{+} + \beta^{*}(l)S_{-},$$
 (37)

and so Wegner's choice

$$\eta^{W}(l) = \beta_{z}(l) \left[\beta(l) S_{+} - \beta^{*}(l) S_{-} \right]$$
(38)

is employed. Therefore, clearly the condition in Eq. (17) is fulfilled. The flow equation for $\beta(l)$ is

$$\frac{d\beta(l)}{dl} = -\beta_z^2(l)\beta(l),\tag{39}$$

from which the phase of $\beta(l)$ is seen to be independent of l, and accordingly Eq. (15) vanishes. Comparison of the above generator, $\eta^W(l)$, with $\left[dU(l)/dl\right]U^\dagger(l)$ yields $\varphi(l)=$ const, which leads to the fact that the variations of the arc-length functional (S, calculated by using the above metric) with respect to $\theta(l)$ and $\varphi(l)$ vanish. Also, the spectrum of $H_{\rm d}(l)=\beta_z(l)S_z$ with $s\geq 1$ is equally spaced, and so Case-C in the preceding section is realized. Thus, Wegner's flow is geodesic (i.e., the great circle).

The third example is the Jaynes-Cummings model [18], which describes a two-level atom interacting with a single-mode radiation field. The Hamiltonian is given by

$$H = \frac{1}{2}\omega_0\sigma_3 + \omega a^{\dagger}a + \kappa(\sigma_+ a + \sigma_- a^{\dagger}). \tag{40}$$

Here, ω_0 , ω , and κ are the transition frequency of the atom, the frequency of the radiation, and the coupling constant, respectively. a^{\dagger} and a are the creation and annihilation operators of the radiation field satisfying the same algebra as in the first example. σ 's are the operators of the atom, which are given in terms of the orthonormal basis, the ground state $|g\rangle$ and the excited state $|e\rangle$, as follows: $\sigma_+ = |e\rangle\langle g|$, $\sigma_- = |g\rangle\langle e|$, $\sigma_3 = |e\rangle\langle e| - |g\rangle\langle g|$. This model is known to be exactly solvable. The associated unitary operator is

$$U(l) = \sum_{n=0}^{\infty} \left[\alpha_n(l) \sigma_+ \sigma_- \mid n \rangle \langle n \mid + \beta_{n-1}(l) \sigma_- \sigma_+ \mid n \rangle \langle n \mid + \gamma_n(l) \sigma_+ \mid n \rangle \langle n + 1 \mid + \delta_n(l) \sigma_- \mid n + 1 \rangle \langle n \mid \right], \quad (41)$$

where $|n\rangle=(n!)^{-1/2}a^{\dagger^n}|0\rangle$ is the *n*-photon state (for the details of this unitary operator, see [13]). The unitarity of this operator leads to the conditions: $|\alpha_n|^2+|\gamma_n|^2=1$, $|\alpha_n|=|\beta_n|$, $|\gamma_n|=|\delta_n|$, and $\alpha_n\delta_n^*+\beta_n^*\gamma_n=0$. From the polar forms, $\alpha_n=|\alpha_n|\exp(i\theta_{\alpha_n})$ and so on, follows the condition on the phases: $\theta_{\alpha_n}+\theta_{\beta_n}-\theta_{\gamma_n}-\theta_{\delta_n}=(2m+1)\pi$ ($m=0,\pm 1,\pm 2,\ldots$). Here, we choose $|e\rangle|n\rangle$ as the initial state. $|\alpha_n|$, θ_{α_n} , θ_{β_n} , and θ_{γ_n} form as a set of independent coordinate variables. The metric is

$$ds^{2} = \frac{d\left|\alpha_{n}\right|^{2}}{1-\left|\alpha_{n}\right|^{2}} + \left|\alpha_{n}\right|^{2} \left(1-\left|\alpha_{n}\right|^{2}\right) \left(d\theta_{\alpha_{n}} - d\theta_{\gamma_{n}}\right)^{2},\tag{42}$$

which does not seem to be the one of a familiar space. The θ_{β_n} -dependence disappears due to the above choice of the initial state. The transformed Hamiltonian is

$$H(l) = \sum_{n=0}^{\infty} \left[A_n(l)\sigma_+\sigma_- \mid n\rangle\langle n\mid + B_{n-1}(l)\sigma_-\sigma_+ \mid n\rangle\langle n\mid + C_n(l)\sigma_+ \mid n\rangle\langle n+1\mid + C_n^*(l)\sigma_- \mid n+1\rangle\langle n\mid \right], \quad (43)$$

and so

$$H_{\text{o-d}}(l) = \sum_{n=0}^{\infty} \left[C_n(l)\sigma_+ \mid n\rangle\langle n+1 \mid + C_n^*(l)\sigma_- \mid n+1\rangle\langle n \mid \right], \tag{44}$$

where the coefficients, $A_n(l)$'s and $B_{n-1}(l)$'s are real, and $C_n(l)$'s turn out to be also real (as can be seen from the flow equations). They are expressed in terms of the physical coefficients contained in the original Hamiltonian as well as the coefficients appearing in U(l) (for details, see [13]). Let us consider Wegner's choice:

$$\eta^{W}(l) = \sum_{n=0}^{\infty} \left[A_{n}(l) - B_{n}(l) \right] C_{n}(l) \left(\sigma_{+} \mid n \rangle \langle n+1 \mid -\sigma_{-} \mid n+1 \rangle \langle n \mid \right). \tag{45}$$

Then, we find that Eq. (15) vanishes. The flow equations give rise to

$$\frac{d\theta_{\alpha_n}(l)}{dl} = 0, (46)$$

$$\frac{d\theta_{\gamma_n}(l)}{dl} = 0, (47)$$

which explicitly make the variations of the arc-length functional with respect to $|\alpha_n(l)|$, $\theta_{\alpha_n}(l)$, and $\theta_{\gamma_n}(l)$ all vanish. In addition, comparing $\eta(l) = [H_{\rm d}(l), H_{\rm 0-d}(l)]$ with $\eta^W(l)$ above, we see that Case-B in the preceding section is realized. Thus, again Wegner's flow is geodesic.

5 Concluding Remarks

To summarize, generalizing Wegner's method of flow equations, we have found a condition, under which the corresponding flow of a quantum state becomes geodesic in a quantum evolution submanifold, independently of specific initial conditions at l=0. We have illustrated this by employing the generalized harmonic oscillator, the spin in an external magnetic field, and the Jaynes-Cummings model.

The present result implies that the method of flow equations is not just a mathematical tool for diagonalizing a Hamiltonian but provides the optimal strategy in quantum state engineering for realizing a stationary state from a given initial state in each of a certain class of systems. In fact, a formal solution, $U(l) = P_l \exp \int_0^l dl' \eta^W(l')$ with P_l being Dyson's "l-ordering" symbol, can be divided into the product of many unitary operators, each of which defines a small translation along the geodesic flow and may represent each quantum operation performed experimentally. In the present work, we have discussed the metric structure and associated geodesic nature of the flow of a state in a quantum evolution submanifold. Other basic quantities such as curvature are not explicitly treated here. However, curvature becomes relevant, for example, when geodesic deviation is considered. This kind of considerations may cast further light on geometry of the flow equations.

All the examples discussed in Section 4 are exactly solvable, and Wegner's method perfectly works, generating the geodesic flows. This observation naturally leads to an anticipation that Wegner's flow may always be geodesic if a system is exactly solvable. Still there does not exist an affirmative/negative proof of this point, and further investigations are needed.

References

- 1. Wegner, F.: Ann. Phys. (Leipzig) 3, 77 (1994)
- 2. Wegner, F.J.: Phys. Rep. 348, 77 (2001);
- Wegner, F.: J. Phys. A: Math. Gen. 39, 8221 (2006);
- Kehrein, S.: The Flow Equation Approach to Many-Particle Systems. Springer-Verlag, Berlin (2006)
- 3. Kehrein, S.K., Mielke, A.: Ann. Phys. (NY) 252, 1 (1996)
- 4. Bylev, A.B., Pirner, H.J.: Phys. Lett. B 428, 329 (1998)
- 5. Raas, C., Löw, U., Uhrig, G.S., Kühne, R.W.: Phys. Rev. B 65, 144438 (2002)
- 6. Hankevych, V., Wegner, F.: Eur. Phys. J. B **31**, 333 (2003)
- 7. Stauber, T.: Phys. Rev. B 67, 205107 (2003)
- 8. Domanski, T., Ranninger, J.: Phys. Rev. B 70, 184513 (2004)
- 9. Dusuel, S., Vidal, J.: Phys. Rev. Lett. 93, 237204 (2004)
- 10. Dusuel, S., Vidal, J., Arias, J.M., Dukelsky, J., García-Ramos, J.E.: Phys. Rev. C 72, 064332 (2005)
- 11. Glazek, S.D.: Nucl. Phys. B (Proc. Suppl.) 161, 59 (2006)
- 12. Aprea, G., Di Castro, C., Grilli, M., Lorenzana, J.: Nucl. Phys. B 744, 277 (2006)
- 13. Itto, Y., Abe, S.: Physica A **387**, 1063 (2008)
- 14. Chu, M.T.: Linear Algebra Appl. 147, 75 (1991)
- 15. Abe, S.: Phys. Rev. A **46**, 1667 (1992);
 - Abe, S.: Phys. Rev. A 48, 4102 (1993)
- 16. Kobayashi, S., Nomizu, K.: Foundations of Differential Geometry, Vol. II. Interscience, New York (1969)
- 17. Page, D.N.: Phys. Rev. A 36, 3479 (1987)
- 18. Jaynes, E.T., Cummings, F.W.: Proc. IEEE 51, 89 (1963)