# SOME CONTROL VARIATES FOR EXOTIC OPTIONS

#### JC NDOGMO

ABSTRACT. There are no known exact formulas for the valuation of a number of exotic options, and this is particularly true for options under discrete monitoring and for American style options. Therefore, one usually recourses to a Monte Carlo Simulation approach, amongst other numerical methods, to estimate the value of these options. The problem which then arises with this method is one of variance reduction. Control variates are often used, and we present some results concerning these control variables, for the valuation of Asian and lookback options. An inequality on functions of correlations useful for comparing estimators in variance reduction procedures is also provided.

## 1. The problem of exotic options valuations

There are two main types of options. Standard European or American put and call options are referred to as vanilla options. These are exchange traded options and their payoffs depend on the price of the underlying asset at exercise time only. Contrary to these, there are derivatives with more complicated payoffs and which are referred to as exotic options. Most exotic options trade in over-thecounter markets and have been designed to meet the particular needs of customers (corporations).

One important class of exotic options consists of path-dependent (or historydependent) derivatives. They have a payoff that depends not just on the final price of the underlying asset, but also on the path followed by this price. Asian options and lookback options are typical examples of path-dependent options. In some instances, numerical methods are the only means available to value these exotic options, and in addition to Monte Carlo methods, other commonly used numerical methods include the finite difference and the finite element methods, the trinomial tree method, and some times the Markov Chain method (see [7, 8, 9, 10, 11] and the references therein). Until the last decade, Monte Carlo method has been considered as costly and unreliable, but based on innovative techniques which are now a topic of current research, they've yielded more promising results. In this context, Barraquand and Martineau [10], amongst others, obtained some interesting results.

Consider an option on a given security having n days to maturity. Let  $S_d(j)$  denote the closing price of the security at the end of day j (for j = 1, ..., n). Assume also that the risk free interest rate is r per year, and denote by N the number of trading days in a year. With these notation, the payoff from an asian option is either given by

$$Y = e^{-rn/N} \left( S_d(n) - \sum_{j=1}^n \frac{s_d(j)}{n} \right)^+,$$
 (1.1)

where the strike price K is taken as the average of all prices up to the expiry date, or

$$Y = e^{-rn/N} \left( \sum_{j=1}^{n} \frac{S_d(j)}{n} - K \right)^+,$$
(1.2)

when the underlying's terminal price is taken as the average of all prices up to maturity. The payoff from a lookback option is given by

$$Y = e^{-rn/N} \left( S_d(n) - \min_{j=1,\dots,n} S_d(j) \right)^+.$$
 (1.3)

Barrier options are another type of frequently traded path-dependent options. These are options for which a barrier value v, which may be a one- or two-dimensional vector, is specified, and the option becomes alive or cease to exist according to the path followed by the price w.r.t. the barrier value. When a barrier option is alive at maturity, its payoff is that of the equivalent vanilla option, i.e. that with the same strike price K, and number of days n to maturity.

The main problem with using Monte Carlo Simulation to value path-dependent derivatives is that the computation time necessary to achieve a reasonable level of accuracy can be excessively high. Control variate techniques are some of the tools available for implementing the necessary variance reduction procedures, which can lead to dramatic savings in computation time.

Our focus in this paper will be on the valuation of Asian and lookback options. In section 2 we discuss Monte Carlo valuation simulations for these options and explain how controlled variates techniques can be gainfully employed to improve the accuracy of the valuation. In section 3, we present some results concerning the control variables for an improvement of the variance reduction procedure, and an inequality on functions of correlation coefficients useful for comparing estimators in variance reduction techniques is also provided.

#### 2. Monte carlo simulations and variance reduction procedures

Suppose that the price S(t) of a given security follows a risk neutral geometric Brownian motion with constant parameters  $\mu$  and  $\sigma^2$ . This means in particular that for all  $y \ge 0$ , and  $t \ge 0$ ,  $\ln(S(t+y)/S(y))$  has a normal distribution with mean  $\mu t$ and standard deviation  $\sigma\sqrt{t}$ , and that  $\mu = r - \sigma^2/2$ , where r represent the risk-free interest rate.

We denote as usual by S(0) the initial price of the security, and by  $S_d(j)$  the price of the security at the end of day j (for j = 1, ..., n), where n is the number of days to maturity. Set

$$X(i) = \ln\left(\frac{S_d(i)}{S_d(i-1)}\right), \quad (i = 1, \dots, n).$$
(2.1)

The random variables  $X(1), \ldots, X(n)$  are independent and identically distributed with mean  $\mu/N$  and variance  $\sigma^2/N$ . That is,  $X(i) \sim \Phi(\frac{\mu}{N}, \frac{\sigma^2}{N})$ . A straightforward calculation using (2.1) shows that

$$S_d(i) = S_d(i-1)e^{X(i)}$$
(2.2)

$$= S(0)e^{X(1) + \dots + X(i)}$$
(2.3)

Values of  $X(1), \ldots, X(n)$  can be generated by a computer and Eq. (2.2)) or (2.3) can be used to sample a random path for the price S of the security.

The valuation simulation for an asian option whose payoff is

$$Y = e^{-rn/N} \left( \sum_{j=1}^{n} \frac{S_d(j)}{n} - K \right)^{-1}$$

as in Eq. (1.2) can be implemented as follows

- (1) Generate n values for the normal random variables  $X(1), \ldots, X(n)$ .
- (2) Calculate the end-of-day prices  $S_d(i) = S(0)e^{X(1)+\dots+X(n)}$ , for  $i = 1, \dots, n$ .
- (3) Calculate the payoff of the derivative  $Y_j = e^{-rn/N} \left( \sum \frac{S_d(i)}{n} K \right)^+$ , for the *j*-th simulation run.
- (4) Compute the average mean  $\overline{Y} = \frac{1}{R} \sum_{j=1}^{R} Y_j$ , where R is the total number of simulation runs, i.e. the number of times that steps (1) through (3) are performed.

The value of  $\bar{Y}$  represents an estimate of the exact cost Y of the derivative. We have

$$\mathbf{E}[\bar{Y}] = \mathbf{E}[Y] \tag{2.4}$$

and

$$\operatorname{Var}(\bar{Y}) = \operatorname{Var}(Y)/R. \tag{2.5}$$

The last equality, Eq. (2.5), shows that the accuracy of  $\overline{Y}$  improves with the number of simulations.

The simulation procedure is similar for lookback and for most of the pathdependent derivatives. Monte carlo simulation presents some advantages. They can be used to value path-dependent derivatives as well as those derivatives whose payoff depend only on the final value S of the security. They can also be used to value barrier options type securities, whose payoffs may occur several times before maturity. More discussions on the advantages of Monte Carlo simulation techniques can be found in [1, P. 408].

The drawbacks with Monte Carlo simulation is that the amount of time necessary to achieve a reasonable accuracy can be unacceptably high. As remedial measures, a number of variance reduction techniques are available, and they can lead to dramatic savings in computation time. These include the antithetic variable technique, the importance sampling, the stratified sampling, and the control variate technique. All these procedures and many others are described in [1, 2, 3].

#### JC NDOGMO

We now move on to explain how control variate can be used to improve the simulation valuations of derivatives. Suppose that in a general setup we want to estimate

$$M = \mathbb{E}[Y] \tag{2.6}$$

by simulation. Suppose furthermore that in the process of generating the random variables that determine Y we also find another random variable V whose mean value is known to be  $\mu_v = E[V]$ . Then rather than using the value of Y as an estimator, we can replace it with another value

$$W = Y + c(V - \mu_v),$$
 (2.7)

where c is a constant to be specified. The value of W is an estimate for M since E[W] = E[Y] = M. We have

$$\operatorname{Var}(W) = \operatorname{Var}(Y) + c^{2} \operatorname{Var}(V) + 2c \operatorname{Cov}(Y, V)$$
(2.8)

The best estimator is obtained by choosing c to be the value that makes Var(W) as small as possible. Elementary calculus shows that the value of c that minimizes Var(W) is given by

$$c^* = -\frac{\operatorname{Cov}(Y, V)}{\operatorname{Var}(V)} \tag{2.9}$$

substituting this value back into Eq. (2.8) yields

$$\operatorname{Var}(W) = \operatorname{Var}(Y) - \frac{\operatorname{Corr}^2(Y, V)}{\operatorname{Var}(V)}$$

That is,

$$\frac{\operatorname{Var}(W)}{\operatorname{Var}(Y)} = 1 - \operatorname{Corr}^2(Y, V)$$
(2.10)

$$=1-r^2,$$
 (2.11)

where the quantity  $r = \operatorname{Corr}(Y, V)$  is the correlation coefficient between Y and V. The number  $r^2$  is often called the coefficient of determination between Y and V. Eq. (2.10) shows that the variance reduction obtained with the control variate V is completely determined by  $\operatorname{Corr}(Y, V)$ . More precisely, the reduction obtained is  $100\% \operatorname{Corr}^2(Y, V)$  and consequently, the higher the correlation between Y and V, the greater the variance reduction obtained will be.

2.1. Control variates. Suppose that the payoff Y from an option on a given security is of any of the forms given by the Eqs. (1.1)-(1.3). With the notation of the previous section, set

$$V = \sum_{i=1}^{n} \alpha_i X(i) \tag{2.12}$$

where  $\alpha_i$  are some weights to be determined, and take

$$W = Y + c(V - E(V))$$
 (2.13)

as the new estimator for Y. We have

$$\mathbf{E}[V] = \sum_{i=1}^{n} \alpha_i \, \mathbf{E}[X(i)] = \sum_{i=1}^{n} \frac{\alpha_i}{N} \left( r - \frac{\sigma^2}{2} \right),$$

so that

$$W = Y + c \left[ \sum_{i=1}^{n} \alpha_i X(i) - \frac{\alpha_i}{N} \left(r - \frac{\sigma^2}{2}\right) \right]$$
$$= Y + \sum_{i=1}^{n} \beta_i \left[ X(i) - \left(r - \frac{\sigma^2}{2}\right) \right]$$

where

$$\beta_i = c \,\alpha_i. \tag{2.14}$$

On the other hand, we have

$$\operatorname{Var}(W) = \operatorname{Var}(Y) + \sum_{i=1}^{n} \beta_i^2 \operatorname{Var}(X(i)) + 2\sum_{i=1}^{n} \beta_i \operatorname{Cov}(Y, X(i))$$

and it readily follows that the values of  $\beta_i$  that minimize W are given by

$$\beta_i^* = -\frac{\operatorname{Cov}(Y, X(i))}{\operatorname{Var}(X(i))} \tag{2.15}$$

In terms of these optimal values of the  $\beta_i$ , the smallest possible value for Var(W) is given by

$$\operatorname{Var}(W) = \operatorname{Var}(Y) + \sum_{i} \frac{\operatorname{Cov}^{2}(Y, X(i))}{\operatorname{Var}(X(i))} + 2\sum_{i} - \frac{\operatorname{Cov}^{2}(Y, X(i))}{\operatorname{Var}(X(i))}$$

That is

$$\operatorname{Var}(W) = \operatorname{Var}(Y) - \sum_{i} \frac{\operatorname{Cov}^{2}(Y, X(i))}{\operatorname{Var}(X(i))}$$
(2.16)

Consequently,

$$\frac{\operatorname{Var}(W)}{\operatorname{Var}(Y)} = 1 - \sum_{i} \operatorname{Corr}^{2}(Y, X(i))$$
(2.17)

We have thus proven the following result

**Theorem 1.** Let  $V = \sum_i \alpha_i X(i)$ , for some arbitrary weights  $\alpha_i$  and fori = 1,...,n. Let W = Y + c(V - E[V]) be an estimator of Y for some constant c.

JC NDOGMO

(a): The optimal variance reduction is achieved with this control variable V for the values of  $\alpha_i$  and c such that

$$c\alpha_i = -\frac{\operatorname{Cov}(Y, X(i))}{\operatorname{Var}(X(i))}, \qquad (i = 1, \dots, n)$$

(b): The optimal variance reduction thus obtained is given by

$$\frac{\operatorname{Var}(W)}{\operatorname{Var}(Y)} = 1 - \sum \operatorname{Corr}^2(Y, X(i))$$

Remark 2.

- (1) It follows from part (a) of the theorem that a control variate of the form  $V = \sum_{i} \alpha_i X(i)$  gives rise to an optimal reduction if and only if  $\alpha_i = -\lambda \frac{\cos(Y,X(i))}{\operatorname{Var}(X(i))}$  for all  $i = 1, \ldots, n$  and for a nonzero constant  $\lambda$ .
- (2) This theorem implies that for all  $\alpha_1, \ldots, \alpha_n$  and for every constant c

$$\frac{\operatorname{Var}(W)}{\operatorname{Var}(Y)} \ge 1 - \sum \operatorname{Corr}^2(Y, X(i))$$

where

$$W = Y + c \left[ \sum_{i} \alpha_i X(i) - \alpha_i (r - \sigma^2/2) \right]$$

In [4, P. 138], the problem of finding the values of  $\alpha_i$  (for i = 1, ..., n) for the best control variate of the form  $V = \sum_i \alpha_i X(i)$ , and alternatively, the problem of finding the values of  $c_1, \ldots, c_n$  for the best estimator of Y of the form

$$Y + \sum_{i=1}^{n} c_i \left( X(i) - \frac{r - \sigma^2/2}{N} \right)$$

are posed. Theorem 1 gives an answer to this question, by determining the values of  $\alpha_i$  and  $c_i$  and by indicating precisely the optimal variance reduction that can be achieved.

**Theorem 3.** Let Y be a random variable and let  $X_1, \ldots, X_n$  be n independent random variables. Then for all numbers  $\alpha_1, \ldots, \alpha_n$ 

$$\operatorname{Corr}^2\left(Y, \sum_i \alpha_i X_i\right) \leq \sum_i \operatorname{Corr}^2(Y, X_i).$$

*Proof.* Let  $V = \sum_{i} \alpha_i X_i$  and let W = Y + c(V - E[V]) be an estimator of Y for some constant c. For any fixed values of the  $\alpha_i$  the optimal variance reduction given by Eq. (2.10) is

$$\frac{\operatorname{Var}(W)}{\operatorname{Var}(Y)} = 1 - \operatorname{Corr}^2(Y, V)$$

This is not smaller than the optimal variance reduction obtained for all possible values of the  $\alpha_i$  and c given by Eq. (2.17). Consequently,  $1 - \sum_i \operatorname{Corr}^2(Y, X_i) \leq 1 - \operatorname{Corr}^2(Y, \sum_i \alpha_i X_i)$ ; that is,  $\operatorname{Corr}^2(Y, \sum_i \alpha_i X_i) \leq \sum_i \operatorname{Corr}^2(Y, X_i)$ .

The result stipulated by this theorem is certainly very important, but it is likely to be unknown. Indeed in statistics books that present the most comprehensive material on the topic of correlation coefficient between random variables, there is rarely any discussion of a relationship between functions of correlations (see [5, 6]).

6

## 2.2. Applications.

- (1) If  $V = \sum_{i} \alpha_{i} X_{i}$  where the  $\alpha_{i}$  are constants independent of  $X_{i}$ , then no estimator of Y of the form  $W = Y + \sum_{i} \alpha_{i} X_{i} E[V]$  using V as control variable can lead to an optimal variance reduction, since by Theorem 1, the  $\alpha_{i}$  must depend on  $X_{i}$ .
- (2) Let  $V = \ln(s_d(n)/S(0))$ . By the definition of the X(i), it readily follows that  $V = \sum_i X(i)$ . Thus by the preceding remark, taking V as control variable cannot give rise to the best estimator.

## 3. Conclusion

We have made use of the variance reduction formula (2.10) obtained for an estimator of the payoff Y, with a given control variable V, to derive the form of the best possible control variable of the form  $V = \sum_{i} \alpha_i X(i)$ , where the random variables X(i) are as in Eq. (2.1). We've also obtained an upper bound for the correlation coefficient between an arbitrary random variable Y and an arbitrary linear combination of independent random variables. This inequality can be useful in comparing estimators in variance reduction procedures.

A generalization of this type of inequalities between functions of correlation coefficients is desirable, to compare for instance an estimator obtained with a control variable of the form  $V = \sum_i \beta_i S_d(i)$ , with that obtained using a control variable of the form  $\sum_i \alpha_i X(i)$ . In this instance, one would need a general relationship between functions of correlation coefficients of the type  $\operatorname{Corr}^2(Y, \sum_i \beta_i S_d(i))$  and  $\operatorname{Corr}^2(Y, \sum_i \beta_i \ln(S_d(i)/S_d(i-1)))$ . As stated in this paper, no result on such type of relationship seems to be available in the literature. Finally, the results obtained in this paper clearly apply to any random variable for which variance reduction is required, and not only to the specific case of exotic options considered in this paper.

### References

- John C. Hull, Options, Futures, & Other derivatives, Fourth edition, Prentice Hall, New Jersey, 2000.
- [2] L. Clewlow and C. Strickland, Implementing derivatives models, John Wiley and Sons, London, 1998.
- [3] P.P. Boyle, Options: A Monte Carlo Approach, Journal of financial economics 4 (1997) 323-38.
- [4] Sheldom M. Ross, An Introduction to Mathematical Finance, Cambridge University Press, Cambridge, 1999.
- [5] B. Ostle, Statistics in research: Basic concepts and techniques for research workers, 2nd Edition, Iowa State University Press, Iowa, 1972.
- [6] D.G. Rees, Essential Statistics, Fouth Edition, Chpman & Hall/CRC, Boca Raton, 2001.
- [7] M. Bertoli, M. Bianchetti, Monte carlo simulation of discrete barrier options, internal report, Financial Engineering - Derivatives Modelling, Caboto SIM S.p.a., Banca Intesa Group, Milan, Italy (2003).
- [8] J. Duan, E. Dudley, G. Gauthier, J. Simonato, Pricing discretely monitored barrier options by a markov chain, J. Derivatives 10 (2003) 9–32.
- [9] M. Broadie, P. Glasserman, S. Kou, A continuity correction for barrier options, Math. Finance 7 (1997) 325–349.
- [10] J. Barraquand, D. Martineau, Numerical valuation of high dimensional multivariate American securities, Journal of Financial and Quantitative analysis 30 (1995) 383–405.
- [11] A. Mayo, High-order accurate implicit finite difference method for evaluating American options, The European Journal of Finance 10 (2004) 212–237.

JC NDOGMO

Department of Mathematics University of the Western Cape P/B X17, Bellville 7535

 ${\it Email: jndogmo@uwc.ac.za}$