Counting paths in Bratteli diagrams for $SU(2)_k$

Toufik Mansour

Department of Mathematics, University of Haifa, Haifa 31905, Israel

Simone Severini

Institute for Quantum Computing and Department of Combinatorics & Optimization, University of Waterloo, Waterloo N2L 3G1, Canada

It is known that the Hilbert space dimensionality for quasiparticles in an $SU(2)_k$ Chern-Simons-Witten theory is given by the number of directed paths in certain Bratteli diagrams. We present an explicit formula for these numbers for arbitrary k. This is on the basis of a relation with Dyck paths and Chebyshev polynomials.

I. INTRODUCTION

The so-called Bratteli diagrams have been introduced by Bratteli in 1972 [\[2\]](#page-5-0) for the classification of some classes of C^* -algebras. For our scope, it is sufficient to give here an "algebra free" description of this notion (see, e.g., [\[1](#page-5-1)] or [\[3\]](#page-5-2)). More specifically, lattice points will be labeled by numbers instead of Young diagrams. A Bratteli diagram for $SU(2)_k$ is defined as a finite digraph $D_k = (V, E)$, where:

• The vertices of D_k are associated to lattice points in the positive quadrant of the Cartesian plane:

$$
(j, i) \in (\mathbb{Z}^{\geq 0})^{\times 2}
$$
 with $j \geq i$.

The vertex set V is $\{(j, i) : 0 \le i \le k, j; i \text{ and } j \text{ have the same parity}\}.$ Here 0 is assumed to be even.

- There is an arc $((j, i), (j', i'))$ in E if $j' = j + 1$ and $i' = i \pm 1$.
- A directed path from a vertex (j, i) to a vertex (j', i') of length n is a sequence of n arcs of the form $(j, i)(j +$ $(1, l_1), (j + 1, l_1)(j + 2, l_2), \ldots, (j + n - 1, l_{n-1})(j', i')$. Let $D_k(x, y)$ be the number of directed paths from the vertex $(0,0)$ to the vertex (j,i) in the Bratteli diagram D_k . In a Bratteli diagram D_k , the vertex (j,i) is labeled by the number $D_k(i, j)$. In our notation, the function f introduced in the general definition of a Bratteli diagram is exactly $D_k(i, j)$. Notice that

$$
D_k(i,j) = D_k(i-1,j-1) + D_k(i+1,j-1)
$$

with the initial conditions $D_k(0, j) = D_k(1, j - 1)$ and $D_k(k, j) = D_k(k - 1, j - 1)$.

Let

$$
d^+(i) = |\{j : (i,j) \in A(D_k)\}|
$$

and

$$
d^{-}(i) = |\{j : (j,i) \in A(D_k)\}|
$$

be the *indegree* and the *outdegree* of a vertex *i*, respectively. Notice that $d^+(0,0) = 1$, $d^-(0,0) = 0$, and $d^+(0,0) = 0$ $d^{\pm}(k, i) = d^{\pm}(l, l) = 1$. All other vertices have indegree and outdegree 2. The figure below illustrates the Bratteli diagram D_2 .

The number $D_k(i, j)$ corresponds to the Hilbert space dimensionality for i q-spin $1/2$ quasiparticles having total qspin j, in an $SU(2)_k$ Chern-Simons-Witten theory (see, e.g., [\[4](#page-5-3)] and the references contained therein). An asymptotic

FIG. 1: Bratteli diagram D_2 .

expression and a recurrence relation for $D_k(i, j)$ have been pointed out by Slingerland and Bais (see [\[9](#page-5-4), Section 2.4]). More precisely, they showed that

$$
D_k(i,j) \approx \left(2\cos\left(\frac{\pi}{k+2}\right)\right)^j,\tag{1}
$$

where $i+j = 0$ mod 2) (From the definitions we have that $D_k(i,j) = 0$ for all $i+j = 1 \pmod{2}$.) We present here an explicit formula for $D_k(i, j)$, which implies the asymptotic expression in [\(1\)](#page-1-0). In particular, we find an explicit formula for the total number of directed paths from the vertex $(0,0)$ to the vertex (i, j) in the Bratteli diagram D_k , for any k, as shown in the next section. Our tools will be Dyck paths and Chebyshev polynomials.

II. MAIN RESULT

Chebyshev polynomials of the second kind are defined by

$$
U_r(\cos\theta) = \frac{\sin\left((r+1)\theta\right)}{\sin\theta},
$$

for $r \geq 0$. Evidently, $U_r(x)$ is a polynomial of degree r in x with integer coefficients. Chebyshev polynomials were invented for the needs of approximation theory, but are also widely used in various other branches of mathematics, including algebra, number theory, and the study of lattice paths in combinatorics (see [\[8](#page-5-5)]). For $k \geq 0$, we define $R_k(x)$ by

$$
R_k(x) = \frac{U_{k-1}\left(\frac{1}{2\sqrt{x}}\right)}{\sqrt{x}U_k\left(\frac{1}{2\sqrt{x}}\right)}.
$$

For example, $R_0(x) = 0$, $R_1(x) = 1$, and $R_2(x) = 1/(1-x)$. It is easy to see that for any k, $R_k(x)$ is a rational function in x .

A Dyck path is a lattice path in the plane integer lattice $(\mathbb{Z}^{\geq 0})^{\times 2}$ consisting of up-steps $u = (1, 1)$ and down-steps $d = (1, -1)$. It follows that a Dyck path never passes below the x-axis. The length of a Dyck path is defined as the number of its up-steps and down-steps. From the definitions we can state the following observation.

Observation 1 The number $D_k(i, j)$ is exactly the number of Dyck paths below the line $y = k + 1$, starting at the origin and ending at the point with x-coordinate j and y-coordinate i .

Each Dyck path P below the line $y = k + 1$, starting at the origin and ending at (j, i) , has the following form:

$$
P = P_1 u P_2 u P_3 \cdots u P_{i+1},
$$

where P_s is a Dyck path of height at most $k + 1 - s$, and the length of P is exactly j (each up-step and down-step is counted as a unit step).

Let us fix a variable x for the generating function for counting the number of up-steps and down-steps in a Dyck path. Using the fact that the generating function for the number of Dyck paths of length 2n with height at most s is given by $R_{s+1}(x^2)$ (see [\[5\]](#page-5-6)), we get that the generating function

$$
D_k(x;i) = \sum_{j\geq 0} D_k(i,j)x^j
$$

is given by

$$
D_k(x; i) = R_{k+1}(x^2) \prod_{r=1}^{i} (xR_{k+1-r}(x^2)),
$$
\n(2)

which is equivalent to

$$
D_k(x; i) = x^i \prod_{r=0}^i \frac{U_{k-r} \left(\frac{1}{2x}\right)}{xU_{k+1-r} \left(\frac{1}{2x}\right)} = \frac{U_{k-i} \left(\frac{1}{2x}\right)}{xU_{k+1} \left(\frac{1}{2x}\right)}.
$$
\n(3)

Using the fact that the roots of $U_k(x)$ are

$$
\cos\left(\frac{r\pi}{k+1}\right),\ r=1,2,\ldots,k
$$

we obtain that the minimal positive pole of the function $D_k(x;i)$ is given by $\left(2\cos\left(\frac{\pi}{k+2}\right)\right)^{-1}$. Therefore, the asymptotic behaviour of the function $D_k(i, j)$ is given by

$$
D_k(i,j) \approx \left(2\cos\left(\frac{\pi}{k+2}\right)\right)^j,
$$

where $i + j = 0 \pmod{2}$, as described in [\[9,](#page-5-4) Equation 26].

In order to find an explicit formula for $D_k(i, j)$ we need the following lemma.

Lemma 2 The generating function

$$
\frac{U_{k-i}(x)}{U_{k+1}(x)}
$$

is given by

$$
\frac{1}{k+2} \sum_{r=1}^{k+1} \frac{(-1)^{r+1} U_{k-i}(\rho_{k+1,r}) \sin^2 \frac{\pi r}{k+2}}{x - \rho_{k+1,r}},
$$

where $\rho_{m,r} = \cos\left(\frac{r\pi}{m+1}\right)$.

Proof. Let us compute the partial fraction decomposition of

$$
\frac{U_{k-i}(x)}{U_{k+1}(x)}.
$$

By general principles, it is

$$
\sum_{r=1}^{k+1} \frac{a_{k+1,r}}{x - \rho_{k+1,r}},
$$

where $\rho_{m,r} = \cos\left(\frac{r\pi}{m+1}\right)$ are the zeros of the m-th Chebyshev polynomials of the second kind. Now,

$$
a_{k+1,r} = \frac{U_{k-i}(\rho_{k+1,r})}{\frac{d}{dx}U'_{k+1}(x)|_{\rho_{k+1,r}}}
$$

and note that

$$
U'_{k+1}(x) = \frac{dU_{k+1}(x)}{dx} = \frac{d}{d\theta} \left(\frac{\sin((k+2)\theta)}{\sin(\theta)} \right) \cdot \frac{d\theta}{dx}.
$$

We work out that

$$
\frac{dU_{k+1}}{d\theta} = \frac{(k+2)\cos((k+2)\theta) \cdot \sin(\theta) - \sin((k+2)\theta) \cdot \cos(\theta)}{\sin^2(\theta)},
$$

and if we plug in $x = \rho_{k+1,r}$ simplification occurs, since certain terms are just zero; we obtain that

$$
\frac{dU_{k+1}}{d\theta}(\arccos \rho_{k+1,r})
$$
\n
$$
= \frac{(k+2)\cos(k+2)\theta \cdot \sin \theta - \sin(k+2)\theta \cdot \cos \theta}{\sin^2 \theta}\Big|_{\theta = \frac{\pi r}{k+2}}
$$
\n
$$
= \frac{(k+2)\cos(\pi r) \cdot \sin \frac{\pi r}{k+2} - \sin(\pi r) \cdot \cos \frac{\pi r}{k+2}}{\sin^2 \frac{\pi r}{k+2}}
$$
\n
$$
= \frac{(k+2)\cos(\pi r) \cdot \sin \frac{\pi r}{k+2}}{\sin^2 \frac{\pi r}{k+2}}
$$
\n
$$
= \frac{(k+2)(-1)^r}{\sin \frac{\pi r}{k+2}}.
$$

Further $\frac{dx}{d\theta} = -\sin \theta$, so together

$$
a_{k+1,r} = \frac{U_{k-i}(\rho_{k+1,r})}{U'_{k+1}(\rho_{k+1,r})}
$$

=
$$
\frac{1}{k+2}(-1)^{r+1}U_{k-i}(\rho_{k+1,r})\sin^2\frac{\pi r}{k+2},
$$

which completes the proof. \blacksquare

Now we are ready to give an explicit formula for $D_k(i, j)$.

Theorem 3 For all k, i, j , the number $D_k(i, j)$ of directed paths from the vertex $(0, 0)$ to the vertex with x-coordinate j and y -coordinate i in the Bratteli diagram D_k is given by

$$
\frac{2}{k+2} \sum_{r=1}^{k+1} (-1)^{r+1} U_{k-i}(\rho_{k+1,r}) \sin^2\left(\frac{\pi r}{k+2}\right) (2\rho_{k+1,r})^j,
$$

where $\rho_{m,r} = \cos\left(\frac{r\pi}{m+1}\right)$.

Proof. Lemma [2](#page-2-0) and Equation [3](#page-2-1) give that the generating function $D_k(x; i)$ is given by

$$
\frac{2}{k+2} \sum_{r=1}^{k+1} \frac{(-1)^{r+1} U_{k-i}(\rho_{k+1,r}) \sin^2\left(\frac{\pi r}{k+2}\right)}{1 - 2\rho_{k+1,r} x},
$$

which implies that the x^j coefficient in the generating function $D_k(x; i)$ is given by

$$
\frac{2}{k+2} \sum_{r=1}^{k+1} (-1)^{r+1} U_{k-i}(\rho_{k+1,r}) \sin^2\left(\frac{\pi r}{k+2}\right) (2\rho_{k+1,r})^j,
$$

as claimed.

For example, the above theorem for $k = 1, 2, 3, 4, 5$ gives the following values:

• $D_1(0, 2j) = 1$ and $D_1(1, 2j + 1) = 1$ (otherwise $D_1(i, j) = 0$).

FIG. 2: Number of paths in the Bratteli diagram D_2 .

- $D_2(0, 2j) = D_2(2, 2j) = 2^{j-1}$ and $D_2(1, 2j + 1) = 2^j$, see Figure [2.](#page-4-0)
- $D_3(0, 2j) = F_{2j-1}, D_3(1, 2j) = F_{2j+1}, D_3(2, 2j+2) = F_{2j+2}$ and $D_3(3, 2j+3) = F_{2j+2}$, where F_m is the m-th Fibonacci number (defined as $F_0 = 0$, $F_1 = 1$, and $F_n = F_{n-1} + F_{n-2}$ for any n), see Figure [3.](#page-4-1) It was pointed out in [\[4\]](#page-5-3) that the theory of q-spin 1 quasiparticles in $SU(2)_3$ is equivalent to $SO(3)_3$, a theory which is in fact also known as the Fibonacci anyon theory [\[7](#page-5-7)].

FIG. 3: Number of paths in the Bratteli diagram D_3 .

- $D_4(0, 2j + 2) = D_4(1, 2j + 1) = (3^{j} + 1)/2$, $D_4(2, 2j + 2) = 3^j$, and $D_4(3, 2j + 3) = D_4(4, 2j + 4) = (3^{j+1} - 1)/2,$ for all $j \geq 0$.
- $D_5(0, 2j) = a_j, D_5(1, 2j + 1) = a_{j+1},$ $D_5(2, 2j + 2) = a_{j+2} - a_{j+1},$ $D_5(3, 2j + 3) = a_{j+3} - 2a_{j+2},$ $D_5(4, 2j + 4) = D_5(5, 2j + 5) = a_{j+4} - 3a_{j+3} + a_{j+2},$

for all $j \geq 0$, where a_j is a sequence satisfies the following recurrence relation $a_m = 5a_{m-1} - 6a_{m-2} + a_{m-3}$ with the initial conditions $a_0 = a_1 = 1$ and $a_2 = 2$. Note that, the generating function for the sequence a_n is given by

$$
\frac{1 - 4x + 3x^2}{1 - 5x + 6x^2 - x^3}
$$

.

Using the continued fraction representation of the generating function $R_k(x)$, namely

$$
R_k(x) = \frac{1}{1 - \frac{x}{1 - \frac{x}{\ddots}}}
$$

with exactly k-levels (see [\[6](#page-5-8), Lemma 3.1]), we obtain that

$$
R_{\infty}(x) = \lim_{k \to \infty} R_k(x) = C(x) := \frac{1 - \sqrt{1 - 4x}}{2x}.
$$

Hence, From [\(2\)](#page-2-2) we obtain that the generating function

$$
D_{\infty}(x;i) = \lim_{k \to \infty} D_k(x;i)
$$

is given by

$$
D_{\infty}(x;i) = x^i C^{i+1}(x^2).
$$

The generating function $C(x)$ satisfies

$$
C(x) = 1 + xC^2(x),
$$

thus by the Lagrange inversion formula [\[10,](#page-5-9) Sec. 5.4] we obtain that

$$
D_{\infty}(i, j) = \frac{i+1}{j+1} \binom{j+1}{\frac{j-i}{2}},
$$

where $i + j = 0 \pmod{2}$, as shown in [\[9,](#page-5-4) Equation 21].

- [1] S. Bezuglyi, J. Kwiatkowski, K. Medynets, B. Solomyak, Invariant Measures on Stationary Bratteli Diagrams, [arXiv:0812.1088v](http://arxiv.org/abs/0812.1088)1 [math.DS]
- [2] O. Bratteli, Inductive limits of finite dimensional C*-algebras, Trans. Amer. Math. Soc. 171 (1972), 195-234.
- [3] F. Durand, B. Host, C. Skau, Substitutional dynamical systems, Bratteli diagrams and dimension groups, Ergodic Theory Dynam. Systems 19 (1999), no. 4, 953–993.
- [4] L. Hormozi, G. Zikos, N.E. Bonesteel, S.H. Simon, Topological Quantum Compiling, Phys. Rev. B 75, 165310 (2007). [arXiv:quant-ph/0610111v](http://arxiv.org/abs/quant-ph/0610111)2
- [5] T. Mansour, Counting peaks at height k in a Dyck path, J. Int. Seq. 5 (2002) Article 02.1.1.
- [6] T. Mansour and A. Vainshtein, Restricted permutations, continued fractions, and Chebyshev polynomials, Elect. J. Combin. 7 (2000) $\#\text{R17}$.
- [7] J. Preskill, Lecture Notes for Physics 219: Quantum Computation – Part III: Topological Quantum Computation. www.theory.caltech.edu/˜preskill/ph219/topological.pdf
- [8] Th. Rivlin, Chebyshev polynomials. From approximation theory to algebra and number theory, John Wiley, New York (1990).
- [9] J. K. Slingerland, F. A. Bais, Quantum groups and nonabelian braiding in quantum Hall systems, Nucl.Phys. B 612 (2001) 229-290. [arXiv:cond-mat/0104035v](http://arxiv.org/abs/cond-mat/0104035)1 [cond-mat.mes-hall]
- [10] R. P. Stanley, Enumerative Combinatorics, Vol. 2, Cambridge University Press, Cambridge, 1999.