

Semiclassical Corrections to the Cardy-Verlinde Formula of Kerr Black Holes

M.R. Setare *

Department of Science, Payame Noor University. Bijar. Iran

Abstract

In this letter, we compute the corrections to the Cardy-Verlinde formula of 4-dimensional Kerr black hole. These corrections are considered within the context of KKW analysis and arise as a result of the self-gravitational effect. Then we show, one can taking into account the semiclassical corrections of the Cardy-Verlinde entropy formula by just redefining the Virasoro operator L_0 and the central charge c .

arXiv:0807.0273v1 [hep-th] 2 Jul 2008

*E-mail: rezakord@ipm.ir

1 Introduction

Concerning the quantum process called Hawking effect [1] much work has been done using a fixed background during the emission process. The idea of Keski-Vakkuri, Kraus and Wilczek (KKW) [2]-[6] is to view the black hole background as dynamical by treating the Hawking radiation as a tunnelling process. The energy conservation is the key to this description. The total (ADM) mass is kept fixed while the mass of the black hole under consideration decreases due to the emitted radiation. The effect of this modification gives rise to additional terms in the formulae concerning the known results for black holes [7]-[12]; a non-thermal partner to the thermal spectrum of the Hawking radiation shows up. The Cardy-Verlinde formula proposed by Verlinde [13], relates the entropy of a certain CFT with its total energy and its Casimir energy in arbitrary dimensions. Using the $\text{AdS}_d/\text{CFT}_{d-1}$ [14]-[17] and $\text{dS}_d/\text{CFT}_{d-1}$ correspondences [18, 19, 20], this formula has been shown to hold exactly for different black holes (see for example [21]-[27]). All cases where the Cardy-Verlinde formula has been shown to hold up in these references had as a necessary ingredient a negative cosmological constant, or, more generally, a certain potential term for supergravity scalars. This guarantees that the theory admits AdS vacua, and thus one has a dual description in terms of a conformal field theory on the boundary of AdS. A natural question is whether the Cardy-Verlinde formula holds in a more general setting, e. g. for black holes that are asymptotically flat rather than approaching AdS space. The authors of [28] have shown that this is indeed the case. In this paper, in the context of KKW analysis we consider the Kerr black hole as a dynamical background and we derive the modified event horizon radius of Kerr black hole. Then we derive the corresponding modified thermodynamic quantities of the Kerr black hole and in the lowest order of the emitted shell of energy. Then we consider the generalized Cardy-Verlinde formula of a 4-dimensional Kerr black hole. We obtain the semiclassical corrections to this entropy formula. Finally we obtain the corrections to the quantities entering the Cardy-Verlinde formula: Virasoro operator and the central charge.

2 Thermodynamical quantities of Kerr black hole

We consider the asymptotically flat Kerr black holes [29]

$$ds^2 = -\frac{\Delta}{\rho^2}[dt - a \sin^2 \theta d\phi]^2 + \frac{\rho^2}{\Delta}dr^2 + \rho^2 d\theta^2 + \frac{\sin^2 \theta}{\rho^2}[adt - (r^2 + a^2)d\phi]^2, \quad (1)$$

where

$$\begin{aligned} \Delta &= (r^2 + a^2) - 2Mr, \\ \rho^2 &= r^2 + a^2 \cos^2 \theta. \end{aligned} \quad (2)$$

The event horizon $r = r_+$ is given by

$$r_+ = M + \sqrt{M^2 - a^2} \quad (3)$$

The temperature, entropy, energy, angular momentum and angular velocity of the horizon are as following[29]

$$T = \frac{r_+^2 - a^2}{4\pi r_+(r_+^2 + a^2)} \quad (4)$$

$$S_{BH} = \frac{V_2}{4G}(r_+^2 + a^2), \quad (5)$$

$$E = \frac{V_2}{8\pi G r_+}(r_+^2 + a^2) \quad (6)$$

$$J = \frac{aV_2}{8\pi G r_+}(r_+^2 + a^2) \quad (7)$$

$$\Omega_H = \frac{a}{r_+^2 + a^2} \quad (8)$$

where V_2 denotes the volume of the unit 2-sphere.

3 Self-Gravitational Corrections to Cardy-Verlind Formula

The entropy of a $(1+1)$ -dimensional CFT is given by the well-known Cardy formula [30]

$$S = 2\pi\sqrt{\frac{c}{6}\left(L_0 - \frac{c}{24}\right)}, \quad (9)$$

where L_0 represent the product ER of the energy and radius, and the shift of $\frac{c}{24}$ is caused by the Casimir effect. After making the appropriate identifications for L_0 and c , the same Cardy formula is also valid for CFT in arbitrary spacetime dimensions d in the form [13]

$$S_{CFT} = \frac{2\pi R}{d-2}\sqrt{E_c(2E - E_c)}, \quad (10)$$

the so called Cardy-Verlind formula, where R is the radius of the system, E is the total energy and E_C is the Casimir energy. The definition of the Casimir energy is derived by the violation of the Euler relation as [13]

$$E_C = 2(E - TS_{BH} + pV - \Omega_H J), \quad (11)$$

the pressure of the CFT is defined as $p = E/2V$, where V is the volume of the system. The total energy may be written as the sum of two terms

$$E(S, V) = E_E(S, V) + \frac{1}{2}E_C(S, V) \quad (12)$$

where E_E is the purely extensive part of the total energy E and the Casimir energy, is the subextensive part of E . Using the equation of state, $pV = E/2$, and substituting Eqs.(4)-(8) into Eq.(11), we get

$$E_C = \frac{V_2}{4\pi G r_+}(r_+^2 + a^2) = 2E. \quad (13)$$

So far, mostly asymptotically AdS and dS black hole solutions have been considered [14]-[27]. In [28], it is shown that even the Schwarzschild and Kerr black hole solutions, which are asymptotically flat, satisfy the modification of the Cardy-Verlinde formula

$$S_{CFT} = \frac{2\pi R}{d-2} \sqrt{2EE_c}. \quad (14)$$

This result holds also for various charged black hole solution with asymptotically flat spacetime [31].

We are interested primarily in the corrections to the entropy (5) that arise in the context of KKW analysis [2]-[6]. Let us remind that the key point to the KKW analysis is that the total energy of the spacetime under study is kept fixed while the black hole mass is allowed to vary. To calculate the emission rate, we should introduce the Painlevé-Ker coordinate system. The line element in the Painlevé-Kerr coordinate system is given in Ref. [32]. Namely,

$$ds^2 = \hat{g}_{00} dt^2 + 2\sqrt{\hat{g}_{00}(1-g_{11})} dt dr + dr^2 + [\hat{g}_{00}G(r,\theta)^2 + g_{22}] d\theta^2 + 2\hat{g}_{00}G(r,\theta) dt d\theta + 2\sqrt{\hat{g}_{00}(1-g_{11})}G(r,\theta) dr d\theta, \quad (15)$$

where

$$\hat{g}_{00} = -\frac{\rho^2 \Delta}{(r^2 + a^2)^2 - \Delta a^2 \sin^2 \theta}, \quad (16)$$

$$g_{11} = \frac{\rho^2}{\Delta}, \quad g_{22} = \rho^2. \quad (17)$$

and $G(r, \theta)$ is given by

$$G(r, \theta) = \int \frac{\partial F(r, \theta)}{\partial \theta} dr + C(\theta), \quad (18)$$

also $F(r, \theta)$ satisfies

$$g_{11} + \hat{g}_{00}F(r, \theta)^2 = 1, \quad (19)$$

here, $C(\theta)$ is an arbitrary analytic function of θ . The outgoing radial null geodesics followed by the massless particles, i.e. the shell of energy, is given by

$$\dot{r} = \frac{\Delta}{2} \sqrt{\frac{\rho^2}{(\rho^2 - \Delta)[(r^2 + a^2)^2 - \Delta a^2 \sin^2 \theta]}}. \quad (20)$$

To calculate the emission rate correctly, we should take into account the self-gravitation of the tunnelling particle with energy ω . That is, we should replace M with $M - \omega$ in (15). The coordinate φ does not appear in the line element expressions (15). That is to say, φ is an ignorable coordinate in the Lagrangian function L . To eliminate this freedom completely, the action for the classically forbidden trajectory should be written as

$$S = \int_{t_i}^{t_f} (L - P_\varphi \dot{\varphi}) dt, \quad (21)$$

which is related to the emission rate of the tunnelling particle by

$$\Gamma \sim e^{-2ImS}. \quad (22)$$

Therefore, the imaginary part of the action is

$$ImS = Im\left(\int_{r_i}^{r_f} \left[P_r - \frac{P_\varphi \dot{\varphi}}{\dot{r}}\right] dr\right) = Im\left(\int_{r_i}^{r_f} \left[\int_{(0,0)}^{(P_r, P_\varphi)} dP'_r - \frac{\dot{\varphi}}{\dot{r}} dP'_\varphi\right] dr\right), \quad (23)$$

where P_φ is the canonical momentum conjugate to φ . If we treat the black hole as a rotating sphere and consider the particle self-gravitation, we have

$$\dot{\varphi} = \Omega'_H, \quad (24)$$

and

$$J' = (M - \omega')a = P'_\varphi, \quad (25)$$

where Ω'_H is the dragged angular velocity of the event horizon. The imaginary part of the action can be rewritten as

$$ImS = Im\left\{\int_{r_i}^{r_f} \left[\int_{(0,J)}^{(P_r, J-\omega a)} dP'_r - \frac{\Omega'_H}{\dot{r}} dJ'\right] dr\right\}. \quad (26)$$

We now eliminate the momentum in favor of energy by using Hamilton's equation

$$\dot{r} = \frac{dH}{dP_r} \Big|_{(r;\varphi,P_\varphi)} = \frac{d(M - \omega')}{dP_r} = \frac{dM'}{dP_r}, \quad (27)$$

Based on similar discussion to [2]-[6], it follows directly that a particle tunnelling across the event horizon sees the effective metric of Eq. (15), although with the replacements $M \rightarrow M - \omega'$. The same substitutions in Eq. (20) yield the desired expression of \dot{r} as a function of ω' . Thus, we can rewrite (26) in the following explicit manner

$$ImS = Im \int_{r_i}^{r_f} \left[\int \frac{2\sqrt{(\rho^2 - \Delta')[(r^2 + a^2)^2 - \Delta' a^2 \sin^2 \theta]}}{\Delta' \sqrt{\rho^2}} (dM' - \Omega'_H dJ') \right] dr. \quad (28)$$

where

$$\Delta' = r^2 + a^2 - 2M'r \quad (29)$$

$$r_i = r_+ = M + \sqrt{M^2 - a^2}, \quad (30)$$

$$r_f = M - \omega + \sqrt{(M - \omega)^2 - a^2}. \quad (31)$$

We see that $r = (M - \omega') + \sqrt{(M - \omega')^2 - a^2}$ is a pole. The integral can be evaluated by deforming the contour around the pole, so as to ensure that positive energy solution decay in time. Doing the r integral first we obtain

$$ImS = -\frac{1}{2} \int_{(M)}^{(M-\omega)} \frac{4\pi(M'^2 + M'\sqrt{M'^2 - a^2})}{\sqrt{M'^2 - a^2}} (dM' - \Omega'_H dJ') dr. \quad (32)$$

Finishing the integration we get

$$ImS = \pi[M^2 - (M - \omega)^2 + M\sqrt{M^2 - a^2} - (M - \omega)\sqrt{(M - \omega)^2 - a^2}] = -\frac{1}{2}\Delta S_{BH}. \quad (33)$$

The tunnelling rate is therefore

$$\Gamma \sim \exp[-2ImS] = e^{\Delta S_{BH}}. \quad (34)$$

The Casimir energy Eq.(11) now will be modified due to the semi-classical corrections as

$$E'_C = 2(3/2E' - T'S'_{BH} - \Omega'_H J'). \quad (35)$$

It is easily seen that

$$2E'_C E' = \frac{V_2^2}{16\pi^2 G^2 r_f^2} (r_f^2 + a^2)^2 \quad (36)$$

We substitute the above expression in the Cardy-Verlinde formula in order that self-gravitational corrections to first order in ω to be considered:

$$S'_{CFT} = S_{CFT} \left[1 - \frac{2\omega r_i^2}{(r_i^2 + a^2)\sqrt{M^2 - a^2}} \right] = S_{CFT} \left[1 - \frac{\omega(2M^2 - a^2 + 2M\sqrt{M^2 - a^2})}{(M^2 + M\sqrt{M^2 - a^2})\sqrt{M^2 - a^2}} \right] \quad (37)$$

For the Kerr black holes, the dual CFT lives on a flat space, and thus the energy has no subextensive part. In the two-dimensional conformal field theory, when the conformal weight of the ground state is zero, we have

$$S_{CFT} = 2\pi \sqrt{\frac{cL_0}{6}}, \quad (38)$$

where $L_0 = ER$ and $\frac{c}{6} = \frac{2S_c}{\pi} = 2E_c R$, here S_c is the Casimir entropy. Then, we can taking into account the self-gravitational corrections of the Cardy-Verlinde entropy formula by just redefining the Virasoro operator and the central charge as following. Using Eqs.(13, 30, 31, 36) one can obtain

$$L'_0 = E'R' = ER - \frac{\omega V_2 r_i^2}{4\pi G \sqrt{M^2 - a^2}} = L_0 - \frac{\omega V_2 (2M^2 - a^2 + 2M\sqrt{M^2 - a^2})}{4\pi G \sqrt{M^2 - a^2}} \quad (39)$$

$$c' = 12E'_c R' = 12E_c R - \frac{6\omega V_2 r_i^2}{\pi G \sqrt{M^2 - a^2}} = c - \frac{6\omega V_2 (2M^2 - a^2 + 2M\sqrt{M^2 - a^2})}{\pi G \sqrt{M^2 - a^2}} \quad (40)$$

In [33] Carlip have computed the logarithmic corrections to the Cardy formula, according to his calculations, logarithmic corrections to the density of states is as

$$\rho(\Delta) \approx \left(\frac{c}{96\Delta^3}\right)^{1/4} \exp\left(2\pi \sqrt{\frac{c\Delta}{6}}\right) \quad (41)$$

where $\Delta = L_0$ the exponential term in (41) gives the standard Cardy formula, but we have now found the lading correction, which is logarithmic. In the other hand as we saw the effect of the self-gravitational corrections to the Cardy-Verlinde formula appear as the redefinition of the c and L_0 only. As Carlip have discussed in [33], the central charge c appearing in (41) is the full central charge of the conformal field theory. In general, c will consist of a classical term which appear in the Poisson brackets of the Virasoro algebra generators, plus a correction due to the quantum (here self-gravitational corrections) effects, that can change the exponent in (41) from its classical value. Moreover we saw

that L_0 take a similar correction as eq.(39) then the similar discussion about L_0 is correct. Therefore the first order corrections to the L_0 and c are given by

$$\Delta L_0 = L'_0 - L_0 = -\frac{\omega V_2(2M^2 - a^2 + 2M\sqrt{M^2 - a^2})}{4\pi G\sqrt{M^2 - a^2}} \quad (42)$$

$$\Delta c = c' - c = -\frac{6\omega V_2(2M^2 - a^2 + 2M\sqrt{M^2 - a^2})}{\pi G\sqrt{M^2 - a^2}} \quad (43)$$

Thus, this redefinition can be considered as a renormalization of the quantities entering in the Cardy formula.

4 Conclusions

Black hole thermodynamic quantities depend on the event horizon radius as Eqs.(4-8). The event horizon radius undergoes corrections from the self-gravitational effect as Eq.(31). Then we have obtained the corrections to the entropy of a dual conformal field theory live on flat space as Eqs.(37). Then we have considered this point that the generalized Cardy-Verlinde formula is the outcome of a striking resemblance between the thermodynamics of CFTs with asymptotically flat dual's and CFTs in two dimensions. After that we have obtained the corrections to the quantities entering the Cardy-Verlinde formula:Virasoro operator and the central charge.

References

- [1] S. W. Hawking, Commun. Math. Phys. **43**, 199, (1975).
- [2] E. Keski-Vakkuri and P. Kraus, Phys. Rev. D **54**, 7407, (1996).
- [3] P. Kraus and F. Wilczek, Nucl. Phys. B **433**, 403, (1995).
- [4] P. Kraus and F. Wilczek, Nucl. Phys. B **437**, 231, (1995).
- [5] E. Keski-Vakkuri and P. Kraus, Nucl. Phys. B **491**, 249, (1997).
- [6] M. K. Parikh and F. Wilczek, Phys. Rev. Lett. **85**, 5042,(2000).
- [7] Y. Kwon, Il Nuovo Cimento B **115**, 469, (2000).
- [8] S. Hemming and E. Keski-Vakkuri, Phys. Rev. D **64**, 044006 (2001).
- [9] M. R. Setare, Phys. Lett. **B602**, 1, (2004).
- [10] M. R. Setare, E. C. Vagenas ,Int. J.Mod. Phys. **A20**, 7219, (2005).
- [11] M. R. Setare, Phys. Lett. **B612**, 100, (2005).
- [12] M. R. Setare, Eur. Phys. J. **C47**, 851, (2006).
- [13] E. Verlinde, *On the Holographic Principle in a Radiation Dominated Universe*, hep-th/0008140.

- [14] J. M. Maldacena, *Adv. Theor. Math. Phys.* **2**, 231, (1998).
- [15] E. Witten, *Adv. Theor. Math. Phys.* **2**, 253 (1998).
- [16] S. Gubser, I. Klebanov and A. Polyakov, *Phys. Lett.* **B428**, 105 (1998).
- [17] O. Aharony, S. Gubser, J. Maldacena, H. Ooguri and Y. Oz, *Phys. Repts.* **323** , 183, (2000).
- [18] A. Strominger, *JHEP* **0110**, 034, (2001).
- [19] A. Strominger, *JHEP* **0111**, 049, (2001).
- [20] M. Spradlin, A. Strominger, A. Volovich, hep-th/0110007.
- [21] S. Nojiri, S.D. Odintsov, *Phys. Lett.***B519**, 145, (2001).
- [22] S. Nojiri, S.D. Odintsov, *JHEP* **0112**, 033, (2001).
- [23] S. Nojiri, S.D. Odintsov, *Phys. Lett.* **B523**, 165, (2001).
- [24] V. Balasubramanian, J. de Boer and D. Minic, *Phys. Rev.* **D65**, 123508 (2002).
- [25] M. R. Setare, *Mod. Phys. Lett.* **A17**, 2089, (2002).
- [26] M. R. Setare and R. Mansouri, *Int. J. Mod. Phys.* **A18**, 4443, (2003).
- [27] M. R. Setare and M. B. Altaie, *Eur. Phys. J.* **C30**, 273, (2003).
- [28] D. Klemm, A. C. Petkou, G. Siopsis, D. Zanon, *Nucl. Phys.* **B620**, 519, (2002).
- [29] R. C. Myers and M. J. Perry, *Annals Phys.* **172**, 304, (1986).
- [30] J. L. Cardy, *Nucl. Phys. B* **270**, 186, (1986).
- [31] D. Youm, *Mod. Phys. Lett.* **A16**, 1263, (2001).
- [32] J. Zhang, Z. Zhao, *Phys. Lett. B* **618**, 14, (2005).
- [33] S. Carlip, *Class. Quant. Grav.* **17**, 4175, (2000).