

# Advances in the measurement of the Lense-Thirring effect with planetary motions in the field of the Sun

Lorenzo Iorio

INFN-Sezione di Pisa. Permanent address for correspondence: Viale Unità di Italia 68,  
70125, Bari (BA), Italy. E-mail: [lorenzo.iorio@libero.it](mailto:lorenzo.iorio@libero.it)

Received \_\_\_\_\_; accepted \_\_\_\_\_

## ABSTRACT

E.V. Pitjeva, by processing more than 400,000 planetary observations of various types with the dynamical models of the EPM2006 ephemerides, recently estimated a correction to the canonical Newtonian-Einsteinian Venus' perihelion precession of  $-0.0004 \pm 0.0001$  arcseconds per century. The prediction of general relativity for the Lense-Thirring precession of the perihelion of Venus is  $-0.0003$  arcseconds per century. It turns out that neither other mismodelled/unmodelled standard Newtonian/Einsteinian effects nor exotic ones, postulated to, e.g., explain the Pioneer anomaly, may have caused the determined extra-precession of the Venus orbit which, thus, can be reasonably attributed to the gravitomagnetic field of the Sun, not modelled in the routines of the EPM2006 ephemerides. However, it must be noted that the quoted error is the formal, statistical one; the realistic uncertainty might be larger. Future improvements of the inner planets' ephemerides, with the inclusion of the Messenger and Venus-Express tracking data, should further improve the accuracy and the consistency of such a test of general relativity which would also benefit of the independent estimation of the extra-precessions of the perihelia (and the nodes) by other teams of astronomers.

*Subject headings:* Experimental tests of gravitational theories; Celestial mechanics; Orbit determination and improvement; Ephemerides, almanacs, and calendars

In the weak-field and slow motion approximation the Einstein field equations of general relativity get linearized resembling the Maxwellian equations of electromagnetism. As a consequence, a gravitomagnetic field arises (Mashhoon 2001, 2007); it is induced by the off-diagonal components  $g_{0i}$ ,  $i = 1, 2, 3$  of the space-time metric tensor related to the mass-energy currents of the source of the gravitational field. It affects orbiting test particles, precessing gyroscopes, moving clocks and atoms and propagating electromagnetic waves (Ruggiero and Tartaglia 2002; Schäfer 2004). The most famous gravitomagnetic effects are, perhaps, the precession of the axis of a gyroscope (Pugh 1959; Schiff 1960), whose detection in the gravitational field of the rotating Earth is the goal of the space-based GP-B experiment<sup>1</sup> (Everitt et al 1974), and the Lense-Thirring<sup>2</sup> precessions (Lense and Thirring 1918) of the orbit of a test particle for which some disputed satellite-based tests in the gravitational fields of the spinning Earth (Ciufolini and Pavlis 2004, 2005; Iorio 2005a, 2006a, 2007a; Lucchesi 2005) and Mars (Iorio 2006b; Krogh 2007; Iorio 2008a) have been reported.

We focus on the detection of the solar gravitomagnetic field through the Lense-Thirring planetary precessions of the longitudes of perihelia<sup>3</sup>  $\varpi = \omega + \cos i \Omega$

$$\frac{d\varpi}{dt} = -\frac{4GS \cos i}{c^2 a^3 (1 - e^2)^{3/2}}, \quad (1)$$

where  $G$  is the Newtonian gravitational constant,  $S$  is the proper angular momentum of the Sun,  $c$  is the speed of light in vacuum,  $a$  and  $e$  are the semimajor axis and the eccentricity,

---

<sup>1</sup>See on the WEB <http://einstein.stanford.edu/>

<sup>2</sup>According to an interesting historical analysis recently performed in (Pfister 2007), it would be more correct to speak about an Einstein-Thirring-Lense effect.

<sup>3</sup>Here  $\omega$  is the argument of pericentre, reckoned from the line of the nodes,  $i$  is the inclination of the orbital plane to the equator of the central rotating mass and  $\Omega$  is the longitude of the ascending node.

respectively, of the planet’s orbit. It may be interesting to know that in (Haas and Ross 1975) it was proposed to measure the solar gravitomagnetic field through the Schiff effect with a drag-free gyroscope orbiting the Sun in a polar orbit.

The impact of the Sun’s rotation on the Mercury’s longitude of perihelion was calculated for the first time with general relativity by de Sitter (1916) who, by assuming a homogenous and uniformly rotating Sun, found a secular rate of  $-0.01$  arcseconds per century ( $'' \text{cy}^{-1}$  in the following). This value is also quoted at pag. 111 of Soffel (1989). Cugusi and Proverbio (1978) yield  $-0.02'' \text{cy}^{-1}$  for the argument of perihelion of Mercury. Instead, recent determinations of the Sun’s proper angular momentum<sup>4</sup>  $S_{\odot} = (190.0 \pm 1.5) \times 10^{39} \text{ kg m}^2 \text{ s}^{-1}$  from helioseismology (Pijpers 1998, 2003), accurate to 0.8%, yield a precessional effect one order of magnitude smaller. The predicted gravitomagnetic precessions of the four inner planets, according to the recent value of the Sun’s angular momentum, are reported in Table 1; they are of the order of  $10^{-3} - 10^{-5}'' \text{cy}^{-1}$ . Due to their extreme smallness it has been believed for a long time, until recently, that the planetary Lense-Thirring effect would have been undetectable; see, e.g., p. 23 of Soffel (1989). A preliminary analysis showing that recent advances in the ephemerides field are making the situation more favorable was carried out in (Iorio 2005b). Pitjeva (2005a) processed more than 317,000 planetary observations of various kinds collected from 1917 to 2003 with the dynamical

---

<sup>4</sup>It could be measured also with a different approach (Ni 2008).

Table 1: Lense-Thirring precessions, in  $'' \text{cy}^{-1}$ , of the longitudes of the perihelion  $\varpi$  of the inner planets of the Solar System induced by the gravitomagnetic field of the Sun. The value  $S_{\odot} = (190.0 \pm 1.5) \times 10^{39} \text{ kg m}^2 \text{ s}^{-1}$  has been assumed for its angular momentum.

Mercury	Venus	Earth	Mars
$-0.0020$	$-0.0003$	$-0.0001$	$-0.00003$

force models of the EPM2004 ephemerides (Pitjeva 2005b) producing a global solution in which she estimated, among many other parameters, also corrections  $\Delta\dot{\varpi}$  to the canonical Newton-Einstein perihelion precessions for all the inner planets; since the gravitomagnetic force was not modelled at all, contrary to the static part of the general relativistic force of order  $\mathcal{O}(c^{-2})$ , such corrections to the usual perihelia evolutions account, in principle, for the Lense-Thirring effect as well, in addition to the mismodelled parts of the standard Newtonian/Einsteinian precessions. Thus, the estimated corrections for the perihelion rates of Mercury, the Earth and Mars have been used in (Iorio 2007b) to perform a first test. The errors  $\delta(\Delta\dot{\varpi})$  released in (Pitjeva 2005a) were slightly larger than the gravitomagnetic precessions whose predicted values, however, were found compatible with the estimated corrections. Venus was not used because of the poor data set used in the estimation of its extra-precession whose value, indeed, turned out too large to be due to a physically plausible effect amounting to  $+0.53 \pm 0.30'' \text{ cy}^{-1}$ : the Lense-Thirring prediction for the Venus perihelion precession was incompatible with such a result at about  $2 - \sigma$  level.

Now the situation for the second planet of the Solar System has remarkably improved allowing for a more stringent test of the Lense-Thirring effect. Indeed, Pitjeva (2007, 2008), in the effort of continuously improving the planetary ephemerides, recently processed more than 400,000 data points (1913-2006) with the EPM2006 ephemerides which encompasses better dynamical models with the exception, again, of the gravitomagnetic force itself. Also in this case she estimated, among more than 230 parameters, the corrections to the usual perihelion precessions for some planets (Pitjeva 2007). In the case of Venus the inclusion of the radiometric data of Magellan (Pitjeva 2008) as well allowed her to obtain<sup>5</sup>

$$\Delta\dot{\varpi}_{\text{Venus}} = -0.0004 \pm 0.0001'' \text{ cy}^{-1}, \quad (2)$$

in which the quoted uncertainty is the formal, statistical one. By looking at Table 1 it turns

---

<sup>5</sup>Personal communication by Pitjeva to the author, June 2008.

out that such an extra-precession can be well accommodated by the general relativistic prediction for the Lense-Thirring rate of the Venus’ perihelion whose existence would, thus, be confirmed at 25%. Somebody may object that the gravitomagnetic force should have been explicitly modelled and an ad-hoc parameter accounting for it should have been inserted in the set of parameters to be estimated. Certainly, it may be an alternative approach which could be implemented in future; in addition, we note that the procedure followed by Pitjeva may be regarded, in a certain sense, as safer for our purposes because it is truly model-independent and, since her goal in estimating  $\Delta\dot{\varpi}$  was not the measurement of the Lense-Thirring effect, there is a priori no risk that, knowing in advance the desired answer, something was driven just towards the expected outcome.

The main question to be asked is, at this point, the following one: Can the result of eq. (2) be explained by other unmodelled/mismodelled canonical or non-conventional dynamical effects? Let us, first, examine some standard candidates like, e.g., the residual precession due to the still imperfect knowledge of the Sun’s quadrupole mass moment  $J_2^\odot$  (Pireaux and Rozelot 2003) whose action was, in fact, modelled by Pitjeva (2005a) by keeping it fixed to  $J_2^\odot = 2 \times 10^{-7}$  in the global solution in which she estimated the corrections to the perihelion precessions. The answer is negative since the Newtonian secular precession due to the Sun’s oblateness<sup>6</sup>, whatever magnitude  $J_2$  may have, is positive. Indeed, it is (Capderou 2005; Iorio 2007b)

$$\dot{\varpi}_{J_2} = \frac{3}{2} \frac{nJ_2}{(1-e^2)^2} \left(\frac{R}{a}\right)^2 \left(1 - \frac{3}{2} \sin^2 i\right), \quad (3)$$

where  $n = \sqrt{GM/a^3}$  is the Keplerian mean motion and  $R$  is the Sun’s mean equatorial radius; the angle  $i$  between the Venus’ orbit and the Sun’s equator amounts to<sup>7</sup> 3.4 deg only. For  $J_2^\odot = 2 \times 10^{-7}$ , the nominal value of the Venus’ perihelion precession induced by it

---

<sup>6</sup>For an oblate body  $J_2 > 0$ .

<sup>7</sup>Indeed, the orbit of Venus is tilted by 3.7 deg to the mean ecliptic of J2000

amounts to  $+0.0026'' \text{ cy}^{-1}$ ; by assuming an uncertainty of about  $\delta J_2 \approx 10\%$  (Fienga et al 2008), if  $\Delta\dot{\omega}_{\text{Venus}}$  was due to such a mismodelled effect it should amount to  $+0.0002'' \text{ cy}^{-1}$ , which is, instead, ruled out at  $6 - \sigma$  level. Concerning the precession due to the solar octupole mass moment  $J_4^\odot$ , it is (Capderou 2005)

$$\dot{\omega}_{J_4} = -\frac{15}{16}nJ_4 \left(\frac{R}{a}\right)^4 \left[ \frac{3}{(1-e^2)^3} + 7\frac{(1+\frac{3}{2}e^2)}{(1-e^2)^4} \right] \left( \frac{7}{4}\sin^4 i - 2\sin^2 i + \frac{2}{5} \right). \quad (4)$$

For Venus it amounts to  $-1.2 J_4^\odot'' \text{ cy}^{-1}$ ; since  $J_4^\odot \approx -4 \times 10^{-9}$  (Roxburgh 2001; Mecheri et al 2004), we conclude that the second even zonal harmonic of the multipolar expansion of the solar gravitational potential cannot be responsible for eq. (2) and, more generally, it does not represent a potentially relevant source of systematic error for the measurement of the Lense-Thirring planetary precessions. Similar arguments hold also for other potential sources of systematic errors like, e.g., the asteroid ring and the Kuiper Belt objects, both modelled in EPM2006: the precessions induced by them are positive. Indeed, a Sun-centered ring of mass  $m_{\text{ring}}$  and inner and outer radius  $R_{\text{min/max}} \gg a$  induces a perihelion precession (Iorio 2007c)

$$\dot{\omega}_{\text{ring}} = \frac{3}{4}\sqrt{\frac{Ga^3(1-e^2)}{M}} \frac{m_{\text{ring}}}{R_{\text{min}}R_{\text{max}}(R_{\text{min}}+R_{\text{max}})} > 0. \quad (5)$$

According to eq. (5), the precession induced by the asteroids' ring on the Venus' perihelion amounts to  $+0.0007 \pm 0.0001'' \text{ cy}^{-1}$  by using  $m_{\text{ring}} = (5 \pm 1) \times 10^{-10}M_\odot$  (Krasinsky et al 2002); the lowest value  $+0.0006'' \text{ cy}^{-1}$  is incompatible with eq. (2) at  $10 - \sigma$  level. In the case of the Kuiper Belt Objects, eq. (5) yields a precession of the order of  $+0.00006'' \text{ cy}^{-1}$  with  $m = 0.052m_\oplus$  (Iorio 2007c). Thus, we can rule out such modelled classical features of the Sun and the Solar System as explanations of  $\Delta\dot{\omega}_{\text{Venus}}$ . General relativistic terms of order  $\mathcal{O}(c^{-4})$  were not modelled by Pitjeva; however, the first correction of order  $\mathcal{O}(c^{-4})$  to

---

([http://ssd.jpl.nasa.gov/txt/aprx\\_pos\\_planets.pdf](http://ssd.jpl.nasa.gov/txt/aprx_pos_planets.pdf)), while the Carrington's angle between the Sun's equator and the ecliptic is  $7.15 \text{ deg}$  (Beck and Giles 2005).

the perihelion precession (Damour and Schäfer 1988) can be safely neglected because for Venus it is

$$\dot{\omega}_{c^4} \propto \frac{n(GM)^2}{c^4 a^2 (1-e^2)^2} \approx 10^{-7} \text{ '' cy}^{-1}. \quad (6)$$

Concerning possible exotic explanations, i.e. due to some modifications of the currently known Newton-Einstein laws of gravity, it may have some interest to check some of the recently proposed extra-forces (Standish 2008) which would be able to phenomenologically accommodate the Pioneer anomaly (Nieto 2006). All of such hypothetical new forces have not been modelled by Pitjeva, so that if they existed in Nature they would affect  $\Delta\dot{\omega}_{\text{Venus}}$ . A central acceleration quadratic in the radial component  $v_r$  of the velocity of a test particle<sup>8</sup> (Jaekel and Reynaud 2005; Standish 2008)

$$A = -v_r^2 \mathcal{H}, \quad \mathcal{H} = 6.07 \times 10^{-18} \text{ m}^{-1} \quad (7)$$

would induce a retrograde perihelion precession according to (Iorio 2008b)

$$\dot{\omega} = \frac{\mathcal{H} n a \sqrt{1-e^2}}{e^2} \left( -2 + e^2 + 2\sqrt{1-e^2} \right) < 0. \quad (8)$$

However, eq. (8) predicts a precession of  $-0.0016 \text{ '' cy}^{-1}$  for Venus, which is ruled out by eq. (2) at  $12 - \sigma$  level. Another possible candidate considered in (Standish 2008) is an acceleration linear in the radial velocity

$$A = -|v_r| \mathcal{K}, \quad \mathcal{K} = 7.3 \times 10^{-14} \text{ s}^{-1} \quad (9)$$

which yields a retrograde perihelion precession (Iorio 2008b)

$$\dot{\omega} = -\frac{\mathcal{K} \sqrt{1-e^2}}{\pi} \left[ \frac{2e - (1-e^2) \ln\left(\frac{1+e}{1-e}\right)}{e^2} \right] < 0. \quad (10)$$

The prediction of eq. (10) for Venus is  $-0.1 \text{ '' cy}^{-1}$ , clearly incompatible with eq. (2). Should one consider a central uniform acceleration with the magnitude of the Pioneer

---

<sup>8</sup>The quoted numerical value of  $\mathcal{H}$  allows to reproduce the Pioneer anomaly.

anomalous one, i.e.  $A = -8.74 \times 10^{-10} \text{ m s}^{-2}$ , the exotic precession induced by it (Iorio and Giudice 2006; Sanders 2006) on the perihelion of Venus would be

$$\dot{\omega}_{\text{Ven}} = A \sqrt{\frac{a(1-e^2)}{GM}} = -16 \text{ '' cy}^{-1}. \quad (11)$$

Another non-conventional effect which may be considered is the precession predicted by Lue and Starkman (2003) in the framework of the DGP multidimensional braneworld model by Dvali, Gabadadze and Porrati (Dvali et al 2000) proposed to explain the cosmic acceleration without invoking dark energy. It is

$$\dot{\omega}_{\text{LS}} = \mp \frac{3c}{8r_0} + \mathcal{O}(e^2) \approx \mp 0.0005 \text{ '' cy}^{-1}, \quad (12)$$

where the plus sign is related to the self-accelerated branch, while the minus sign is for the standard, Friedmann-Lemaître-Robertson-Walker (FLRW) branch;  $r_0 \approx 5 \text{ Gpc}$  is a threshold characteristic of the DGP model after which gravity would experience neat deviations from the Newtonian-Einsteinian behavior. As can be noted, the self-accelerated branch is ruled out at  $9 - \sigma$  level by eq. (2), while the FLRW case is still compatible with eq. (2) ( $1 - \sigma$  discrepancy). By the way, apart from the fact that there are theoretical concerns with the DGP model (see, e.g., (Izumi et al 2007) and references therein), the existence of both the Lue-Starkman FLRW precession and the Lense-Thirring one, implying a total unmodelled effect of  $-0.0008 \text{ '' cy}^{-1}$ , would be ruled out by eq. (2) at  $4 - \sigma$  level. As a consequence, we can conclude not only that the examined exotic modifications of the standard laws of gravity, not modelled by Pitjeva, are not responsible for the estimated  $\Delta \dot{\omega}_{\text{Venus}}$ , but also that their existence in the inner regions of the Solar System is falsified by the observations. Moreover, given the magnitudes of the hypothetical effects with the negative sign, it is not possible that reciprocal cancelations with the positive classical mismodelled precessions can explain eq. (2). Indeed, the sum of the latter ones is  $+0.0004 \text{ '' cy}^{-1}$ ; the sum of, e.g., eq. (8) and eq. (12) (FLRW) is  $-0.0021 \text{ '' cy}^{-1}$ , while the sum of eq. (8) and eq. (12) (self-accelerated branch) is  $-0.0011 \text{ '' cy}^{-1}$ .

Thus, we conclude that the most likely explanation for eq. (2) is just the general relativistic Lense-Thirring effect. However, caution is in order in assessing the realistic uncertainty in such a test because, as already stated, the released error of  $0.0001'' \text{ cy}^{-1}$  is the formal, statistical one; the realistic uncertainty might be larger. By the way, we can at least firmly conclude that now also in the case of Venus the general relativistic predictions for the Lense-Thirring effect on  $\dot{\varpi}$  are compatible with the observational determinations for the unmodelled perihelion precessions, contrary to the case of (Iorio 2007b). Moreover, future modelling of planetary motions should take into account the relativistic effects of the rotation of the Sun as well. The steady improvement in the planetary ephemerides, which should hopefully benefit of the radiometric data from Messenger and Venus-Express as well, should allow for more accurate and stringent test in the near-mid future. Of great significance would be if also other teams of astronomers would estimate their own corrections to the canonical perihelion (and also node) precessions in order to enhance the statistical significance and robustness of this important direct test of general relativity.

### Acknowledgments

I thank E.V. Pitjeva for useful and important communications.

## REFERENCES

- Beck J G and Giles P 2005 *Astroph. J.* **621** L153
- Capderou M 2005 *Satellites. Orbits and missions* (Paris: Springer) p. 136
- Ciufolini I and Pavlis E C 2004 *Nature* **431** 958
- Ciufolini I and Pavlis E C 2005 *New Astron.* **10** 636
- Cugusi L and Proverbio E 1978 *Astron. Astrophys.* **69** 321
- Damour T and Schäfer G 1988 *Nuovo Cimento B* **101** 127
- de Sitter W 1916 *Mon. Not. R. Astron. Soc.* **76**, 699
- Dvali G, Gabadadze G and Porrati M 2000 *Phys. Lett. B* **485** 208
- Everitt C W F 1974 The Gyroscope Experiment I. General Description and Analysis of Gyroscope Performance in *Proc. Int. School Phys. “Enrico Fermi” Course LVI* ed B Bertotti (New York: New Academic Press) pp 331–60
- Fienga A, Manche H, Laskar J and Gastineau M 2008 *Astron. Astrophys.* **477**, 315
- Haas M R and Ross D K 1975 *Astrophys. Space Sci.* **32** 3
- Iorio L 2005a *New Astron.* **10** 616
- Iorio L 2005b *Astron. Astrophys.* **431**, 385
- Iorio L 2006a *J. Geod.* **80** 128
- Iorio L 2006b *Class. Quantum Grav.* **23** 5451
- Iorio L 2007a *Planet. Space Sci.* **55** 503

Iorio L 2007b *Planet. Space Sci.* **55** 1290

Iorio L 2007c *Mon. Not. Roy. Astron. Soc.* **375** 1311

Iorio L 2008a *J. Gravit. Phys.* at press, <http://arxiv.org/abs/gr-qc/0701146>

Iorio L 2008b *Int. J. Mod. Phys. D*, at press, <http://arxiv.org/abs/0806.3011>

Iorio L and Giudice G 2006 *New Astron.* **11** 600

Izumi K, Koyama K, Pujolàs O and Tanaka T 2007 *Phys. Rev. D* **76** 104041

Jaekel M-T and Reynaud S 2005 *Mod. Phys. Lett. A* **20** 1047

Krasinky G A, Pitjeva E V, Vasilyev M V and Yagudina E I 2002 *Icarus* **158** 98

Krogh K 2007 *Class. Quantum Grav.* **24** 5709

Lense J and Thirring H 1918 *Phys. Z.* **19** 156 Translated and discussed in Mashhoon B, Hehl F W and Theiss D S 1984 *Gen. Relativ. Gravit.* **16** 711. Reprinted in Ruffini R J and Sigismondi C (eds) 2003 *Nonlinear Gravitodynamics* (Singapore: World Scientific) pp 349–88

Lue A and Starkman G 2003 *Phys. Rev. D* **67** 064002

Lucchesi D M 2005 *Int. J. Mod. Phys. D* **14** 1989

Mashhoon B 2001 *Gravitoelectromagnetism* in Pascual-Sánchez J F, Floría L, San Miguel A and Vicente F (eds.) *Reference Frames and Gravitomagnetism* (Singapore: World Scientific) pp. 121-132

Mashhoon B 2007 *Gravitoelectromagnetism: A Brief Review* in Iorio L (ed.) *The Measurement of Gravitomagnetism: A Challenging Enterprise* (Hauppauge, New York: NOVA publishers) pp. 29-39

- Mecheri R, Abdelatif T, Irbah A, Provost J and Berthomieu G 2004 *Solar Phys.* **222** 191
- Ni W-T 2008 *Int. J. Mod. Phys. D* **17** 921
- Nieto M M 2006 *Europhysics News* **37**/6, 30
- Pfister H 2007 *Gen. Relativ. Gravit.* **39** 1735
- Pijpers F P 1998 *Mon. Not. R. Astron. Soc.* **297**, L76
- Pijpers F P 2003 *Astron. Astrophys.* **402**, 683
- Pitjeva E V 2005a *Astron. Lett.* **31**, 340
- Pitjeva E V 2005b *Sol. Syst. Res.* **39**, 176
- Pitjeva E V 2007 *Proceedings of IAA RAS* **17**, pp. 42–59.
- Pitjeva E V 2008 in Jin W J, Platais I and Perryman M A C (eds) *A Giant Step: from Milli- to Micro-arcsecond Astrometry Proceedings IAU Symposium No. 248, 2007*, (Cambridge: Cambridge University Press), pp. 20–22.
- Pugh G E 1959 *WSEG Research Memorandum No. 11* Reprinted in Ruffini R J and Sigismondi C (eds) 2003 *Nonlinear Gravitodynamics. The Lense–Thirring Effect* (Singapore: World Scientific) pp. 414–26
- Roxburgh I W 2001 *Astron. Astrophys.* **377** 688
- Pireaux S and Rozelot J P 2003 *Astrophys. Space Sci.* **284** 1159
- Ruggiero M L and Tartaglia A 2002 *N. Cim. B* **117** 743
- Sanders R H 2006 *Mon. Not. Roy. Astron. Soc.* **370** 1519
- Schiff L 1960 *Phys. Rev. Lett.* **4** 215

Schäfer G 2004 *Gen. Relativ. Gravit.* **36** 2223

Soffel M 1989 *Relativity in Astrometry, Celestial Mechanics and Geodesy* (Berlin: Springer)

Standish E M 2008 in Macías A, Lämmerzahl C and Camacho A (eds) 2008 *AIP Conference Proceedings Volume 977. RECENT DEVELOPMENTS IN GRAVITATION AND COSMOLOGY: 3rd Mexican Meeting on Mathematical and Experimental Physics* (American Institute of Physics), pp. 254–263.