Symmetry Reduction of Lane-Emden Equation for Polytropes

BABUR M. MIRZA

Department of Mathematics, Quaid-i-Azam University, Islamabad 45320, Pakistan E-mail: bmmirza2002@yahoo.com

October 16, 2005

Abstract

We describe an ansatz for symmetry reduction of the Lane-Emden equation for an arbitrary polytropic index n, admitting only one symmetry generator. For the reduced first order differential equation it is found that standard reduction procedure do not admit any non-trivial Lie point symmetry. However some special solutions for the differential equation are obtained.

The simplest model of a gravitationally bound spherically symmetric polytrope with a power law distribution function is determined by the Lane-Emden equation [1, 2]

$$\frac{1}{r^2}\frac{d}{dr}(r^2\frac{d\psi(r)}{dr}) = -\psi(r)^n\tag{1}$$

where $\psi(r)$ is the dimensionless gravitational potential, r is the dimensionless radius of the sphere, and n is the polytropic index. For the second order differential equation the symmetry generator has the form[3]

$$\mathbf{X} = \xi(r, \psi) \frac{\partial}{\partial r} + \eta(r, \psi) \frac{\partial}{\partial \psi} + \eta'(r, \psi) \frac{\partial}{\partial \psi'}$$
 (2)

Here the prime denotes differentiation with respect to the independent variable r. For the differential equation the symmetry condition is given by

$$\mathbf{X}\omega(r,\psi,\psi') = \eta''(r,\psi), \quad \text{mod } \psi'' = \omega(r,\psi,\psi'). \tag{3}$$

This gives an identity in the dependent and independent variables (r, ψ) . From (1) we have

$$\omega(r,\psi,\psi') = -\psi(r)^n - \frac{2}{r}\psi'. \tag{4}$$

The infinitesimals (ξ, η) are now determined by the identity

$$(\psi^{n} + \frac{2}{r}\psi')[\eta_{,\psi} - 2\xi_{,r} - 3\psi'\xi_{,\psi}] + \frac{2}{r^{2}}\psi'\xi - n\psi^{n-1}\eta - \frac{2}{r}[\eta_{,r} + \psi'(\eta_{,\psi} - \xi_{,r})]$$

$$-(\psi')^{2}\xi_{,\psi}] = \eta_{,rr} + \psi'[2\eta_{,r\psi} - \xi_{,rr}] + (\psi')^{2}[\eta_{,\psi\psi} - 2\xi_{,r\psi}] - (\psi')^{3}\xi_{,\psi\psi}$$
 (5)

We can determine the infinitesimals ξ and η by an iterative procedure. First equating coefficients of $(\psi')^3$ and $(\psi')^2$ in (5), we find that $\xi \neq \xi(\psi)$ and $\eta_{,\psi\psi} = 0$. Therefore we assume that $\xi = \alpha(r)$ and $\eta = \beta(r)\psi + \gamma(r)$. The symmetry condition then reduces to

$$(\psi^{n} + \frac{2}{r}\psi')[\beta - 2\alpha'] + \frac{2}{r^{2}}\psi'\alpha - n\psi^{n-1}(\beta\psi + \gamma) - \frac{2}{r}[\beta'\psi + \gamma' + \psi'(\beta - \alpha')]$$

$$= \beta''\psi + \gamma'' + \psi'[2\beta' - \alpha''] \tag{6}$$

Equating the coefficients of ψ' and ψ^n we obtain the following two conditions

$$2\alpha - 2r\alpha' - [2\beta' - \alpha'']r^2 = 0, \quad [\beta - 2\alpha'] - n\beta = 0$$
 (7)

The coefficients of r^2 equated to zero leave for the first two terms in the first condition $\alpha = kr$ where k does not depend on r. The second equation gives $\beta = 2\alpha'/(1-n)$. Substitution back into the determining equation yields

$$n\gamma\psi^{n-1} + \frac{2}{r}\gamma' + \gamma'' \equiv 0 \tag{8}$$

Comparing coefficients of ψ^{n-1} we find that $\gamma = 0$. Thus the infinitesimals are

$$(\xi, \eta) = (kr, \frac{-2k}{n-1}\psi), \quad n \neq 1.$$

$$(9)$$

We have here a one-parameter group G_1 of scaling transformation with a parameter k. Thus reduction in the order of the equation by one is possible [4]. For convenience we take k to be unity.

Let (t, s) be the canonical variables for the differential equation (1) with infinitesimals (9). Then

$$s = \int \frac{dx}{\xi}, \quad t = \int (\xi dy - \eta dx); \tag{10}$$

gives

$$s = \ln x, \quad t = y \exp(\frac{2}{n-1}); \quad n \neq 1.$$
 (11)

We have for s = s(t) and s' = ds/dt:

$$dr = d(e^s) = e^s ds, \quad d\psi = d(te^{2s/(1-n)}) = e^{2s/(1-n)} dt + \frac{2t}{1-n} e^{2s/(1-n)} ds.$$
(12)

which implies that

$$\psi' = \frac{d\psi}{dr} = \frac{e^{(\frac{1+n}{1-n})s}}{s'} \left[1 + \frac{2}{1-n}ts'\right],\tag{13}$$

and

$$\psi'' = \frac{d}{dt}(\frac{d\psi}{dr}) = \frac{e^{(\frac{2n}{1-n})s}}{(s')^3} \left[s' \left\{ \frac{2(ts'' + s')}{1-n} + \frac{1+n}{1-n} (1 + \frac{2ts'}{1-n})s' \right\} - s'' (1 + \frac{2ts'}{1-n}) \right]$$
(14)

With these substitutions we obtain from equation (1) the following reduced form of the Lane-Emden equation

$$u'(t) = \frac{n-5}{n-1}u^2 + \left\{\frac{2(3-n)}{(n-1)^2}t + t^n\right\}u^3, \quad n \neq 1.$$
 (15)

Where u(t) = s'(t) and u'(t) = s''(t). This highly nonlinear first order differential equation has solutions for n = 0 and 5. These correspond to the well known solutions for ψ namely $-(r^2/6) + (1/6x)$ and $(1 + r^2/3)^{-1/2}$ respectively. The only other known solution to (15) is the spherical Bessel function corresponding to n = 1[5, 6]. To investigate the solution for other

permissible values of n we first transform the variable u(t) as -1/y(t). This substitution weakens the nonlinearity of the equation as

$$y'(t) = a - \frac{bt + t^n}{y(t)}, \quad n \neq 1,$$
 (16)

where a = (n-5)/(n-1) and $b = 2(3-n)/(n-1)^2$. For the infinitesimals (ς, ϕ) The symmetry condition for equation (16) gives

$$\phi_{,t} + (\phi_{,y} - \varsigma_{,t})(a - \frac{bt + t^n}{y(t)}) - \varsigma_{,y}\left(a - \frac{bt + t^n}{y(t)}\right)^2 = \varsigma\left(-\frac{b + nt^{n-1}}{y}\right) + \phi\left(\frac{bt + t^n}{y^2}\right)$$
(17)

For the infinitesimals of this equation we find that no non-trivial solution to (16) exists for the above ansatz.

However one immediate consequence of the infinitesimals in (9) is an invariant particular solution for the Lane-Emden equation. We first have

$$\psi'(r) = \frac{\eta}{\xi} = \frac{-2}{n-1} (\frac{\psi}{r}), \quad \psi''(r) = \frac{2(n+1)}{(n-1)^2} \frac{\psi}{r^2}$$
 (18)

Substitution into equation (1) we obtain the particular solution in an implicit form

$$\psi^n = \frac{2(n-3)}{(n-1)^2} \frac{\psi}{r^2}, \quad n \neq 1.$$
 (19)

For n less than one this solution is bounded for all values of r. For n > 1 the solution has a singularity at r = 0, which becomes stronger as n increases.

References

- [1] Liu F K, Polytropic gas spheres: An approximate analytic solution of the Lane-Emden equation, MNRAS 281 (1996), 1197.
- [2] Honda M and Honda Y S, On Exact Polytropic Equilibria of Self-Gravitating Gaseous and Radiative Systems: Their Application to Molecular Cloud Condensation, MNRAS **341** (2003), 164.
- [3] Stephani H, Differential Equations: Thier Solution using Symmetries, University Press, Cambridge, 1989, p28.

- [4] Bluman G W and Kumei S, Symmetries and Differential Equations. Applied Mathematical Sciences 81, Springer-Verlag, New York, 1989, p110.
- [5] Rose W K, Advanced Stellar Astrophysics. University Press, Cambridge, 1998, p447.
- [6] McKee C F and Holliman J H II, Multi-Pressure Polytropes as Models for the Structure and Stability of Molecular Clouds. I. Theory. ApJ. 522 (1999), 313.