

Correction to “Knotted Hamiltonian cycles in spatial embeddings of complete graphs”

Joel Foisy

ABSTRACT. We state and prove a correct version of a theorem presented in [1].

Professor Masakazu Teragaito has recently pointed out that Theorem 3.3 of [1] is incorrect as stated. The fact that $\mu_f(G, \Gamma; 6) = 3$ is independent of embedding of K_8 does not necessarily imply that there are at least 3 knotted Hamiltonian cycles. For example, there could be exactly one knotted Hamiltonian cycle with $a_2(K) = 3$.

Professor Kouki Taniyama has further pointed out that Lemma 2, in [2], has a gap in its proof. He has proposed a rigorous proof of a weaker version Lemma 2. In this short paper, we will state and prove this weaker version of Lemma 2 in [2], and then apply it to obtain a weaker version of Theorem 3.3 of [1]. For definitions of terms, see [2] and [1]. Here is the modified version of Shimabara’s Lemma 2 that we will prove:

Lemma 0.1. *Let Γ be a set of cycles in an undirected graph G . The invariant $\mu_f(G, \Gamma; n)$ does not depend on the spatial embedding f of G if the following two conditions hold:*

- (1) *For any edges A, B, E such that A is adjacent to B , $\nu_1(\Gamma; A, B, E) \equiv 0 \pmod{n}$.*
- (2) *For any pairs of non-adjacent edges (A, B) and (E, F) , $\nu_2(\Gamma; A, B; E, F) \equiv 0 \pmod{2n}$.*

The difference between this new lemma and the original comes in the second condition, where the equivalence is mod $2n$, not n . For the proof of Lemma 2, case 2, on p. 410 of [2], the definition of linking number used does not work. For the version of linking number used, $\zeta(A, B)$ depends on the order of A and B , whereas the equality $\sum_{E, F} \sum_{\gamma \in \Gamma_1} \epsilon(c) \zeta(f_\gamma(E), f_\gamma(F)) = \sum_{E, F} (n_3 - n_4) \zeta(f(E), f(F))$ implicitly uses the assumption that $\zeta(A, B) = \zeta(B, A)$.

A proof of Lemma 0.1 is possible if one uses a different (but equivalent) version of ζ and linking number. For A and B two disjoint oriented arcs or circles in \mathbb{R}^3 , define $\zeta'(A, B) = (1/2) \sum_c \epsilon(c)$, with the summation being over all crossings

Mathematics Subject Classification. 57M15, 57M25.

Key words and phrases. spatial graph, embedded graph, intrinsically knotted.

between A and B . If A and B are circles, then $\zeta'(A, B)$ gives the linking number of A and B , $lk(A, B)$. It then follows immediately that $\zeta'(A, B) = \zeta'(B, A)$. The proof of Lemma 0.1 proceeds as in the proof of Lemma 2 in [2], but now ends with:

$$\delta(u) = \sum_{E, F} \sum_{\gamma \in \Gamma_1} \epsilon(c) \zeta(f_\gamma(E), f_\gamma(F)) = \sum_{E, F} (n_3 - n_4) \zeta'(f(E), f(F)).$$

It then follows, since each $\zeta'(f(E), f(F))$ is either an integer or an integer divided by 2, that, $\delta(u)$ is congruent to 0 mod n if $|n_3 - n_4| \equiv 0 \pmod{2n}$.

Now, in [1], it was shown that $\nu_2 \equiv 0 \pmod{6}$. It was also shown that there exists an embedding of K_8 with exactly 21 knotted Hamiltonian cycles, each with Arf invariant 1. One can also verify that each of these knotted cycles is a trefoil, with $a_2 = 1$. This embedding, together with Lemma 0.1, implies that $\mu_f(K_8, \Gamma, 3) \equiv 0$ for every spatial embedding of K_8 . By Theorem 2.2 of [1], there is at least one Hamiltonian cycle with Arf invariant 1, in every spatial embedding of K_8 .

We thus have the following corrected version of Theorem 3.3 from [1].

Theorem 0.2. *Given an embedding of K_8 , at least one of the following must occur in that embedding:*

1. *At least 3 knotted Hamiltonian cycles.*
2. *Exactly 2 knotted Hamiltonian cycles C_1 and C_2 , with $a_2(C_1) \equiv 1 \pmod{3}$ and $a_2(C_2) \equiv 2 \pmod{3}$, or $a_2(C_1) \equiv 0 \pmod{3}$ and $a_2(C_2) \equiv 0 \pmod{3}$. Either C_1 or C_2 has non-zero Arf invariant.*
3. *Exactly 1 knotted Hamiltonian cycle, C with $a_2(C) \equiv 1 \pmod{2}$ and $a_2(C) \equiv 0 \pmod{3}$. (Equivalently: $a_2(C) \equiv 3 \pmod{6}$.)*

It thus remains an open question to determine if 1 is the best lower bound for the minimum number of knotted Hamiltonian cycles in every spatial embedding of K_8 .

Acknowledgments The author would like to thank Professors Masakazu Teragaito, Kouki Taniyama and Ryo Nikkuni for valuable comments and suggestions.

References

- [1] P. Blain, G. Bowlin, J. Hendricks, J. LaCombe, J. Foisy, *Knotted Hamiltonian cycles in spatial embeddings of complete graphs*, New York Journal of Mathematics, **13** (2007), 11-16.
- [2] M. Shimabara *Knots in Certain Spatial Graphs* Tokyo J. Math. Vol. 11, No 2, 1988.

DEPARTMENT OF MATHEMATICS, SUNY POTSDAM, POTSDAM, NY 13676
foisyjs@potdam.edu